



HAL
open science

Effect of rapid response systems on hospital mortality: a systematic review and meta-analysis

Audrey de Jong, Boris Jung, Aurelien Daurat, Gerald Chanques, Martin Mahul, Marion Monnin, Nicolas Molinari, Samir Jaber

► To cite this version:

Audrey de Jong, Boris Jung, Aurelien Daurat, Gerald Chanques, Martin Mahul, et al.. Effect of rapid response systems on hospital mortality: a systematic review and meta-analysis. *Intensive Care Medicine*, 2016, 42 (4), pp.615 - 617. 10.1007/s00134-016-4263-1 . hal-01780710

HAL Id: hal-01780710

<https://hal.umontpellier.fr/hal-01780710>

Submitted on 3 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of rapid response systems on hospital mortality: a systematic review and meta-analysis

Audrey De Jong^{1†}, Boris Jung^{1,2†}, Aurelien Daurat¹, Gerald Chanques^{1,2}, Martin Mahul¹, Marion Monnin¹, Nicolas Molinari³ and Samir Jaber^{1,2*}

Dear Editor,

This article reports a meta-analysis and systematic review of studies of rapid response system (RRS) compared to no RRS including the recent study by Jung et al. [1].

We made a quantitative synthesis selecting the following study designs: randomized controlled trials, prospective observational studies (before and after treatment), and retrospective observational studies [see also Electronic Supplemental Material (ESM) 1 for selection criteria and data collection and analysis] [2]. The primary endpoint of this meta-analysis was the incidence of overall hospital mortality. The secondary endpoint was the incidence of unexpected mortality.

Data were extracted as they were reported in the original paper or based on the authors' answers to our queries (see ESM for further details). Odds ratios (ORs) were used as the summary measure for dichotomous outcomes. Statistical heterogeneity was quantified by the Q-Cochrane heterogeneity test [Q statistic with degree of freedom (df)] and the I^2 statistic [3]. In case of heterogeneity, a random effect model was performed. Predefined subgroups analysis and exclusion of outlying studies were performed to reduce heterogeneity for the primary outcome [4, 5]. To assess the effect of potential confounding factors we performed a meta-regression (see also ESM). A funnel plot (plot of treatment effect against trial precision) was created to determine the presence of

publication bias, true heterogeneity, data irregularities, and choice of effect measure in the meta-analysis.

The updated literature search provided a total of 32 studies and 10,563,083 patients (see ESM Fig. S1). Of those, nine studies (1,045,364 patients) provided data on unexpected mortality and 32 studies (9,517,719 patients) reported overall mortality rate. The meta-regression did not reveal any significant effect of potential confounding factors (see ESM) for overall hospital mortality.

RRSs were associated with a significant decrease in both overall hospital mortality [OR 0.89; 95 % CI 0.85–0.93] (Fig. 1a) and unexpected mortality [OR 0.51; 95 % CI (0.35–0.76)] (Fig. 1b). Funnel plots did not reveal publication bias for overall mortality or unexpected mortality (see ESM Figs. S2 and S3). Subgroups analysis performed according to the multiple center or single center design (Fig. S4), the prospective or retrospective design (Fig. S5), and the location of the studies (Fig. S6) showed consistent results, with a significant decrease of overall hospital mortality. After exclusion of outlying studies (Fig. S7), heterogeneity became low and a fixed model could be applied, showing identical results with a decreased incidence of overall hospital mortality [OR 0.92; 95 % CI (0.90–0.94)].

In this updated systematic review and meta-analysis, medical emergency team (MET) implementation was associated with a significant decrease in overall and unexpected mortality of hospitalized patients. One limitation was the large amount of statistical heterogeneity. Nevertheless, we considered that to combine data using a random effect meta-analysis would allow a more useful result than to include a small number of homogeneous studies as presented in Fig. S7 [5]. Moreover, subgroups analysis and exclusion of outlying studies allowing

*Correspondence: s-jaber@chu-montpellier.fr

†A. De Jong and B. Jung contributed equally to this work.

¹ Intensive Care Unit, Department of Anesthesia and Critical Care Medicine, University of Montpellier, Saint Eloi Teaching Hospital, 80 Avenue Augustin Fliche, 34295 Montpellier Cedex 5, France
Full author information is available at the end of the article




Fig. 1 Forests plots of **a** overall mortality and **b** unexpected hospital mortality. Study-specific odds ratios (95% CI) are denoted by *black boxes* (*black lines*) for unexpected mortality. The combined odds ratios estimate for all studies is represented by a *black diamond*. Overall diamond width corresponds to 95% CI bounds. Heights of boxes and diamonds are inversely proportional to the precision of the odds ratio estimate. The *p* value for heterogeneity of the odds ratio by study is shown (Q statistic with degree of freedom) and I² statistic

reduced heterogeneity showed similar results. The present updated meta-analysis strongly suggests a positive effect of RRS on overall and unexpected mortality.

Author details

¹ Intensive Care Unit, Department of Anesthesia and Critical Care Medicine, University of Montpellier, Saint Eloi Teaching Hospital, 80 Avenue Augustin Fliche, 34295 Montpellier Cedex 5, France. ² Centre National de la Recherche Scientifique (CNRS 9214), Institut National de la Santé et de la Recherche Medicale (INSERM U-1046), Montpellier University, Montpellier, France. ³ Department of Statistics, University of Montpellier Lapeyronie Hospital, UMR 729 MISTEA, Montpellier, France.

Compliance with ethical standards

Conflicts of interest

Boris Jung reports personal fees from Merck (Whitehouse Station, NJ) and Astellas (Tokyo, Japan) not related to the present study. Aurelien Daurat, Nicolas Molinari, Audrey De Jong, Martin Mahul, Marion Monnin, and Gerald Chanques have nothing to disclose. Samir Jaber reports personal fees from Maquet, Draeger, Hamilton Medical, Fisher Paykel, and Abbott not related to the present study.

Source of funding

This study was supported by the University Hospital of Montpellier.

References

1. Jung B, Daurat A, De Jong A, Chanques G, Mahul M, Monnin M, Molinari N, Jaber S (2016) Rapid response team and hospital mortality in hospitalized patients. *Intensive care Med*. doi:10.1007/s00134-016-4254-2
2. De Jong A, Molinari N, Conseil M, Coisel Y, Pouzeratte Y, Belafia F, Jung B, Chanques G, Jaber S (2014) Video laryngoscopy versus direct laryngoscopy for orotracheal intubation in the intensive care unit: a systematic review and meta-analysis. *Intensive Care Med* 40:629–639
3. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 339:b2700
4. Higgins JPT, Green S (2011) *Cochrane handbook for systematic reviews of interventions* version 5.1.0. The Cochrane Collaboration. <http://www.cochrane-handbook.org>. Accessed 1 Oct 2013
5. Hole J, Hirsch M, Ball E, Meads C (2015) Music as an aid for postoperative recovery in adults: a systematic review and meta-analysis. *Lancet* 386:1659–1671