



HAL
open science

Dual Molecular Effects of Dominant RORA Mutations Cause Two Variants of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia

Claire Guissart, Xenia Latypova, Paul Rollier, Tahir Khan, Hannah Stamberger, Kirsty Mcwalter, Megan Cho, Susanne Kjaergaard, Sarah Weckhuysen, Gaetan Lesca, et al.

► **To cite this version:**

Claire Guissart, Xenia Latypova, Paul Rollier, Tahir Khan, Hannah Stamberger, et al.. Dual Molecular Effects of Dominant RORA Mutations Cause Two Variants of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia. *American Journal of Human Genetics*, 2018, 102 (5), pp.744 - 759. 10.1016/j.ajhg.2018.02.021 . hal-01796580

HAL Id: hal-01796580

<https://hal.umontpellier.fr/hal-01796580>

Submitted on 13 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dual molecular effects of dominant *RORA* mutations cause two variants of syndromic intellectual disability with either autistic features or cerebellar ataxia

Claire Guissart,^{1,36} Xenia Latypova,^{2,3,4,37} Paul Rollier,^{5,37} Tahir N Khan,^{3,37} Hannah Stamberger,^{6,7,8} Kirsty McWalter,⁹ Megan T. Cho,⁹ Susanne Kjaergaard,¹⁰ Sarah Weckhuysen,^{6,7,8} Gaetan Lesca,^{11,12} Thomas Besnard,^{2,4} Katrin Öunap,¹³ Lynn Schema,¹⁴ Andreas G. Chiocchetti,¹⁵ Marie McDonald,¹⁶ Julitta de Bellescize,¹⁷ Marie Vincent,^{2,4} Hilde Van Esch,¹⁸ Shannon Sattler,¹⁹ Irman Forghani,²⁰ Isabelle Thiffault,^{21,22,23} Christine M. Freitag,¹⁵ Deborah Sara Barbouth,²⁰ Maxime Cadieux-Dion,²¹ Rebecca Willaert,⁹ Maria J. Guillen Sacoto,⁹ Nicole P. Safina,^{23,24,25} Christèle Dubourg,²⁶ Lauren Grote,^{23,24,25} Wilfrid Carré,²⁶ Carol Saunders,^{21,22,23} Sander Pajusalu,¹³ Emily Farrow,^{21,23,24} Anne Boland,²⁷ Danielle Hays Karłowicz,¹⁶ Jean-François Deleuze,²⁷ Monica H. Wojcik,²⁸ Rena Pressman,²⁰ Bertrand Isidor,^{2,4} Annick Vogels,¹⁸ Wim Van Paesschen,²⁹ Lihadh Al-Gazali,³⁰ Aisha Mohamed Al Shamsi,³¹ Mireille Claustres,¹ Aurora Pujol,³² Stephan Sanders,³³ François Rivier,³⁴ Nicolas Leboucq,³⁵ Benjamin Cogné,^{2,4} Souphatta Sasorith,¹ Damien Sanlaville,^{11,12} Kyle Retterer,⁹ Sylvie Odent,^{5,36} Nicholas Katsanis,³ Stéphane Bézieau,^{2,4} Michel Koenig,^{1,38} Erica E Davis,^{3,38,*} Laurent Pasquier,^{5,38} Sébastien Küry^{2,38,*}

Affiliations:

¹ EA7402 Institut Universitaire de Recherche Clinique, and Laboratoire de Génétique Moléculaire, CHU and Université de Montpellier, 34093 Montpellier, France;

² Service de Génétique Médicale, CHU Nantes, 9 quai Moncoussu, 44093 Nantes cedex 1, France;

³ Center for Human Disease Modeling, Duke University Medical Center, Durham, NC 27701, USA;

⁴ l'institut du thorax, INSERM, CNRS, UNIV Nantes, 44007 Nantes, France;

⁵ Service de Génétique Clinique, Centre Référence “Déficiences Intellectuelles de causes rares” (CRDI), Centre de référence anomalies du développement CLAD-Ouest, CHU Rennes, 35203 Rennes, France;

⁶ Division of Neurology, University Hospital Antwerp (UZA), 2610 Antwerp, Belgium;

⁷ Neurogenetics Group, Center for Molecular Neurology, VIB, 2650Antwerp, Belgium;

- ⁸ Laboratory of Neurogenetics, Institute Born-Bunge, University of Antwerp, 2650 Antwerp, Belgium;
- ⁹ GeneDx, 207 Perry Parkway, Gaithersburg, Maryland 20877, USA;
- ¹⁰ Chromosome Laboratory, Department of Clinical Genetics, Copenhagen University Hospital, Rigshospitalet, 2100 Copenhagen, Denmark;
- ¹¹ Service de génétique, Groupement Hospitalier Est, Hospices Civils de Lyon, Lyon, France;
- ¹² INSERM U1028, CNRS UMR5292, Centre de Recherche en Neurosciences de Lyon, Université Claude Bernard Lyon 1, Lyon, France;
- ¹³ Department of Clinical Genetics, United Laboratories, Tartu University Hospital and Institute of Clinical Medicine, University of Tartu, 2 L.Puusepa street, Tartu 51014, Estonia;
- ¹⁴ University of Minnesota -Fairview Minneapolis, MN 55454, USA;
- ¹⁵ Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, JW Goethe University Frankfurt, Deutschordenstraße 50, Frankfurt am Main, D-60528, Germany;
- ¹⁶ Division of Medical Genetics, Department of Pediatrics, Duke University, Durham, NC 27710, USA;
- ¹⁷ Epilepsy, Sleep and Pediatric Neurophysiology Department, Hospices Civils, Lyon, 69677 Bron, France;
- ¹⁸ Center for Human Genetics, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium;
- ¹⁹ Carle Physician Group, Urbana, IL 61801, USA;
- ²⁰ Dr. John T. Macdonald Foundation Department of Human Genetics, University of Miami, Miller School of Medicine, 1501 NW 10th Avenue, BRB, room 359 (M-860), Miami, FL 33136, USA;
- ²¹ Center for Pediatric Genomic Medicine, Children's Mercy Hospital, Kansas City, MO, 64108, USA;
- ²² Department of Pathology and Laboratory Medicine, Children's Mercy Hospitals, Kansas City, MO, 64108, USA;
- ²³ University of Missouri Kansas City, School of Medicine, Kansas City, MO, 64108, USA;
- ²⁴ Division of Clinical Genetics, Children's Mercy Hospital, Kansas City, MO, 64108, USA;
- ²⁵ Department of Pediatrics, Children's Mercy Hospital, Kansas City, MO, 64108, USA;
- ²⁶ Laboratoire de Génétique Moléculaire & Génomique, CHU de Rennes, 35033, Rennes, France;

²⁷ Centre National de Recherche en Génomique Humaine (CNRGH), Institut de Biologie François Jacob, DRF, CEA, Evry, France;

²⁸ The Broad Institute of MIT and Harvard, Cambridge, MA, 02142, USA;

²⁹ Department of Neurology, University Hospitals Leuven, Herestraat 49, 3000, Leuven, Belgium;

³⁰ Department of Paediatrics, College of Medicine and Health Sciences, United Arab Emirates University, PO box 17666, Al Ain, United Arab Emirates;

³¹ Department of Paediatrics, Tawam Hospital, PO box 15258, Al-Ain, United Arab Emirates;

³² Neurometabolic Diseases Laboratory, IDIBELL, Gran Via, 199, L'Hospitalet de Llobregat, 08908 Barcelona, and CIBERER U759, Center for Biomedical Research on Rare Diseases, 08908 Barcelona, Spain; Catalan Institution of Research and Advanced Studies (ICREA), 08010 Barcelona, Spain;

³³ Department of Psychiatry, UCSF Weill Institute for Neurosciences, University of California, San Francisco, San Francisco, California 94158, USA;

³⁴ Department of Neuropaediatrics and CR Maladies Neuromusculaires, CHU Montpellier, PhyMedExp, INSERM, CNRS, University of Montpellier, Montpellier, France;

³⁵ Neuroradiologie, CHU de Montpellier, 34090 Montpellier, France;

³⁶ CNRS UMR 6290, Université de Rennes, 2 Avenue du Professeur Léon Bernard, 35043 Rennes, France;

³⁷ equally contributed to this work;

³⁸ equally contributed to this work.

* Correspondence: Dr. Sébastien Küry, Service de Génétique Médicale, CHU Nantes, 9 quai Moncoussu, 44093 Nantes cedex 1, France. Tel: +33 2 44 76 82 85. Fax: +33 2 40 08 40 26. E-mail: sebastien.kury@chu-nantes.fr; Dr. Erica Davis, Center for Human Disease Modeling, Duke University Medical Center, 300 North Duke Street, Carmichael Building, Room 47-104, Durham, NC 27701, USA. E-mail: erica.davis@duke.edu.

Key words

RORA; neurodevelopmental disorder; intellectual disability; autistic features; cerebellar ataxia; epilepsy; dual molecular effects

Abstract

ROR α , the RAR-related orphan nuclear receptor alpha, is essential for cerebellar development. The spontaneous mutant mouse, *staggerer*, with an ataxic gait caused by neurodegeneration of cerebellar Purkinje cells, was discovered two decades ago to result from homozygous intragenic *Rora* deletions. However, *RORA* mutations were hitherto undocumented in humans. Through a multi-centric collaboration, we identified three copy number variant deletions (two *de novo*, one dominantly inherited in three generations); one *de novo* disrupting duplication; and eight *de novo* point mutations (three truncating, one canonical splice site and four missense mutations) involving *RORA* in 16 individuals from 13 families with variable neurodevelopmental delay and intellectual disability (ID)-associated autistic features, cerebellar ataxia and epilepsy. Consistent with the human and mouse data, disruption of the *D. rerio* ortholog, *rora*, causes significant reduction in the size of the developing cerebellum. Systematic *in vivo* complementation studies showed that, whereas wild-type human *RORA* mRNA could complement the cerebellar pathology, missense variants had two distinct pathogenic mechanisms of either haploinsufficiency or a dominant toxic effect according to their localization in the ligand-binding or DNA-binding domains, respectively. This dichotomous direction of effect is likely relevant to the phenotype in humans: individuals with loss of function variants leading to haploinsufficiency show ID with autistic features, while individuals with *de novo* dominant toxic variants present with ID, ataxia and cerebellar atrophy. Our combined genetic and functional data highlight the complex mutational landscape at the human *RORA* locus and suggest that dual mutational effects likely determine phenotypic outcome.

INTRODUCTION

Nuclear receptors appeared in the metazoan lineage as an adaptation to multicellular organization requiring distant cellular signaling through non-peptidic growth and differentiation factors.¹ The nuclear receptor superfamily is a group of transcription factors regulated by small hydrophobic hormones, such as retinoic acid, thyroid hormone, and steroids.² Mutations in nuclear receptors cause a diverse range of disorders, including central nervous system (CNS) pathologies, cancer, and metabolic disorders. For example, haploinsufficiency of *RORB* (OMIM: 601972), encoding the nuclear receptor ROR β , results in behavioral and cognitive impairment and epilepsy,³ while biallelic mutations in *RORC* (OMIM: 602943), encoding ROR γ , result in immunodeficiency (OMIM: 616622,⁴). Nuclear Receptor Subfamily 0, Group B, Member 1 (NR0B1, OMIM: 300473) is an orphan member of the nuclear receptor superfamily, and has been implicated in sex reversal (OMIM: 300018,⁵) and congenital adrenal hypoplasia (OMIM: 300200,⁶). ROR α (*RORA*) is most closely related to Retinoic Acid Receptor (*RAR*) yet functions differently; *RAR* acts as a ligand responsive heterodimer with retinoid X receptor (*RXR*). However, ROR α isoforms 1 and 2 constitutively activate transcription and bind DNA as monomers at responsive elements which consist of 6-bp AT-rich sequences.⁷ The amino-terminal domain of the ROR α 1 isoform determines its affinity and specific DNA-binding properties by acting in concert with the zinc finger domain.⁷

ROR α deficiency is known to cause the mouse *staggerer* (*sg*) phenotype, a cerebellar degenerative model.⁸ In humans, microdeletions overlapping *RORA* on 15q22.2 have been reported in affected individuals as part of a contiguous gene syndrome,⁹ with the smallest deletion involving two genes, NMDA receptor-regulated 2 (*NARG2/ICE2*) and *RORA*. All reported individuals with 15q22.2 microdeletion share epileptic seizures, mild intellectual disability (ID) and dysmorphic features, with variable ataxia.⁹ Here, we report 16 affected individuals from 13 syndromic ID families with intergenic or intragenic deletions; truncating mutations; or missense changes in *RORA*. We modeled these genetic findings in zebrafish larvae through endogenous *rora* ablation or heterologous expression of *RORA*, followed by relevant cerebellar phenotyping. In zebrafish models, we

recapitulated the neuroanatomical features of affected humans as well as the *staggerer* mouse, and we show that loss of *roraa* leads to reduction of both the Purkinje and granule compartments of the cerebellum. Further, our *in vivo* data indicate that missense variants in the DNA binding domain confer a dominant toxic effect, while a missense change in the ligand binding domain results in a loss of function effect. Together, our data highlight how different mutation effects at the *RORA* locus can produce overlapping but distinct phenotypic outcomes.

INDIVIDUALS AND METHODS

Genetic studies and ethics statement

Human genetic studies conducted in research laboratories were approved by local ethics committees from participating centers (Antwerp, Belgium; Lyon, France; Frankfurt, Germany; Copenhagen, Denmark; Montpellier, France; Tartu, Estonia; Minneapolis, Kansas City, USA). Written informed consent was obtained from all study participants. All 16 affected individuals underwent extensive clinical examination by at least one expert clinical geneticist. Routine genetic testing was performed whenever clinically relevant, including copy number variation (CNV) analysis by high-resolution array-based comparative genomic hybridization (aCGH) using: (1) 180k CytoSure ISCA v2 array (Oxford Gene Technology; individuals 10A-D); (2) 180k SurePrint G3 CGH microarray as described¹⁰ (Agilent; individual 11), or (3) 400k SurePrint G3 CGH microarray, as described¹¹ (Agilent, individuals 12 and 13). Affected individuals with a negative aCGH result underwent whole exome sequencing (WES) on an Illumina HiSeq platform according to the following paradigms: (1) trio-based clinical diagnostic WES (individuals 1, 2, 6, 7 and 9); (2) trio-based WES in a research laboratory (individuals 2, and 3); or (3) WES of an affected individual followed by single site testing in parental DNA samples (individuals 5 and 8; see Table S1 for further details). Point mutations were confirmed by Sanger sequencing of DNA sample from all available family members, when possible.

Establishment and culture of primary fibroblasts

We conducted biochemical studies in primary cultures of skin fibroblasts from individuals 2, 3 and 6 and from two control individuals (WT1 and WT2). Fibroblasts were grown in RPMI 1640 medium, containing 5% fetal calf serum (FCS, ThermoFisher, Waltham), 2 mM L-glutamine (ThermoFisher), 1% Ultrosor G (Pall, Cortland), 1X Antibiotic-Antimycotic (ThermoFisher) and were maintained in a humidified incubator at 37°C/5% CO₂. Both RT-PCR and Western blotting studies were performed using individual fibroblasts plated in 6-well plates (200,000 cells/well).

RT-PCR studies in primary fibroblasts

Total RNA was extracted using the RNeasy Plus Mini kit (Qiagen) and reverse transcribed with the Moloney Murine Leukemia Virus Reverse Transcriptase (ThermoFisher). PCR amplification was performed with primers located in exons 3 and 4 (Table S2) and the Promega MasterMix (Promega). PCR products were separated on a 1.5% agarose gel and fragments were cut from the gel. After extraction from gel slices, the PCR products were purified with ExoSAP-IT (ThermoFisher) and sequenced by standard cycle-sequencing reactions with Big Dye terminators (ThermoFisher) with the PCR forward and reverse primers in an ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems, Foster City). Mutation detection analysis was performed using 4Peaks.

RORA immunoblotting

Total protein lysate was extracted from primary fibroblasts using 1X Laemmli buffer. Proteins were separated on 10% SDS-PAGE gels and transferred to PVDF membrane (Westran® Clear Signal Whatman®; Dominique Dutscher, Brumath). The membranes were incubated overnight with 1:200 diluted anti-ROR α (sc-6062, Santa Cruz biotechnology, Dallas) primary antibody in 5% skim milk. Protein levels of the housekeeping protein GAPDH were assayed for internal control of protein loading with 1:40,000 diluted GAPDH antibody (sc-25778, Santa Cruz Biotechnology).

Three-dimensional (3D) protein modeling

Missense changes located in the DNA-binding domain of RORA isoform a were studied by 3D modeling. Wild type (WT) and mutated RORA DNA binding domain homology models were

generated according to the crystal structure of the RXR-RAR DNA-binding complex on the retinoic acid response element DR1 using Modeller¹² software with standard parameters.

RORA expression analysis in neuronal cell types

There are four RefSeq transcripts of *RORA* (GenBank), referred to as: NM_134261.2 (*RORA* 1); NM_134260.2 (*RORA* 2); NM_002943.3 (*RORA* 3); NM_134262.2 (*RORA* 4). Relative mRNA levels of the four *RORA* transcripts were assessed in three different commercially-obtained human cDNAs by quantitative (q)RT-PCR: (i) cerebellar cDNA from 26 year-old male (Amsbio); (ii) whole brain cDNA pooled from two males of Northern European descent aged 43-55, Human multiple tissue cDNA (MTC) panel I, (Clontech); (iii) Human Universal QUICK-Clone II, (Clontech) using transcript-specific primers (Table S2) and SYBR Green PCR Master Mix (Thermo Fisher) on an ABI 7900HT real-time PCR system. We determined relative gene expression levels in quadruplicate samples according to the Δ Ct method. After normalization to β -actin levels, $-\log^2$ of Δ Ct for each of the four transcripts was added together to determine total expression levels from the *RORA* locus for each cDNA. Relative *RORA* transcript expression levels expressed in $-\log^2$ of Δ Ct were obtained by calculating the percentage of each transcript compared to total *RORA* expression.

Zebrafish lines and husbandry

All zebrafish work was performed in accordance with protocols approved by the Duke University Institutional Animal Care and Use Committee. Zebrafish embryos were obtained by natural matings of WT (ZDR strain, Aquatica BioTech) or transgenic (*neurod:egfp*) adults¹³ and maintained on a 14h/10h light-dark cycle. Embryos were reared in embryo media (0.3 g/L NaCl, 75 mg/L CaSO₄, 37.5 mg/L NaHCO₃, 0.003% methylene blue) at 28°C until processing for phenotypic analyses at 3 days post-fertilization (dpf).

CRISPR/Cas9 genome editing of roraa in zebrafish

We designed two guide (g) RNAs targeting *roraa* (GRCz10: ENSDARG00000031768) using the CHOPCHOP v2 tool,¹⁴ and synthesized them *in vitro* with the GeneArt precision gRNA synthesis kit

(Table S2; Thermo Fisher) as described.¹⁵⁻¹⁷ We targeted the *rora* locus by microinjection into the cell of zebrafish embryos with 1 nl of cocktail containing 100 pg gRNA and 200 pg Cas9 protein (PNA Bio) at the one cell stage. We harvested individual embryos ($n=8$) at 1 dpf for DNA extraction to assess targeting efficiency. We PCR-amplified the region flanking the targeted site (Table S2), denatured the resulting product, and reannealed it slowly to form heteroduplexes (95°C for 5 min, ramped down to 85°C at 1°C/s and then to 25°C at 0.1°C/s). We performed polyacrylamide gel electrophoresis (PAGE) on a 20% precast 1 mm gel (Thermo Fisher) to visualize heteroduplexes. To estimate mosaicism of F0 mutants, PCR products were cloned into a TOPO-TA vector (Thermo Fisher) and individual colonies ($n=24$) were sequenced ($n=3$ larvae/gRNA).

Transient rora suppression, in vivo complementation and heterologous expression experiments

We designed splice blocking (sb) morpholinos (MO) targeting either the splice donor site of exon 2 (e2i2) or exon 3 (e3i3) of *rora* (GeneTools, LLC; Table S2), and injected 1 nl of MO into the yolk of zebrafish embryos at one-to-four cell stage. To confirm expression of the two annotated *rora* transcripts (GRCz10: ENSDART00000148537.2 and ENSDART00000121449.2) and to determine MO efficiency, we harvested uninjected control and MO-injected larvae in Trizol (Thermo Fisher) at 3 dpf, extracted total RNA, and conducted first-strand cDNA synthesis with the QuantiTect Reverse Transcription kit (Qiagen). The targeted region of *rora* was PCR-amplified using primers complementary to sites in flanking exons (Table S2), and migrated by electrophoresis on a 1 % agarose gel; bands were excised, gel purified using QIAquick gel extraction kit (Qiagen) and resulting clones were Sanger sequenced. The optimal MO dose for *in vivo* complementation experiments was determined by injection of three concentrations of MO (2, 3, 4 ng of e2i2; or 6, 7, 8 ng of e3i3). To generate mRNA for injections, we purchased Gateway-compatible open reading frame (ORF) clones from Genecopoeia (*RORA* 1; *RORA* 2; and *RORA* 3) or Thermo Fisher (*RORA* 4) and transferred the ORFs to a pCS2+ vector by LR clonase II-mediated recombination (Thermo Fisher). We performed site-directed mutagenesis of *RORA* 4 according to the QuikChange protocol (Table S2; Agilent), using described methodology;¹⁸ sequences were validated by Sanger sequencing. Linearized pCS2+ vectors

containing WT or mutant ORFs were transcribed *in vitro* with the mMessage mMachine SP6 Transcription kit (Ambion).

Whole-mount immunostaining

We stained axonal tracts of the cerebellum with monoclonal anti-acetylated tubulin antibody produced in mouse (Sigma-Aldrich, T7451, 1:1000), as described.¹⁹ We fixed larvae in Dent's solution (80% methanol and 20% DMSO) and carried out primary antibody detection overnight, and secondary detection for one hour with Alexa Fluor 488 goat anti-mouse IgG (A11001, Invitrogen; 1:1000). We visualized the Purkinje compartment of the zebrafish cerebellum with anti-zebrin II antibody (a gift from Dr. Richard Hawkes) in *neurod:egfp* transgenic larvae. Briefly, zebrafish larvae were fixed in 4% paraformaldehyde and incubated overnight in mouse anti-zebrin II antibody (1:100); and secondary antibody was applied for one hour (Alexa Fluor 594 goat anti-mouse IgG, A11005, Invitrogen; 1:1000). Fluorescent signal was imaged manually on dorsally positioned larvae using an AxioZoom.V16 microscope and AxioCam 503 monochromatic camera, using Zen Pro 2012 software (Zeiss). Cerebellar structures of interest or optic tecta were measured using ImageJ.²⁰ Total cerebellar area was measured on acetylated tubulin stained larvae by outlining structures with fluorescent signal; regions comprised of Purkinje cells were measured on zebrin II stained regions; region comprised of granule cells were measured on GFP positive regions. Statistical analyses were performed using a two-tailed parametric *t*-test (GraphPad software).

RESULTS

Identification of point mutations or copy number variants disrupting RORA

As part of our ongoing studies to understand the molecular basis of neurodevelopmental disorders, we identified a total of 16 individuals with rare variants suspected to alter *RORA* function (Figure 1; Table 1; Table S3; Table S4). The first individual under investigation was a female with severe syndromic ID, multifocal seizures, mild cerebellar hypoplasia and hypotonia (individual 6, Table 1; Figure 2A). Upon performing WES, we identified a *de novo* frameshifting mutation in *RORA* (GenBank ID:

NM_134261.2, c.1019delG; p.(Arg340Profs*17)) that was not present in the genome aggregation database (gnomAD; >246,000 chromosomes) or the NHLBI Exome Variant Server (EVS; >13,000 alleles). Information exchange on community data sharing platforms including GeneMatcher²¹, the DatabasE of genomic variation and Phenotype in Humans using Ensembl Resources (DECIPHER), and the Broad Institute matchbox repository, facilitated the identification of an additional 15 affected individuals with *RORA* variants who displayed overlapping phenotypes. These variants were present *de novo* in 11 simplex families with affected individuals and segregated in one three-generation pedigree under a dominant paradigm, bolstering the candidacy of this locus. Of note, segregation analysis was impossible in individual 1's family, since the child had been adopted.

Four of the 16 affected individuals harbored likely pathogenic SNVs predicted to alter *RORA* protein sequence or dose. In addition to individual 6, three affected individuals had *de novo* SNVs or small indels in *RORA*; these included one frameshifting mutation; one nonsense mutation; and one mosaic canonical splice acceptor site change (Figure 1A; Table 1; Table S3). These three *RORA* variants are predicted to result in protein sequence with either complete (c.804_805delGT; p.Ser269Hisfs*13), or partial truncation (c.1019delG; p.Arg340Profs*17 and c.1498C>T; p.Arg500*) of the ligand binding domain at the C-terminus and/or nonsense-mediated mRNA decay (NMD). Notably, analyses of fibroblast protein lysates from individual 6 (p.Arg340Profs*17) showed that *RORA* protein levels were not decreased compared to controls, suggesting that related mRNA harboring the premature stop codon would not be eliminated by NMD. According to Alamut software (compiling five different prediction tools: Splicesite Finder-like, MaxEntScan, NNSPLICE, GeneSplicer and Human Splicing Finder), the splice acceptor site mutation, c.425-1G>A, is predicted to cause in-frame exon skipping and deletion of 77 amino acids (p.Ala142_Leu273del); however, cell lines were unavailable from the affected individual to confirm this prediction; mosaicism was estimated at about 20% from exome and Sanger sequencing data.

An additional seven affected individuals had non-recurrent CNVs impacting the *RORA* locus detected by high-resolution aCGH (Table 1; Figure 1A-C; Figure S1; Table S4). We identified one ~63 kb

duplication interrupting *RORA* exons 3-6, which is predicted to result in a premature termination of the protein (p.Gln315Leufs*51); one intragenic CNV that resulted in a ~27 kb deletion of *RORA* exon 3 and its flanking intronic regions to produce a putative frameshifting truncation; and one ~3.7 Mb intergenic deletion impacting 17 genes on 15q22.3q22.2 including *RORA*. Segregation in each of the three families showed that all three CNVs occurred *de novo*. Furthermore, we detected a ~1.5 Mb intergenic deletion in four individuals segregating neurological phenotypes in an autosomal dominant pedigree (Figure 1B); this CNV encompasses nine genes, including *RORA*.

Finally, an additional five affected individuals had missense mutations predicted to be deleterious (Table 1; Table S3; Figure S2). All five alleles were absent from all available public databases queried (gnomAD, ExAC, EVS). Two of the five variants, c.1225A>C (p.Ser409Arg) and c.1385G>A (p.Arg462Gln), map to the ligand binding domain; both residues are conserved in vertebrate orthologues of ROR α but not in the human paralogous nuclear receptor proteins, indicating that these positions are specific to the ROR sub-family. The other three variants c.269C>G (p.Cys90Ser), c.275G>C (p.Gly92Ala) and c.281A>G (p.Lys94Arg) are located in the conserved zinc-finger DNA-binding domain of ROR α . The three residues Cys90, Gly92 and Lys94, together with Cys93, belong to the P box motif that is part of the alpha-helix of the first zinc-finger domain of the nuclear receptors and interacts directly with DNA (Figure S3A).²² Because the c.281A>G variant lies in the donor splice site of exon 3, we examined whether it impacts mRNA splicing in primary skin fibroblasts from individual 3; RT-PCR showed neither abnormal sized product nor semi-quantitative differences in cDNA amounts (Figure S4A, B). Additionally, Western blotting of lysates from primary fibroblasts harboring the c.275G>C change (individual 2) showed a modest increase of ROR α levels when compared to a matched control (Figure S4C).

Clinical features of individuals with RORA variants

The 16 affected individuals in our cohort display complex phenotypes with regards to their cognition, motor function, and electrophysiology (Table 1). The predominant phenotype was ID. However, cognitive function was variable among the 16 individuals, and ranged from mild to moderate (13/16)

to severe (2/16), while one individual had a low intelligence quotient (IQ) without ID. We noted IQ regression in one affected individual (individual 4) who had cognitive decline at ~10 years of age. Developmental milestones were also delayed. The mean age of walking was delayed in 8/12 individuals; 11/12 walked by 3 years of age. Further, 11/13 individuals had speech delay and/or poor verbal communication abilities, and 2 were not speaking in sentences by the age of 6. We diagnosed epilepsy in 11/16 individuals (Table 1); the predominant seizure semiology was that of a generalized epilepsy with absences, drop attacks and tonic-clonic seizure sub-types. None of the patients with *RORA* intragenic mutation had overt dysmorphic facial features. Of the nine individuals who underwent brain MRI, six had normal results and three were diagnosed with cerebellar hypoplasia, which predominantly affected the vermis (individuals 2 and 3; Figure 2B,C). These individuals developed early onset ataxia and hypotonia by age 1.

Modeling RORA disruption in zebrafish

To corroborate the *Rora*^{sg} mouse mutant phenotype data,^{8,23} and to determine the effect of missense variants identified in affected individuals, we developed zebrafish models of *RORA* ablation and expression. We and others have shown previously that zebrafish is a robust model of neuroanatomical phenotypes observed in humans.^{19,24–26} In particular, assay of cerebellar defects in zebrafish has provided crucial insights toward understanding underlying pathomechanism,^{27–29} especially given the high conservation of granule and Purkinje cell types between mammals and teleost species.³⁰ First, we aimed to model *RORA* disruption *in vivo* by targeting and knocking down the relevant zebrafish ortholog. Through reciprocal BLAST of the zebrafish genome and four annotated human *RORA* transcripts (NM_134261, NM_134260, NM_002943 and NM_134262; Figure S5A), we identified two zebrafish orthologs: *roraa* (Ensembl ID: ENSDARG00000031768; GRCz10; 88%, 91%, 91%, and 91% identity to proteins encoded by *RORA* 1, *RORA* 2, *RORA* 3, *RORA* 4, respectively) and *rorab* (Ensembl ID: ENSDARG00000001910; 69%, 72%, 72%, 73% identity to *RORA* 1, *RORA* 2, *RORA* 3, *RORA* 4, respectively). Next, we considered endogenous expression data to determine the most appropriate *D. rerio* transcript(s) to modulate *in vivo*. RNA *in situ* hybridization studies in zebrafish larvae have documented *roraa* expression patterns in the developing cerebellum as well as other

anterior structures (optic tecta, hindbrain and retina). However, *rorab* expression is restricted to the hindbrain.^{31,32} Considering both the amino acid conservation and also the spatiotemporal expression of *roraa* in the developing cerebellum, we deemed *roraa* as the most relevant *D. rerio* locus to test the activity of *RORA* mutations in humans.

Haploinsufficiency or knockout of *Rora* in mice lead to abnormal cerebellar layer morphology.^{8,23} Therefore, to assess the effect of loss of *RORA* function consistent with deletion or truncating mutations identified in affected individuals, we compared the size of the developing cerebellum between *roraa* knock-down and uninjected controls. The zebrafish *roraa* locus has two annotated transcripts (GRCz10: [*roraa*-201] ENSDART00000121449.2 and [*roraa*-202] ENSDART00000148537.2), the latter of which has an incomplete 5' coding sequence; RT-PCR using cDNA originating from 3 dpf embryos confirmed that both transcripts are detectable in this time window (Figure S6A, B). Next, we generated CRISPR/Cas9-based zebrafish F0 mutant models by targeting exon 5 or exon 8 of *roraa*-201 (corresponding to exon 4 or exon 7 of *roraa*-202) to produce mosaic mutants with >90% mosaicism (Figure S7). Immunostaining of the central nervous system (CNS) with anti-acetylated tubulin antibody and measurements of neuroanatomical structures showed that *roraa* F0 mutants display a reduced cerebellar area and smaller optic tecta area compared to either control larvae or larvae injected with gRNA alone ($P < 0.0001$, for gRNA 1 and gRNA 2, n=33-43 and 41-45 larvae/batch respectively, repeated, masked scoring; Figure 3A-C; Figure S8A-C).

To assess phenotype specificity and to determine the effect of *RORA* missense variants, we performed MO-mediated transient suppression of *roraa*. We designed two sb MOs targeting the splice donor site of exons 2 and 3 of *roraa*-201, which we injected into embryo batches, generated cDNA at 3 dpf and performed RT-PCR to determine efficiency; subsequent Sanger sequencing confirmed a frameshifting deletion of *roraa* exon 2 or exon 3 (Figure S6C, D). Next, we injected increasing concentrations of either MO (2, 3, 4 ng e2i2; or 6, 7, 8 ng e3i3). Consistent with our CRISPR F0 mutant data, we observed a dose-dependent reduction in cerebellar size for both reagents ($P < 0.0001$, 30 to 94 larvae/condition, Figure S6E). Co-injection of e2i2 sb MO with each of the four RefSeq annotated WT

human *RORA* mRNAs rescued cerebellar and optic tecta defects, indicating MO specificity ($P=0.0060$, 0.0387 , <0.0001 , <0.0001 for *RORA* 1, 2, 3 and 4, respectively vs. MO, $n=31-60$ larvae/batch, repeated, Figure 3B and D; Figure S8A and D). In parallel, heterologous expression of each of the four WT *RORA* mRNAs did not lead to cerebellar phenotypes that differed from uninjected controls ($n=40-52$ larvae/batch, Figure S9). To investigate further the relative expression of the four annotated *RORA* transcripts and to identify the most relevant isoform for *in vivo* modeling in zebrafish, we performed qRT-PCR on commercial human adult cDNA; we identified *RORA* 4 as the most abundantly expressed transcript in the human CNS (Figure S5A and B), potentially explaining the observation that WT *RORA* 4 mRNA produced the most significant restoration of cerebellar size when co-injected with MO.

Altered Purkinje and granule layers in roraa zebrafish models

Similar to mammals, Purkinje and granule cell crosstalk is vital for the function of the zebrafish cerebellum.³⁰ Cerebellar cell subpopulations have been characterized previously in zebrafish with lineage specific markers and transgenic lines. In *neurod:egfp* transgenic zebrafish, granule cells are marked with GFP.²⁹ Furthermore, zebrin II is a specific marker of the Purkinje compartment,²⁷ which develops between 2.3 and 4 dpf.³³ To assess whether *roraa* suppression affects granule or Purkinje cell layers of zebrafish cerebellum, we conducted whole mount zebrin II immunostaining on *neurod:egfp* transgenic zebrafish larvae at 3 dpf. F0 *roraa* mutants presented with a significantly decreased size of Purkinje and granule cell layers compared to controls ($P < 0.0001$ for both cell types; Purkinje cells, $n=26-55$; granule cells, $n=11-55$, repeated; Figure 3E-H; Figure S10A-C). Importantly, gRNA injected alone was indistinguishable from controls (Figure 3G, H). Injection of e2i2 MO recapitulated these observations ($P < 0.0001$ or $P < 0.001$ for Purkinje [$n=29-55$] and granule cells [$n=30-55$], respectively in morphants vs. controls; repeated; Figure 3E-H; Figure S10). Further, co-injection of e2i2 MO with WT human *RORA* 4 mRNA, rescued the measured area of each of the granule and Purkinje compartments ($P < 0.0001$ or $P < 0.001$ vs MO alone for Purkinje or granule cells, respectively; Figure 3E-H; Figure S10). Together, our data confirm that in F0 mutant or transient

knockdown zebrafish models, *rora* disruption leads to defects of Purkinje and granule cell layers, consistent with the *Rora*^{sg} mouse model.³⁴⁻³⁶

In vivo complementation studies indicate dual direction of RORA allele effect

To determine the pathogenicity of *RORA* missense variants (c.275G>C p.(Gly92Ala), c.281A>G p.(Lys94Arg) and c.1385G>A p.(Arg462Gln)), we assessed the effect of mutant *RORA* 4 mRNA in the presence or absence of MO. First, we tested a loss of function hypothesis by co-injecting *rora* e2i2 MO with WT or variant *RORA* 4 mRNA. *In vivo* complementation of *rora* MO with p.(Gly92Ala) or p.(Lys94Arg) encoding mRNA resulted in early developmental defects including a reduction in the size of anterior structures and tail extension failure (65% and 69%, respectively) and a high mortality rate (19% and 10%, respectively), while co-injection of MO with p.(Arg462Gln), WT, or with a potentially benign variant p.(Thr476Ala) (rs190933482, minor allele frequency 0.001 in gnomAD) induced no appreciable morphological defects (Figure 4A). The gross morphological phenotype observed by co-injection of MO with mutant mRNAs located in the DNA binding domain was more severe than the morphant phenotype, and suggested either a dominant negative or a gain of function effect. To investigate these possibilities, we injected 200 pg of each mutant mRNA alone, and again observed notable mortality at 3 dpf (2% for controls (1 dead larva/54) versus 31% for p.(Gly92Ala) (14/45); 9% for p.(Lys94Arg) (4/46); and 7% for p.(Arg462Gln) (3/41)). Of the larvae that survived to 3 dpf, we performed anti-acetylated immunostaining and measurement of the cerebellar area; p.(Gly92Ala) encoding mRNA induced a reduction in the mean cerebellar area compared to WT ($p < 0.0001$, 31-53 larvae/batch, Figure 4B). To determine whether WT mRNA could rescue the phenotypes induced by p.(Gly92Ala), we titrated increasing amounts of p.(Gly92Ala) mRNA together with *RORA* 4 WT mRNA and observed a dose-dependent effect that correlated with the injected dose of p.(Gly92Ala) (Figure 4C-D). These data suggested a dominant toxic effect as the likely mechanism for the p.(Gly92Ala) variant. Although expression of p.(Lys94Arg) induced morphological defects similar to p.(Gly92Ala) and we observed a broadened distribution of cerebellar measurements at the highest dose of mutant mRNA that was consistent with p.(Gly92Ala), these results did not reach statistical significance (Figure 4E, F). Finally, *in vivo* complementation of *rora*

MO with p.(Arg462Gln) encoding mRNA did not rescue the size of the cerebellum (Figure 4G-H, $p < 0.001$, $n = 40-41$), suggesting a loss of function effect as the possible disease mechanism for this change impacting the ligand binding domain (Figure 4G, H).

DISCUSSION

Here, we describe a cohort of 16 affected individuals who harbor 13 different rare variants disrupting *RORA*. We observe a clinical spectrum of neurodevelopmental delay with at least two different presentations: (1) a cognitive and motor phenotype; and (2) a cognitive and behavioral phenotype. The first sub-phenotype is characterized by a moderate to severe ID with a marked ataxic component, severe cerebellar vermis hypoplasia and epilepsy with predominant generalized seizures as already reported in *RORB*³; whereas the hallmark of the second sub-phenotype is ASD with mild ID or normal cognition frequently associated with epilepsy. In addition to these two groups, three individuals exhibit only mild ID without behavioral problems.

Although the size of our cohort limits the delineation of an unambiguous genotype-phenotype correlation, we can infer pathomechanisms related to specific phenotypes from our *in vivo* complementation data. Individuals 2 and 3, who harbor the two dominant toxic mutations of the *RORα* DNA binding domain display severe ID and motor phenotypes likely due to cerebellar hypoplasia. A similar severe phenotype is coincident with the truncating deletion (individual 6; p.(Arg340Profs*17)), raising the possibility that this mutation could also result in a dominant toxic effect by encoding a truncated protein; immunoblotting studies from fibroblast protein lysates showed *RORA* protein levels to be similar to that of controls, arguing against haploinsufficiency (Figure S4C). Individual 6 also displayed autistic traits, however these phenotypes occurred secondary to established ID, potentially excluding her from the Autism Diagnosis Interview (ADI) criteria for idiopathic ASD.^{37,40}

By contrast, we note variability in cognitive function and behavioral phenotypes in individuals with

likely haploinsufficiency. One individual displays an ASD phenotype in the absence of ID (individual 8), and *in vivo* complementation testing in zebrafish indicate that the p.(Arg462Gln) mutation confers a loss of function effect. Further, 6/12 individuals from simplex families who likely have a reduction in protein dosage display mild to moderate ID but with normal behavior, while the remaining 5/12 affected individuals from simplex families present with both ID and autistic features. Notably, there is phenotypic variability among the individuals within multiplex family 10; all four individuals display mild ID, but only two display behavioral anomalies. However, we cannot exclude the possibility of *trans* effects from elsewhere in the genome, especially from *DISC1* affected by the CNVs harbored by individuals 10A-C, or individual 11 (Figure S1; Table S4). Phenotypic variability is not uncommon for neurodevelopmental disorders,³⁸ and here we account, in part, for variability due to allelism at the *RORA* locus by elucidating the direction of allele effect for missense changes.

The sub-group of our cohort who show a constellation of cognitive and motor defects are reminiscent of the homozygous *staggerer* (*sg/sg*) mouse, an early reported animal model of *Rora* ablation. The *staggerer* mutation consists of an intragenic CNV that results in a 122-bp frameshifting deletion that truncates the ligand binding domain, leading to the loss of ROR α activity.⁸ This phenotype is similar to *Rora*^{-/-} mice³⁹ in which the most obvious symptom is an ataxic gait associated with defective Purkinje cell development leading to an abnormal cerebellar size.⁴¹ Heterozygous *Rora*^{+/-} mice present with a comparable phenotype, although they display a late onset of neuronal loss and reduced phenotypic severity.⁴² Further studies will be required to understand the precise molecular mechanisms of p.(Gly92Ala) and p.(Lys94Arg) to account for their apparent toxic effects. We speculate that these changes in the DNA binding domain might hamper access of WT ROR α to its natural target sites (Figure S3B), thereby leading to a phenotype resembling to the *staggerer* and *Rora*^{-/-} homozygous mutants.

The autistic signs observed in two individuals with truncating mutations (Individuals 6 and 13) and two individuals with missense mutations altering the ligand binding domain of *RORA* (Individuals 7

and 8), are in agreement with recent reports suggesting that *RORA* is a candidate gene for ASD.⁴³ ChIP-on-chip analysis has revealed that ROR α can be recruited to the promoter regions of 2,544 genes across the human genome, with a significant enrichment in biological functions including neuronal differentiation, adhesion and survival, synaptogenesis, synaptic transmission and plasticity, and axonogenesis, as well as higher level functions such as development of the cortex and cerebellum, cognition, memory, and spatial learning.⁴⁴ Independent ChIP-quantitative PCR analyses confirmed binding of RORA to promoter regions of selected ASD-associated genes, including *A2BPI*, *CYP19A1*, *ITPR1*, *NLGNI*, and *NTRK2*, whose expression levels are also decreased in *RORA*-repressed human neuronal cells and in prefrontal cortex tissues from individuals with ASD.⁴⁴ Additionally, two *RORA* polymorphisms (rs11639084 and rs4774388) have been associated with ASD risk in Iranian individuals.⁴⁵ Consistent with these data, treatment with a synthetic ROR α/γ agonist, SR1078, reduced repetitive behavior in the BTBR mouse model of autism, suggesting that *RORA* upregulation could be a viable therapeutic option for ASD.⁴⁶

In summary, our data implicate a diverse series of disruptive mutations in *RORA* with neurological phenotypes hallmarked by ID and either severe motor phenotypes or behavioral anomalies. Through combined clinical, genetic, and functional studies, we expand the genetic basis of rare neurodevelopmental syndromes, and show how *in vivo* modeling can reveal dual molecular mutational effects.

Supplemental Data

Supplemental Data include ten figures and four tables.

Acknowledgments

We are grateful to Dr. R. Hawkes and Dr. C. Armstrong (University of Calgary) for providing the zebrin II antibody. We also thank Mr. Z. Kupchinsky for zebrafish husbandry; Mr. D. Morrow for assisting with reagents for the *in vivo* modeling study; and members of the Center for Human Disease Modeling for helpful comments. We are also grateful to Ms. J. A. Rosenfeld (Mokry) for her precious

help in our search for similar affected individuals and for her reviewing of the manuscript. This work was supported by funds from the Agence Nationale pour la Recherche/E-rare Joint-Transnational-Call 2011 (2011-RARE-004-01 “Euro-SCAR”) to MK; from the National Human Genome Research Institute with supplemental funding provided by the National Heart, Lung, and Blood Institute under the Trans-Omics for Precision Medicine (TOPMed) program and the National Eye Institute (to the Broad Center for Mendelian Genomics (UM1 HG008900)); from the NIH grant T32 HD07466 (to M.H.W.); US NIH grant R01 MH106826 to E.E.D.; and from the Estonian Research Council, grant PUT355 (to S.P. and K.Õ.). We also acknowledge the Deciphering Developmental Disorders (DDD) Study.

Conflict of interest:

Kirsty McWalter, Megan T. Cho, Rebecca Willaert, Maria J. Guillen Sacoto, and Kyle Retterer are employees of GeneDx, Inc., a wholly owned subsidiary of OPKO Health, Inc. N.K. is a paid consultant for and holds significant stock of Rescindo Therapeutics, Inc.

Web Resources

1000 Genomes, <http://www.1000genomes.org/>

CHOPCHOP v2, <http://chopchop.cbu.uib.no/>

Database of Genomic Variants, <http://dgv.tcag.ca/dgv/app/home/>

DECIPHER, <https://decipher.sanger.ac.uk/>

dbSNP, <http://www.ncbi.nlm.nih.gov/projects/SNP/>

Ensembl Genome Browser, <https://www.ensembl.org/>

ExAC Browser, <http://exac.broadinstitute.org/>

Fiji, <https://fiji.sc/>

GenBank, <http://www.ncbi.nlm.nih.gov/genbank/>

GeneMatcher, <https://genematcher.org/>

gnomAD browser, <http://gnomad.broadinstitute.org/>

The Human Protein Atlas, <http://www.proteinatlas.org/>

Matchbox (Broad Institute), <https://seqr.broadinstitute.org/>

NHLBI Exome Sequencing Project (ESP) Exome Variant Server, <http://evs.gs.washington.edu/EVS/>

OMIM, <http://www.omim.org/>

PLINK, <http://pngu.mgh.harvard.edu/~purcell/plink/>

UCSC Genome Browser, <http://genome.ucsc.edu>

UniProt, <http://www.uniprot.org/>

ClustalW, <http://www.clustal.org/>

QuickChange, <http://www.genomics.agilent.com/primerDesignProgram.jsp>

Zfin, <https://zfin.org>

References

1. Escriva, H., Langlois, M.C., Mendonça, R.L., Pierce, R., and Laudet, V. (1998). Evolution and diversification of the nuclear receptor superfamily. *Ann. N. Y. Acad. Sci.* 839, 143–146.
2. Sladek, F.M. (2011). What are nuclear receptor ligands? *Mol. Cell. Endocrinol.* 334, 3–13.
3. Rudolf, G., Lesca, G., Mehrjouy, M.M., Labalme, A., Salmi, M., Bache, I., Bruneau, N., Pendziwiat, M., Fluss, J., de Bellescize, J., et al. (2016). Loss of function of the retinoid-related nuclear receptor (RORB) gene and epilepsy. *Eur. J. Hum. Genet. EJHG* 24, 1761–1770.
4. Okada, S., Markle, J.G., Deenick, E.K., Mele, F., Averbuch, D., Lagos, M., Alzahrani, M., Al-Muhsen, S., Halwani, R., Ma, C.S., et al. (2015). IMMUNODEFICIENCIES. Impairment of immunity to *Candida* and *Mycobacterium* in humans with bi-allelic RORC mutations. *Science* 349, 606–613.
5. Bardoni, B., Zanaria, E., Guioli, S., Floridaia, G., Worley, K.C., Tonini, G., Ferrante, E., Chiumello, G., McCabe, E.R., and Fraccaro, M. (1994). A dosage sensitive locus at chromosome Xp21 is involved in male to female sex reversal. *Nat. Genet.* 7, 497–501.
6. Muscatelli, F., Strom, T.M., Walker, A.P., Zanaria, E., Récan, D., Meindl, A., Bardoni, B., Guioli, S., Zehetner, G., and Rabl, W. (1994). Mutations in the DAX-1 gene give rise to both X-linked adrenal hypoplasia congenita and hypogonadotropic hypogonadism. *Nature* 372, 672–676.
7. Giguère, V., Tini, M., Flock, G., Ong, E., Evans, R.M., and Otulakowski, G. (1994). Isoform-specific amino-terminal domains dictate DNA-binding properties of ROR alpha, a novel family of orphan hormone nuclear receptors. *Genes Dev.* 8, 538–553.
8. Hamilton, B.A., Frankel, W.N., Kerrebrock, A.W., Hawkins, T.L., FitzHugh, W., Kusumi, K., Russell, L.B., Mueller, K.L., van Berkel, V., Birren, B.W., et al. (1996). Disruption of the nuclear hormone receptor RORalpha in staggerer mice. *Nature* 379, 736–739.
9. Yamamoto, T., Mencarelli, M.A., Di Marco, C., Mucciolo, M., Vascotto, M., Balestri, P., Gérard, M., Mathieu-Dramard, M., Andrieux, J., Breuning, M., et al. (2014). Overlapping microdeletions involving 15q22.2 narrow the critical region for intellectual disability to NARG2 and RORA. *Eur. J. Med. Genet.* 57, 163–168.
10. Boutry-Kryza, N., Labalme, A., Till, M., Schluth-Bolard, C., Langue, J., Turleau, C., Edery, P., and Sanlaville, D. (2012). An 800 kb deletion at 17q23.2 including the MED13 (THRAP1) gene,

revealed by aCGH in a patient with a SMC 17p. *Am. J. Med. Genet. A* 158A, 400–405.

11. Grønborg, S., Kjaergaard, S., Hove, H., Larsen, V.A., and Kirchoff, M. (2015). Monozygotic twins with a de novo 0.32 Mb 16q24.3 deletion, including TUBB3 presenting with developmental delay and mild facial dysmorphism but without overt brain malformation. *Am. J. Med. Genet. A* 167A, 2731–2736.

12. Sali, A., and Blundell, T.L. (1993). Comparative protein modelling by satisfaction of spatial restraints. *J. Mol. Biol.* 234, 779–815.

13. Obholzer, N., Wolfson, S., Trapani, J.G., Mo, W., Nechiporuk, A., Busch-Nentwich, E., Seiler, C., Sidi, S., Söllner, C., Duncan, R.N., et al. (2008). Vesicular glutamate transporter 3 is required for synaptic transmission in zebrafish hair cells. *J. Neurosci. Off. J. Soc. Neurosci.* 28, 2110–2118.

14. Montague, T.G., Cruz, J.M., Gagnon, J.A., Church, G.M., and Valen, E. (2014). CHOPCHOP: a CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Res.* 42, W401–407.

15. Küry, S., Besnard, T., Ebstein, F., Khan, T.N., Gambin, T., Douglas, J., Bacino, C.A., Craigen, W.J., Sanders, S.J., Lehmann, A., et al. (2017). De Novo Disruption of the Proteasome Regulatory Subunit PSMD12 Causes a Syndromic Neurodevelopmental Disorder. *Am. J. Hum. Genet.* 100, 352–363.

16. Stankiewicz, P., Khan, T.N., Szafranski, P., Slattery, L., Streff, H., Vetrini, F., Bernstein, J.A., Brown, C.W., Rosenfeld, J.A., Rednam, S., et al. (2017). Haploinsufficiency of the Chromatin Remodeler BPTF Causes Syndromic Developmental and Speech Delay, Postnatal Microcephaly, and Dysmorphic Features. *Am. J. Hum. Genet.* 101, 503–515.

17. Ta-Shma, A., Khan, T.N., Vivante, A., Willer, J.R., Matak, P., J alas, C., Pode-Shakked, B., Salem, Y., Anikster, Y., Hildebrandt, F., et al. (2017). Mutations in TMEM260 Cause a Pediatric Neurodevelopmental, Cardiac, and Renal Syndrome. *Am. J. Hum. Genet.* 100, 666–675.

18. Niederriter, A.R., Davis, E.E., Golzio, C., Oh, E.C., Tsai, I.-C., and Katsanis, N. (2013). In vivo modeling of the morbid human genome using *Danio rerio*. *J. Vis. Exp. JoVE* e50338.

19. Margolin, D.H., Kousi, M., Chan, Y.-M., Lim, E.T., Schmahmann, J.D., Hadjivassiliou, M., Hall, J.E., Adam, I., Dwyer, A., Plummer, L., et al. (2013). Ataxia, dementia, and hypogonadotropism caused by disordered ubiquitination. *N. Engl. J. Med.* 368, 1992–2003.

20. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675.
21. Sobreira, N., Schiettecatte, F., Valle, D., and Hamosh, A. (2015). GeneMatcher: a matching tool for connecting investigators with an interest in the same gene. *Hum. Mutat.* 36, 928–930.
22. Ruff, M., Gangloff, M., Marie Wurtz, J., and Moras, D. (2000). Estrogen receptor transcription and transactivation: Structure-function relationship in DNA- and ligand-binding domains of estrogen receptors. *Breast Cancer Res* 2, 353–359.
23. Sidman, R.L., Lane, P.W., and Dickie, M.M. (1962). Staggerer, a new mutation in the mouse affecting the cerebellum. *Science* 137, 610–612.
24. Marin-Valencia, I., Novarino, G., Johansen, A., Rosti, B., Issa, M.Y., Musaev, D., Bhat, G., Scott, E., Silhavy, J.L., Stanley, V., et al. (2017). A homozygous founder mutation in TRAPPC6B associates with a neurodevelopmental disorder characterised by microcephaly, epilepsy and autistic features. *J. Med. Genet.* 55, 48–54.
25. Schaffer, A.E., Eggens, V.R.C., Caglayan, A.O., Reuter, M.S., Scott, E., Coufal, N.G., Silhavy, J.L., Xue, Y., Kayserili, H., Yasuno, K., et al. (2014). CLP1 founder mutation links tRNA splicing and maturation to cerebellar development and neurodegeneration. *Cell* 157, 651–663.
26. Borck, G., Hög, F., Dentici, M.L., Tan, P.L., Sowada, N., Medeira, A., Gueneau, L., Thiele, H., Kousi, M., Lepri, F., et al. (2015). BRF1 mutations alter RNA polymerase III-dependent transcription and cause neurodevelopmental anomalies. *Genome Res.* 25, 155–166.
27. Akizu, N., Cantagrel, V., Zaki, M.S., Al-Gazali, L., Wang, X., Rosti, R.O., Dikoglu, E., Gelot, A.B., Rosti, B., Vaux, K.K., et al. (2015). Biallelic mutations in SNX14 cause a syndromic form of cerebellar atrophy and lysosome-autophagosome dysfunction. *Nat. Genet.* 47, 528–534.
28. Frosk, P., Arts, H.H., Philippe, J., Gunn, C.S., Brown, E.L., Chodirker, B., Simard, L., Majewski, J., Fahiminiya, S., Russell, C., et al. (2017). A truncating mutation in CEP55 is the likely cause of MARCH, a novel syndrome affecting neuronal mitosis. *J. Med. Genet.* 54, 490–501.
29. Anttonen, A.-K., Laari, A., Kousi, M., Yang, Y.J., Jääskeläinen, T., Somer, M., Siintola, E., Jakkula, E., Muona, M., Tegelberg, S., et al. (2017). ZNHIT3 is defective in PEHO syndrome, a severe encephalopathy with cerebellar granule neuron loss. *Brain A J. Neurol.* 140, 1267–1279

30. Hibi, M., and Shimizu, T. (2012). Development of the cerebellum and cerebellar neural circuits. *Dev. Neurobiol.* 72, 282–301.
31. Bertrand, S., Thisse, B., Tavares, R., Sachs, L., Chaumot, A., Bardet, P.-L., Escrivà, H., Duffraisse, M., Marchand, O., Safi, R., et al. (2007). Unexpected novel relational links uncovered by extensive developmental profiling of nuclear receptor expression. *PLoS Genet.* 3, e188.
32. Katsuyama, Y., Oomiya, Y., Dekimoto, H., Motooka, E., Takano, A., Kikkawa, S., Hibi, M., and Terashima, T. (2007). Expression of zebrafish ROR alpha gene in cerebellar-like structures. *Dev. Dyn. An Off. Publ. Am. Assoc. Anat.* 236, 2694–2701.
33. Hamling, K.R., Tobias, Z.J.C., and Weissman, T.A. (2015). Mapping the development of cerebellar Purkinje cells in zebrafish. *Dev. Neurobiol.* 75, 1174–1188.
34. Gold, D.A., Gent, P.M., and Hamilton, B.A. (2007). ROR alpha in genetic control of cerebellum development: 50 staggering years. *Brain Res.* 1140, 19–25.
35. Vogel, M.W., Sinclair, M., Qiu, D., and Fan, H. (2000). Purkinje cell fate in staggerer mutants: agenesis versus cell death. *J. Neurobiol.* 42, 323–337.
36. Yoon, C.H. (1972). Developmental mechanism for changes in cerebellum of “staggerer” mouse, a neurological mutant of genetic origin. *Neurology* 22, 743–754.
37. Tammimies, K., Marshall, C.R., Walker, S., Kaur, G., Thiruvahindrapuram, B., Lionel, A.C., Yuen, R.K.C., Uddin, M., Roberts, W., Weksberg, R., et al. (2015). Molecular Diagnostic Yield of Chromosomal Microarray Analysis and Whole-Exome Sequencing in Children With Autism Spectrum Disorder. *JAMA* 314, 895–903.
38. Hu, W.F., Chahrour, M.H., and Walsh, C.A. (2014). The diverse genetic landscape of neurodevelopmental disorders. *Annu. Rev. Genomics Hum. Genet.* 15, 195–213.
39. Doulazmi, M., Frédéric, F., Capone, F., Becker-André, M., Delhay-Bouchaud, N., and Mariani, J. (2001). A comparative study of Purkinje cells in two RORalpha gene mutant mice: staggerer and RORalpha(-/-). *Brain Res. Dev. Brain Res.* 127, 165–174.
40. Chérot, E., Keren, B., Dubourg, C., Carré, W., Fradin, M., Lavillaureix, A., Afenjar, A., Burglen, L., Whalen, S., Charles, P., et al. (2017). Using medical exome sequencing to identify the causes of neurodevelopmental disorders: Experience of 2 clinical units and 216 patients. *Clin. Genet.* doi:

10.1111/cge.13102. [Epub ahead of print].

41. Herrup, K., Shojaeian-Zanjani, H., Panzini, L., Sunter, K., and Mariani, J. (1996). The numerical matching of source and target populations in the CNS: the inferior olive to Purkinje cell projection. *Brain Res. Dev. Brain Res.* *96*, 28–35.
42. Doulazmi, M., Capone, F., Frederic, F., Bakouche, J., Boukhtouche, J., Lemaigre-Dubreuil, Y., and Mariani, J. (2006). Cerebellar purkinje cell loss in heterozygous *rora*^{+/-} mice: a longitudinal study. *J. Neurogenet.* *20*, 1–17.
43. Nguyen, A., Rauch, T.A., Pfeifer, G.P., and Hu, V.W. (2010). Global methylation profiling of lymphoblastoid cell lines reveals epigenetic contributions to autism spectrum disorders and a novel autism candidate gene, RORA, whose protein product is reduced in autistic brain. *FASEB J.* *24*, 3036–3051.
44. Sarachana, T., and Hu, V.W. (2013). Genome-wide identification of transcriptional targets of RORA reveals direct regulation of multiple genes associated with autism spectrum disorder. *Mol. Autism* *4*, 14.
45. Sayad, A., Noroozi, R., Omrani, M.D., Taheri, M., and Ghafouri-Fard, S. (2017). Retinoic acid-related orphan receptor alpha (RORA) variants are associated with autism spectrum disorder. *Metab. Brain Dis.* *32*, 1595–1601.
46. Wang, Y., Billon, C., Walker, J.K., and Burris, T.P. (2016). Therapeutic Effect of a Synthetic ROR α / γ Agonist in an Animal Model of Autism. *ACS Chem. Neurosci.* *7*, 143–148.

Figure legends




Figure 1. *RORA* variants cause intellectual disability with autistic features or cerebellar hypoplasia. (A) Schematic of ROR α isoform a (GenBank: NP_599023.1; encoded by *RORA* transcript 1) depicting exons (black outlined boxes with numbers 1-10) and domains; green, DNA binding domain (DBD); orange, ligand binding domain (LBD). Missense variants (top) and truncating variants (bottom) are shown. (B) Autosomal dominant pedigree (family 10) in which a ~1.5 Mb intergenic deletion containing *RORA* segregates with disease. Filled shapes, affected individuals; unfilled shapes, healthy individuals. Individuals 10A, 10B, 10C, 10D, and the spouse of 10C were tested. (C) Schematic depicting the *RORA* locus at 15q22. Zoomed region shows two intergenic deletions (individuals 10A-D and individual 11), one intragenic deletion (individual 12) and an intragenic duplication (individual 13) involving *RORA*. Genes are indicated by black bars; copy number loss, red; copy number gain, blue.




Figure 2. Brain MRI of individuals with cerebellar hypoplasia. MRI of individual 6 done at 6.5 years old (A), individual 2 done at 2 years and 1 month old (B) and individual 3 done at 2 years old (C) showing cerebellar hypoplasia, predominant on the vermis (white arrow). T1-weighted sequence, sagittal section through cerebellum.




Figure 3. Disruption of *roraa* in zebrafish larvae results in cerebellar hypoplasia driven by Purkinje and granule cell loss. (A) Schematic of neuroanatomical structures painted with anti-acetylated tubulin antibody at 3 days post fertilization (dpf); IT, intertectal. (B) Representative dorsal images of acetylated tubulin immunostained larvae show that *roraa* ablation causes cerebellar defects in CRISPR/Cas9 F0 mutants and morphants. Cerebellar size was measured as indicated by the dashed red outline on inset panels. (C) Quantification of cerebellar area in larval batches is shown for two guide (g)RNAs targeting either *roraa* exon 5 (gRNA 1) or exon 8 (gRNA 2) (D) *roraa* morphants (injected with 3 ng morpholino; MO) display a cerebellar phenotype that can be rescued by four

different wild type *RORA* mRNA transcripts: co-injection of *rora* e2i2 splice-blocking MO with *RORA* splice variants (*RORA* 1: NM_134261, *RORA* 2: NM_134260, *RORA* 3: NM_002943 and *RORA* 4: NM_134262). **(E)** Schematic of cerebellar cell types assessed in 3 dpf larvae using either a *neurod:egfp* transgene (green) or anti-zebrin II immunostaining (red). Orientation is indicated with A, anterior; P, posterior; L, left; R, right. **(F)** Representative dorsal images show that reduction of Purkinje and granule cells contributes to cerebellar defects induced by *rora* targeting. Transgenic *neurod:egfp* larvae were fixed, immunostained with anti-zebrin II antibody (red), and the area comprised of each cell type was measured (as indicated in the schematic). Dashed white lines indicate measured area. **(G)** Quantification of granule layer cells (GFP-positive). **(H)** Quantification of Purkinje cells (zebrin II-positive). Both cell populations are reduced significantly in *rora* F0 mutants as well as morphant zebrafish. Scale bar in **(B)** and **(F)**: 100 μ m. AU, arbitrary units. Stars indicate *P* value compared to uninjected controls (CRISPR/Cas9 and MO) or to morphants (MO + *RORA*). ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$; ns, not significant. Error bars in **(C)**, **(D)**, **(G)** and **(H)** represent 5th and 95th Percentile.




Figure 4. *RORA* missense variant located in the DNA binding domain and ligand binding domain confer a dominant toxic and loss-of-function effect, respectively. (A) Complementation of *rora* morpholino (MO) with variant mRNA results in severe gross morphological defects or larval death before 3 days post-fertilization (dpf). Left, representative embryo classes are shown; right, quantification of outcomes. p.Gly92Ala (G92A), p.Lys94Arg (K94R) are located in the DNA binding domain; p.Arg462Gln (R462Q) localizes to the ligand-binding domain; T476A (rs190933482), is a negative control for the assay with an allele frequency of 0.001091 in gnomAD). **(B)** Quantification of

cerebellar area of larval batches with ectopic expression of *RORA* mRNA. G92A mRNA confers a significant reduction in cerebellar size. **(C)** Representative dorsal images of 3 dpf larvae injected with gradually modified doses of WT and G92A RNA (top, amount in pg) and stained with anti-acetylated tubulin antibody. **(D)** Quantification of cerebellar measurements from the batches shown in **(C)**. **(E)** Representative dorsal images of 3 dpf larvae injected with gradually modified doses of WT and K94R RNA (top, amount in pg) and stained with anti-acetylated tubulin antibody. **(F)** Quantification of cerebellar measurements from the batches shown in **(E)**. **(G)** Representative dorsal images of 3 dpf larvae injected either with e2i2 splice-blocking MO, or MO co-injected with WT *RORA* mRNA, R462Q or population control T476A variant *RORA* mRNA. **(H)** Quantification of cerebellar measurements from the batches shown in **(E)**. AU, arbitrary units. Stars indicate *P* value compared to WT. ****, $p < 0.0001$; ns, not significant. Error bars in **(B)**, **(D)**, **(F)** and **(H)** represent 5th and 95th Percentile.

Table 1. Molecular and clinical data from the 15 individuals with *RORA* variants.

Individual ID	1	2	3	4	5	6	7	8	9	10-D	10-C	10-B	10-A	11	12	13
DECIPHER ID	-	-	-	-	-	-	-	-	-	-	-	-	289628		255754	293939
Sex	female	male	female	female	female	female	male	male	male	female	female	male	female	male	male	male
Geographic origin	USA	Estonia	France	USA	USA	France	USA	Germany	USA	Belgium	Belgium	Belgium	Belgium	France	Denmark	Denmark
Age at last investigation (years)	4	3.5	6	28	3	10	9	4	14	25	50	44	80	15	8	6,5
Mutation type	missense	missense	missense	splice site	frameshift	frameshift	missense	missense	nonsense	15q22 deletion including <i>RORA</i>	15q22.3q22.2 deletion including <i>RORA</i>	15q22 intragenic deletion of <i>RORA</i>	15q22 disruptive duplication of <i>RORA</i>			
Mutation (GRCh37) ^a	c.269G>C	c.275G>C	c.281A>G	c.425-1G>A	c.804_805delGT	c.1019delG	c.1225A>C	c.1385G>A	c.1498C>T	15q22.2 (59,641,986-61,104,231)x1	15q22.2 (59,641,986-61,104,231)x1	15q22.2 (59,641,986-61,104,231)x1	15q22.2 (59,641,986-61,104,231)x1	15q21.3q22.2 (58,622,268-62,320,616)x1	15q22.2 (60,809,984-60,837,029)x1	15q22.2 (60,797,691-60,860,668)x3
Protein variant ^b	p.(Cys90Ser)	p.(Gly92Ala)	p.(Lys94Arg)	p.(Ala142_Leu273del)	p.(Ser269Hisfs*13)	p.(Arg340Profs*17)	p.(Ser409Arg)	p.(Arg462Gln)	p.(Arg500*)	-	-	-	-	-	p.(Ser66Trpfs*24)	p.(Gln315Leufs*51)
Mode of inheritance	unknow (adopted)	<i>de novo</i>	<i>de novo</i>	<i>de novo</i> (mosaic, 20% in blood) ^c	<i>de novo</i>	<i>de novo</i>	<i>de novo</i>	<i>de novo</i>	<i>de novo</i>	Inherited from mother (ID 10-C)	Inherited from mother (ID 10-A)	Inherited from mother (ID 10-A)	dominant	<i>de novo</i>	<i>de novo</i>	<i>de novo</i>
Functional effect ^d	NA	dominant toxic	dominant toxic	NA	NA	NA	NA	loss-of-function	NA	NA	NA	NA	NA	NA	NA	NA
Growth parameters																
Birth weight (grams/SD)	NA	3500/0	3310/0	4238/+2	4054/+1.4	2655/-1.5	3487/0	NA	3340/-0.4	NC/+0.7	NA	NA	NA	3200/-0.5	4050/+1	NC; unremarkable
Birth length (cm/SD)	NA	52/+1	49/0	NA	54.6/+1.9	46/-2	52/+1	NA	48.3/-0.7	NC/+0.7	NA	NA	NA	49.5/0	56/+2.5	NC; unremarkable
Birth head circumference (cm/SD)	NA	38/+2.5	34.5/0	NA	NA	33/-1	34/-0.5	NA	46.5/+2 (6 mo)	NC/+0.7	NA	NA	NA	34/-0.5	NA	NC; unremarkable
Height at age last investigation (cm/SD)	102/0	102/+0.5	104/-2	167.8/+0.5	89/-1.5	132/-1	120.8/-2 (8y 5 mo)	NA	150.4/-0.7 (13 y)	166/+0.7	NA	NA	NA	170/0	NC/+2.5	NC/-1
Weight at age last investigation (kg/SD)	17.2/+0.5	17.8/+1	16/-1.5	69.4/+1	14.5/+0.3	26.2/-1.3	31/+1.3 (9y 3 mo)	NA	44.4/-0.2 (13 y)	74.3/+1.3	NA	NA	NA	58/+0.7	NC/+1	NC/-1
Head circumference at age last investigation (cm/SD)	49/0	50.6/0	50/0	58.4/+1.9	NA	53/0	51.7/-0.3 (8y 4 mo)	NA	58/+2.6 (13 y)	54/-0.7	NA	NA	NA	56/+0.7	NA	NA

Degree of developmental delay or ID	mild	mild-moderate	severe	mild (regression at 10 years)	mild	severe	mild	no ID (IQ 85)	moderate	mild	mild	mild	mild	moderate	mild	mild
Age of walking	2 years	3 years	6 years	14 months	20 months	3 years	15 months	NA	24 months	16 months	NA	NA	NA	21 months	normal	mild delay
Age of first words	2 years	delayed	before 1 year	14-15 months	12-13 months	5 years	>2 years	NA	delayed	14 months	NA	NA	NA	2 years	delayed	delayed
Current language ability	speaking with sentences	speaking with sentences	no phrase	normal	around 20 words	no phrase	delayed	normal	rudimentary sentences	delayed	NA	NA	NA	normal	delayed	delayed
Behavioral anomalies	no	no	no	no	ASD	ASD	ASD	ASD	no	behavioral problems	no	borderline personality disorder	no	hyperactivity	no	ASD
Neurological examination																
Seizures (age at onset/type)	no	no	yes (4 years / absences)	yes (8 years / tonic-clonic seizures)	yes (ND / 1 episode of absence)	yes (5 years / multifocal)	yes (neonatal / myoclonic seizures)	no	yes (ND / generalized seizures)	yes (4 years / absences, drop attacks)	yes	yes (ND / generalized, status epilepticus)	no	yes (ND / generalized, febrile seizures)	no	yes
Neurological examination	tremor, hypotonia, coordination disorder	tremor, hypotonia, ataxia	tremor, hypotonia, ataxia, pyramidal syndrome	tremor, hypotonia, ataxia	hypotonia, poor coordination	tremor, hypotonia, ataxia	occasional fine tremor	NA	mild hand tremor, hypotonia	NA	NA	tremor	no	palpebral myoclonia	no	no
Brain imaging	normal	cerebellar hypoplasia (mainly vermis)	pontocerebellar atrophy (mainly vermis)	NA	normal	mild global cerebellar hypoplasia	corpus callosal hypoplasia	NA	CSF spaces mildly prominent	normal	NA	normal	NA	NA	NA	NA
Global examination																
Eye anomalies	strabismus, esotropia, nystagmus, amblyopia	no	mild oculomotor apraxia, strabismus	visual processing deficits	no	no	esotropia, hyperopia	NA	strabismus, hyperopia	strabismus	strabismus	NA	NA	strabismus	no	no
Other	-	spina bifida occulta	recurrent vomiting, constipation	hypoglycemia in childhood	bilateral hydronephrosis	hypercholesterolemia	popliteal contractures	-	cryptorchidism, renal cysts	-	-	-	-	-	atopic dermatitis	-
Other genetic findings	VUS in <i>NR4A2</i>	-	maternally inherited heterozygous p.(Arg662*) in <i>DNM1L</i>	-	maternally inherited deletions of 3p26.3 and 10q21.3 (both classified as VUS), mother reports no history of developmental delays	-	HSPG2 c.7006+1G>A (maternally inherited), HSPG2 c.4391G>A (paternally inherited), GSK3B c.625dupC (de novo), CACNA1A c.5883G>A (maternally inherited)	-	NGLY1 c.1516C>T, p.Arg506* (maternally inherited), PKD2 c.2143delC, p.Leu715* (1q42.2(231,834,554-231,885,282)x1)	intragenic <i>DISC1</i> deletion (1q42.2(231,834,554-231,885,282)x1)	-	-	-			

Initial diagnostic hypotheses	none	ID and disorders with cerebellar involvement	congenital ataxia with pontocerebellar atrophy	disorder of mitochondrial metabolism	none	none	speech delay, toe walking	idiopathic ASD F84.0	<i>CUL4B</i> -related disorder	GGE and ID	ID and possible epilepsy	ID?	epilepsy (GGE?) and ID	Deletion of <i>RORB</i>	none	none
-------------------------------	------	--	--	--------------------------------------	------	------	---------------------------	----------------------	--------------------------------	------------	--------------------------	-----	------------------------	-------------------------	------	------

^a nomenclature HGVS V2.0 according to mRNA reference sequence NM_134261.2. Nucleotide numbering uses +1 as the A of the ATG translation initiation codon in the reference sequence, with the initiation codon as codon 1

^b inferred from bioinformatic predictions but not verified from the individual's mRNA.

^c mosaicism was inferred and estimated initially from raw exome sequencing data and confirmed by Sanger sequencing.

^d effect inferred from *in vivo* tests in zebrafish or analyses from patient fibroblasts; *in vitro* experiments; LoF: loss of function, GoF: gain of function, ND: not determined. ASD: autism spectrum disorder; ID: intellectual disability; IQ: intelligence quotient; GGE: genetic generalized epilepsy; NA: not analyzed; NC: not communicated; ND: not determined; y: years