*Reply to the Editor—Carnitine and
ventricular repolarization: A complex link?*

Dr Acampa discussed the link between QT interval abnormalities, ventricular arrhythmias, the autonomic nervous system (ANS), and metabolic disorders associated with the low level of carnitine. First, in Rett syndrome the level of carnitine is low but not null as in primary carnitine deficiency (PCD). Rett syndrome (RS) is a severe neurodevelopmental disorder associated with a high frequency of seizure (68%) in contrast to PCD. It is known that the frequency of sudden death is high in the epileptic population.¹ Second, Rasmussen et al² and Guideri et al³ did not report abnormalities of the QT interval in PCD and RS. De Biase⁴ and Schimmenti⁵ noted one patient with prolonged QT interval and ventricular arrhythmias, but a genetic mutation in the main genes involved in long QT syndrome was not ruled out. Third, short QT syndrome published by Brugada is rare and probably underestimated. Recently, several abnormalities of the T wave were reported in PCD.⁶ The effect of the ANS on ventricular repolarization and ventricular arrhythmias is well known. All conditions associated with ANS dysfunction favor cardiac arrhythmias. However, in the Guideri study³ the low frequency/high frequency ratio was unchanged after carnitine supplementation, demonstrating a limited effect of carnitine supplementation on heart rate variability. Finally, the electrophysiological effects of acyl-L-carnitine and L-carnitine are different⁷: in vitro carnitine has no effect in contrast to acyl-L-carnitine, which increases the potassium current I_{Kr} that plausibly decreases the QT interval. These effects should be studied in the intact heart.

Julien Roussel, PhD*

Jérôme Thireau, PhD*

Charlotte Farah, PhD*

Jean Yves Le Guennec, PhD*

Dominique Babuty, PhD,[†]

d.babuty@chu-tours.fr

*INSERM U1046, CNRS UMR 9214,

Université de Montpellier

Montpellier, France

[†]Cardiologie, CHRU de Tours

Université François Rabelais

Tours, France

References

1. Rugg-Gunn FJ, Simister RJ, Squirell M, Holdright DR, Duncan JS. Cardiac arrhythmias in focal epilepsy: a prospective long-term study. *Lancet* 2004;364:2212–2219.
2. Rasmussen J, Kober L, Lund AM, Nielsen. Primary carnitine deficiency in the Faroe Islands: health and cardiac status in 76 adult patients diagnosed by screening. *J Inherit Metab Dis* 2014;37:223–230.
3. Guideri F, Acampa M, Hayek Y, Zappella M. Effects of acetyl-L-carnitine on cardiac dysautonomia in Rett syndrome: prevention of sudden death? *Pediatr Cardiol* 2005;26:574–577.
4. De Biase I, Champaigne NL, Schroer R. Primary carnitine deficiency presents atypically with long QT syndrome: a case report. *JIMD Rep* 2012;2:87–90.
5. Schimmenti LA, Crombez EA, Schwahn BC. Expanded newborn screening identifies maternal primary carnitine deficiency. *Mol Genet Metabol* 2007;90:441–445.
6. Fu L, Huang M, Chen S. Primary carnitine deficiency and cardiomyopathy. *Korean Circ J* 2013;43:785–792.
7. Ferro F, Ouille A, Tran TA, Fontanaud P, Bois P, Babuty D, Labarthe F, Le Guennec JY. Long-chain acylcarnitines regulate the hERG channel. *PLoS One* 2012;7:e41686.