



**HAL**  
open science

## **3D models related to the publication: Neotropics provide insights into the emergence of New World monkeys: new dental evidence from the late Oligocene of Peruvian Amazonia**

Laurent Marivaux, Rodolfo Salas-Gismondi, Pierre-Olivier Antoine

### ► To cite this version:

Laurent Marivaux, Rodolfo Salas-Gismondi, Pierre-Olivier Antoine. 3D models related to the publication: Neotropics provide insights into the emergence of New World monkeys: new dental evidence from the late Oligocene of Peruvian Amazonia. *MorphoMuseum*, 2017, 3 (1), 10.18563/m3.3.1.e1 . hal-01813187

**HAL Id: hal-01813187**

**<https://hal.umontpellier.fr/hal-01813187>**

Submitted on 13 Jun 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 3D models related to the publication: Neotropics provide insights into the emergence of New World monkeys: new dental evidence from the late Oligocene of Peruvian Amazonia

Marivaux Laurent<sup>1\*</sup>, Salas-Gismondi Rodolfo<sup>2</sup>, Antoine Pierre-Olivier<sup>1</sup>

<sup>1</sup>Laboratoire de Paléontologie, Institut des Sciences de l'Évolution de Montpellier (ISE-M, UMR 5554, CNRS/UM/IRD/EPHE), c.c. 064, Université de Montpellier, place Eugène Bataillon, 34095 Montpellier Cedex 05, France

<sup>2</sup>Departamento de Paleontología de Vertebrados, Museo de Historia Natural - Universidad Nacional Mayor San Marcos (MUSM), Av. Arenales 1256, Lima 11, Peru

\*Corresponding author: laurent.marivaux@umontpellier.fr

## Abstract

This contribution contains the 3D models of the isolated teeth of *Canaanimico amazonensis*, a new stem platyrrhine primate, described and figured in the following publication: Marivaux et al. (2016), Neotropics provide insights into the emergence of New World monkeys: new dental evidence from the late Oligocene of Peruvian Amazonia. Journal of Human Evolution. <http://dx.doi.org/10.1016/j.jhevol.2016.05.011>

**Keywords:** Homunculidae, Paleogene, Peru, Soriacebinae, South America

Submitted:2017-08-14, published online:2017-09-13. <https://doi.org/10.18563/m3.3.1.e1>

## INTRODUCTION

We present here the 3D digital models of the fossils documenting *Canaanimico amazonensis* Marivaux et al. 2016, a medium-sized platyrrhine monkey, recently found in Peruvian Amazonia (Contamana, Loreto Department) from fluvial deposits (CTA-61) dating from the late Oligocene (ca. 26.5 Ma; i.e., Deseadan South American Land Mammal Ages [SALMA]; Antoine et al., 2016). This anthropoid primate is known by only two isolated upper teeth (Fig. 1; Table 1) recovered after wet-screening of more than 800 kg of sediment. MUSM-2500 is the lingual part of a pristine left M1, which was probably broken during the fossilization process. MUSM-2499 is a complete left M2, which is particularly well-preserved aside from minor wear to the apices of the main cusps due to occlusal attrition during the life of the animal. From our comparative dental analysis, we have seen that *Canaanimico* exhibits a few upper molar characters that are otherwise found only in a limited group of platyrrhines. However, most other characters may be found individually in extinct or/and living taxa belonging either to the Pitheciidae, Cebidae or Atelidae. In order to investigate and formalize the phylogenetic position of *Canaanimico* in a high-level platyrrhine phylogeny, we have performed a cladistic assessment of the dental evidence. The results strengthen support for the soriacebine affinities of *Canaanimico* among the early Miocene 'Homunculidae', a group of stem platyrrhines placed outside the crown platyrrhine radiation. *Canaanimico* therefore extends back the soriacebine lineage to 26.5 Ma. This discovery adds to the evidence that platyrrhines were well established in western Amazonia during mid-Cenozoic times (see Marivaux et al., 2016).

Model IDs	Taxon	Description
MUSM-2499	<i>Canaanimico amazonensis</i>	Left upper M2 (Holotype)
MUSM-2500	<i>Canaanimico amazonensis</i>	left upper M1 (lingual part)


**Table 1.** List of models. All specimens come from the Museo de Historia Natural - Universidad Nacional Mayor San Marcos (MUSM), Lima, Peru.

## METHODS

AVIZO 7.1 (Visualization Sciences Group) software was used for visualization, segmentation and 3D rendering. The teeth were prepared within a "labelfield" module of AVIZO, using the segmentation threshold selection tool. The 3D models are provided in .ply format, and then can be opened with a wide range of freeware.

## ACKNOWLEDGEMENTS

3D data presented in this work were produced through the technical facilities of the Montpellier RIO Imaging (MRI) platform and of the LabEx CeMEB. We particularly thank R. Lebrun (ISE-M, Université de Montpellier) for his help and advices during micro-CT scan acquisitions and treatments. This work (paleontological fieldwork and laboratory post-field analyses) was financially supported by the L. S. B. Leakey Foundation. This work has also benefited from an "Investissements d'Avenir" grant managed by the Agence Nationale de la Recherche, France (CEBA, ANR-10-LABX-0025-01).



**Figure 1.** Fossil dental specimens of *Canaanimico amazonensis* Marivaux et al. 2006, from the late Oligocene (Deseadan SALMA) of Contamana locus n°61 (CTA-61), Peruvian Amazonia. A) MUSM-2499, left M2 in occlusal view; B) MUSM-2500, left M1 in occlusal view. The white arrow situated on the left indicate the orientation of the teeth on the jaws (i.e., mesiolingual). Scale bar = 2 mm.

## BIBLIOGRAPHY

- Antoine, P.-O., Abello, M. A., Adnet, S., Altamirano-Sierra, A. J., Baby, P., Billet, G., Boivin, M., Calderón, Y., Candela, A. M., Chabain, J., Corfu, F., Croft, D. A., Ganerød, M., Jaramillo, C., Klaus, S., Marivaux, L., Navarrete, R. E., Orliac, M. J., Parra, F., Pérez, M. E., Pujos, F., Rage, J.-C., Ravel, A., Robinet, C., Roddaz, M., Tejada-Lara, J. V., Vélez-Juarbe, J., Wesselingh, F. P., Salas-Gismondi, R., 2016. A 60-million year Cenozoic history of western Amazonian ecosystems in Contamana, Eastern Peru. *Gondwana Research* 31, 30-59. <http://dx.doi.org/10.1016/j.gr.2015.11.001>
- Marivaux, L., Adnet, S., Altamirano-Sierra, A. J., Boivin, M., Pujos, F., Ramdarshan, A., Salas-Gismondi, R., Tejada-Lara, J. V., Antoine, P.-O., 2016. Neotropics provide insights into the emergence of New World monkeys: new dental evidence from the late Oligocene of Peruvian Amazonia. *Journal of Human Evolution* 97, 159-175. <http://dx.doi.org/10.1016/j.jhevol.2016.05.011>