



HAL
open science

New remains of the very small cuckoo, *Chambicuculus pusillus* (Aves, Cuculiformes, Cuculidae) from the late Early or early Middle Eocene of Djebel Chambi, Tunisia

Cécile Mourer-Chauviré, El Mabrouk Essid, Hayet Khayati Ammar, Laurent Marivaux, Wissem Marzougui, Rim Temani, Monique Vianey-Liaud, Rodolphe Tabuce

► **To cite this version:**

Cécile Mourer-Chauviré, El Mabrouk Essid, Hayet Khayati Ammar, Laurent Marivaux, Wissem Marzougui, et al.. New remains of the very small cuckoo, *Chambicuculus pusillus* (Aves, Cuculiformes, Cuculidae) from the late Early or early Middle Eocene of Djebel Chambi, Tunisia. *Palaeovertebrata*, 2016, 40 (1), 10.18563/pv.40.1.e2 . hal-01813293

HAL Id: hal-01813293

<https://hal.umontpellier.fr/hal-01813293>

Submitted on 13 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New remains of the very small cuckoo, *Chambicuculus pusillus* (Aves, Cuculiformes, Cuculidae) from the late Early or early Middle Eocene of Djebel Chambi, Tunisia

CÉCILE MOURER-CHAUVIRÉ^a, EL MABROUK ESSID^b, HAYET KHAYATI^b, LAURENT MARIVAUX^c, WISSEM MARZOUGUI^b, RIM TEMANI^b, MONIQUE VIANEY-LIAUD^c & RODOLPHE TABUCE^c

^aLaboratoire de Géologie de Lyon, Université Claude Bernard – Lyon 1, UMR-CNRS 5276 Terre, Planètes, Environnement, 2 rue Dubois, 69622 Villeurbanne Cedex, France

^bOffice National des Mines de Tunis (Service géologique National), 24 rue 8601, 2035 BP215, ZI La Charguia, 2035 Ariana, 1080 Tunis, Tunisia

^cLaboratoire de Paléontologie, Institut des Sciences de l'Évolution de Montpellier (ISE-M, UMR 5554, CNRS/IRD/EPHE), c.c. 064, Université de Montpellier, Place Eugène Bataillon, F-34095 Montpellier Cedex 05, France

* Corresponding author: cecile.mourer@univ-lyon1.fr

Abstract: A very tiny cuckoo, *Chambicuculus pusillus*, was recently described after a few fragments of tarsometatarsi from the late Early/early Middle Eocene of Djebel Chambi, Tunisia. New remains, notably a coracoid, confirm the attribution of this genus to the family Cuculidae. This coracoid shows a very elongate and strap-like processus procoracoideus. This morphological feature is present in some recent or extinct taxa but it is more developed in the family Cuculidae. The characteristics of the coracoid and tarsometatarsi show that *Chambicuculus* is morphologically more advanced over the other stem cuculids described in Europe and North America. *Chambicuculus* is the oldest Cuculidae known so far.

Keywords: Fossil bird, Eocene, Cuckoos

Submitted 6 November 2015, Accepted 26 January 2016

Published Online 15 February 2016, doi: [10.18563/pv.40.1.e2](https://doi.org/10.18563/pv.40.1.e2)

© Copyright Cécile Mourer-Chauviré February 2015

INTRODUCTION

The Paleogene avifauna is still very poorly known in the whole of the African continent. Marine birds are known in the Late Paleocene and the Early Eocene of the phosphate Ouled Abdoun Basin in Morocco. They belong to the Odontopterygiformes (pseudo-toothed birds), the Prophaethontidae and possibly the Phaethontidae (tropicbirds) (Bourdon, 2006; Bourdon *et al.*, 2005, 2008a, 2008b, 2010). Odontopterygiformes are also known in the Middle Eocene of Nigeria (Andrews, 1916) and Togo (Bourdon & Cappetta, 2012).

As far as terrestrial birds are concerned, the locality of Glib Zegdou in Algeria, dated from the late Early to early Middle Eocene, has yielded the femur of a bird related to the family Phorusrhacidae, *Lavocatavis africana* (Mourer-Chauviré *et al.*, 2011b). The Phorusrhacidae were giant flightless birds mainly known in South America. In Namibia, the locality of Silica South, previously dated from the early Middle Eocene, now considered as late Middle Eocene (Bartonian) (Pickford *et al.*, 2014), has yielded the remains of a stem galliform, *Namaortyx sperrgebietensis* (Mourer-Chauviré *et al.*, 2011a). Also situated in Namibia, and also dated from the late Middle Eocene, the locality of Eocliff has yielded several remains of a galliform, *Scopelortyx klinghardensis*, and of a psittaciform, *Namapsitta praeruptorum* (Mourer-Chauviré *et al.*, 2015). Unlike the galliforms from Chambi and Silica South, the familial position of which is unknown, *Scopelortyx* belongs to the extinct family Paraortygidae, which was widespread in the northern hemisphere during the Late Eocene to the Oligocene in Europe

and North America (Mourer-Chauviré *et al.*, 2015).

In the Late Eocene and Early Oligocene localities of the Fayum, Egypt, the avifauna is more diversified and includes birds that can be attributed to 14 orders or families, including two extinct endemic African families, Eremopezidae and Xenorodiopidae (Rasmussen *et al.*, 1987, 2001; Stidham & Smith, 2015). Most of the families present in the Fayum deposits correspond to aquatic birds such as cormorants, shoebills, herons, storks, jacanas, rails, cranes, and flamingos.

In Tunisia the localities of Chambi 1 and Chambi 2, dated from the late Early to early Middle Eocene as the locality of Glib Zegdou, have yielded the remains of a small stem galliform, family incertae sedis, *Chambiortyx cristata*, and of a representative of the family Cuculidae, *Chambicuculus pusillus* (Mourer-Chauviré *et al.*, 2013). New bird material has recently been extracted from the lacustrine limestones of Chambi 1 and can be ascribed to the same very small cuculiform. This form was previously known only by three distal fragments of tarsometatarsi. The new material not only includes two very incomplete distal fragments of tarsometatarsi, but also an omal part of coracoid and a proximal part of femur. The size of the coracoid and femur, which are very small, matches the size of the tarsometatarsi, which correspond to a very tiny bird. The presence of the coracoid is crucial and substantiates the attribution of the genus *Chambicuculus* to the family Cuculidae.

MATERIAL AND METHODS

The anatomical terminology follows Baumel & Witmer (1993).

The fossil material is deposited in the collection of the “Office national des Mines de Tunis”, Tunisia. Marivaux *et al.* (2015) have recently synthesized our current knowledge of the Chambi vertebrate fauna.

SYSTEMATIC PALEONTOLOGY

Order Cuculiformes Wagler, 1830

Family Cuculidae Leach, 1820

Chambicuculus pusillus Mourer-Chauviré *et al.*, 2013

The new material includes the following remains: CBI-1-561 Left coracoid, omal part; CBI-1-562 Right femur, proximal part; CBI-1-563 Distal fragment of shaft of right tarsometatarsus, with trochlea metatarsi IV; CBI-1-564 Distal fragment of shaft of left tarsometatarsus, with trochlea metatarsi III.

Description and comparisons

The coracoid (Fig. 1)

In the members of the family Cuculidae, the coracoid shows a processus procoracoideus that is wide and flattened in dorsoventral direction, and strongly developed in mediolateral direction. This processus is particularly developed in the recent genus *Cuculus*, but also in the other genera of this family (Fig. 2). It tends to come closer to the processus acrocoracoideus. Two families were previously included in the order Cuculiformes, the Cuculidae (cuckoos) and the Musophagidae (turacos), but phylogenetic analyses based on genomic sequence data show that these two families belong to two different orders, the Cuculiformes and the Musophagiformes (Jarvis *et al.*, 2014). In the musophagiform coracoid the processus procoracoideus fuses with the processus acrocoracoideus, thereby forming a ring which surrounds the sulcus m. supracoracoidei.

On the Chambi coracoid the foramen nervi supracoracoidei is absent, the cotyla scapularis is a shallow depression, the processus acrocoracoideus ends in sternal direction in a small point, and there is no pneumatic foramen under the medial ledge of the processus acrocoracoideus. All these morphological characteristics are present in the family Cuculidae. The processus procoracoideus ends in medial direction in two lobes, a small one in omal direction, and a wider one in medial direction.

The femur (Fig. 1)

The femur CBI-1-562 does not show many particular morphological characteristics. The femoral head is relatively small and medially projecting. The crista trochanteris does not run beyond the facies articularis antitrochanterica in proximal direction, but it is projecting in dorsal direction. These characteristics are present in the Cuculiformes and differ from the Musophagiformes. In the recent turacos the femoral head is wider, stouter, and less projecting medially, and the crista trochanteris, slightly projecting proximally, shows a pneumatic foramen on the dorsal surface.

The fragments of tarsometatarsi

These fragments show the same characteristics as the fragments

CBI-1-541 and CBI-2-1001, upon which the description of *Chambicuculus pusillus* was based (Mourer-Chauviré *et al.*, 2013).

Measurements (in mm)

Coracoid: length as preserved, 5.5; proximal width from the facies scapularis to the tip of processus procoracoideus, 2.6; width of shaft, 0.9.

Femur: length as preserved, 7.3; proximal width, 2.9.

Comparisons with stem Cuculidae and related forms

The genus *Eocuculus* Chandler, 1999, has been attributed to Cuculidae by Chandler (1999) and considered as ?Cuculidae by Mayr (2006). It is known by the species *Eocuculus cherpinae*, from the Late Eocene of North America (USA; Chandler, 1999) and *Eocuculus cf. cherpinae*, from the Early Oligocene of France (Mayr, 2006). The coracoid is known in a specimen on a slab, which is tentatively referred to *Eocuculus*, and which comes from the same area as *Eocuculus cf. cherpinae* (Mayr, 2008). On this specimen the processus procoracoideus is long and narrow, but on the slab its medial extremity is covered by the proximal part of the humerus and it is not possible to see whether the processus was as elongate as in *Chambicuculus* (Mayr, 2008, fig. 4, A and D). The coracoid is unknown in *Neococcyx mcorquodalei*, from the Late Eocene (Mayr, 2009) of Canada (Weigel, 1963), but it is known in *Cursoriococcyx geraldinae*, from the Early Miocene of Colorado (Martin & Mengel, 1984). In this species the processus procoracoideus is long but not as extensive as in *Chambicuculus*. The shape of the processus acrocoracoideus is similar to that of the members of the recent subfamily Neomorphinae, while in *Chambicuculus* this shape is identical to that of the members of the recent subfamily Cuculinae.

The very tiny bird, *Pumiliornis tessellatus*, from the Early Eocene (Lenz *et al.*, 2015) of Messel, Germany, was previously considered as related to the genus *Eocuculus* (Mayr, 2008; 2009;



Figure 1. *Chambicuculus pusillus*, a-b: left coracoid, omal part, CBI-1-561, dorsal view (a) and ventral view (b); c-d: right femur, proximal part, CBI-1-562, dorsal view (c) and ventral view (d). Scale bars = 1 mm.

2011). But, according to new data, *Pumiliornis* is no longer considered to be close to Cuculiformes but it is considered as a stem group representative of Psittacopasseres, the clade including Psittaciformes and Passeriformes (Mayr, 2015). The coracoid of *Pumiliornis* shows a processus procoracoideus wide in omosternal direction, and medially elongate, but not as elongate as in *Chambicuculus* (Mayr, 1999, fig. 4, b-c; 2008, fig. 4, E; 2011, fig. 2, C). In addition it shows on the ventral side of the omal part “an elevation of approximately triangular shape” (Mayr, 1999: 78) a character which is not observed in *Chambicuculus*.

DISCUSSION

Chambicuculus is older than the other extinct forms of Cuculiformes known so far. Its tarsometatarsus shows a canalis interosseus distalis covered by a roof on the dorsal surface, which is a plesiomorphic characteristic (Mayr *et al.*, 2003). In the modern Cuculidae, this canalis is not roofed, but dorsally open. However this tarsometatarsus also shows an apomorphic characteristic, i.e. the presence of a sehnenhalter, or trochlea accessoria, on the trochlea metatarsi IV. This sehnenhalter, well-developed, plantarly and slightly medially oriented, indicates that *Chambicuculus*, like the present-day Cuculidae, was zygodactyl. With the exception of its plesiomorphic canalis interosseus distalis, *Chambicuculus* appears more advanced than the more recent genus *Eocuculus*. In *Eocuculus*, indeed, the foot was not zygodactyl but probably only semizygodactyl. In addition, in *Eocuculus* the trochlea metatarsi IV extends as far distally as the trochlea metatarsi II, and it reaches the mid-section of the trochlea metatarsi III, while in the living Cuculidae, and in *Chambicuculus*, the trochlea metatarsi IV is situated more proximally than the base of the trochlea metatarsi III and it is much shorter than the trochlea metatarsi II.

The shape of the coracoid, especially the strap-like and medially elongate processus procoracoideus (very similar to that of the modern Cuculinae) is also a derived characteristic of the family Cuculidae (Fig. 2). In some other groups of extinct or recent birds there is also a processus procoracoideus strongly developed and extending medially, as for example in some Zygodactylidae (Mayr, 1998, pl. 7, fig. 12), in some Upupiformes (Phoeniculidae), in some Alcediniformes (Meropidae), in the




Figure 2. Left coracoids of recent Cuculidae, in dorsal view. **a:** *Cuculus canorus*; **b:** *Centropus senegalensis*; **c:** *Clamator glandarius*.

Coraciiformes s. s. (Coraciidae and Brachypteraciidae) or in some Piciformes (Galbulae), but it is more medially elongate in the Cuculidae than in any of these groups. This indicates that, as early as in the Early or Middle Eocene, the Cuculidae had acquired some of their morphological characteristics, and that the here described African form was more derived than the more recent forms known in Europe and North America. *Chambicuculus* is the earliest member of the family Cuculidae known so far.

Cuculiformes represent a very ancient group of birds. Phylogenetic analyses based on genomic sequence data suggest that Cuculiformes belong to the same clade as Musophagiformes and Otidiformes (Jarvis *et al.*, 1914; Prum *et al.*, 2015). The lineages leading to these three groups would have diverged shortly after the Cretaceous/Palaeogene boundary (Jarvis *et al.*, 2014), or during the Palaeocene (Prum *et al.*, 2015). The fossil record of the Cuculidae is poor but the occurrence of a cuculid in the Early to Middle Eocene confirms the great antiquity of this family.

ACKNOWLEDGMENTS

The authors thank Jean Vannier for his help in taking the photographs of Figure 1. We are also highly indebted to Anne-Lise Charrault and Sandra Unal (ISE-M, Montpellier) for their substantial contribution in the extraction and preparation of the fossil specimens from Chambi 1. This research was supported by the French ANR-ERC PALASIAFRICA Program (ANR-08-JCIC-0017), a grant from the Conseil Scientifique (CS) of the Université Montpellier 2 (UM2), and by the ONM of Tunis (AC-1785). Finally the authors thank the two reviewers, Gerald Mayr and Nikita Zelenkov, whose comments have greatly improved the manuscript.

BIBLIOGRAPHY

- Andrews, C.W., 1916. Note on the sternum of a large carinate bird from the (?) Eocene of Southern Nigeria. *Proceedings of the Zoological Society of London*, 519-524.
- Baumel, J. J., Witmer, L. M., 1993. Osteologia. In: Baumel, J. J. (Ed.), *Handbook of Avian Anatomy. Nomina Anatomica Avium*, 2nd edition. Nuttall Ornithological Club 23, 45-132.
- Bourdon, E., 2006. L'avifaune du Paléogène des phosphates du Maroc et du Togo : diversité, systématique et apports à la connaissance de la diversification des oiseaux modernes (Néornithes). Ph. D. dissertation, Muséum national d'Histoire naturelle, Paris.
- Bourdon, E., Amaghaz, M., Bouya, B., 2008a. A new seabird (Aves, cf. Phaethontidae) from the Lower Eocene phosphates of Morocco. *Geobios* 41, 455-459. doi: 10.1016/j.geobios.2007.11.002
- Bourdon, E., Amaghaz, M., Bouya, B., 2010. Pseudotoothed Birds (Aves, Odontopterygiformes) from the Early Tertiary of Morocco. *American Museum Novitates* 3704, 71 pp.
- Bourdon, E., Bouya, B., Iarochène, M., 2005. Earliest African neornithine bird: a new species of Prophaethontidae (Aves) from the Paleocene of Morocco. *Journal of Vertebrate Paleontology* 25, 157-170.
- Bourdon, E., Cappelletta, H., 2012. Pseudo-toothed birds (Aves, Odontopterygiformes) from the Eocene phosphate deposits of Togo, Africa. *Journal of Vertebrate Paleontology* 32 (4), 965-970.
- Bourdon, E., Mourer-Chauviré, C., Amaghaz, M., Bouya, B., 2008b. New specimens of *Lithoptila abdounensis* (Aves, Prophaethontidae) from the Lower Paleogene of Morocco. *Journal of Vertebrate Paleontology* 28 (3), 751-761.

- Chandler, R., 1999. Fossil birds of Florissant, Colorado: with a description of a new genus and species of cuckoo. Geologic Resources Division, Technical Report, NPS/NRGRD/GRDTR-99, 49-53.
- Jarvis, E. D. et alii (105 authors), 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science* 146 (6215), 1320-1331. doi: [10.1126/science.1253451](https://doi.org/10.1126/science.1253451)
- Lenz, O. K., Wilde, V., Mertz, D. F., Riegel, V., 2015. New palynology-based astronomical and revised $^{40}\text{Ar}/^{39}\text{Ar}$ ages for the Eocene maar lake of Messel (Germany). *International Journal of Earth Sciences* 104, 873-889. doi: [10.1007/s00531-014-1126-2](https://doi.org/10.1007/s00531-014-1126-2)
- Marivaux, L., Essid, E.M., Marzougui, W., Khayati Ammar, H., Merzeraud, G., Tabuce, R., Vianey-Liaud, M., 2015. The early evolutionary history of anomaluroid rodents in Africa: new dental remains of a zegdomyid (Zegdomyidae, Anomaluroidea) from the Eocene of Tunisia. *Zoologica Scripta* 44 (2), 117-134. doi: [10.1111/zsc.12095](https://doi.org/10.1111/zsc.12095)
- Martin, L. D., Mengel, R. M., 1984. A new cuckoo and a chachalaca from the early Miocene of Colorado. *Special Publication Carnegie Museum of Natural History* 9, 171-177.
- Mayr, G., 1998. "Coraciiforme" und "piciforme" Kleinvögel aus dem Mittel-Eozän der Grube Messel (Hessen, Deutschland). *Courier Forschungsinstitut Senckenberg* 205, 1-101.
- Mayr, G., 1999. *Pumiliornis tessellatus* n. gen. n. sp., a new enigmatic bird from the Middle Eocene of Grube Messel (Hessen, Germany). *Courier Forschungsinstitut Senckenberg* 216, 75-83.
- Mayr, G., 2006. A specimen of *Eocuculus* Chandler, 1999 (Aves, ? Cuculidae) from the early Oligocene of France. *Geobios* 39, 865-872. doi: [10.1016/j.geobios.2005.10.007](https://doi.org/10.1016/j.geobios.2005.10.007)
- Mayr, G., 2008. *Pumiliornis tessellatus* MAYR, 1999 revisited – new data on the osteology and possible phylogenetic affinities of an enigmatic Middle Eocene bird. *Paläontologische Zeitschrift* 82/3, 247-253.
- Mayr, G., 2009. *Paleogene Fossil Birds*. Springer Verlag, Berlin Heidelberg, xiii + 262 pp. doi: [10.1007/978-3-540-89628-9](https://doi.org/10.1007/978-3-540-89628-9)
- Mayr, G., 2011. On the osteology and phylogenetic affinities of *Morsoravis sedilis* (Aves) from the early Eocene Fur Formation of Denmark. *Bulletin of the Geological Society of Denmark* 59, 23-35.
- Mayr, G., 2015. A reassessment of Eocene parrotlike fossils indicates a previously undetected radiation of zygodactyl stem group representatives of passerines (Passeriformes). *Zoologica scripta* 44 (6), 587-602. doi: [10.1111/zsc.12128](https://doi.org/10.1111/zsc.12128)
- Mayr, G., Manegold, A., Johansson, U. S., 2003. Monophyletic groups within "higher land birds" – Comparison of morphological and molecular data. *Journal of Zoological Systematics and Evolutionary Research* 41, 233-248.
- Mourer-Chauviré, C., Pickford, M., Senut, B., 2011a. The first Palaeogene galliform from Africa. *Journal of Ornithology* 152 (3), 617-622. doi: [10.1007/s10336-010-0630-9](https://doi.org/10.1007/s10336-010-0630-9)
- Mourer-Chauviré, C., Pickford, M., Senut, B., 2015. Stem group galliform and stem group psittaciform birds (Aves, Galliformes, Paraortygidae, and Psittaciformes, family incertae sedis) from the Middle Eocene of Namibia. *Journal of Ornithology* 156 (1), 275-286. doi: [10.1007/s10336-014-1124-y](https://doi.org/10.1007/s10336-014-1124-y)
- Mourer-Chauviré, C., Tabuce, R., Essid, E. M., Marivaux, L., Khayati, H., Vianey-Liaud, M., Ben Haj Ali, M., 2013. A new taxon of stem group Galliformes and the earliest record for stem group Cuculidae from the Eocene of Djebel Chambi, Tunisia. In Göhlich, U. B., Kroh, A. (Eds.), *Palaeornithological Research 2013*. Proceeding of the 8th International Meeting of the Society of Avian Paleontology and Evolution, pp. 1-15.
- Mourer-Chauviré, C., Tabuce, R., Mahboubi, M., Adaci, M., Bensalah, M., 2011b. A Phororhacoid bird from the Eocene of Africa. *Naturwissenschaften* 98, 815-823. doi: [10.1007/s00114-011-0829-5](https://doi.org/10.1007/s00114-011-0829-5)
- Pickford, M., Senut, B., Mocke, H., Mourer-Chauviré, C., Rage, J.-C., Mein, P., 2014. Eocene aridity in southwestern Africa: timing of onset and biological consequences. *Transactions of the Royal Society of South Africa* 69 (3), 139-144. doi: [10.1080/0035919X.2014.933452](https://doi.org/10.1080/0035919X.2014.933452)
- Prum, R. O., Berv, J.S., Dornburg, A., Field, D. J., Townsend, J. P., Lemmon, E. M., Lemmon, A. R., 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature*. doi: [10.1038/nature15697](https://doi.org/10.1038/nature15697)
- Rasmussen, D. T., Olson, S. L., Simons, E. L., 1987. Fossil birds from the Oligocene Jebel Qatrani Formation, Fayum Province, Egypt. *Smithsonian Contribution to Paleobiology* 62, iv + 20 pp.
- Rasmussen, D. T., Simons, E. L., Hertel, F., Judd, A., 2001. Hindlimb of a giant terrestrial bird from the Upper Eocene, Fayum, Egypt. *Palaeontology* 44 (2), 325-337.
- Stidham, T. A., Smith, N. A., 2015. An ameghinornithid-like bird (Aves, Cariamae ?Ameghinornithidae) from the early Oligocene of Egypt. *Palaeontologia Electronica* 18.1.5A, 1-8.
- Weigel, R. D., 1963. Oligocene birds from Saskatchewan. *Quarterly Journal of the Florida Academy of Sciences* 26/3, 257-262.