



HAL
open science

Targeting the complement cascade: novel treatments coming down the pike

Joshua Thurman, Moglie Le Quintrec

► **To cite this version:**

Joshua Thurman, Moglie Le Quintrec. Targeting the complement cascade: novel treatments coming down the pike. *Kidney International*, 2016, 90 (4), pp.746-752. 10.1016/j.kint.2016.04.018. hal-01838465

HAL Id: hal-01838465

<https://hal.umontpellier.fr/hal-01838465>

Submitted on 17 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HHS Public Access

Author manuscript

Kidney Int. Author manuscript; available in PMC 2017 October 01.

Published in final edited form as:

Kidney Int. 2016 October ; 90(4): 746–752. doi:10.1016/j.kint.2016.04.018.

Targeting the Complement Cascade: Novel Treatments Coming down the Pike

Joshua M. Thurman, MD* and Moglie Le Quintrec, MD†

*Department of Medicine, University of Colorado School of Medicine, Anschutz Medical Campus, Aurora, CO

†Department of Nephrology and renal Transplantation, Lapeyronnie hospital and INSERM U1183, IRMB, Montpellier, France

Abstract

The complement cascade is a vital component of both the innate and adaptive immune systems. Complement activation also contributes to the pathogenesis of many diseases, however, and the kidney is particularly susceptible to complement-mediated injury. Drugs that block complement activation can rapidly reduce tissue inflammation and also attenuate the adaptive immune response to foreign and tissue antigens. Eculizumab is a monoclonal antibody that prevents the cleavage of C5. It has been approved for the treatment of atypical hemolytic uremic syndrome (aHUS), and it has been used in selected patients with other kidney diseases. Many additional drugs are also in development for blocking the complement cascade, including new monoclonal antibodies, recombinant proteins, small molecules, and small interfering RNA (siRNA) agents. Validation of these new drugs as effective treatments for kidney diseases faces several challenges. Many complement-mediated kidney diseases are rare, so it is not feasible to test all of the new drugs in numerous different rare diseases. The onset and course of the diseases are heterogeneous and many of these diseases also carry a life-long risk of recurrence, and it is not clear how long complement inhibition must be maintained. In spite of these challenges, new therapeutic options for targeting the complement system will likely become available in the near future and may prove useful for treating patients with kidney disease.

Keywords

complement; glomerulonephritis; inflammation

Introduction

The kidney is a common target of immune-mediated injury. Several kidney diseases are caused by autoimmunity against antigens expressed within the glomeruli, and the innate

Corresponding author: Joshua M. Thurman, M.D., Division of Nephrology and Hypertension, B-115, 1775 Aurora Court, M20-3103, Aurora, CO 80045, Phone: 303-724-7584, FAX: 303-724-7581, Joshua.Thurman@ucdenver.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

immune system also frequently causes renal injury. Furthermore, kidney failure causes dysregulation of the immune system. Chronic kidney disease (CKD) is associated with a reduced ability to fight infection, for example, yet patients with CKD also have evidence of chronic systemic inflammation.¹ Thus, there is a delicate interrelationship between the kidney and the immune system (Figure 1), and immunomodulatory drugs may be beneficial for treating a many different kidney diseases and their complications.

The complement cascade is a vital component of both the innate and adaptive immune systems, making it an important therapeutic target. Drugs that block complement activation can rapidly reduce tissue inflammation and also attenuate the adaptive immune response to foreign and tissue antigens. Although the specific mechanisms vary, complement activation contributes to the pathogenesis of almost every kidney disease.² This protein cascade is amenable to many different pharmacologic approaches, and anti-complement drugs could play a larger role in the treatment of kidney disease in the years to come.

The complement system

The complement system is comprised of more than 30 plasma and membrane-bound proteins. Activation of the system proceeds in a cascade fashion via three initiation pathways: the classical (CP), lectin (LP), and alternative (AP). During activation the proteins C2, C4, C3, and C5 are cleaved. The resultant protein fragments bind to nearby tissues or enter the systemic circulation, eliciting both local and systemic responses. The complement system mediates detection and removal of pathogens, local inflammatory reactions, the recruitment and activation of phagocytes, direct cell lysis, and the removal of apoptotic cells and immune-complexes.

These downstream effects are primarily mediated by C3a, C5a, C3b, and C5b-9 (Figure 2). C3a and C5a (the “anaphylatoxins”) are small peptides released during complement activation that bind to transmembrane spanning G protein coupled receptors (C3aR and C5aR). C5a also binds to a non-G protein coupled receptor (C5L2). The anaphylatoxin receptors are expressed on myeloid and non-myeloid cells. They induce vasodilation, cytokine and chemokine release, the recruitment of immune cells, and they induce an oxidative burst by macrophages, eosinophils and neutrophils. C5a also contributes to T-cell and antigen-presenting cell activation, expansion, and survival.

During complement activation, C3b is fixed to nearby cells where it amplifies AP activation and contributes to formation of the C5 convertase (activating enzyme). C3 fragments bound to the surface of cells are also ligands for four different complement receptors (CR1–4). C5b-9 (also referred to as the terminal complement complex [TCC] and the membrane attack complex [MAC]) is a multimer that forms pores in the outer membranes of target cells. The flux of fluid and ions through C5b-9 pores can cause cell activation, proliferation, apoptosis, or lysis.

Complement and the kidney

Given its promiscuous involvement in the both the innate and adaptive immune responses, the complement system may provide a convenient “node” for treating a variety of distinct

renal diseases. IgM and IgG containing immune-complexes are strong activators of the CP, which is implicated in many forms of glomerulonephritis including lupus nephritis and cryoglobulinemia. The CP is also activated in antibody mediated transplant rejection (AMR). The LP is activated when MBL proteins or ficolin bind to carbohydrates present on bacteria surfaces. The MBLs and ficolin also bind to molecules displayed on damaged cells, and detection of MBL proteins in the glomeruli of patients with IgA nephropathy suggests involvement the LP.³

AP activation is involved in the pathogenesis of many different types of kidney disease. The AP is continually activated in plasma by the conversion of C3 to its hydrolyzed form, C3(H₂O), which forms part of an initiation C3 convertase. This convertase generates more C3b, self-amplifying AP activation. Spontaneous AP activation provides a rapid response to pathogens, but it must be tightly regulated on host tissues. This balance is maintained by a group of complement regulatory proteins.⁴ Congenital and acquired defects in complement regulation are associated with inflammation, and the kidney is particularly susceptible in this setting. Two severe forms of kidney disease are strongly associated with impaired AP regulation: aHUS and C3 glomerulopathy (C3G).^{5, 6}

The AP also amplifies activation that is initiated through the CP and LP, and can produce more than 80% of the C5b-9 that is generated. As a result, blockade of the AP is protective in immune-complex glomerulonephritis⁷ and ANCA associated vasculitis (AAV).⁸ Factor D is a key component of the AP, and the concentration of factor D increases in patients with CKD due to reduced clearance.⁹ This increases AP activation and may be a cause of systemic inflammation. The AP is also activated by contact of the plasma with artificial surfaces during hemodialysis and cardiopulmonary bypass, contributing to the pro-inflammatory nature of these treatments. Because of the unique and complex relationship between the AP and the kidney, agents that selectively block AP activation may be particularly useful for patients with kidney disease.

Use of eculizumab in patients with kidney disease

Eculizumab is a humanized hybrid IgG2/IgG4 monoclonal antibody directed against human C5. It prevents production of C5a and C5b-9 (Figure 2). Eculizumab was first approved for the treatment of paroxysmal nocturnal hemoglobinuria (PNH), and the efficacy and safety of eculizumab for treating aHUS were demonstrated in two prospective phase II studies in patients who were either unresponsive to or dependent on plasma exchange.¹⁰ Another study showed that eculizumab reduces the recurrence of aHUS after renal transplantation.¹¹ Many patients with aHUS have underlying mutations in complement-associated genes, and the disease can recur after eculizumab is stopped.¹² Consequently, the optimal duration of treatment is not known.

Eculizumab is now the standard of care for PNH and aHUS. There is also compelling evidence that complement activation plays an important role in the pathogenesis of AMD and AMR, and clinical trials of complement inhibitors in these diseases are underway (for example, clinicaltrials.gov NCT01399593 and NCT01567085). Complement activation also plays a direct role in the pathogenesis of C3G. The results of eculizumab in patients with

C3G have been mixed,^{13, 14} however, and further studies will be needed to determine whether eculizumab is beneficial in this disease. The high cost of eculizumab has limited its off-label use in other diseases, but there are reports of its efficacy in lupus nephritis,¹⁵ membranoproliferative glomerulonephritis (MPGN),¹⁶ AMR,¹⁷ and IgA nephropathy.¹⁸

Biologic considerations in the development of complement therapeutics

Mitigation of infection risk

Patients with congenital complement deficiencies are at increased risk of infection. Early complement pathway component deficiencies (C1q, C2, C4, and C3) are associated with recurrent bacterial infections, and terminal complement deficiencies (C5, C6, C7, C8, and C9) are associated with increased risk of infection with encapsulated bacteria, particularly *Neisseria* species.¹⁹ Because eculizumab blocks the terminal complement pathway it carries a similar risk, and has a black box warning recommending that patients receiving the drug should be immunized with polyvalent meningococcal vaccine. Patients living in regions with prevalent serotypes not covered by the vaccine should be empirically treated with appropriate antibiotics. Even patients immunized against meningococcus are at risk of infection during treatment, because antibody-mediated killing of bacteria involves complement activation. In patients with paroxysmal nocturnal hemoglobinuria (PNH) who have been immunized against meningococcus and are treated with eculizumab, the incidence of infection is ~0.9 cases per hundred patient years,²⁰ and infections were reported in three of 180 (~2%) immunized aHUS patients treated with eculizumab. In some centers, empiric antibiotics are now administered to all patients receiving the drug, although this is opinion-based and the risk/benefit of this approach is uncertain.²¹ Children treated with eculizumab should also be vaccinated against *Streptococcus pneumoniae* and *Haemophilus influenzae* type b.

Complement inhibitors must be used cautiously in patients with active infections. Ironically, infections can trigger recurrences of aHUS^{22, 23} and C3G.²⁴ Infections and inflammation increase complement activity in plasma,²⁵ causing breakthrough in patients who were previously completely inhibited.^{26, 27} Thus, infections can be a complication of treatment with complement inhibitory drugs, but may also increase the dose requirement of these drugs. The decision to discontinue, maintain, or increase complement inhibition depends on the severity of the infection and the underlying renal disease. In patients with infections who are continued on therapy, the CH50 and “free anti-C5” should be monitored to ensure full inhibition. The CH50 measures the ability of diluted patient serum to lyse sheep erythrocytes coated with antibody. Lysis of the erythrocytes involves the formation of C5b-9 on the cell surface, and a value of 0 indicates that even with undiluted serum no lysis is seen. Typically the CH50 of patients treated with eculizumab should be 0, or below the lower limit of detection for the lab.¹¹ Eculizumab binds to C5 in the plasma, preventing its cleavage. The detection of anti-C5 antibody that has not bound C5 (free anti-C5) ensures that there is sufficient drug to block all of the available C5. Free anti-C5 levels > 35 µg/mL are associated with complete inhibition of lytic activity.²⁸ In the study by Legendre et al., trough levels were maintained at levels of at least 50–100 µg/mL in treated aHUS patients.¹⁰

Other physiologic functions of the complement system

The complement cascade has physiologic functions beyond its role in fighting infection, including the removal of apoptotic and necrotic cells and solubilization of immune-complexes.²⁹ Classical pathway deficient mice have an impaired ability to remove apoptotic cells.³⁰ Patients with deficiency of early CP proteins are at increased risk of autoimmune diseases, possibly because apoptotic nuclear debris is not efficiently cleared.³¹ In a mouse model of lupus nephritis, deficiency of C3 did not protect the mice from nephritis.³² More glomerular IgG was detected in these mice even though the level of circulating immune complexes was the same as in C3 sufficient mice, suggesting that complement helps with the removal of immune complexes in this model. There are also case reports linking congenital deficiency of complement proteins, including C3, with MPGN-like disease, possibly due to an impaired ability to remove glomerular ICs.³³ It is not known whether complement inhibition at the level of C3 could similarly increase deposition of ICs in the glomeruli, but the findings mentioned above suggest that this is a possibility.

Complete complement blockade versus blockade of specific pathways or fragments

The complement activation pathways converge at the C3 and C5 convertases (Figure 2). These enzymatic complexes are critical for complement activation and are logical targets for complement inhibitors. The advantage of using agents that target the C3 convertase is that this shuts down production of all pro-inflammatory fragments (Figure 2). Targeting the complement cascade at the level of C3 may be more effective for preventing tissue inflammation than targeting it downstream, but this approach blocks physiologic as well as pathologic functions of the cascade.

A related question is whether C3a contributes to the pathogenesis of renal disease. The C3a receptor directly triggers NF- κ B activation in renal tubular epithelial cells and causes the cells to release pro-inflammatory chemokines.³⁴ C3a may also have anti-inflammatory effects,³⁵ however, and C3a receptor deficiency was associated with more severe autoimmunity in a lupus-prone strain of mice.³⁶ In spite of its complicated effects, there is experimental evidence that C3a contributes to injury in several models of renal disease, including a rat model of diabetic nephropathy³⁷ and a model of focal segmental glomerulosclerosis.³⁸ Blockade of the complement system at the level of C3 may therefore be advantageous.

Does one need to completely inhibit the complement system?

Complement activation is not a binary process. There is continual low-level activation in the plasma, and in health there is a delicate balance between activating and inhibitory factors. Patients with heterozygous deficiencies do not seem to be susceptible to infection,³¹ so partial complement inhibition could reduce the cost and the infectious risk of therapy. On the other hand, prolongation of the period between doses has been associated with increased disease activity in post-transplant aHUS patients.¹¹ The published studies studying the efficacy of eculizumab in aHUS achieved full inhibition of the complement system, however, and current dosing guidelines for eculizumab are designed to fully suppress the CH50.

Complement inhibition in patients with different underlying molecular complement defects

A large number of genetic variants and autoantibodies have been identified in patients with complement-mediated diseases. These underlying defects may make some patients unresponsive to particular drugs. For example, a C5 polymorphism has been identified in Japanese patients who are unresponsive to treatment with eculizumab.³⁹ Given the large number of autoantibodies, loss of function genetic variants, and gain of function genetic variants that are associated with kidney disease, there probably are subsets of patients who will not respond to the complement inhibitory drugs in development.

Practical considerations in the development of complement therapeutics

The complement inhibitory drugs currently in development fall into several different classes of molecules. Some are protein therapeutics: purified proteins, monoclonal antibodies, and recombinant proteins. Small molecule complement antagonists are in development, and siRNA agents are being developed to “knock-down” complement protein production in the liver. Each of these strategies has practical advantages and disadvantages.

Cost

Eculizumab has been one of the most expensive drugs on the market since its release, in part because of the cost of manufacturing monoclonal antibodies.⁴⁰ The average cost for small molecules, in contrast, is estimated to be less than 1/20th the cost of biologic drugs.⁴¹

Route of administration

Monoclonal antibodies, recombinant proteins, and SiRNAs must be administered intravenously or subcutaneously. In contrast, small molecule drugs can be orally administered. Although directly observed intravenous administration may improve compliance,⁴² it adds to the cost and inconvenience of treatment. Patients can be trained to perform subcutaneous injections, so drugs that can be delivered by this route can be administered at home.

Pharmacokinetics/pharmacodynamics

The half-life of monoclonal antibodies and protein therapeutics is longer than that of small molecules. SiRNA therapies may provide prolonged complement inhibition. A long half-life is advantageous in terms of compliance and convenience but is a disadvantage if treatment needs to be discontinued due to side effects or infection.

Toxicity

Because the complement system is comprised of interacting proteins, it lends itself to protein therapeutics. Monoclonal antibodies bind to protein targets with high specificity and affinity, and there are minimal off-target effects. Protein therapies can be immunogenic, however, particularly when used chronically. Small molecule therapeutics, in contrast, generally have more off-target effects than biologic agents.⁴²

Novel complement therapeutics in development

Purified plasma proteins

Plasma infusion can provide patients with soluble proteins, such as factor H, that are abundant in the plasma of healthy donors. The administration of a purified complement regulatory protein is a logical method to suppress complement activation, particularly in patients with a factor H deficiency. The administration of a purified protein may not be effective, however, in patients with auto-antibodies or circulating inhibitors of the protein.

Purified factor H—Recombinant factor H was developed by Ophtherion (now licensed to Baxter) for the treatment of age-related macular degeneration (AMD). Although purified factor H may be beneficial in patients with genetic mutations in factor H, this agent is not currently being developed for use in patients with kidney diseases.

C1 inhibitor—Hereditary angioedema is a disease that affects patients with deficiency or acquired deficiencies of C1 inhibitor (C1inh), and purified C1inh has been approved for the treatment of hereditary angioedema.⁴³ A phase I/II study was conducted in which highly sensitized renal transplant recipients were randomized to C1inh or placebo.⁴⁴ None of the 10 patients in the C1inh group developed AMR during the study, although only 1 patient in the placebo group developed AMR. Recently, the C1inh was used as rescue therapy for AMR and seemed to improve kidney graft function.⁴⁵ A randomized double-blind study to evaluate the efficacy and safety of C1inh for the treatment of acute AMR in kidney transplantation is currently underway (clinicaltrials.gov NCT02052141).

Monoclonal antibodies

Monoclonal antibodies can bind target proteins with high affinity and high specificity. This can cause depletion of the target protein, or it can block the biologic activity of the target. Monoclonal antibodies have been developed against many of the complement proteins, including C5, factor B, factor D, C1s, the mannose associated serine protease-2 (MASP2), properdin, and C3b.⁴⁶ None of these are currently being tested in patients with renal disease, although some have shown promise in patients with other diseases.

Anti-C1s—A monoclonal antibody to C1s (TNT009, True North Therapeutics) selectively blocks classical pathway activation and has been developed for use in antibody-mediated diseases. It is currently being tested in a Phase 1a/1b clinical trial (clinicaltrials.gov NCT02502903). This study includes patients with several diseases believed to be complement-mediated, including patients with end stage renal disease (ESRD) who also have anti-HLA antibodies. By measuring biomarkers of complement activity, the investigators may accumulate evidence that this agent effectively blocks complement activation in these diseases.

Anti-factor D—A monoclonal antibody to factor D (Lampalizumab, Genentech) selectively blocks AP activation in the eye when injected intra-vitreally.⁴⁷ In an 18-month phase II study it prevented progression of AMD (clinicaltrials.gov NCT01229215, results

not yet published), and it is being tested in a phase III clinical trial (clinicaltrials.gov NCT02247479).

Engineered proteins

In addition to purifying complement inhibitory proteins from plasma, proteins can also be synthesized using recombinant technology. These proteins can be designed to incorporate the complement regulatory regions of the endogenous proteins, and can also be engineered to have other useful properties.

Complement receptor 1—CR1 is a potent inhibitor of the CP, LP, and AP, and a soluble form of CR1 was developed for therapeutic use.⁴⁸ A clinical trial tested whether this drug is protective in patients undergoing cardiac surgery, but the drug did not significantly improve the primary endpoint.⁴⁹ In a single patient with C3G, this drug reduced complement consumption, as evidenced by increased total C3 levels and decreased sC5b-9 levels.⁵⁰

Targeted complement regulatory proteins—Several strategies have been used to deliver complement inhibitors specifically to sites of inflammation. These agents may have fewer systemic side effects than untargeted inhibitors. One such inhibitor was developed in which the complement regulatory region of CR1 was attached to a membrane associating peptide and a hydrophobic myristoyl group that inserts itself into cell membranes (APT070).⁵¹ Treatment of rat kidney allografts with this agent reduced tubular injury and improved allograft survival. A phase II study is currently underway to test whether this agent will reduce the incidence of delayed graft function in transplant patients.

Complement receptor 2 selectively binds C3d, and engineered proteins have been developed that use the C3d binding region of CR2 to deliver the complement inhibitory regions of factor H to sites of complement activation (TT30, Alexion Pharmaceuticals). A phase I study in patients with PNH demonstrated that intravenous and subcutaneous injections of the drug reduced plasma complement activity and reduced LDH levels.⁵²

Small molecules

Small organic molecules (typically <1 kD) provide another class of molecules that can affect biologic processes.⁴² Large libraries of these molecules can be screened to identify candidates that interfere with receptors or enzymatic processes, and several small molecules have been developed to target the complement cascade.

C3 blockade—Compstatin (Potentia Pharmaceuticals) is a cyclic tridecapeptide that blocks the cleavage of C3.⁵³ Compstatin was tested intra-vitreally in a Phase I study of patients with AMD. It also blocked complement activation by serum from patients with C3G in vitro, including patients with C3 nephritic factors and anti-factor H autoantibodies.⁵⁴

Factor D blockade—Small molecule inhibitors that bind factor D and block AP activation are currently under development by Achillion Pharmaceuticals (ACH-3856, ACH-4100, ACH-4471).

C5a receptor blockade—A small molecule inhibitor of the C5aR (CCX168, ChemoCentryx) protected mice in a model of AAV and is currently being tested in a phase II trial ([clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02222155) NCT02222155). Preliminary results indicate that the drug improves overall disease activity, including renal manifestations.⁵⁵ Trials are also underway to test the efficacy of the drug in IgA nephropathy ([Clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02384317) NCT02384317) and in patients with aHUS who have reached ESRD ([Clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02464891) NCT02464891).

Small interfering RNA

siRNA therapeutics suppress the production of target proteins. A siRNA agent that prevents production of C5 (ALN-CC5, Alnylam Pharmaceuticals) has been developed. In a Phase I study in healthy adults, a single subcutaneous dose of the drug reduced complement hemolytic activity by more than 90% for more than 2 months.⁵⁶ The PK/PD of siRNA are advantageous for patients who require chronic complement inhibition, although complement inhibition is difficult to reverse in patients undergoing treatment.

Future directions

The kidney is perhaps more susceptible to complement-mediated injury than any other organ,² and the development of new complement inhibitory drugs could improve the care of many different kidney diseases. As these drugs become available, however, a major challenge will be to design clinical trials that can adequately test these numerous different agents in patients with rare or slowly progressive kidney diseases. The difficulty of achieving approval for new treatments for the glomerular diseases is not unique to complement inhibitors, and some features of the complement system may actually be advantageous for testing new therapies. There are good animal models (e.g. “knockout mice”) for studying complement biology and there is a long history of measuring complement proteins in the plasma and in biopsy tissue from patients with glomerular diseases.

Even though aHUS is a very rare disease, it may provide the easiest pathway for new complement inhibitory drugs into the clinic. Clinical trials of eculizumab have demonstrated the efficacy of complement blockade in aHUS, and the response to treatment can be very rapid – measured in weeks, not years.¹⁰ Complement activation also appears to contribute to the pathogenesis of more common diseases, such as lupus nephritis and IgA nephropathy,⁵⁷ but the role for the new complement inhibitory drugs in these diseases is not yet clear. Most forms of glomerulonephritis can be slowly progressive, adding to the length and cost of clinical trials. Better biomarkers of complement activation within tissues may be useful for rapidly confirming the biologic effects of new complement inhibitory drugs. The currently available biomarkers of complement activation, such as plasma C3 and C4 levels, are imperfect indicators of complement activation within the kidney. More sensitive biomarkers of intra-renal complement activation/inhibition could provide rapid evidence that a new drug is working as expected. Nevertheless, rigorous clinical studies will still be required to confirm that the new drugs improve patient outcomes.

Will the new complement inhibitors transform the way we treat glomerulonephritis or simply provide additional treatments for aHUS? There is reason to be optimistic that

complement inhibitory drugs will have a role in the treatment of a wide range of renal diseases. To achieve this, however, will require carefully designed multi-center clinical trials and a broad-based commitment of physicians and patients to testing the new therapies.

Acknowledgments

This work was supported by National Institutes of Health Grants R01 DK076690 (JMT).

Disclosure.

JMT receive royalties from Alexion Pharmaceuticals, Inc. MLQ has received consulting fees from Alexion Pharmaceuticals, Inc.

References

1. Betjes MG. Immune cell dysfunction and inflammation in end-stage renal disease. *Nature reviews Nephrology*. 2013; 9:255–265. [PubMed: 23507826]
2. Thurman JM. Complement in kidney disease: core curriculum 2015. *Am J Kidney Dis*. 2015; 65:156–168. [PubMed: 25441433]
3. Endo M, Ohi H, Ohsawa I, et al. Glomerular deposition of mannose-binding lectin (MBL) indicates a novel mechanism of complement activation in IgA nephropathy. *Nephrol Dial Transplant*. 1998; 13:1984–1990. [PubMed: 9719152]
4. Zipfel PF, Skerka C. Complement regulators and inhibitory proteins. *Nat Rev Immunol*. 2009; 9:729–740. [PubMed: 19730437]
5. Le Quintrec M, Roumenina L, Noris M, et al. Atypical hemolytic uremic syndrome associated with mutations in complement regulator genes. *Seminars in thrombosis and hemostasis*. 2010; 36:641–652. [PubMed: 20865641]
6. Servais A, Noel LH, Roumenina LT, et al. Acquired and genetic complement abnormalities play a critical role in dense deposit disease and other C3 glomerulopathies. *Kidney Int*. 2012; 82:454–464. [PubMed: 22456601]
7. Watanabe H, Garnier G, Circolo A, et al. Modulation of renal disease in MRL/lpr mice genetically deficient in the alternative complement pathway factor B. *J Immunol*. 2000; 164:786–794. [PubMed: 10623824]
8. Xiao H, Schreiber A, Heeringa P, et al. Alternative complement pathway in the pathogenesis of disease mediated by anti-neutrophil cytoplasmic autoantibodies. *Am J Pathol*. 2007; 170:52–64. [PubMed: 17200182]
9. Volanakis JE, Barnum SR, Giddens M, et al. Renal filtration and catabolism of complement protein D. *N Engl J Med*. 1985; 312:395–399. [PubMed: 3844050]
10. Legendre CM, Licht C, Muus P, et al. Terminal complement inhibitor eculizumab in atypical hemolytic-uremic syndrome. *N Engl J Med*. 2013; 368:2169–2181. [PubMed: 23738544]
11. Zuber J, Quintrec ML, Krid S, et al. Eculizumab for Atypical Hemolytic Uremic Syndrome Recurrence in Renal Transplantation. *Am J Transplant*. 2012
12. Ardissino G, Testa S, Possenti I, et al. Discontinuation of eculizumab maintenance treatment for atypical hemolytic uremic syndrome: a report of 10 cases. *Am J Kidney Dis*. 2014; 64:633–637. [PubMed: 24656451]
13. Le Quintrec M, Lionet A, Kandel C, et al. Eculizumab for treatment of rapidly progressive C3 glomerulopathy. *Am J Kidney Dis*. 2015; 65:484–489. [PubMed: 25530108]
14. Vivarelli M, Emma F. Treatment of C3 glomerulopathy with complement blockers. *Seminars in thrombosis and hemostasis*. 2014; 40:472–477. [PubMed: 24799307]
15. Pickering MC, Ismajli M, Condon MB, et al. Eculizumab as rescue therapy in severe resistant lupus nephritis. *Rheumatology (Oxford)*. 2015
16. Radhakrishnan S, Lunn A, Kirschfink M, et al. Eculizumab and refractory membranoproliferative glomerulonephritis. *N Engl J Med*. 2012; 366:1165–1166. [PubMed: 22435384]

17. Stegall MD, Diwan T, Raghavaiah S, et al. Terminal complement inhibition decreases antibody-mediated rejection in sensitized renal transplant recipients. *Am J Transplant*. 2011; 11:2405–2413. [PubMed: 21942930]
18. Rosenblad T, Rebetz J, Johansson M, et al. Eculizumab treatment for rescue of renal function in IgA nephropathy. *Pediatr Nephrol*. 2014; 29:2225–2228. [PubMed: 24924752]
19. Skattum L, van Deuren M, van der Poll T, et al. Complement deficiency states and associated infections. *Mol Immunol*. 2011; 48:1643–1655. [PubMed: 21624663]
20. Kelly RJ, Hill A, Arnold LM, et al. Long-term treatment with eculizumab in paroxysmal nocturnal hemoglobinuria: sustained efficacy and improved survival. *Blood*. 2011; 117:6786–6792. [PubMed: 21460245]
21. Zuber J, Fakhouri F, Roumenina LT, et al. Use of eculizumab for atypical haemolytic uraemic syndrome and C3 glomerulopathies. *Nature reviews Nephrology*. 2012; 8:643–657. [PubMed: 23026949]
22. Licht C, Greenbaum LA, Muus P, et al. Efficacy and safety of eculizumab in atypical hemolytic uremic syndrome from 2-year extensions of phase 2 studies. *Kidney Int*. 2015; 87:1061–1073. [PubMed: 25651368]
23. Zuber J, Le Quintrec M, Morris H, et al. Targeted strategies in the prevention and management of atypical HUS recurrence after kidney transplantation. *Transplant Rev (Orlando)*. 2013; 27:117–125. [PubMed: 23937869]
24. Vernon KA, Goicoechea de Jorge E, Hall AE, et al. Acute presentation and persistent glomerulonephritis following streptococcal infection in a patient with heterozygous complement factor H-related protein 5 deficiency. *Am J Kidney Dis*. 2012; 60:121–125. [PubMed: 22503529]
25. Schutte M, DiCamelli R, Murphy P, et al. Effects of anesthesia, surgery and inflammation upon host defense mechanisms. I. Effects upon the complement system. *International archives of allergy and applied immunology*. 1975; 48:706–720. [PubMed: 805770]
26. Nester C, Stewart Z, Myers D, et al. Pre-emptive eculizumab and plasmapheresis for renal transplant in atypical hemolytic uremic syndrome. *Clin J Am Soc Nephrol*. 2011; 6:1488–1494. [PubMed: 21617085]
27. Weitz M, Amon O, Bassler D, et al. Prophylactic eculizumab prior to kidney transplantation for atypical hemolytic uremic syndrome. *Pediatr Nephrol*. 2011; 26:1325–1329. [PubMed: 21556717]
28. Hill A, Hillmen P, Richards SJ, et al. Sustained response and long-term safety of eculizumab in paroxysmal nocturnal hemoglobinuria. *Blood*. 2005; 106:2559–2565. [PubMed: 15985537]
29. Ricklin D, Hajishengallis G, Yang K, et al. Complement: a key system for immune surveillance and homeostasis. *Nat Immunol*. 2010; 11:785–797. [PubMed: 20720586]
30. Taylor PR, Carugati A, Fadok VA, et al. A hierarchical role for classical pathway complement proteins in the clearance of apoptotic cells in vivo. *J Exp Med*. 2000; 192:359–366. [PubMed: 10934224]
31. Pettigrew HD, Teuber SS, Gershwin ME. Clinical significance of complement deficiencies. *Ann N Y Acad Sci*. 2009; 1173:108–123. [PubMed: 19758139]
32. Sekine H, Reilly CM, Molano ID, et al. Complement component C3 is not required for full expression of immune complex glomerulonephritis in MRL/lpr mice. *J Immunol*. 2001; 166:6444–6451. [PubMed: 11342671]
33. Coleman TH, Forristal J, Kosaka T, et al. Inherited complement component deficiencies in membranoproliferative glomerulonephritis. *Kidney Int*. 1983; 24:681–690. [PubMed: 6663990]
34. Thurman JM, Lenderink AM, Royer PA, et al. C3a is required for the production of CXC chemokines by tubular epithelial cells after renal ischemia/reperfusion. *J Immunol*. 2007; 178:1819–1828. [PubMed: 17237432]
35. Coulthard LG, Woodruff TM. Is the Complement Activation Product C3a a Proinflammatory Molecule? Re-evaluating the Evidence and the Myth. *J Immunol*. 2015; 194:3542–3548. [PubMed: 25848071]
36. Wenderfer SE, Wang H, Ke B, et al. C3a receptor deficiency accelerates the onset of renal injury in the MRL/lpr mouse. *Mol Immunol*. 2009; 46:1397–1404. [PubMed: 19167760]

37. Li L, Yin Q, Tang X, et al. C3a receptor antagonist ameliorates inflammatory and fibrotic signals in type 2 diabetic nephropathy by suppressing the activation of TGF-beta/smad3 and IKKalpha pathway. *PLoS one*. 2014; 9:e113639. [PubMed: 25422985]
38. Tang Z, Lu B, Hatch E, et al. C3a mediates epithelial-to-mesenchymal transition in proteinuric nephropathy. *J Am Soc Nephrol*. 2009; 20:593–603. [PubMed: 19158354]
39. Nishimura J, Yamamoto M, Hayashi S, et al. Genetic variants in C5 and poor response to eculizumab. *N Engl J Med*. 2014; 370:632–639. [PubMed: 24521109]
40. Shaughnessy AF. Monoclonal antibodies: magic bullets with a hefty price tag. *BMJ*. 2012; 345:e8346. [PubMed: 23236036]
41. McCamish M, Woollett G. Worldwide experience with biosimilar development. *MABs*. 2011; 3:209–217. [PubMed: 21441787]
42. Mocsai A, Kovacs L, Gergely P. What is the future of targeted therapy in rheumatology: biologics or small molecules? *BMC Med*. 2014; 12:43. [PubMed: 24620738]
43. Cicardi M, Aberer W, Banerji A, et al. Classification, diagnosis, and approach to treatment for angioedema: consensus report from the Hereditary Angioedema International Working Group. *Allergy*. 2014; 69:602–616. [PubMed: 24673465]
44. Vo AA, Zeevi A, Choi J, et al. A phase I/II placebo-controlled trial of C1-inhibitor for prevention of antibody-mediated rejection in HLA sensitized patients. *Transplantation*. 2015; 99:299–308. [PubMed: 25606785]
45. Viglietti D, Gosset C, Loupy A, et al. C1-Inhibitor in Acute Antibody-Mediated Rejection Non-Responsive to Conventional Therapy in Kidney Transplant Recipients: A Pilot Study. *Am J Transplant*. 2015
46. Ricklin D, Lambris JD. Progress and Trends in Complement Therapeutics. *Adv Exp Med Biol*. 2013; 735:1–22.
47. Le KN, Gibiansky L, van Lookeren Campagne M, et al. Population Pharmacokinetics and Pharmacodynamics of Lampalizumab Administered Intravitreally to Patients With Geographic Atrophy. *CPT Pharmacometrics Syst Pharmacol*. 2015; 4:595–604. [PubMed: 26535160]
48. Weisman HF, Bartow T, Leppo MK, et al. Soluble human complement receptor type 1: in vivo inhibitor of complement suppressing post-ischemic myocardial inflammation and necrosis. *Science*. 1990; 249:146–151. [PubMed: 2371562]
49. Lazar HL, Bokesch PM, van Lenta F, et al. Soluble human complement receptor 1 limits ischemic damage in cardiac surgery patients at high risk requiring cardiopulmonary bypass. *Circulation*. 2004; 110:II274–II279. [PubMed: 15364875]
50. Zhang Y, Nester CM, Holanda DG, et al. Soluble CR1 therapy improves complement regulation in C3 glomerulopathy. *J Am Soc Nephrol*. 2013; 24:1820–1829. [PubMed: 23907509]
51. Patel H, Smith RA, Sacks SH, et al. Therapeutic strategy with a membrane-localizing complement regulator to increase the number of usable donor organs after prolonged cold storage. *J Am Soc Nephrol*. 2006; 17:1102–1111. [PubMed: 16510763]
52. Risitano, AM.; Storek, M.; Sahelijo, L., et al., editors. Safety and Pharmacokinetics of the Complement Inhibitor TT30 in a Phase I Trial for Untreated PNH Patients; *Proceedings of the Conference Name*; Date Year of Conference; Conference Location|. Publisher|; Place Published|, Year Published|.
53. Qu H, Ricklin D, Bai H, et al. New analogs of the clinical complement inhibitor compstatin with subnanomolar affinity and enhanced pharmacokinetic properties. *Immunobiology*. 2013; 218:496–505. [PubMed: 22795972]
54. Zhang Y, Shao D, Ricklin D, et al. Compstatin analog Cp40 inhibits complement dysregulation in vitro in C3 glomerulopathy. *Immunobiology*. 2015; 220:993–998. [PubMed: 25982307]
55. Jayne, DRW.; Bruchfeld, A.; Schaier, M., et al., editors. PHASE 2 RANDOMISED TRIAL OF ORAL C5A RECEPTOR ANTAGONIST CCX168 IN ANCA-ASSOCIATED RENAL VASCULITIS; *Proceedings of the Conference Name*; Date Year of Conference; Conference Location|. Publisher|; Place Published|, Year Published|.
56. Hill, A.; Taubel, J.; Bush, J., et al., editors. Subcutaneously Administered Investigational RNAi Therapeutic (ALN-CC5) Targeting Complement C5 for Treatment of PNH and Complement-

Mediated Diseases: Interim Phase 1 Study Results; *Proceedings of the Conference Name*; Date Year of Conference; Conference Location|. Publisher|; Place Published|, Year Published|.

57. Maillard N, Wyatt RJ, Julian BA, et al. Current Understanding of the Role of Complement in IgA Nephropathy. *J Am Soc Nephrol*. 2015

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. The Complement System and Kidney Disease

Complement activation contributes to the pathogenesis of acute and chronic kidney injury. Damage to the kidney, in turn, increases local and systemic complement activation. The complement cascade may link kidney disease with an increased susceptibility to infection and systemic inflammation. Complement inhibitory drugs hold the promise of blocking many forms of immune-mediated kidney injury and reducing the systemic effects of kidney disease.




Figure 2. Overview of drugs that target the complement cascade

Complement activation is initiated through three pathways: the classical pathway, alternative pathway, and lectin pathway. Full activation leads to the generation of several biologically active fragments, namely C3a, C5a, C3b, and C5b-9. Drugs are currently being developed to selectively block the classical pathway, the alternative pathway, activation at the level of C3, activation at the level of C5, and C5a. ¹Purified proteins, ²Monoclonal antibodies, ³Engineered proteins, ⁴Small molecules, ⁵siRNA.