



HAL
open science

Two Novel, Distantly Related Papillomaviruses Isolated from Healthy Skin of the Timor Deer (*Rusa timorensis*)

Beatriz Mengual-Chuliá, Gudrun Wibbelt, Marc Gottschling, Ignacio G Bravo

► To cite this version:

Beatriz Mengual-Chuliá, Gudrun Wibbelt, Marc Gottschling, Ignacio G Bravo. Two Novel, Distantly Related Papillomaviruses Isolated from Healthy Skin of the Timor Deer (*Rusa timorensis*). *Genome Announcements*, 2018, 6 (24), pp.505-523. 10.1128/genomeA.00505-18 . hal-01898177

HAL Id: hal-01898177

<https://hal.umontpellier.fr/hal-01898177>

Submitted on 18 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Two Novel, Distantly Related Papillomaviruses Isolated from Healthy Skin of the Timor Deer (*Rusa timorensis*)

Beatriz Mengual-Chuliá,^a Gudrun Wibbelt,^b Marc Gottschling,^c  Ignacio G. Bravo^d

^aInfections and Cancer Laboratory, Catalan Institute of Oncology (ICO), Barcelona, Spain

^bDepartment of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

^cDepartment Biologie I, Systematic Botany and Mycology, GeoBio-Center, Ludwig-Maximilians-Universität, Munich, Germany

^dCentre National de la Recherche Scientifique (CNRS) Laboratory MIVEGEC (UMR CNRS IRD UM), Montpellier, France

ABSTRACT We report the complete genome sequences of *Rusa timorensis papillomavirus 1* (RtimPV1) and *Rusa timorensis papillomavirus 2* (RtimPV2), isolated from hair follicles of asymptomatic skin from the same Timor deer specimen. RtimPV1 and RtimPV2 are evolutionarily only distantly related. RtimPV1 lacks a canonical E2-binding site, and RtimPV2 does not carry an E6 gene.

The genome of papillomaviruses (PVs) is a single double-stranded DNA (dsDNA) circular molecule of approximately 8 kbp, carrying genes involved in cell replication, viral replication, and viral encapsidation. Most PVs cause epithelial asymptomatic infections in amniotes and probably also in bony fishes, while some of them induce proliferative benign lesions and different cancers (1, 2).

The Timor deer *Rusa timorensis* (de Blainville, 1822) is native to the island of Java and is classified as “vulnerable” by the International Union for Conservation of Nature (IUCN) Red List of Threatened Species (<http://www.iucnredlist.org>) (3). A biopsy specimen from asymptomatic skin (isolate IZW 39/08) was collected from a female deer, about 19 years old, living in captivity in a zoo in Berlin, Germany. The animal did not present papilloma-like lesions and was euthanized because of a severe internal infection that was unrelated to any PV infection. Total DNA was extracted from hair bulbs and PCR tested for the presence of PV DNA using broad-spectrum primers targeting the E1 and L1 genes (4). Using this partial information, we designed tail-to-tail primers to amplify the full-length genomes using long-range PCR, which were then Sanger sequenced in both strands through primer walking and cloned. Phylogenetic relationships for the E1E2, L2L1, and E1E2L2L1 gene concatenates were inferred under a maximum-likelihood framework (5).

We retrieved two complete PV genomes from the same skin sample of the *R. timorensis* individual, here named *Rusa timorensis papillomavirus 1* (RtimPV1) (isolate IZW 39/08) and *Rusa timorensis papillomavirus 2* (RtimPV2) (isolate IZW 39/08). Both viral genomes display the standard features of PVs, with a long control region; the early genes E7, E1, and E2; and the late genes L2 and L1. Remarkably, RtimPV2 does not carry an E6 gene. This gene may have been lost independently in several PV lineages through time (6, 7). The RtimPV1 L1 gene displays maximum nucleotide similarity values around 65% with very divergent PVs, while RtimPV2 L1 displays identity values around 65% with divergent PVs in the 5' end, and only above 50% in the 3' end. Therefore, this criterion alone does not suffice for taxonomic assignment following the current rules (8), which need to be revised (7, 9). Phylogenetic reconstruction confidently places RtimPV1 as a sister lineage to *Capra hircus papillomavirus 1* (ChPV1), isolated from healthy skin of a female goat (10), a *Phipapillomavirus* in the Beta + Xi PV supergroup

Received 2 May 2018 Accepted 7 May 2018 Published 14 June 2018

Citation Mengual-Chuliá B, Wibbelt G, Gottschling M, Bravo IG. 2018. Two novel, distantly related papillomaviruses isolated from healthy skin of the Timor deer (*Rusa timorensis*). Genome Announc 6:e00505-18. <https://doi.org/10.1128/genomeA.00505-18>.

Copyright © 2018 Mengual-Chuliá et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ignacio G. Bravo, ignacio.bravo@ird.fr.

(11). Neither ChPV1 nor RtimPV1 displays a canonical E2-binding site in the long control region, thus suggesting alternative regulation of early gene expression. Phylogenetic inference unambiguously places RtimPV2 basal to *Epsilonpapillomavirus* and *Deltapapillomavirus*, within the Delta + Zeta PV supergroup, which includes PVs isolated from giraffes and camels, as well as from different bovid and deer species (11). The RaPV genome isolated from *Rusa alfredi* (Sclater, 1870), with a host closely related to *R. timorensis*, branches in a derived position within the Delta + Zeta PV supergroup (12) and is only distantly related to RtimPV2.

Our results exemplify the diversity of PVs asymptotically infecting similar cells within a single host individual and highlight the genomic plasticity within *Papillomaviridae*.

Accession number(s). The complete genome sequences for RtimPV1 and for RtimPV2 are available in GenBank under the accession no. [KP757765](#) and [KT852571](#), respectively.

ACKNOWLEDGMENTS

B.M.-C. was the recipient of an IDIBELL PhD fellowship. This work was partly funded by the former Spanish Ministry for Science and Innovation (CGL2010-16713). I.G.B. is funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (CODOVIREVOL, grant agreement 647916). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1. Bravo IG, Féllez-Sánchez M. 2015. Papillomaviruses: viral evolution, cancer and evolutionary medicine. *Evol Med Public Health* 2015:32–51. <https://doi.org/10.1093/emph/eov003>.
2. Van Doorslaer K. 2013. Evolution of the *Papillomaviridae*. *Virology* 445: 11–20. <https://doi.org/10.1016/j.virol.2013.05.012>.
3. International Union for Conservation of Nature. 2015. *Rusa timorensis*, p e.T41789A22156866. In Hedges S, Duckworth JW, Timmins R, Semiadi G, Dryden G (ed), The IUCN Red List of Threatened Species. International Union for Conservation of Nature, Gland, Switzerland.
4. Mengual-Chuliá B, García-Pérez R, Gottschling M, Nindl I, Bravo IG. 2012. Novel animal papillomavirus sequences and accurate phylogenetic placement. *Mol Phylogenet Evol* 65:883–891. <https://doi.org/10.1016/j.ympev.2012.08.011>.
5. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>.
6. Van Doorslaer K, McBride AA. 2016. Molecular archeological evidence in support of the repeated loss of a papillomavirus gene. *Sci Rep* 6:33028. <https://doi.org/10.1038/srep33028>.
7. Bravo IG, de Sanjosé S, Gottschling M. 2010. The clinical importance of understanding the evolution of papillomaviruses. *Trends Microbiol* 18: 432–438. <https://doi.org/10.1016/j.tim.2010.07.008>.
8. Bernard H-U, Burk RD, Chen Z, van Doorslaer K, zur Hausen H, de Villiers E-M. 2010. Classification of papillomaviruses (PVs) based on 189 PV types and proposal of taxonomic amendments. *Virology* 401:70–79. <https://doi.org/10.1016/j.virol.2010.02.002>.
9. Bravo IG, de Sanjosé S, Gottschling M. 2011. Papillomaviruses and Darwinian classification: response to Van Doorslaer et al. *Trends Microbiol* 19:49–50. <https://doi.org/10.1016/j.tim.2010.11.008>.
10. Van Doorslaer K, Rector A, Vos P, Van Ranst M. 2006. Genetic characterization of the *Capra hircus* papillomavirus: a novel close-to-root artiodactyl papillomavirus. *Virus Res* 118:164–169. <https://doi.org/10.1016/j.virusres.2005.12.007>.
11. Gottschling M, Göker M, Stamatakis A, Bininda-Emonds ORP, Nindl I, Bravo IG. 2011. Quantifying the phylodynamic forces driving papillomavirus evolution. *Mol Biol Evol* 28:2101–2113. <https://doi.org/10.1093/molbev/msr030>.
12. Fux R, Langenmayer MC, Jörgens D, Schubert C, Heckel J-O, Sutter G. 2016. *Rusa alfredi* papillomavirus 1—a novel *Deltapapillomavirus* inducing endemic papillomatosis in the endangered Visayan spotted deer. *J Gen Virol* 97:128–133. <https://doi.org/10.1099/jgv.0.000340>.