



HAL
open science

Opposing Patterns of Seasonal Change in Functional and Phylogenetic Diversity of Tadpole Assemblages

Axel Strauss, Francois Guilhaumon, Roger Daniel Randrianiaina, Katharina Wollenberg Valero, Miguel Vences, Julian Glos

► To cite this version:

Axel Strauss, Francois Guilhaumon, Roger Daniel Randrianiaina, Katharina Wollenberg Valero, Miguel Vences, et al.. Opposing Patterns of Seasonal Change in Functional and Phylogenetic Diversity of Tadpole Assemblages. PLoS ONE, 2016, 11 (3), pp.e0151744. 10.1371/journal.pone.0151744 . hal-01923907

HAL Id: hal-01923907

<https://hal.umontpellier.fr/hal-01923907>

Submitted on 26 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Opposing Patterns of Seasonal Change in Functional and Phylogenetic Diversity of Tadpole Assemblages

Axel Strauß^{1,2*}, François Guilhaumon³, Roger Daniel Randrianiaina^{1,4}, Katharina C. Wollenberg Valero⁵, Miguel Vences¹, Julian Glos⁶

1 Zoologisches Institut, Technische Universität Braunschweig, Mendelssohnstr. 4, 38106, Braunschweig, Germany, **2** Institut für Genetik, Ludwig-Maximilians-Universität München, Großhaderner Str. 2–4, 82152 Martinsried, München, Germany, **3** IRD, MARBEC, Université de Montpellier, Montpellier, France, **4** Département de Biologie Animale, Université d'Antananarivo, Antananarivo, 101, Madagascar, **5** College of Science, Engineering and Mathematics, Department of Natural Sciences, Bethune-Cookman University, 640 Dr. Mary McLeod Bethune Blvd., Daytona Beach, Florida, 32114, United States of America, **6** Zoologisches Institut, Universität Hamburg, Martin-Luther-King Platz 3, 20146, Hamburg, Germany

* axel@tsiboboka.de



OPEN ACCESS

Citation: Strauß A, Guilhaumon F, Randrianiaina RD, Wollenberg Valero KC, Vences M, Glos J (2016) Opposing Patterns of Seasonal Change in Functional and Phylogenetic Diversity of Tadpole Assemblages. *PLoS ONE* 11(3): e0151744. doi:10.1371/journal.pone.0151744

Editor: Carlos A Navas, University of Sao Paulo, BRAZIL

Received: July 21, 2015

Accepted: March 3, 2016

Published: March 25, 2016

Copyright: © 2016 Strauß et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Sequencing data are deposited in GenBank (for accession numbers see paper). Data used for statistical analyses are accessible at figshare (DOI [10.6084/m9.figshare.2059941](https://doi.org/10.6084/m9.figshare.2059941)).

Funding: Financial support was granted by the Volkswagen Foundation to MV and RDR, by the Deutsche Forschungsgemeinschaft (grant VE247/2-1) to MV, AS and JG, and grant DFG Glos 665/1-1 to JG, and by the Deutscher Akademischer Austauschdienst to RDR. The funders had no role in

Abstract

Assemblages that are exposed to recurring temporal environmental changes can show changes in their ecological properties. These can be expressed by differences in diversity and assembly rules. Both can be identified using two measures of diversity: functional (FD) and phylogenetic diversity (PD). Frog communities are understudied in this regard, especially during the tadpole life stage. We utilised tadpole assemblages from Madagascan rain-forest streams to test predictions of seasonal changes on diversity and assemblage composition and on diversity measures. From the warm-wet to the cool-dry season, species richness (SR) of tadpole assemblages decreased. Also FD and PD decreased, but FD less and PD more than expected by chance. During the dry season, tadpole assemblages were characterised by functional redundancy (among assemblages—with increasing SR), high FD (compared to a null model), and low PD (phylogenetic clustering; compared to a null model). Although mutually contradictory at first glance, these results indicate competition as tadpole community assembly driving force. This is true during the limiting cool-dry season but not during the more suitable warm-wet season. We thereby show that assembly rules can strongly depend on season, that comparing FD and PD can reveal such forces, that FD and PD are not interchangeable, and that conclusions on assembly rules based on FD alone are critical.

Introduction

The properties of species assemblages vary in time. On an ecological time scale, these variations might be anthropogenically caused, e.g., by drastic single events (such as fire), or recurring due to annual climatic changes [1–4]. Most obviously, changes in species assemblages are reflected by changes in species richness (SR). However, SR as measure of diversity is likely to miss

study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

relevant information because ecological traits and phylogenetic relatedness of species in an assemblage are not independent. Ecological traits of species (and therefore the functional diversity of assemblages, FD) can determine their ability to exploit resources, influence potential of coexistence, and may reflect the ecological impact of assemblages (i.e., ecosystem functions such as biomass production [5–7]). The degree of relatedness (summarised as the phylogenetic diversity of an assemblage, PD), is often used as alternative measure of assemblage structure and its influence on ecosystem processes [8]. The information needed to measure PD (i.e., a phylogenetic tree usually based on molecular data) is easier and less expensive to obtain, and often less ambiguous, than complex ecological data on species, and PD has been proposed as a proxy for FD [9, 10]. PD can be [11] but is not necessarily [12] a better determinant of diversity than FD. Whether PD indeed well reflects ecological functions in species assemblages has been questioned [13, 14] and exceptions have been found [11, 15, 16].

Such non-congruency between FD and PD can be caused by different factors. It is often assumed that closely related species show a high degree of morphological and ecological similarity [17]. PD covers more dimensions than the number of traits typically used to calculate FD [10]; however, it does not provide information on what these dimensions are [10]. Furthermore, PD is based on the assumption that relevant traits are conserved across phylogenies but there are cases of traits missing a phylogenetic signal (e.g., [18, 19]), or of a strong variation in phylogenetic signal of traits [20], usually due to extensive homoplasy. Since PD is easy to access and includes evolutionary history as well as unmeasured traits [21] it is still considered valuable if handled (and interpreted) with care [22] and can be a good predictor of ecosystem functioning [10].

The influence exerted by competition on FD and PD of species assemblages is usually tested by comparison of observed with randomly assembled "null assemblages"—consisting of members of the local species pool (e.g., [9, 15, 23–25]). If observed assemblages have higher FD (simply called "high FD") than the null assemblages, this is understood as an indication of competition as a relevant factor in shaping these assemblages, whereas low FD is an indicator of environmental filtering [23, 24]. The competition-relatedness hypothesis [26] transfers this interpretation to PD. However, the role of competition and environmental filtering in shaping assemblages has continuously been debated and alternative outcomes on FD and PD have been suggested [27]. Studies of PD combined with quantitative ecological traits will help interpreting relatedness in assemblages and environmental conditions shaping them.

If environmental conditions change, subsequent compositional changes of species assemblages may affect FD and PD in a different way than it would affect SR. Two different processes have been characterized by which such effects can lead to functional redundancy. *Intrinsic functional redundancy* (sensu [24]) occurs for instance if an assemblage, as a starting point, contains a high proportion of functionally similar species. In such a case, *random* decreases of species numbers will have a lower effect on FD than on SR [24]. *Extrinsic functional redundancy* instead originates by a process of *non-random* change in a species assemblage, e.g., when species disappearing from an assemblage are mostly functionally unique. These concepts can be extended to intrinsic or extrinsic phylogenetic redundancy leading to phylogenetic overdispersion or clustering regarding relatedness [28, 29]. Redundancy obviously is low in opposite situations, i.e., assemblages consist of high proportions of unique species (intrinsic), or species disappearing from the communities are mostly similar (extrinsic). Intrinsic redundancy has been observed in a range of disturbed ([19, 30] but see [23, 24]) and undisturbed [25] ecosystems across several taxa regarding species function (i.e., FD), and in urban plant assemblages [31] regarding relatedness (i.e., PD). Examples for extrinsic redundancy are observed in directly human influenced systems across several animal and plant taxa [30, 32, 33].

Tropical anuran assemblages represent an appropriate model to study seasonal changes and their impact on different measures of diversity as they are known to be remarkably rich but still

can be completely assessed and taxonomically handled [34]. Seasonal changes in SR, i.e., seasonal changes in frog (reproductive) activity, have been observed for adult amphibians [35–37], and to some extent for amphibian larval assemblages [38].

In this study we focus on the world’s most species rich stream tadpole assemblages in the rainforests of Madagascar [25] to evaluate seasonal patterns of SR, FD and PD. We expect SR to be lower in the dry season than in the wet season, following the pattern observed for adults [35]. Subsequently, we analyze according seasonal changes in FD and PD, and compare these changes against null models to identify high or low FD and PD, and functional redundancy and phylogenetic clustering or overdispersion, respectively.

Methods

Study sites

We conducted fieldwork in the wet season (January and February) and the dry season (July) of 2008. Study sites were located in one of the centres of amphibian species richness in Madagascar, Ranomafana National Park (RNP; 21°16'S; 47°25'E). RNP covers over 40,000 ha of rain forest from ca. 500 m up to ca. 1.500 m a.s.l. and harbours over 100 frog species [34]. Seasonality in this area is characterised by clear differences in precipitation and temperature (Fig 1). Activity patterns of adult frogs in Ranomafana are different between seasons [35].




Fig 1. Annual changes in precipitation and temperature. Annual changes in precipitation and temperature from the RNP region from January 2007 to December 2008. The total monthly precipitation is given in grey bars. The curves represent the mean monthly temperature retrieved from daily minimum-maximum-measurements of air temperature (red triangles) and the stream water temperature (blue circles; data from Fompohonina River). Data of rainfall and air temperature were provided by ValBio research station, water temperature was retrieved from iButton temperature loggers.

doi:10.1371/journal.pone.0151744.g001

During the austral winter dry season, the precipitation in RNP can reach values as low as <100 mm in some months, and temperatures are distinctly cooler than in summer. The wet season in the austral summer is generally characterised by high precipitation but the maximum can vary between months (see e.g. January 2007 vs. January 2008 and February 2007 vs. February 2008; see also Wollenberg et al. [39]).

Species sampling

We sampled tadpole assemblages by intensive capture in 30 m sections of 12 streams (in a mid-elevational area ranging from 910 m to 1.130 m a.s.l.). Sampling details are provided in Strauß et al. [40]. We kept tadpoles alive and carried them in water containers into the laboratory. They were euthanized by immersion in MS222 solution, and immediately sorted into series based on their morphology. From each series, we identified one specimen by DNA barcoding based on a fragment of the mitochondrial *16S* rRNA gene [41, 42]. All newly determined DNA sequences have been deposited in Genbank under accession numbers KF609548-KF611386. Sampling was carried out once in the wet and once in the dry season, applying the same sampling methods to the same stream sections. All analyses are based exclusively on tadpole data which are a reliable representation of the frog species actually breeding in a single stream; sightings of adults were not considered.

We tracked water temperature during the seasons by placing temperature loggers (Thermochron® iButton, Dallas Semiconductor) in the streams one year before this study. They conducted measurements every 255 min covering about one year (early 2007 to early 2008). Data on aerial temperature and rainfall were provided by the ValBio research station, Ranomafana (J. C. Razafimahaimodison).

Ethics Statement

No experiments were conducted using living animals.

All field research, including collection of specimens using dip nets, carriage to the lab in water containers, anaesthesia, and euthanasia of specimens, were approved by the Madagascan Ministère de l'Environnement, des Eaux et des Forêts (Direction des Eaux et Forêts, DEF) under the following permits: 300/06/MINENV.EF/SG/DGEF/DPB/SCBLF/RECH dated 22 December 2006; 003/08-MEEFT/SG/DGEF/DSAP/SSE dated 4 January 2008; 004/08-MEEFT/SG/DGEF/DSAP/SSE dated 4 January 2008; and MEEFT/SG/DGEF/DSAP/SSE dated 30 January 2008. Export of specimens was approved by the DEF under permit: 052N-EA02/MG08 dated 28 February 2008. Sampling for this study was conducted in January, February, and July 2008. No species protected by the Convention of the International Trade in Endangered Species CITES were concerned by this research. Voucher specimens were euthanized using approved methods (e.g., anaesthesia with MS222, followed by overdose of the same substance and 95% ethanol fixation). These are standard methods included in the permits above that do not require approval by an ethics committee. Killings were recorded and reported to Madagascan authorities as requested in the respective authorisations (included in the permits stated above).

Statistical analyses

We assessed species richness (SR) of tadpole assemblages in dry and wet season based on molecular identification of tadpoles sampled. We first confirmed the expectation that species richness (SR) of tadpole assemblages differs between wet and dry season by paired t-test (all software details are stated and referenced at the end of this section). Before testing whether changes in FD or PD are similar to changes in SR or whether species loss and turnover depend




Fig 2. Scheme illustrating process of setting null assemblages. Scheme illustrating process of setting null assemblages based on observations in the wet and the dry season. (A)—all species found are pooled. (B)—This species pool is randomly ordered a thousand times. (C)—null assemblages for wet season are retrieved: the first nine species if there have been nine species observed in the wet season. (D)—for the null assemblages of the dry season, first the remaining (here two) species that have not been used for the respective wet season assemblage are taken and (E)—filled up by randomly chosen species from the respective wet season assemblage (here four) until the original species richness of the dry season assemblage is reached.

doi:10.1371/journal.pone.0151744.g002

on the ecological function of species, we developed a null model. For this null model, we remodelled random assemblages ("null assemblages") of wet and dry season. Modelling was based on observed SR of assemblages in both seasons and therefore the observed changes in SR and the observed species turnover, as well as the observed species pool available for each stream. In detail, for each stream we pooled all species found in wet and/or dry season in this stream (= species pool available). For example, if nine species were found in the wet season and six in the dry season of which two were not sampled in the wet season, the species pool consisted of 9+2 (11) species (Fig 2A). We reordered them randomly 1000 times (Fig 2B). Out of these 1000 species pools, we each picked the first x species (with x being the number of species observed for this stream in the wet season; nine in our example) and thus achieved null assemblages for the wet season (Fig 2C). To achieve null assemblages for the dry season, we first picked each the remaining y species that were not already included in wet season assemblages (with y being the number of species that were found in the dry but not in the wet season in this stream; two in our example; Fig 2D) and then restocked with randomly chosen species from the respective random wet season assemblage (four in our example; Fig 2E) to reach the final number of species for the random dry season assemblage as observed (six in our example). This was repeatedly done for each stream based on the species that we observed in the stream. For each stream and season we first calculated FD of the tadpole null assemblages in the seasons and then the change in FD from the wet to the dry season, also based on the null assemblages. To calculate the relative change of diversity from the wet season ("wet") to the dry season ("dry") we developed an approach using $-1 \cdot (1 - \text{dry/wet})$ resulting in positive values indicating an increase and negative values indicating a decrease in the respective diversity measure.

FD was calculated following Petchey & Gaston [43, 44]. This method applies a three-step dendrogram based classification function in which a species trait matrix is used to calculate a pair-wise species distance matrix. This matrix is used to construct dendrograms of specific species assemblages. The total branch length needed to connect all species in the assemblage

represents the respective FD. Best distance measure (Gower's distance) and cluster method (UPGMA) were identified following Mouchet et al. [45]. Our trait matrix consisted of categorical and continuous morphological trait variables (oral disc, body shape) of ecological relevance (i.e. feeding, microhabitat choice; [46]) for all species (for a list of traits used, see [25]).

We calculated FD and the relative changes in FD for the observed assemblages, and for null assemblages (see above). We compared relative changes of observed and null assemblage FD using paired t-test. Accordingly, significant differences show that seasonal changes in FD are not random.

Non randomness can be caused by differences in redundancy patterns, or by high or low FD in one or both seasons [24]. We therefore analysed data of the wet and the dry season separately. We applied polynomial regression with observed FD as dependent and SR (linear term) and SR^2 (polynomial term) as independent variables to reveal possible patterns of functional redundancy in each of both seasons (i.e., if nonlinearity is found). To test whether the full polynomial model or a simplified (i.e., linear) model performs better we used stepwise deletion (polynomial term first) and compared the models based on Akaike's Information Criterion (AIC, [47, 48]) until the minimum adequate models were reached. Residuals were checked using diagnostic plots. We used Moran's I autocorrelation coefficient [49] to prove that there is no spatial autocorrelation of the study sites regarding SR, FD, and PD, and their respective changes between wet and dry season. In this process no functional redundancy is indicated if these analyses show only a linear relationship between FD and SR (i.e., if the linear term in the polynomial model happens to be significant and the polynomial term to be non significant). Functional redundancy is indicated if the slope of the relationship between FD and SR decreases with increasing SR (i.e., if the polynomial term in the model is significant), caused by a stronger overlap of ecological traits with increasing number of syntopic tadpole species. We compared observed FD data with the respective null model data using paired t-tests, separately for each season. This allowed identifying possible patterns of low or high FD in the tadpole assemblages if differences were significant. All t-test were two-sided Welch t-tests.

We conducted the same procedure to analyse patterns of phylogenetic diversity (PD) and its respective changes from wet to dry season. Our analyses are based on a time-calibrated phylogeny of mantellid species [50] as all tadpoles sampled in the streams belonged to this family [40]. We pruned from this tree all taxa not represented in our Ranomafana tadpole sampling using TreeEdit version v1.0a10. This reduced the number of species by retaining branch lengths. For a few so far undescribed species, genetic data were insufficient to include them in the tree. In these cases, other species that are known to be in the same clade (according to molecular data of Vieites et al. [34]) were used as replacement for the purpose of PD calculation. Similar to FD calculations [43, 44, 51], we extracted branch lengths from the tree for the assemblages [52] by using the function `treedive` included in the R package `vegan`. The sum of branch lengths needed to connect all species of an assemblage represents the assemblage's PD.

All analyses were run using the statistical software R 2.15.1 [53]. Packages used for FD calculations include `car` R package version 2.0–18 [54], `gtools` R package version 2.7.0. [55], `cluster` R package version 1.14.2 [56] and `clue` R package version 0.3–45 [57, 58]. For PD calculations we used the R packages `ape` version 3.0–5 [59, 60] and `vegan` version 2.0–4 [61]. Moran's I was also calculated using the package `ape`.

Results

Summarizing data over wet and dry seasons, we found tadpoles of a total of 31 species in all twelve stream sections (= assemblages). In the wet season we found five to 15 species, in the dry season two to twelve species per assemblage. Mean SR in the dry season was about 27% lower than in the wet season (paired t-test, $t = 3.44$, $df = 11$, $p = 0.006$). Beside this decrease in




Fig 3. Change from the wet to the dry season: Deviation of observed loss of FD and PD from the null models. The dashed lines represent the null models for the change from the wet to the dry season, the box-whisker-plots show the observed deviations from the null models. Values below the dashed line show a smaller loss, values above the line show a higher loss of FD and PD than predicted by the null models. FD of tadpoles decreases significantly less than predicted whereas PD decreases significantly more than predicted.

doi:10.1371/journal.pone.0151744.g003

SR (i.e., probably due to reduced reproductive activity), a species turnover from wet to dry season was observed in eight streams, i.e., an assemblage in the dry season harboured on average one or more species that were not present in the assemblage of the same stream during the wet season. However, no species was exclusively found in the dry season.

Functional diversity

As observed for SR, FD decreased from the wet to the dry season. However, the observed relative loss of FD (8%) differs from the loss predicted by the null models (17%) based on the observed loss in SR (paired t-test, $t = -2.34$, $df = 11$, $p = 0.04$; Fig 3). This difference suggests




Fig 4. Comparing null models and observed FD from tadpole assemblages from RNP. Given are observed (dark grey filled circles, continuous regression line) and null model FD (light grey circles with dark margins, dashed line) along the observed SR gradient for the wet (A) and the dry season (B). Each symbol represents one tadpole assemblage. (C) and (D) show the differences between null model (dashed line) and observed values of FD (grey circles) of the wet and dry season, respectively. Values above or below the line show observed values being higher (high FD) or lower (low FD) than predicted, respectively. As graphical summary, the respective box-whisker plots are provided next to the scatter plot with outliers indicated as asterisks. Differences were not significant in the wet season; in the dry season assemblages show significantly higher FD than expected.

doi:10.1371/journal.pone.0151744.g004

that species loss and/or species turnover from the wet to the dry season is non random with respect to the species traits, and it is explained by the structure of the tadpole assemblages of the dry season. Firstly, tadpole assemblages do not show functional redundancy in the wet season, as both observed and null model FD show a linear relationship with SR (Fig 4A; linear regressions; null model: $R^2 = 0.78$, $F_{1,10} = 35.8$, $p_{\text{model}} < 0.001$, $p_{\text{intercept}} = 0.012$, $p_{\text{SR}} < 0.001$; observed: $R^2 = 0.63$, $F_{1,10} = 17$, $p_{\text{model}} = 0.002$, $p_{\text{intercept}} = 0.015$, $p_{\text{SR}} = 0.002$). Also, there is no

difference between observed and null model FD (Fig 4C; paired t-test, $t = -1.43$, $df = 11$, $p = 0.18$). In the dry season, however, the tadpole assemblages are characterised by functional redundancy as indicated by curvilinear relationship of FD with SR (Fig 4B; polynomial regressions; null model: $R^2 = 0.96$, $F_{2,9} = 114$, $p_{\text{model}} < 0.001$, $p_{\text{SR}} < 0.001$, $p_{\text{SR}^2} < 0.001$, observed: $R^2 = 0.91$, $F_{2,9} = 45.8$, $p_{\text{model}} < 0.001$, $p_{\text{SR}} < 0.001$, $p_{\text{SR}^2} < 0.005$). Furthermore, these assemblages show high functional diversity, i.e. observed FD values are higher than predicted by the null model (Fig 4D; paired t-test; $t = 2.99$, $df = 11$, $p = 0.012$).

In a nutshell, the loss and/or turnover of species in tadpole assemblages in RNP from the wet to the dry season is non random with respect to species traits with patterns of high FD (compared to the null model) and higher functional redundancy (with increasing SR among sites) in the dry season, whereas in the wet season FD does not provide any more or different information than SR.

Phylogenetic diversity

The observed relative loss of PD (28%) of tadpole assemblages was stronger than predicted (23%) by null model assemblages (Fig 3; paired t-test; $t = 3.08$, $df = 11$, $p = 0.011$). To identify the reason for this deviation of the observed data from the null model, we focused on PD in the wet and the dry season separately. In the wet season, both observed and null model PD show a linear relationship with SR indicating no phylogenetic redundancy (Fig 5A; linear regressions; null model: $R^2 = 0.96$, $F_{1,10} = 267.7$, $p_{\text{model}} < 0.001$, $p_{\text{intercept}} = 0.022$, $p_{\text{SR}} < 0.001$; observed: $R^2 = 0.93$, $F_{1,10} = 123.8$, $p_{\text{model}} < 0.001$, $p_{\text{intercept}} = 0.008$, $p_{\text{SR}} < 0.001$). The observed PD of tadpole assemblages in the wet season does not differ from the predicted values (Fig 5C; paired t-test; $t = -0.88$, $df = 11$, $p = 0.4$). Therefore, there is neither phylogenetic clustering nor overdispersion in tadpole assemblages in the wet season. In the dry season, null assemblages predict that PD will be highly related to SR with a trend to curvilinearity (Fig 5B; polynomial regression; $R^2 = 0.99$, $F_{2,9} = 378.2$, $p_{\text{model}} < 0.001$, $p_{\text{intercept}} = 0.82$, $p_{\text{SR}} < 0.001$, $p_{\text{SR}^2} = 0.06$). Indeed, a similar relationship of SR and PD is shown for the observed assemblages. However, as our data set includes highly influential data points to be considered in regression analysis we consider our model conservative and do not refer it to the polynomial model (Fig 5B; simple linear regression; $R^2 = 0.96$, $F_{1,10} = 227.5$, $p_{\text{model}} < 0.001$, $p_{\text{intercept}} = 0.63$, $p_{\text{SR}} < 0.001$). More of relevance, however, is that PD of tadpole assemblages in the dry season is significantly lower than predicted by null models (Fig 5D; paired t-test, $t = -0.64$, $df = 11$, $p = 0.004$). Therefore, in the dry season tadpole assemblages show phylogenetic clustering, i.e., they are assembled by species that are more closely related to each other than expected by chance.

Summarising the results concerning PD, species loss and/or turnover from the wet to the dry season are non-random with respect to the degree of relatedness of the species in an assemblage. This non-randomness is expressed in phylogenetic clustering in the tadpole assemblages in the dry season (Fig 5B and 5D).

There was no spatial autocorrelation of the diversity measures in any of the seasons (all p range from 0.22 to 0.93).

Discussion

Seasonality as observed in the tropical rainforest of Madagascar causes changes in SR, FD, and PD of tadpole assemblages within stream habitats. All of these measures of diversity were observed to decrease in the dry season compared to the wet season. However, FD and PD do neither change congruently to SR nor to each other. Furthermore, changes as observed differ significantly from predictions by null models. The loss of FD from the wet to the dry season was smaller than predicted, while PD changed more strongly than predicted. Whereas in the

wet season all three measures (SR, FD, PD) provide similar information on tadpole assemblages, strong differences are found in the dry season. In the dry season the assemblages are characterised by functional redundancy (among assemblages—with increasing SR), high FD (compared to a null model), and low PD (phylogenetic clustering; compared to a null model). This suggests that both tadpole functional traits and tadpole relatedness are important determinants in the change of assemblage composition following seasonal climatic changes.




Fig 5. Comparing null models and observed PD from tadpole assemblages from RNP. Given are observed (dark grey filled circles, continuous regression line) and null model PD (light grey circles with dark margins, dashed line) along the observed SR gradient for the wet (A) and the dry season (B). Each symbol represents one tadpole assemblage. (C) and (D) show the differences between null model (dashed line) and the observed values of PD (grey circles) of the wet and dry season, respectively. Values above or below the line show observed values being higher (phylogenetic overdispersion) or lower (phylogenetic clustering) than predicted by null models, respectively. As graphical summary, the respective box-whisker plots are provided next to the scatter plot with outliers indicated as asterisks. Differences were not significant in the wet season; in the dry season, assemblages show significant phylogenetic clustering.

doi:10.1371/journal.pone.0151744.g005

Species richness decreases from wet to dry season

From the warm and wet to the climatically less suitable, cooler dry season [38] species richness in tadpole assemblages at Ranomafana National Park (RNP) decreased significantly. Tropical seasonality has been shown to affect frog activity and accordingly tadpole SR at ponds [36, 38] but not always at streams [35, 62], indicating often more stable conditions in streams than in ponds [63]. In Madagascar, frogs in the dry season are much less active [37] and might also be more susceptible to infection with pathogens such as the amphibian chytrid fungus [64]. Beside the loss of SR, assemblages can change in composition by species turnover [32], as it is the case in the observed RNP tadpole assemblages. This is often due to species that are seasonal specialists [65]. However, no species in this study can be classified as being strictly specialised on one season. We assume that in most RNP stream breeding species, reproduction takes part throughout the year, at least occasionally. Also, some larvae observed in the dry season (i.e., species with long larval development) might derive from clutches deposited in the wet season, blurring possible patterns of specialists reproducing in one of the seasons only.

Functional diversity decreases less than expected from wet to dry season

As expected from the observed decrease of SR from wet to dry season, tadpole assemblages show a decrease of FD. This decrease of FD is lower than predicted by a null model, and accordingly assemblages are presumed to have a relatively higher FD in the dry season. This study therefore shows that a seasonal loss of species from assemblages is trait-related, as it has also been observed in tropical beetles [66], and mainly such species remain in the assemblages that are functionally different [24]. This is an indication that interspecific competition might play a role in shaping these assemblages ([24] but see [discussion](#) of PD below) in the dry season, but less so in the wet season. So far, interspecific competition was assumed to not being important at all in shaping stream tadpole assemblages [25, 63, 67]. These studies, however, were all conducted during the wet season, and agree with the wet season data presented herein. Evidence for competition in tadpole assemblages came so far only from temporary ponds [68–71]. In the dry season in Malagasy rainforest streams reduced food availability, e.g., due to slower algal growth at low temperatures, or lower influx of nutrients, may induce or increase competition among tadpole species despite lower SR in this time, i.e., less potentially competing species.

Additionally, these assemblages show functional redundancy (when comparing FD of assemblages within the dry season): the more species occur in an assemblage, the more they overlap in their ecological functions. We here find this functional redundancy pattern in the dry season assemblages, and it previously has been observed also in the wet season [25]. The lack of redundancy in the wet season assemblages analysed herein might be due to a lower range of SR compared to the previous study [25] which included 29 assemblages of partly much higher SR, illustrating that the probability of detecting redundancy depends on the level of SR [25, 72]. We interpret the missing signal of wet season functional redundancy in the smaller data set analysed herein as an indication of a weaker redundancy effect as compared to the dry season assemblages.

Phylogenetic diversity decreases stronger than expected from wet to dry season

Phylogenetic diversity of stream tadpole assemblages also decreases with a loss of SR. Contrary to the observed pattern for FD, the loss of PD is stronger than predicted by a null model and

assemblages show phylogenetic clustering [29] in the dry season (but not in the wet season). Hence, those species that are less closely related to the remaining species disappear from the assemblages during the shift from the wet to the dry season. Such phylogenetic clustering has also been observed in several bacterial [28], insect [73] and plant assemblages [74, 75].

Following the arguments of the competition-relatedness hypothesis [17, 26] the species within the dry season assemblages are more closely related than expected and therefore they should also be more functionally similar. This would indicate an influence of environmental filtering in species assembly [9, 28]. However, the inverse pattern observed for FD (i.e., lower decrease than expected of FD vs. higher decrease than expected of PD) is contradictory to this hypothesis. High FD in Malagasy stream tadpoles actually indicates that competition plays a more significant role than environmental filtering, and this is not necessarily in conflict with the PD data. In fact, the observed pattern of phylogenetic clustering (low PD) might arise under competition if certain clades are stronger competitors than others irrespective of other traits [27]. Also, the niche conservatism hypothesis [76] might provide an alternative explanation for the phylogenetic clustering in the dry season. Accordingly, the phylogenetically related species occurring during the dry season might have similar ecological traits allowing them to cope with the harsher conditions in the dry season where other species not having these traits are excluded. During evolutionary time, these phylogenetically related species might have conserved their niches more similar to each other than the other distant related species leading to a closely related dry season assembly, i.e. to phylogenetic clustering.

An additional point needs to be considered: life history variables such as larval developmental time and breeding phenology might be phylogenetically conserved traits, and accordingly those clades, e.g., with quick larval development and breeding at the onset of the rainy season would have completed metamorphosis and left the streams in the dry season while the clades with slow larval development would still remain in the streams throughout the dry season. The remaining, accordingly more closely related species then would face high competition. Obtaining more reliable natural and life history data for Madagascan frogs will be paramount to better understand such assembly processes of tadpole communities at specific points in time.

Opposing patterns of functional and phylogenetic diversity are not necessarily contradictory

The morphological and therefore functional similarity of members of RNP tadpole assemblages does not fully reflect their relatedness. This is evident during the dry season, when assemblages have lower SR, FD, and PD than in the wet season, and tadpoles are more closely related but functionally more different than expected by chance. This opposing pattern of FD and PD is mutually contradictory only at first glance, and several factors might explain why FD, PD and SR are not necessarily interchangeable.

Firstly, FD and PD of assemblages may be partially decoupled from SR, e.g., if a decrease in SR of assemblages is associated with species turnover [32, 77], with species being replaced by others that are less closely related but functionally more similar than the remaining species in the assemblage, and vice versa. Although Madagascan tadpole assemblages do show species turnover and tadpoles may have been replaced by species closely related to remaining species but showing a different morphology, we see a clear link of both FD and PD to SR, and the loss of SR had clearly a stronger effect than the species turnover. In fact, Flynn et al. [10] argues that both FD and PD may be linked to SR, but not directly to each other. However, there are several studied systems with a clear phylogenetic signal in functional traits (e.g., [20, 78] and others cited above). Additionally, if genetic markers are linked to a function (e.g., protein coding genes in microorganisms) PD and FD may be even more strongly correlated to each other

than to SR [79]. In Madagascan tadpoles, some closely related groups indeed show a broader range of traits than others which is not necessarily linked to their species richness (e.g., [80–82]). There are several accounts of homoplasy in morpho-functional traits in Madagascan tadpoles [83] although in general, tadpole morphology largely fits phylogeny ([84] and references cited above). As the mentioned exceptions are randomly distributed between the taxonomic groups included in this study, they may not necessarily have influenced the results.

Secondly, a mismatch of FD and PD may appear if some local assemblages comprise species with comparable origin while others comprise species with different origin [15, 16, 85]. The Madagascan anuran fauna originated by five independent colonization events [86] but all species in our study belong to one of these clades, i.e., the Madagascar-Comoroan endemic Mantellidae [40]. The species in the dry and wet season assemblages all underwent comparable evolutionary histories in the eastern rainforest [50], and a bias caused by different origins is therefore unlikely.

Thirdly, it is evident that FD is based on a defined set of ecological traits that may only be a part of the traits covered by phylogeny. The additional traits included in PD, however, may be of relevance, or not, for species assembly. If too many ecologically non-relevant traits are covered by PD they might, as they are included in the analysis as describing factors, mask the information of the relevant traits [21]. Additionally, some functional traits of relevance for species assembly may lack a phylogenetic signal [85, 87] or vary in this signal [20]. Also, PD may include one or a few key traits that simply have not been considered for FD. If so, environmental filtering (or competition) may act and select on these traits and will cause phylogenetic clustering; the remaining species are rather similar regarding these traits (covered by PD but not FD) but also rather different regarding the remaining traits included in FD analysis. If so, a pattern as observed in our study may appear.

Fourthly, Swenson [21] related the applicability of PD to the phylogenetic scale. Which scale is appropriate surely depends on the system studied. In terms of their general morphology and ecology, tadpoles are rather similar to each other (compared to, e.g., plant assemblages with grasses, herbs, and legumes) and adaptive evolution may play a strong role in our study system. However, most tadpoles can easily be assigned to a specific phylogenetic group just by their general appearance indicating an appropriate phylogenetic scale in this study. Additionally, we do see a pattern in PD analysis indicating the relevance of phylogeny. The question of scale is of course also of relevance for FD. This study and data of Strauß et al. [25] do show FD patterns and therefore confirm the scale also appropriate for FD studies.

Conclusions

The harshness of environmental conditions may influence whether traits cluster (more stressful conditions) or overdisperses (less stressful conditions) [19]. Our results illustrate the necessity of observing the diversity of assemblages at different environmental conditions, i.e., in different seasons. Assemblages without seasonal specialists and persisting at the same site during the year vary strongly in their functional and phylogenetic diversity, and obviously their assembly rules. In general, FD and PD provide more information than SR without being interchangeable. Both measures contribute differently to describe diversity. The appropriate selection of traits for FD on the one hand and the missing selection of traits in PD on the other hand remains a major challenge in studies based on these measures. However, if FD is used to indicate assembly rules, the underlying traits must be of relevance (i.e., for competition and environmental filtering) and allow interpretation, respectively. Beside the mentioned advantages of using PD in diversity studies, however, the interpretation of observed patterns is diverse and should be related to FD. Our data, however, are unambiguous in suggesting that

the loss of anuran larvae from the more suitable (i.e., warm-wet) season to the harsher (i.e., cool-dry) season is not random. Although there are no data supporting whether differences in assemblage composition are either based on the duration of larval development or phenological differences in reproductive activity, our results indicate that competition influences the composition of these assemblages at least periodically.

Acknowledgments

We thank S. Ndriantsoa, E. Rajeriarison, T. Rajoafiarison, H. Rasolonjatovo, D. Razafindrabe, and J. Solo for their help during the fieldwork, ValBio, MICET/ICTE, and Madagascar National Parks for logistic support, and G. Keunecke, M. Kondermann, and E. Saxinger for their great work in the lab. This study was carried out in the framework of a cooperation accord between the Département de Biologie Animale of the University of Antananarivo, Madagascar and the Technische Universität Braunschweig, Germany. The Malagasy authorities kindly issued research and export permits.

Author Contributions

Conceived and designed the experiments: AS JG MV. Performed the experiments: AS RDR JG. Analyzed the data: AS KCWV FG. Contributed reagents/materials/analysis tools: FG. Wrote the paper: AS JG MV FG RDR KCWV. Performed molecular or morphological labwork: RDR KCWV.

References

1. Rosenzweig ML. Species diversity in space and time: Cambridge University Press; 1995. 436 p.
2. Huston MA. Biological Diversity—The coexistence of species on changing landscapes: Cambridge University Press; 1995. 681 p.
3. Moretti M, De Bello F, Roberts SPM, Potts SG. Taxonomical vs. functional responses of bee communities to fire in two contrasting climatic regions. *J Anim Ecol.* 2009; 78(1):98–108. doi: [10.1111/j.1365-2656.2008.01462.x](https://doi.org/10.1111/j.1365-2656.2008.01462.x) PMID: [18705629](https://pubmed.ncbi.nlm.nih.gov/18705629/)
4. Dinnage R. Disturbance alters the phylogenetic composition and structure of plant communities in an old field system. *PLoS ONE.* 2009; 4(9).
5. Joshi J, Matthies D, Schmid B. Root hemiparasites and plant diversity in experimental grassland communities. *J Ecol.* 2000; 88(4):634–44.
6. Scherer-Lorenzen M. Functional diversity affects decomposition processes in experimental grasslands. *Funct Ecol.* 2008; 22(3):547–55.
7. Hoehn P, Tscharntke T, Tylianakis JM, Steffan-Dewenter I. Functional group diversity of bee pollinators increases crop yield. *Proceedings of the Royal Society B: Biological Sciences.* 2008; 275(1648):2283–91. doi: [10.1098/rspb.2008.0405](https://doi.org/10.1098/rspb.2008.0405) PMID: [18595841](https://pubmed.ncbi.nlm.nih.gov/18595841/)
8. Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW. The merging of community ecology and phylogenetic biology. *Ecol Lett.* 2009; 12(7):693–715. doi: [10.1111/j.1461-0248.2009.01314.x](https://doi.org/10.1111/j.1461-0248.2009.01314.x) PMID: [19473217](https://pubmed.ncbi.nlm.nih.gov/19473217/)
9. Webb CO, Ackerly DD, McPeck Ma, Donoghue MJ. Phylogenies and community ecology. *Annu Rev Ecol Syst.* 2002; 33:475–505. doi: [10.1146/annurev.ecolsys.33.010802.150448](https://doi.org/10.1146/annurev.ecolsys.33.010802.150448)
10. Flynn DFB, Mirotnick N, Jain M, Palmer MI, Naeem S. Functional and phylogenetic diversity as predictors of biodiversity—ecosystem-function relationships. *Ecology.* 2011; 92:1573–81. PMID: [21905424](https://pubmed.ncbi.nlm.nih.gov/21905424/).
11. Cadotte MW, Cavender-Bares J, Tilman D, Oakley TH. Using phylogenetic, functional and trait diversity to understand patterns of plant community productivity. *PLoS ONE.* 2009; 4(5).
12. Cadotte MW, Cardinale BJ, Oakley TH. Evolutionary history and the effect of biodiversity on plant productivity. *Proc Natl Acad Sci USA.* 2008; 105(44):17012–7. doi: [10.1073/pnas.0805962105](https://doi.org/10.1073/pnas.0805962105) PMID: [18971334](https://pubmed.ncbi.nlm.nih.gov/18971334/)
13. Mouquet N, Devictor V, Meynard CN, Munoz F, Bersier L-F, Chave J, et al. Ecophylogenetics: advances and perspectives. *Biol Rev Camb Philos Soc.* 2012; 87:769–85. doi: [10.1111/j.1469-185X.2012.00224.x](https://doi.org/10.1111/j.1469-185X.2012.00224.x) PMID: [22432924](https://pubmed.ncbi.nlm.nih.gov/22432924/).

14. Winter M, Devictor V, Schweiger O. Phylogenetic diversity and nature conservation: where are we? *Trends Ecol Evol.* 2013; 28:199–204. doi: [10.1016/j.tree.2012.10.015](https://doi.org/10.1016/j.tree.2012.10.015) PMID: [23218499](https://pubmed.ncbi.nlm.nih.gov/23218499/).
15. Prinzing A, Reiffers R, Braakhekke WG, Hennekens SM, Tackenberg O, Ozinga WA, et al. Less lineages—more trait variation: phylogenetically clustered plant communities are functionally more diverse. *Ecol Lett.* 2008; 11(8):809–19. doi: [10.1111/j.1461-0248.2008.01189.x](https://doi.org/10.1111/j.1461-0248.2008.01189.x) PMID: [18445034](https://pubmed.ncbi.nlm.nih.gov/18445034/)
16. Devictor V, Moullot D, Meynard C, Jiguet F, Thuiller W, Mouquet N. Spatial mismatch and congruence between taxonomic, phylogenetic and functional diversity: the need for integrative conservation strategies in a changing world. *Ecol Lett.* 2010; 13:1030–40. doi: [10.1111/j.1461-0248.2010.01493.x](https://doi.org/10.1111/j.1461-0248.2010.01493.x) PMID: [20545736](https://pubmed.ncbi.nlm.nih.gov/20545736/).
17. Harvey PH, Pagel MD. *The Comparative Method in Evolutionary Biology.* Oxford, UK: Oxford University Press; 1991. 248 p.
18. Fritz SA, Purvis A. Phylogenetic diversity does not capture body size variation at risk in the world's mammals. *Proc R Soc B.* 2010; 277:2435–41. doi: [10.1098/rspb.2010.0030](https://doi.org/10.1098/rspb.2010.0030) PMID: [20375051](https://pubmed.ncbi.nlm.nih.gov/20375051/)
19. Kluge J, Kessler M. Phylogenetic diversity, trait diversity and niches: species assembly of ferns along a tropical elevational gradient. *J Biogeogr.* 2011; 38:394–405. doi: [10.1111/j.1365-2699.2010.02433.x](https://doi.org/10.1111/j.1365-2699.2010.02433.x)
20. Best RJ, Stachowicz JJ. Phylogeny as a proxy for ecology in seagrass amphipods: which traits are most conserved? *PloS ONE.* 2013; 8:e57550. doi: [10.1371/journal.pone.0057550](https://doi.org/10.1371/journal.pone.0057550) PMID: [23505433](https://pubmed.ncbi.nlm.nih.gov/23505433/).
21. Swenson NG. The assembly of tropical tree communities—the advances and shortcomings of phylogenetic and functional trait analyses. *Ecography.* 2013; 36:264–76. doi: [10.1111/j.1600-0587.2012.00121.x](https://doi.org/10.1111/j.1600-0587.2012.00121.x)
22. Srivastava DS, Cadotte MW, MacDonald AAM, Marushia RG, Mirotnich N. Phylogenetic diversity and the functioning of ecosystems. *Ecol Lett.* 2012; 15:637–48. doi: [10.1111/j.1461-0248.2012.01795.x](https://doi.org/10.1111/j.1461-0248.2012.01795.x) PMID: [22583836](https://pubmed.ncbi.nlm.nih.gov/22583836/).
23. Micheli F, Halpern BS. Low functional redundancy in coastal marine assemblages. *Ecol Lett.* 2005; 8(4):391–400.
24. Petchey OL, Evans KL, Fishburn IS, Gaston KJ. Low functional diversity and no redundancy in British avian assemblages. *J Anim Ecol.* 2007; 76(5):977–85. PMID: [17714276](https://pubmed.ncbi.nlm.nih.gov/17714276/)
25. Strauß A, Reeve E, Randrianiaina R-D, Vences M, Glos J. The world's richest tadpole communities show functional redundancy and low functional diversity: ecological data on Madagascar's stream-dwelling amphibian larvae. *BMC Ecol.* 2010; 10(1):12. doi: [10.1186/1472-6785-10-12](https://doi.org/10.1186/1472-6785-10-12)
26. Cahill JF Jr., Kembel SW, Lamb EG, Keddy PA. Does phylogenetic relatedness influence the strength of competition among vascular plants? *Perspect Plant Ecol Evol Syst.* 2008; 10:41–50. doi: [10.1016/j.ppees.2007.10.001](https://doi.org/10.1016/j.ppees.2007.10.001)
27. Mayfield MM, Levine JM. Opposing effects of competitive exclusion on the phylogenetic structure of communities. *Ecol Lett.* 2010; 13:1085–93. doi: [10.1111/j.1461-0248.2010.01509.x](https://doi.org/10.1111/j.1461-0248.2010.01509.x) PMID: [20576030](https://pubmed.ncbi.nlm.nih.gov/20576030/).
28. Horner-Devine MC, Bohannon BJM. Phylogenetic clustering and overdispersion in bacterial communities. *Ecology.* 2006; 87(sp7):100–8. doi: [10.1890/0012-9658\(2006\)87\[100:PCAOIB\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[100:PCAOIB]2.0.CO;2)
29. Emerson BC, Gillespie RG. Phylogenetic analysis of community assembly and structure over space and time *Trends Ecol Evol.* 2008; 23(11):619–30. doi: [10.1016/j.tree.2008.07.005](https://doi.org/10.1016/j.tree.2008.07.005) PMID: [18823678](https://pubmed.ncbi.nlm.nih.gov/18823678/)
30. Flynn DFB, Gogol-Prokurat M, Nogeire T, Molinari N, Richers BT, Lin BB, et al. Loss of functional diversity under land use intensification across multiple taxa. *Ecol Lett.* 2009; 12(1):22–33. doi: [10.1111/j.1461-0248.2008.01255.x](https://doi.org/10.1111/j.1461-0248.2008.01255.x) PMID: [19087109](https://pubmed.ncbi.nlm.nih.gov/19087109/)
31. Knapp S, Kühn I, Schweiger O, Klotz S. Challenging urban species diversity: contrasting phylogenetic patterns across plant functional groups in Germany. *Ecol Lett.* 2008; 11(10):1054–64. doi: [10.1111/j.1461-0248.2008.01217.x](https://doi.org/10.1111/j.1461-0248.2008.01217.x) PMID: [18616547](https://pubmed.ncbi.nlm.nih.gov/18616547/)
32. Ernst R, Linsenmair KE, Rödel MO. Diversity erosion beyond the species level: dramatic loss of functional diversity after selective logging in two tropical amphibian communities. *Biol Conserv.* 2006; 133(2):143–55.
33. Sakai AK, Allendorf FW, Holt JS, Lodge DM, Molofsky J, With KA, et al. The population biology of invasive species. *Annu Rev Ecol Syst.* 2001; 32:305–32.
34. Vieites DR, Wollenberg KC, Andreone F, Köhler J, Glaw F, Vences M. Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory. *Proc Natl Acad Sci USA.* 2009; 106:8267–72. doi: [10.1073/pnas.0810821106](https://doi.org/10.1073/pnas.0810821106) PMID: [19416818](https://pubmed.ncbi.nlm.nih.gov/19416818/)
35. Andreone F. Seasonal variations of the amphibian communities in two rainforests of Madagascar. In: Lourenço WR, editor. *Biogéographie de Madagascar.* Paris: Editions de l'ORSTOM; 1996. p. 397–402.
36. Kopp K, Eterovick PC. Factors influencing spatial and temporal structure of frog assemblages at ponds in southeastern Brazil. *J Nat Hist.* 2006; 40(29–31):1813–30.

37. Heinemann J, Rodríguez A, Segev O, Edmonds D, Dolch R, Vences M. Year-round activity patterns in a hyperdiverse community of rainforest amphibians in Madagascar. *J Nat Hist*. 2015;1–19. doi: [10.1080/00222933.2015.1009513](https://doi.org/10.1080/00222933.2015.1009513)
38. Both C, Sole M, dos Santos TG, Cechin SZ. The role of spatial and temporal descriptors for neotropical tadpole communities in southern Brazil. *Hydrobiologia*. 2009;1–14.
39. Wollenberg KC, Vieites DR, Vences M. Annual temperature data for two Malagasy sites of high anuran diversity. In: Andreone F, editor. *A Conservation Strategy for the Amphibians of Madagascar—Monografie XLV*. Torino: Museo Regionale di Scienze Naturali; 2008. p. 85–94.
40. Strauß A, Randrianiaina RD, Vences M, Glos J. Species distribution and assembly patterns of frog larvae in rainforest streams of Madagascar. *Hydrobiologia*. 2013; 702(1):27–43.
41. Vences M, Thomas M, van der Meijden A, Chiari Y, Vieites DR. Comparative performance of the 16S rRNA gene in DNA barcoding of amphibians. *Frontiers in Zoology*. 2005; 2:5. PMID: [15771783](https://pubmed.ncbi.nlm.nih.gov/15771783/)
42. Vences M, Chiari Y, Teschke M, Randrianiaina R-D, Raharivololoniaina L, Bora P, et al. Which frogs are out there? A preliminary evaluation of survey techniques and identification reliability of Malagasy amphibians. In: Andreone F, editor. *A Conservation Strategy for the Amphibians of Madagascar—Monografie XLV*. Torino: Museo Regionale di Scienze Naturali; 2008. p. 233–52.
43. Petchey OL, Gaston KJ. Functional diversity (FD), species richness and community composition. *Ecol Lett*. 2002; 5(3):402–11.
44. Petchey OL, Gaston KJ. Functional diversity: back to basics and looking forward. *Ecol Lett*. 2006; 9(6):741–58. PMID: [16706917](https://pubmed.ncbi.nlm.nih.gov/16706917/)
45. Mouchet M, Guilhaumon F, Villéger S, Mason NWH, Tomasini JA, Moullot D. Towards a consensus for calculating dendrogram-based functional diversity indices. *Oikos*. 2008; 117(5):794–800.
46. Alford RA. Ecology: Resource use, Competition, and Predation. In: McDiarmid RW, Altig R, editors. *Tadpoles: The Biology of Anuran Larvae*. Chicago and London: Chicago University Press; 1999. p. 240–78.
47. Burnham KP, Anderson DR. *Model selection and inference: a practical information-theoretic approach*. 1st ed. Heidelberg: Springer Verlag; 1998. 353 p.
48. Johnson JB, Omland KS. Model selection in ecology and evolution. *Trends Ecol Evol*. 2004; 19(2):101–8. PMID: [16701236](https://pubmed.ncbi.nlm.nih.gov/16701236/)
49. Moran PAP. Notes on continuous stochastic phenomena. *Biometrika*. 1950; 37:17–23. PMID: [15420245](https://pubmed.ncbi.nlm.nih.gov/15420245/)
50. Wollenberg KC, Vieites DR, Glaw F, Vences M. Speciation in little: the role of range and body size in the diversification of Malagasy mantellid frogs. *BMC Evol Biol*. 2011; 11:217. doi: [10.1186/1471-2148-11-217](https://doi.org/10.1186/1471-2148-11-217) PMID: [21777445](https://pubmed.ncbi.nlm.nih.gov/21777445/); PubMed Central PMCID: PMC3199771.
51. Podani J, Schmera D. On dendrogram-based measures of functional diversity. *Oikos*. 2006; 115(1):179–85.
52. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv*. 1992; 61:1–10. doi: [10.1016/0006-3207\(92\)91201-3](https://doi.org/10.1016/0006-3207(92)91201-3)
53. R Core Team. *R: A language and environment for statistical computing*. 2.15.1 ed. Vienna, Austria: R Foundation for Statistical Computing; 2012.
54. Fox J, Weisberg S. *An R Companion to Applied Regression*. 2nd ed. Thousand Oaks, CA: Sage; 2011.
55. Warnes GR. *gtools: Various R programming tools*. R package version 2.7.0. 2012.
56. Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. *cluster: Cluster Analysis Basics and Extensions*. R package version 1.14.2. 2012.
57. Hornik K. A CLUE for CLUster Ensembles. *Journal of Statistical Software*. 2005; 14(12).
58. Hornik K. *clue: Cluster ensembles*. R package version 0.3–45. 2012.
59. Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R language. R package version 3.0–5. *Bioinformatics*. 2004; 20:289–90. PMID: [14734327](https://pubmed.ncbi.nlm.nih.gov/14734327/)
60. Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R language. *Bioinformatics*. 2004; 20:289–90. PMID: [14734327](https://pubmed.ncbi.nlm.nih.gov/14734327/)
61. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchi PR, O'Hara RB, et al. *vegan: Community Ecology Package*. R package version 2.0–4. 2012.
62. Afonso LG, Eterovick PC. Spatial and temporal distribution of breeding anurans in streams in south-eastern Brazil. *J Nat Hist*. 2007; 41(13–16):949–63.
63. Eterovick PC, Barata IM. Distribution of tadpoles within and among Brazilian streams: the influence of predators, habitat size and heterogeneity. *Herpetologica*. 2006; 62(4):365–77.

64. Bletz MC, Rosa GM, Andreone F, Courtois EA, Schmeller DS, Rabibisoa NHC, et al. Widespread presence of the pathogenic fungus *Batrachochytrium dendrobatidis* in wild amphibian communities in Madagascar. *Scientific Reports*. 2015; 5. doi: [10.1038/srep08633](https://doi.org/10.1038/srep08633) <http://www.nature.com/srep/2015/150305/srep08633/abs/srep08633.html—supplementary-information>.
65. Choi W, Choi K-S, Lyu D-P, Lee J-S, Lim J, Lee S, et al. Seasonal changes of functional groups in coleopteran communities in pine forests. *Biodivers Conserv*. 2010; 19(8):2291–305.
66. Grimbacher PS, Stork NE. Seasonality of a diverse beetle assemblage inhabiting lowland tropical rain forest in Australia. *Biotropica*. 2009; 41(3):328–37.
67. Inger RF, Voris HK, Frogner KJ. Organization of a community of tadpoles in rain forest streams in Borneo. *J Trop Ecol*. 1986; 2(3):193–205.
68. Morin PJ. Competitive and predatory interactions in natural and experimental populations of *Notophthalmus viridescens dorsalis* and *Ambystoma tigrinum*. *Copeia*. 1983; 1983(3):628–39.
69. Morin PJ, Johnson EA. Experimental studies of asymmetric competition among anurans. *Oikos*. 1988; 53(3):398–407.
70. Werner EE, Glennemeier KS. Influence of forest canopy cover on the breeding pond distributions of several amphibian species. *Copeia*. 1999; 1999(1):1–12.
71. Wells KD. *The Ecology and Behavior of Amphibians*. Chicago: The University of Chicago Press; 2007. 1148 p.
72. Halpern BS, Floeter SR. Functional diversity responses to changing species richness in reef fish communities. *Mar Ecol Prog Ser*. 2008; 364:147–56.
73. Weiblen GD, Webb CO, Novotny V, Basset Y, Miller SE. Phylogenetic dispersion of host use in a tropical insect herbivore community. *Ecology*. 2006; 87(sp7):62–75. doi: [10.1890/0012-9658\(2006\)87\[62:PDOHUI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[62:PDOHUI]2.0.CO;2)
74. Webb CO. Exploring the phylogenetic structure of ecological communities: an example for rain forest trees. *Am Nat*. 2000; 156(2):145–55. PMID: [10856198](https://pubmed.ncbi.nlm.nih.gov/10856198/)
75. Brunbjerg AK, Borchsenius F, Eiserhardt WL, Ejrnæs R, Svenning J-C. Disturbance drives phylogenetic community structure in coastal dune vegetation. *Journal of Vegetation Science*. 2012; 23:1082–94. doi: [10.1111/j.1654-1103.2012.01433.x](https://doi.org/10.1111/j.1654-1103.2012.01433.x)
76. Wiens JJ, Ackerly DD, Allen AP, Anacker BL, Buckley LB, Cornell HV, et al. Niche conservatism as an emerging principle in ecology and conservation biology. *Ecol Lett*. 2010; 13(10):1310–24. doi: [10.1111/j.1461-0248.2010.01515.x](https://doi.org/10.1111/j.1461-0248.2010.01515.x) PMID: [20649638](https://pubmed.ncbi.nlm.nih.gov/20649638/).
77. Swenson NG, Erickson DL, Mi X, Bourg NA, Forero-Montaña J, Ge X, et al. Phylogenetic and functional alpha and beta diversity in temperate and tropical tree communities. *Ecology*. 2012; 93:S112–S25. doi: [10.1890/11-0402.1](https://doi.org/10.1890/11-0402.1)
78. Baraloto C, Hardy OJ, Paine CET, Dexter KG, Cruaud C, Dunning LT, et al. Using functional traits and phylogenetic trees to examine the assembly of tropical tree communities. *J Ecol*. 2012; 100:690–701. doi: [10.1111/j.1365-2745.2012.01966.x](https://doi.org/10.1111/j.1365-2745.2012.01966.x)
79. Bryant JA, Stewart FJ, Eppley JM, DeLong EF. Microbial community phylogenetic and trait diversity declines with depth in a marine oxygen minimum zone. *Ecology*. 2012; 93:1659–73. PMID: [22919912](https://pubmed.ncbi.nlm.nih.gov/22919912/).
80. Grosjean S, Strauß A, Glos J, Randrianiaina R-D, Ohler A, Vences M. Morphological uniformity in the surface-feeding tadpoles of Malagasy litter frogs, subgenus *Chonomantis*. *Zool J Linn Soc*. 2011; 162(1):149–83.
81. Grosjean S, Randrianiaina R-D, Strauß A, Vences M. Sand-eating tadpoles in Madagascar: morphology and ecology of the unique larvae of the treefrog *Boophis picturatus*. *Salamandra*. 2011; 47:63–76.
82. Randrianiaina R-D, Strauß A, Glos J, Glaw F, Vences M. Diversity, external morphology and reverse taxonomy in the specialized tadpoles of Malagasy river bank frogs of the subgenus *Ochthomantis* (genus *Mantidactylus*). *Contributions to Zoology*. 2011; 80:17–65.
83. Randrianiaina RD, Strauß A, Glos J, Vences M. Diversity of the strongly rheophilous tadpoles of Malagasy tree frogs, genus *Boophis* (Anura, Mantellidae), and identification of new candidate species via larval DNA sequence and morphology. *Zookeys*. 2012; 178:59–124. doi: [10.3897/zookeys.178.1410](https://doi.org/10.3897/zookeys.178.1410) PMID: [22539880](https://pubmed.ncbi.nlm.nih.gov/22539880/)
84. Haas A. Phylogeny of frogs as inferred from primarily larval characters (Amphibia: Anura). *Cladistics*. 2003; 19(1):23–89.
85. Losos JB. Phylogenetic niche conservatism, phylogenetic signal and the relationship between phylogenetic relatedness and ecological similarity among species. *Ecol Lett*. 2008; 11:995–1003. doi: [10.1111/j.1461-0248.2008.01229.x](https://doi.org/10.1111/j.1461-0248.2008.01229.x) PMID: [18673385](https://pubmed.ncbi.nlm.nih.gov/18673385/).
86. Crottini A, Madsen O, Poux C, Strauß A, Vieites DR, Vences M. Vertebrate time-tree elucidates the biogeographic pattern of a major biotic change around the K-T boundary in Madagascar. *Proc Natl Acad*

Sci U S A. 2012; 109(14):5358–63. doi: [10.1073/pnas.1112487109](https://doi.org/10.1073/pnas.1112487109) PMID: [22431616](https://pubmed.ncbi.nlm.nih.gov/22431616/); PubMed Central PMCID: [PMC3325728](https://pubmed.ncbi.nlm.nih.gov/PMC3325728/).

87. Best RJ, Caulk NC, Stachowicz JJ. Trait vs. phylogenetic diversity as predictors of competition and community composition in herbivorous marine amphipods. *Ecol Lett.* 2013; 16:72–80. doi: [10.1111/ele.12016](https://doi.org/10.1111/ele.12016) PMID: [23066869](https://pubmed.ncbi.nlm.nih.gov/23066869/).