



HAL
open science

A global database on freshwater fish species occurrence in drainage basins

Pablo A Tedesco, Olivier Beauchard, Rémy Bigorne, Simon Blanchet, Laetitia Buisson, Lorenza Conti, Jean-François Cornu, Murilo Dias, Gaël Grenouillet, Bernard Hugueny, et al.

► **To cite this version:**

Pablo A Tedesco, Olivier Beauchard, Rémy Bigorne, Simon Blanchet, Laetitia Buisson, et al.. A global database on freshwater fish species occurrence in drainage basins. *Scientific Data* , 2017, 4, pp.170141. 10.1038/sdata.2017.141 . hal-01928506

HAL Id: hal-01928506

<https://hal.umontpellier.fr/hal-01928506>

Submitted on 2 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SCIENTIFIC DATA

OPEN Data Descriptor: A global database on freshwater fish species occurrence in drainage basins

Pablo A. Tedesco¹, Olivier Beauchard^{2,3}, Rémy Bigorne¹, Simon Blanchet^{1,4}, Laëtitia Buisson⁵, Lorenza Conti¹, Jean-François Cornu⁶, Murilo S. Dias⁷, Gaël Grenouillet¹, Bernard Hugué¹, Céline Jézéquel¹, Fabien Leprieur⁸, Sébastien Brosse^{1,*} & Thierry Oberdorff^{1,*}

Received: 19 April 2017

Accepted: 1 August 2017

Published: 3 October 2017

A growing interest is devoted to global-scale approaches in ecology and evolution that examine patterns and determinants of species diversity and the threats resulting from global change. These analyses obviously require global datasets of species distribution. Freshwater systems house a disproportionately high fraction of the global fish diversity considering the small proportion of the earth's surface that they occupy, and are one of the most threatened habitats on Earth. Here we provide complete species lists for 3119 drainage basins covering more than 80% of the Earth surface using 14953 fish species inhabiting permanently or occasionally freshwater systems. The database results from an extensive survey of native and non-native freshwater fish species distribution based on 1436 published papers, books, grey literature and web-based sources. Alone or in combination with further datasets on species biological and ecological characteristics and their evolutionary history, this database represents a highly valuable source of information for further studies on freshwater macroecology, macroevolution, biogeography and conservation.

Design Type(s)	data integration objective • database creation objective • species comparison design
Measurement Type(s)	biodiversity assessment objective
Technology Type(s)	digital curation
Factor Type(s)	geographic location
Sample Characteristic(s)	Earth • drainage basin

¹UMR5174 EDB (Laboratoire Evolution et Diversité Biologique), CNRS, IRD, UPS, ENFA, 118 Route de Narbonne, Université Paul Sabatier, F-31062 Toulouse, France. ²Flanders Marine Institute (VLIZ), Wandelaarkaai 7, 8400 Oostende, Belgium. ³Ecosystem Management Research Group, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium. ⁴Station d'Ecologie Théorique et Expérimentale, UMR 5321, 09200 Moulis, France. ⁵UMR 5245 EcoLab (Laboratoire Ecologie Fonctionnelle et Environnement), CNRS, INP, UPS, 118 Route de Narbonne, Université Paul Sabatier, F-31062 Toulouse, France. ⁶Institut des Sciences de l'Evolution (UMR ISEM, CNRS-IRD-UM2), Université de Montpellier, 34000 Montpellier, France. ⁷Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade de Brasília (UnB), Campus Darcy Ribeiro, 70910-900, Brasília-DF, Brazil. ⁸UMR MARBEC, (CNRS, IRD, IFREMER, UM), cc 093, Place E. Bataillon, FR-34095 Montpellier, France. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to P.A.T. (email: pablo.tedesco@ird.fr).

Background & Summary

With c. 126,000 already described animal species, freshwater systems host around 10% of all animals described to date^{1–3} while occupying only 0.8% of the Earth's surface and 0.02% of available aquatic habitable volume⁴. Among aquatic organisms, fishes are a good example of this paradox (the 'freshwater fish paradox' *sensu* Tedesco *et al.*⁵), with c. 40% of all described species inhabiting freshwaters, while the remaining 60% inhabiting marine habitats that comprise >99% of available aquatic habitat⁶. Besides housing a disproportionately high fraction of the global animal diversity considering the small proportion of the earth's surface that they occupy, freshwater ecosystems are also one of the most threatened habitats on Earth^{7,8}. Extinction risk for freshwater fishes, for instance, is thought to be higher than that of terrestrial organisms⁹ and recent extinction rate estimates are 112 to 855 times higher than natural extinction rates^{10–12}.

Describing global scale freshwater fish diversity patterns, understanding the environmental drivers and evolutionary processes shaping such diversity and revealing the major human-related threats were the major goals that motivated the compilation of the present database. Indeed, global scale datasets allowing for biogeographical, macroecological, macroevolutionary and conservation studies were available for only a few well-documented animal taxa such as birds, mammals and amphibians^{13–15}. The present database increases this list of taxa by providing occurrence data by drainage basin worldwide for the most diverse group of vertebrates (i.e. fishes), with more than 33500 species described to date (FishBase; <http://www.fishbase.org>), from which c. 40% inhabit permanently freshwater systems.

We conducted an extensive survey of freshwater fish species distribution based on 1436 published papers, books, grey literature, databases and web-based sources, resulting in species lists for 3119 drainage basins covering more than 80% of the Earth surface (Fig. 1). Two important survey efforts were conducted, respectively completed in 2008 (ref. 16) and 2013 (ref. 17). To date, these databases have been used in several studies that have increased our understanding of freshwater fish species distributions. These studies allowed to accurately map global patterns of native¹⁸, endemic¹⁹ and non-native²⁰ freshwater fish species richness and to reveal their environmental and human-related determinants. The databases were also used to evaluate non-native species influence on native communities structure²¹, to forecast climate change effects on species extinction processes¹¹ and to analyse effects of current and future scenarios of species introductions on fish faunas homogenization processes^{22–25}. Recent studies also focused on analysing the influence of past river connections on the present distribution of native fish species¹⁷, on analysing geographical and trait-based differences in diversification rates and origin of actinopterygian fish families⁵, and on evaluating human-related extinction drivers¹².

Although the database has already provided a lot of insightful knowledge, it still represents a valuable source of information for further studies on freshwater macroecology, macroevolution, biogeography and conservation. For instance, the present dataset could serve to identify diversity hotspots and to generate a global map of ichthyogeographic regions by combining data on the distributions and phylogenetic relationships of species, allowing *in fine* the identification of geographic areas harbouring distinct evolutionary histories. Furthermore, in association with data on the time and place of origin of species or on species functional traits, the global occurrence dataset could provide new insights on the macroevolution of freshwater fishes or approach the functional characteristics of communities. Those




Figure 1. Global map indicating the drainage basins included in the database with different colours by biogeographic realm²⁷. The 3119 drainage basins cover more than 80% of the Earth surface (excluding deserts), ranging from 70% for the Indo-Malay region to over 90% for the Afrotropical region.

forthcoming approaches would surely help designing large scale conservation priorities for freshwater fishes.

The database is organised in three sub-datasets and one shapefile. The first dataset contains the species occurrence records by drainage basin along with their native or non-native status and the corresponding FishBase species code and valid name. The second dataset, which is simply the export of the shapefile attributes table, contains geographic information on the drainage basins (e.g. geographic coordinates, surface area). The third dataset contains the list of references that were used to build the species lists for each of the drainage basins. This reference list is obviously not definitive and updates of the database will be performed regularly to include new occurrence records, the distribution of newly described species, species lists of new drainage basins and nomenclature changes in the always moving taxonomy.

Methods

Information sources

This global database of freshwater fish species distribution results from a joint collaboration between three French research institutes, i.e. the University Paul Sabatier in Toulouse (UPS), the National Museum of Natural History (MNHN) and the Research Institute for Development (IRD). The financial support necessary to build this database mainly came from two projects: the 'Freshwater Fish Diversity' (National Agency for Research: ANR-06-BDIV-010) and 'BioFresh' (7th Framework European program, Contract N°226874) projects. Starting in 2003, we conducted an extensive survey of literature published from 1960 to 2014 on native and non-native freshwater fish species at the drainage basin grain. This survey was complemented with web-based sources from national and international biodiversity inventory initiatives compiling either or both collection and field sampling data.

Our efforts were mainly devoted to find information sources providing complete fish species lists of a given drainage basin, except for some large basins (e.g. the Amazon basin) where we cumulated sub-drainage basin species lists and point sampling locations to obtain the most complete possible coverage of the entire drainage basin. We also used local or regional check lists such as local inventories of stream reaches or inventories based solely on a given family or genus to complement our species lists and for cross-checking available information at the drainage basin scale. The resulting database was gathered from 1436 sources including published papers, books, grey literature and web-based sources that included museum collections, national or regional initiatives compiling monitoring data (mainly for developed countries), continental scale atlases of species distribution and international biodiversity initiatives. When published information was found in languages not handled by any of the team members (e.g. national inventory reports or books), a translator kindly helped us to ensure the collection of correct information on river basins, species lists and location of the species.

Species, taxonomy and status

Sub-species were not considered due to limited data availability and all occurrences not identified to species level were discarded (i.e. occurrences giving only genus names commonly abbreviated to sp., species *affinis* commonly abbreviated to: sp. aff., aff., or affin. or species *confer* abbreviated to cf.). Species migrating between both marine and freshwater environments were systematically included in the database. Concerning marine and estuarine species occasionally occurring in freshwaters, these species may be reported in the database but their distribution information should not be considered as exhaustive in any case, as these systems are not the focus of the database.

All species scientific names are reported in the database as given in each information source. These species names were then carefully checked for typing errors and misspellings. Because taxonomy is a 'moving target', species names were standardized based on valid species names and their synonyms reported in FishBase using the 'rfishbase' package²⁶ from the R environment (<http://www.R-project.org>). For those species names that did not match with any synonym or valid name from FishBase, a manual search was applied in the Catalogue of Fishes (<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). This last step allowed finding valid species names and species recently described that are still not included in FishBase. For recently described species not yet validated by FishBase or species only considered valid by the Catalogue of Fishes, a temporary code was created starting by 'x' (0.04% of all valid species). The remaining species names were considered as invalid and excluded from the database (only 0.6% of all species names). The final standardized species list has 14953 valid names avoiding biases due to synonyms and uncertain identifications (see 'Technical Validation'). According to FishBase, from this list of 14953 valid names our database contains the distribution of 13721 species inhabiting fresh or brackish waters, the remaining being marine species also entering fresh or brackish waters, but not recorded as such by FishBase. As a whole, the database harbours 101779 occurrence records (i.e. single species-drainage basin couples).

A native or exotic status was assigned to each species occurrence record based on the information provided by the sources and further checked (see 'Technical Validation'). An exotic species is defined as a directly or indirectly (e.g. via artificial channels) introduced species that established in the considered drainage basin. Exotic species included in the database are supposed to complete all their life cycle in the considered basins and to present self-sustaining populations in those basins. When an exotic species occurrence was acknowledged to be unsuccessful (i.e. failure of establishment of that species in the drainage basin) or needing regular release of new individuals to maintain the presence of the introduced

species (i.e. stocking), the species was not included in the basin's species list. Species that might be globally extinct or extirpated from a drainage basin were considered as native in the database because the inventory of freshwater fish diversity loss was not targeted (see for instance Dias *et al.*¹² for a compilation of extirpated freshwater fish species for Western Europe and North America).

Drainage basin location and names

Each drainage basin was assigned a unique name that can be used as an identifier and was characterized by its location in one of the eight terrestrial biogeographic realms (as described by Olson *et al.*²⁷; Fig. 1), the country (or main country for shared drainage basins), its endorheic or exorheic type of water flow, its geographic coordinates at the river mouth (for exorheic drainage basins), the geographic coordinates of its centroid and its drainage surface area.

A specific geographic referential (Fig. 1) was built by modifying the 30 sec HydroSheds layer²⁸ to improve the delimitation and accuracy of drainage basins. For instance, some small coastal drainage basins were included in one single HydroSheds polygon but were considered as separate basins because having distinct outlets to the sea. Some drainage basins from oceanic islands have no HydroSheds code simply because not considered in the HydroSheds shapefile. Maps and geographic information available in the compiled literature and web-based sources were used to locate, name and improve our drainage basins layer, complemented by country and continental scale geographic data (e.g. Faunafri project for the African continent; <http://www.poissons-afrique.ird.fr/faunafri/>) and local topographic maps. This new geographic referential is provided as a shapefile to facilitate future uses of the database.

Updates and limitations

Species are continuously being discovered and freshwater fishes are no exception, even in well-known regions²⁹. Rivers are also continuously being explored and re-explored by freshwater scientists. The database is obviously not complete and definitive, and we aim to support the database with regular updates, ideally with bi-annual steps, depending on the resources and funding. Three main factors will be considered in future updates: (1) new or previously non available data sources with species lists or records for additional drainage basins or drainage basins already present in the database; (2) distribution of newly described species; and (3) nomenclature changes in the taxonomic classification. The technical validation procedures described below will also be applied to any new information included in the database. Researchers having access to new data that want this information to be included in the database can send the references or data to the corresponding author PAT. This information will be included, after validation, in the next update release. The resulting new versions of the database will be released through *Figshare* and also through the more specialized *Freshwater Biodiversity Data Portal* (<http://data.freshwaterbiodiversity.eu/>) to ensure the long-term availability of the database.

All biogeographic realms are well represented in terms of surface coverage (Fig. 1). There are however some regional gaps that will be gradually filled in the next updates of the database. For instance, Indonesian islands, coastal rivers of Peru and Northeast Brazil are regions where only few drainage basins are informed in the database. In these regions the scarce existing information is not easily available. Southeast Asia is the less well represented region in terms of surface coverage (Fig. 1), which is certainly related to the low number of freshwater taxonomists working in this highly diverse region³⁰. All these spatial gaps in the database will be prioritized in future updates through literature and web-based sources monitoring.

Data Records

The database is organised in three datasets and one shapefile: the species occurrence records, the drainage basins and the information sources table. The three tables are in csv format (columns separated by commas) and the shapefile in ArcGis shp format (Data Citation 1). The drainage basins table is given in.csv and shapefile formats. Both formats can be linked to the species occurrence table using the unique drainage basin names to visualize and analyse species distribution using any adapted software (e.g. R or QGIS, <http://qgis.osgeo.org>).

- A. The species occurrence records table has six columns: (1) the name of the drainage basin, (2) the scientific name of the fish species according to the information source, (3) the native or exotic status of the occurrence records, (4) the taxonomic serial number (TSN) from the Integrated Taxonomic Information System (ITIS, <https://www.itis.gov/>) when available, (5) the FishBase code when available, (6) the FishBase or Catalogue of Fishes valid scientific name at the time of releasing the database, (7) the occurrence status which can be either 'valid' or 'questionable' (see Technical Validation).
- B. The geographic information on drainage basins is organised in nine columns given in table and shapefile formats: (1) the unique drainage basin name, (2) the main country where it belongs, (3) the corresponding biogeographic region, (4) the endorheic or exorheic status, (5) and (6) the longitude and latitude coordinates of the drainage basin outlet to the sea (only for exorheic drainages), (7) and (8) the centroid longitude and latitude coordinates of the drainage basin, (9) the surface area of the drainage basin.

- C. The information sources table has three columns: (1) the drainage basin names, (2) the type of information sources (e.g. published paper, book, report, online database, PhD Thesis), (3) the references used to build the freshwater fish species list for the corresponding drainage basin.

Technical Validation

Taxonomic validation

Each species name found in a given information source was confronted to the valid and synonym species names lists from FishBase and the Catalogue of Fishes to ensure the validity of the identifications provided in the information source. After taxonomic validation, 103 invalid (unknown) species names were excluded from the database.

Species distribution and status validation

Occurrence records were carefully reviewed by the database' contributors. When several information sources were used to compile the species list of a given drainage basin, particular attention was given to cross check the occurrence records and ensure a good spatial representation of the drainage basin to avoid (or at least minimize) incomplete species lists. Because occurrence data available in FishBase is often incomplete, FishBase was only used as a secondary source to collect species distribution data and to check that the resulting species distributions from our database corresponded to the broad information given in FishBase.

Because only a few (mostly migratory) freshwater species can occur in more than one biogeographic realm, the distribution of every species occurring in more than one realm was carefully verified. Similarly, for species distributed in a single realm, when one or more occurrences were inconsistent with the actual known distribution of a species (i.e. the presence in a drainage located far away from a group of drainages where the species is known to occur), these occurrences were qualified as 'questionable'. Particular care was taken with the occurrences of all species considered exotic at least in one drainage basin. The native and exotic distributions of those species were carefully checked to avoid any status error.

References

- Balian, E. V., Segers, H., Lévêque, C. & Martens, K. The freshwater animal diversity assessment: an overview of the results. *Hydrobiologia* **595**, 627–637 (2008).
- Mora, C., Tittensor, D. P., Adl, S., Simpson, A. G. B. & Worm, B. How many species are there on Earth and in the ocean? *PLoS Biol.* **9**, e1001127 (2011).
- Wiens, J. J. Faster diversification on land than sea helps explain global biodiversity patterns among habitats and animal phyla. *Ecol. Lett.* **18**, 1234–1241 (2015).
- Dawson, M. N. research letter: Species richness, habitable volume, and species densities in freshwater, the sea, and on land. *Front. Biogeogr.* **4**, 105–116 (2012).
- Tedesco, P. A., Paradis, E., Lévêque, C. & Huguency, B. Explaining global-scale diversification patterns in actinopterygian fishes. *J. Biogeogr.* **44**, 773–783 (2017).
- Lévêque, C., Oberdorff, T., Paugy, D., Stiassny, M. L. J. & Tedesco, P. A. Global diversity of fish (Pisces) in freshwater. *Hydrobiologia* **595**, 545–567 (2008).
- Jenkins, M. Prospects for biodiversity. *Science* **302**, 1175–1177 (2003).
- Vörösmarty, C. J. *et al.* Global threats to human water security and river biodiversity. *Nature* **467**, 555–561 (2010).
- Ricciardi, A. & Rasmussen, J. B. Extinction rates of north American freshwater fauna. *Conserv. Biol.* **13**, 1220–1222 (1999).
- Burkhead, N. M. Extinction Rates in North American Freshwater Fishes, 1900–2010. *BioScience* **62**, 798–808 (2012).
- Tedesco, P. A. *et al.* A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. *J. Appl. Ecol.* **50**, 1105–1115 (2013).
- Dias, M. S. *et al.* Anthropogenic stressors and riverine fish extinctions. *Ecol. Indic.* **79**, 37–46 (2017).
- Orme, C. D. L. *et al.* Global hotspots of species richness are not congruent with endemism or threat. *Nature* **436**, 1016–1019 (2005).
- Grenyer, R. *et al.* Global distribution and conservation of rare and threatened vertebrates. *Nature* **444**, 93–96 (2006).
- Dyer, E. E., Redding, D. W. & Blackburn, T. M. The global avian invasions atlas, a database of alien bird distributions worldwide. *Sci. Data* **4**, 170041 (2017).
- Brosse, S. *et al.* Fish-SPRICH: a database of freshwater fish species richness across the World. *Hydrobiologia* **700**, 343–349 (2013).
- Dias, M. S. *et al.* Global imprint of historical connectivity on freshwater fish biodiversity. *Ecol. Lett.* **17**, 1130–1140 (2014).
- Oberdorff, T. *et al.* Global and Regional Patterns in Riverine Fish Species Richness: A Review. *Int. J. Ecol.* **2011**, 1–12 (2011).
- Tedesco, P. A. *et al.* Patterns and processes of global riverine fish endemism. *Glob. Ecol. Biogeogr.* **21**, 977–987 (2012).
- Leprieux, F., Beauchard, O., Blanchet, S., Oberdorff, T. & Brosse, S. Fish Invasions in the World's River Systems: When Natural Processes Are Blurred by Human Activities. *PLoS Biol.* **6**, e28 (2008).
- Blanchet, S. *et al.* Non-native species disrupt the worldwide patterns of freshwater fish body size: implications for Bergmann's rule. *Ecol. Lett.* **13**, 421–431 (2010).
- Toussaint, A., Beauchard, O., Oberdorff, T., Brosse, S. & Villéger, S. Historical assemblage distinctiveness and the introduction of widespread non-native species explain worldwide changes in freshwater fish taxonomic dissimilarity. *Glob. Ecol. Biogeogr.* **23**, 574–584 (2014).
- Toussaint, A., Beauchard, O., Oberdorff, T., Brosse, S. & Villéger, S. Worldwide freshwater fish homogenization is driven by a few widespread non-native species. *Biol. Invasions* **18**, 1295–1304 (2016).
- Villéger, S., Blanchet, S., Beauchard, O., Oberdorff, T. & Brosse, S. Homogenization patterns of the world's freshwater fish faunas. *Proc. Natl. Acad. Sci.* **108**, 18003–18008 (2011).
- Villéger, S., Blanchet, S., Beauchard, O., Oberdorff, T. & Brosse, S. From current distinctiveness to future homogenization of the world freshwater fish faunas. *Divers. Distrib.* **21**, 223–235 (2015).
- Boettiger, C., Lang, D. T. & Wainwright, P. C. fishbase: exploring, manipulating and visualizing FishBase data from R. *J. Fish Biol.* **81**, 2030–2039 (2012).
- Olson, D. M. *et al.* Terrestrial ecoregions of the worlds: A new map of life on Earth. *BioScience* **51**, 933–938 (2001).

28. Lehner, B., Verdin, K. & Jarvis, A. New global hydrography derived from spaceborne elevation data. *EOS Trans. Am. Geophys. Union* **89**, 93–94 (2008).
29. Tedesco, P. A. *et al.* Estimating How Many Undescribed Species Have Gone Extinct: Estimating Undescribed Species Extinctions. *Conserv. Biol.* **28**, 1360–1370 (2014).
30. Kottelat, M. & Whitten, A. J. Freshwater biodiversity in Asia with special reference to fish. World Bank Tech. Pap. No 343 (1996).

Data Citation

1. Tedesco, P. A. *et al.* *Figshare* <https://doi.org/10.6084/m9.figshare.c.3739145> (2017).

Acknowledgements

The construction of this database was mainly supported by the National Agency for Research (ANR) Freshwater Fish Diversity (ANR-06-BDIV-010) and by the EU BIOFRESH project (7th Framework European program, Contract N°226874). EDB laboratory was also supported by 'Investissement d'Avenir' grants (CEBA, ANR-10-LABX-0025; TULIP, ANR-10-LABX-41). BOREA laboratory was also supported by the National Agency for Research (ANR) FISHLOSS project (ANR-09-PEXT-008). We are grateful to the AMAZONFISH project (ERANet-LAC: ELAC2014/DCC-0210; www.amazon-fish.com) for validating our list of fish species for the Amazon River basin. M.S.D. received a PhD grant from the Brazilian government (Science without Borders program, CNPq/GDE n° 201167/2012-3).

Author Contributions

P.A.T. wrote the first draft of the manuscript, and all authors contributed substantially to finalising this manuscript. P.A.T. and O.B. entered occurrence data, revised the information sources and compiled the database, with contributions of all authors. C.J. and J.-F.C. handled the geographic data related to the location and delimitation of the drainage basins, and all authors contributed to checking the information on distribution and status of the species. T.O. and S.B. initiated and designed the Database.

Additional information

Competing interests: The authors declare no competing financial interests.

How to cite this article: Tedesco, P. A. *et al.* A global database on freshwater fish species occurrence in drainage basins. *Sci. Data* 4:170141 doi: 10.1038/sdata.2017.141 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files made available in this article.

© The Author(s) 2017