



Development of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) larvae feeding on the plant material contained in the water

Frédéric Darriet

► To cite this version:

Frédéric Darriet. Development of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) larvae feeding on the plant material contained in the water. *Annals of Community Medicine and Practice*, 2016, 2 (1). hal-01937618

HAL Id: hal-01937618

<https://hal.umontpellier.fr/hal-01937618>

Submitted on 28 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short Communication

Development of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) Larvae Feeding on the Plant Material Contained in the Water

Frédéric Darriet*

Institut de Recherche pour le Développement (IRD), UMR MIVEGEC (UM1-UM2-CNRS 5290-IRD 224) Maladies Infectieuses et Vecteurs, France

*Corresponding author

Frédéric Darriet, Institut de Recherche pour le Développement (IRD), UMR MIVEGEC (UM1-UM2-CNRS 5290-IRD 224) Maladies Infectieuses et Vecteurs, Ecologie, Génétique, Evolution et Contrôle, 911 avenue Agropolis, BP 64501, 34394 Montpellier, France, Tel: 33-(0)4-67-41-63-95; Email: frederic.darriet@ird.fr

Submitted: 28 October 2016

Accepted: 26 November 2016

Published: 28 November 2016

Copyright

© 2016 Darriet

OPEN ACCESS

Keywords

- *Aedes aegypti*
- *Aedes albopictus*
- Plant material
- Log-probit analysis
- Trophic performances

Abstract

Background: In this laboratory study we measured the trophic performances of *Aedes aegypti* and *Aedes albopictus* larvae (Diptera: Culicidae), both being vectors of the dengue fever, chikungunya and zika virus in the world.

Methods: Depending on the quantities of plant material contained in the breeding sites, the bioassays enabled to assess the times it took for 50% of imaginal emergence to occur (IE_{50}). They also enabled to determine the amounts of plant material needed for 50% of adult emergence (IE_{50}).

Results: Water containing 3.3 g of plant material per liter allowed 50% of *Ae. albopictus* adults to emerge within 8 days (IE_{50}), against 60 days at 0.5g/liter. As for *Ae. aegypti*, the IE_{50} took 8 days at 3.3g/l against 29 days at 1.7g/l. The IE_{50} also revealed that 0.61g of plant material were needed for a 50% of adult emergence of *Ae. albopictus*. To reach the same survival rate among *Ae. aegypti*, the larvae must grow in an environment twice as rich in food supply.

Conclusion: This research work has revealed that the *Ae. albopictus* larvae can develop in water collections where the shortage of organic material hinders or compromises the development and survival of *Ae. aegypti*. The outstanding trophic performances of *Ae. albopictus* would thus partly account for the invasive character of *Ae. albopictus*, as well as the dying out of its competitor *Ae. aegypti* in the regions of the world shared by both species.

ABBREVIATIONS

PM: Plant Material; IE_{50} : Imaginal Emergence Time 50%; IE_{50} : Imaginal Emergence 50%

INTRODUCTION

The distribution of mosquito populations mostly depends on the physico-chemical and biological characteristics of the breeding sites. *Aedes aegypti* (L.) and *Aedes albopictus* (Skuse) develop in clear domestic and peridomestic water collections, such as earthenware vases, barrels, cisterns, gutters, cans, tires and plant saucers [1-2]. These two mosquitoes are the major vectors of the dengue fever, chikungunya and zika virus in the world [3-10]. It is now clearly stated that the population growth of *Ae. aegypti* and *Ae. albopictus* larvae is closely associated with the nature of the

resources available [11-13]. If the source of food mainly consists in plant detritus (deciduous and coniferous leaves, flowers and grass) it remains quite rich in cellulose and hard to digest for mosquito larvae [14-15]. In this laboratory study, the bioassays consisted in a follow-up of larvae batches until the emergence of adults, whose larvae grew up in environments with different contents of plant material (dry grasses). The follow-up allowed to assess the times taken by the larvae to develop, depending on the availability plant material (IE_{50}). The quantities of dry grasses needed for each species to reach a 50% adult emergence were also determined (IE_{50}).

MATERIALS AND METHODS

The Bora strain of *Ae. aegypti*, originally from French Polynesia, has been reared for more than 20 years in our laboratory. The

Ae. albopictus strain used in this study comes from the village of Perols located south of the Hérault department (France). This strain of *Ae. albopictus* has been raised in the Institute of Research for Development (IRD) insectarium in Montpellier (France) since September 2014. The plant material (PM) used for the preparation of the larval environment consisted in commercial rodent food (hay) that can be found in pet shops (Zolux®). The quantities of PM assessed on *Ae. aegypti* and *Ae. albopictus* were of 0.5g/l - 1g/l - 2.5g/l and 3.3g/l. A 0.25g/l content was tested on *Ae. albopictus* only and 1.7g/l on *Ae. aegypti* so as to characterize the trophic performances of these two species of mosquitoes more precisely. PM was prepared in 0.0042 m³ plastic trays (length: 0.30m; width: 0.20m; depth: 0.07m), each containing one liter of reverse osmotic water. 24 hr after the preparation of the larval environments, one hundred first instars larvae (L1) of *Ae. aegypti* or *Ae. albopictus* were counted and placed in a tray. Each artificial milieu was evaluated on a total of three replicates. Throughout the duration of the experiment, the trays were maintained at a temperature of 27 ± 2°C in the laboratory.

Female and male adults were counted in each environment to establish the averages of imaginal emergence with a 95% confidence interval [16]. The larval environments allowing more than 50% of adult emergences were analyzed using the log-probit software [17] so as to determine the duration of preimaginal developments leading to 50% of imaginal emergence (IE_{50}). Based on this same log-probit analysis, the quantity of plant material needed for 50% adult emergence was assessed as well (IE_{50}).

RESULTS

Figure 1 shows the averages of male and female emergences in the different larval environment. The imaginal emergences of *Ae. albopictus* were above 50% with only 0.5g of PM per liter. As much as 1.7g/l of PM are needed to reach the same survival rate of *Ae. aegypti*. Except for the 2.5g/l quantity of PM, for which the averages of male and female *Ae. albopictus* proved statistically different ($P = 0.03$), the other quantities tested on both strains did not display significant differences ($0.094 < P < 0.89$). The times needed for 50% of adult *Ae. albopictus* and *Ae. aegypti* to emerge (IE_{50}) (Table 1) were 8 and 11 days respectively, with 3.3g and 2.5g of PM per liter. IE_{50} exceeded a period of one month for 1g/l and equaled two months for 0.5g/l. At the concentration of 1.7g/l, 50% of *Ae. aegypti* adults emerged after 29 days of study. Whatever the species studied, it is worth noticing the regression line slopes steepened as the quantities of PM increased. This observation simply shows that the more food in the environment, the faster the larval growth.

A quantity of 0.61g of PM per liter is enough for 50% (IE_{50}) of the *Ae. albopictus* larvae to develop all the way to imaginal stage (Table 2). For a similar survivorship with *Ae. aegypti* larvae, twice as much organic material is needed in the water. The difference of trophic performances translates in log-probit regression lines whose slopes almost double depending on the species. *Ae. albopictus* larvae's ability to grow in environments that can be either poor or rich in organic matter (Figure 1) will produce a more gradual response in adult with a gentler slope of the regression line.




Figure 1 Averages (95% CI) of males and females mosquitoes whose larvae lived in environments containing different quantities of plant material.

Table 1: Determination for each mosquito species of the value of imaginal emergence time 50% (IE_{50}), estimated from the log-probit analysis.

Species	Plant material quantities (g/l)	IE_{50} (in day) (95% CI)	Slope (± SE)
<i>Aedes albopictus</i>	0.5	60.2 (56.7-64.6)	2.01 (± 0.13)
	1	35.7 (34.7-36.8)	2.6 (± 0.10)
	2.5	10.7 (10.3-11.1)	4.9 (± 0.3)
	3.3	7.7 (6.4-9.1)	5.9 (± 0.6)
<i>Aedes aegypti</i>	1.7	28.8 (27.3-30.6)	1.9 (± 0.14)
	2.5	11.1 (10.7-11.4)	5.4 (± 0.3)
	3.3	8.2 (7.9-8.4)	5.9 (± 0.3)

DISCUSSION

Mosquito larvae breed in water collections whose physico-chemical and biological characteristics differ a lot from one

Species	IE_{50} (in g/l of plant material) (95% CI)	Slope (± SE)
<i>Aedes albopictus</i>	0.61 (0.49-0.76)	1.98 (± 0.23)
<i>Aedes aegypti</i>	1.3 (1.0-1.7)	3.4 (± 0.7)

This research work undoubtedly shows that *albopictus* develops in water collections where the shortage of organic material hinders or compromises the development and survival of *Ae. aegypti*. The outstanding trophic performances would therefore partly account for the invasive character of *Ae. albopictus*, as well as the dying out of its competitor *Ae. aegypti* in the regions of the world shared by both species [23-25].

Special thanks to H  l  ne Darriet for kindly translating this French paper into English.

1. Cordellier R, Germain M, Hervy JP, Mouchet J. Guide pratique pour l'étude des vecteurs de fièvre jaune en Afrique et méthode de lutte. ORSTOM Editions, Initiation, Documents techniques. Paris, France. 1977.
2. Delatte H, Dehecq J, Thiria J, Domerg C, Paupy C, Fontenille D. Geographic distribution and developmental sites of *Aedes albopictus* (Diptera: Culicidae) during a chikungunya epidemic event. Vector Borne Zoonotic Dis. 2008; 8: 25-34.
3. WHO. Regional office for South-East Asia. Guidelines on clinical management of Chikungunya fever. World Health Organization, New Delhi, India. 2008.
4. WHO. Special programme for research and training in tropical diseases. Dengue guidelines for diagnosis treatment, prevention and control. World Health Organization, Geneva, Switzerland. 2009; 147.
5. Hayes EB. Zika virus outside Africa. Emerg Infect Dis. 2009; 15: 1347-1350.
6. Reiter P. Yellow fever and dengue: a threat to Europe? Euro Surveill. 2010; 15: 19509.
7. Rohani A, Adil Azahary AR, Malinda M, Zurainee MN, Rozilawati H, Wan Najdah WM, et al. Eco-virological survey of *Aedes* mosquito larvae in selected dengue outbreak areas in Malaysia. J Vector Borne Dis. 2014; 51: 327-332.
8. Kraemer MU, Sinka ME, Duda KA, Mylne AQ, Shearer FM, Barker CM. et al. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae albopictus*. Elife. 2015; 4.
9. kiner MM, Demirci B, Babuadze G, Robert V, Schaffner F. Spread of the Invasive Mosquitoes *Aedes aegypti* and *Aedes albopictus* in the Black Sea Region Increases Risk of Chikungunya, Dengue, and Zika Outbreaks in Europe. PLoS Negl Trop Dis. 2016; 10.
10. Hennessey M, Fischer M, Staples JE. Zika Virus Spreads to New Areas - Region of the Americas, May 2015-January 2016. MMWR Morb Mortal Wkly Rep. 2016; 65: 55-58.
11. Juliano SA. Species interactions among larval mosquitoes: Context dependence across habitats gradients. Annu Rev Entomol. 2009; 54: 37-56.
12. Murrell EG, Damal K, Lounibos LP, Juliano SA. Distributions of Competing Container Mosquitoes Depend on Detritus Types, Nutrient Ratios, and Food Availability. Ann Entomol Soc Am. 2011; 104: 688-698.
13. Daugherty MP, Alto BW, Juliano SA. Invertebrate carcasses as a resource for competing *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae). J Med Entomol. 2000; 37: 364-372.
14. Clements AN. The biology of mosquitoes: Development, nutrition and reproduction. CABI Publishing, Eastbourne, UK. 2000.
15. Darriet F. Des moustiques et des hommes. Chronique d'une pullulation annoncée. IRD Éditions, collection Didactiques, Marseille, France. 2014.
16. Statistica. Windows statistical software, version 10, Stat Soft France. 2011.
17. Raymond M, Prato G, Ratsira D. Probit and Logit Analysis Program, version 2.0. Praxme, Biometric, Centre National de la Recherche Scientifique, Montpellier, France. 1997.
18. Darriet F, Corbel V. *Aedes aegypti* oviposition in response to NPK fertilizers. Parasite. 2008; 15: 89-92.
19. Darriet F, Zumbo B, Corbel V, Chandre F. [Influence of plant matter and

- NPK fertilizer on the biology of *Aedes aegypti* (Diptera: Culicidae)]. Parasite. 2010; 17: 149-154.
20. Mouchet J, Giacomini T, Julvez J. [Human diffusion of arthropod disease vectors throughout the world]. Sante. 1995; 5: 293-298.
21. Dommergues Y. La biologie des sols. Presses Universitaires de France, Paris, France. 1968.
22. Minard G, Mavingui P, Moro CV. Diversity and function of bacterial microbiota in the mosquito holobiont. Parasit Vectors. 2013; 20: 146.
23. Juliano SA, Lounibos LP. Ecology of invasive mosquitoes: effects on resident species and on human health. Ecol Lett. 2005; 8: 558-574.
24. Bagny L, Delatte H, Quilici S, Fontenille D. Progressive decrease in *Aedes aegypti* distribution in Reunion Island since the 1900s. J Med Entomol. 2009; 46: 1541-1545.
25. Bagny-Beilhe L, Arnoux S, Delatte H, Lajoie G, Fontenille D. Spread of invasive *Aedes albopictus* and decline resident *Aedes aegypti* in urban areas of Mayotte 2007-2010. Biol Invasions. 2012; 14: 1623-1633.

Cite this article

Darriet F (2016) Development of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) Larvae Feeding on the Plant Material Contained in the Water. Ann Community Med Pract 2(1): 1014.