



HAL
open science

Remote Effect of Insecticide-Treated Nets and the Personal Protection against Malaria Mosquito Bites

Nicolas Moiroux, Fabrice Chandre, Jean-Marc Hougard, Vincent Corbel,
Cédric Pennetier

► **To cite this version:**

Nicolas Moiroux, Fabrice Chandre, Jean-Marc Hougard, Vincent Corbel, Cédric Pennetier. Remote Effect of Insecticide-Treated Nets and the Personal Protection against Malaria Mosquito Bites. PLoS ONE, 2017, 12 (1), pp.e0170732. 10.1371/journal.pone.0170732 . hal-01937675

HAL Id: hal-01937675

<https://hal.umontpellier.fr/hal-01937675>

Submitted on 28 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Remote Effect of Insecticide-Treated Nets and the Personal Protection against Malaria Mosquito Bites

Nicolas Moiroux^{1,2*}, Fabrice Chandre¹, Jean-Marc Hougard¹, Vincent Corbel¹, Cédric Pennetier^{1,3}

1 MIVEGEC (IRD 224 - CNRS 5290 - Université de Montpellier), Institut de Recherche pour le Développement (IRD), Montpellier, France, **2** Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso, **3** Institut Pierre Richet (IPR), Bouaké, Côte d'Ivoire

* nicolas.moiroux@ird.fr



OPEN ACCESS

Citation: Moiroux N, Chandre F, Hougard J-M, Corbel V, Pennetier C (2017) Remote Effect of Insecticide-Treated Nets and the Personal Protection against Malaria Mosquito Bites. PLoS ONE 12(1): e0170732. doi:10.1371/journal.pone.0170732

Editor: John Vontas, University of Crete, GREECE

Received: November 17, 2016

Accepted: January 10, 2017

Published: January 27, 2017

Copyright: © 2017 Moiroux et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data of the trials listed in Table 1 and used in the present study were performed under Technical Service Agreements (TSA) between IRD and WHO. According to these TSAs, IRD should not disclose the data without the authorization of WHO. Readers may contact WHOPES office to request the data at the following email address: whopes@who.int.

Funding: Trials listed in Table 1 and used in the analysis received financial support from WHO. All authors are employed by Institut de Recherche pour le Développement (IRD). There was no

Abstract

Experimental huts are part of the WHO process for testing and evaluation of Insecticide Treated Nets (ITN) in semi-field conditions. Experimental Hut Trials (EHTs) mostly focus on two main indicators (i.e., mortality and blood feeding reduction) that serve as efficacy criteria to obtain WHO interim recommendation. However, several other outputs that rely on counts of vectors collected in the huts are neglected although they can give useful information about vectors' behavior and personal protection provided by ITNs. In particular, EHTs allow to measure the deterrent effect and personal protection of ITNs. To provide a better assessment of ITNs efficacy, we performed a retrospective analysis of the deterrence and the personal protection against malaria transmission for 12 unwashed and 13 washed ITNs evaluated through EHTs conducted in West Africa. A significant deterrent effect was shown for six of the 12 unwashed ITNs tested. When washed 20 times, only three ITNs had significant deterrent effect (Rate Ratios (RR)<1; p<0.05) and three showed an apparent "attractiveness" (RR>1; p<0.01). When compared to the untreated net, all unwashed ITNs showed lower number of blood-fed *Anopheles* indicating a significant personal protection (RR<1, p<0.05). However, when washed 20 times, three ITNs that were found to be attractive did not significantly reduce human-vector contact (p>0.05). Current WHO efficacy criteria do not sufficiently take into account the deterrence effect of ITNs. Moreover, the deterrence variability is rarely discussed in EHT's reports. Our findings highlighted the long-range effect (deterrent or attractive) of ITNs that may have significant consequences for personal/community protection against malaria transmission. Indicators measuring the deterrence should be further considered for the evaluation of ITNs.

Background

Between 2000 and 2015, the scale-up of malaria control interventions helped to reduce malaria mortality by 60% globally, and by 66% in sub-Saharan Africa (SSA). However, malaria is still a

additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

major cause of death with 438 000 deaths (uncertainty range: 236 000–635 000) of which 90% occur in SSA [1]. A recent study showed that about 70% of malaria cases were averted since 2000 due to the deployment of insecticide treated net (ITN) [2] hence underlying the need to achieve wide coverage of core interventions in all transmission settings. The ownership of ITNs increased from 2% in 2000 to 56% in 2015 but is still far from the universal coverage objective of WHO [1].

According to WHO [1], National Malaria Control Programs (NMCPs) and global malaria partners should only distribute ITNs that have been recommended by the WHO Pesticide Evaluation Scheme (WHOPES). Sixteen products are currently recommended by WHOPES [3]. WHOPES evaluation scheme is a 3 steps process (1. laboratory—2. small- and 3. large-scale field studies) undertaken to determine the efficacy and operational acceptability of ITNs [4]. The objectives of laboratory testing (phase I) are to determine the efficacy and wash-resistance of an ITN and to study the dynamics of the insecticide on the netting fibre. Candidate ITNs that meet the requirements of phase I testing should subsequently be tested in phase II studies in experimental huts, where the efficacy of ITNs against wild free-flying mosquitoes is investigated. Candidate ITNs that reach the efficacy thresholds of phase I and phase II studies receive an interim recommendation for use as Long Lasting Insecticidal Nets (LLIN) (limited to four years of duration). To get the full recommendation, the net survivorship and attrition, fabrics physical integrity and insecticidal efficacy must be monitored and must reach WHOPES criteria during 3 years under field conditions (phase III large-scale field study) [5].

Experimental huts used in phase II studies allow evaluation of ITNs under controlled conditions that mirror those in which mosquitoes enter a human dwelling and face an ITN in normal use. Results from Experimental Hut Trials (EHTs) usually focus on two main indicators that are criteria for granting the WHO interim recommendation: the blood feeding inhibition (BFI, i.e. the reduction in blood-feeding rates relative to the control) and the mortality rates (proportion of dead mosquitoes). However, several other outputs that rely on counts of vectors collected in the huts are often neglected or analyzed with inappropriate statistical methods although they can provide useful information about vectors' behavior and personal protection provided by ITNs. In particular, EHTs allow to measure the deterrent effect of ITNs. The deterrence is defined as the reduction in the number of mosquitoes entering the treated hut relative to the control hut (untreated nets) [4]. This indicator is measured because some insecticides (e.g. the pyrethroids) are expected to repel malaria vectors at distance preventing their entrance in the dwellings. It is therefore expected that the deterrence will be null or positive. Although it is true for most of EHTs, negative deterrence values (i.e. more malaria *Anopheles* were collected in the treated hut than in the control hut) occurred sometimes. In a recent review studying the impact of pyrethroid resistance in malaria vectors on the efficacy of ITN [6], the authors provide 55 values of deterrence from 17 articles reporting results of EHTs. Thirteen (24%) of these values (from 7 articles) were negatives. In this latter review, in the concerned articles [7–13] and in a recently published study [14], the authors did not discuss much about the cause or origin of these surprising “attractiveness” of treated huts. This phenomenon may have significant consequences on the efficacy of ITNs in term of personal protection against malaria transmission.

In EHT, the personal protection is defined as the reduction in the number of blood-fed mosquitoes in the treatment hut relative to the number of blood-fed mosquitoes in the control hut [4]. However, this outcome that is greatly driven by the deterrent effect of ITNs, is almost totally overlooked with the current Phase II efficacy indicators analysis process. Indeed, although current guidelines recommend calculating the personal protection of ITNs, no statistical guidance is provided to state on its significance. To illustrate the importance of the deterrence on the estimation of the personal protection of a LLIN product, we address the relationships among the




Fig 1. Relationships among the blood-feeding inhibition (BFI), the deterrence and the personal protection as measured in experimental hut trials. See [Methods](#) for details on the mathematical relationship among BFI, deterrence and personal protection. Values of BFI and deterrence from studies cited in the review by Strode *et al.* [6] have been plotted when both indicators can be extracted from Fig 2 and Table 12 of this review article.

doi:10.1371/journal.pone.0170732.g001

BFI, the deterrence and the personal protection (see Fig 1): for a given value of BFI, the personal protection provided by an ITN could be either positive, null, or negative depending on the deterrence (see [Methods](#) for details on the mathematical relationship between the BFI, the deterrence and the personal protection).

Renewed interest for malaria eradication has placed greater emphasis on the development of new tools to target residual transmission (transmission that escape the control by conventional tools such as ITNs and IRS) and mosquito behavioral study are now in the spotlight [15–17]. The study of the remote effect (deterrence) and the personal protection confers by ITNs is of great importance as it might help identify weaknesses of ITNs that should be targeted by complementary vector control tools.

Therefore to provide a better assessment of ITNs used for malaria control, we performed a retrospective analysis of the deterrence and the personal protection against malaria transmission for 13 ITNs evaluated through EHTs. Trials were conducted in West Africa by Institut de Recherche pour le Développement (IRD) in the framework of the West African Anopheles, Biology and Control (ABC) network for testing and evaluation of pesticide products.

Methods

Calculation of the deterrence, the blood-feeding inhibition, and the personal protection (used to draw Fig 1)

The deterrence (D) is the reduction in hut entry relative to control huts (untreated nets):

$$D = 1 - \frac{N_t}{N_c} \tag{1}$$

With N_t the total number of mosquitoes collected in the treatment hut and veranda/exit traps and N_c the total number of mosquitoes in the control hut and veranda/exit traps.

The blood-feeding inhibition (BFI) is defined as “the reduction in blood-feeding in comparison with the control huts” [4]. Although it is not very clear from this definition, “blood-feeding” must be understood as “blood-feeding rate” (i.e. the proportion of blood-fed mosquitoes in the huts) but not as absolute number of blood-fed mosquitoes collected in the huts. The formula commonly used to calculate the BFI is:

$$BFI = 1 - \frac{P_t}{P_c} = 1 - \frac{B_t \times N_c}{N_t \times B_c} \tag{2}$$

With P_t the proportion of blood-fed mosquitoes in the treatment hut, P_c the proportion of blood-fed mosquitoes in the control hut, B_t the number of blood-fed mosquitoes in the treatment hut and B_c the number of blood-fed mosquitoes in the control hut.

The personal protection (PP) against transmission provided by a treatment in an experimental hut study is determined by the reduction in the number of blood-fed mosquitoes in the treatment hut relative to the number of blood-fed mosquitoes in the control hut:

$$PP = 1 - \frac{B_t}{B_c} \tag{3}$$

Relationship among PP, BFI, and D. From [expression 1](#), we deduce [expression 4](#):

$$\frac{N_c}{N_t} = \frac{1}{D - 1} \tag{4}$$

Given [expression 2](#), we can compute BFI by solving for [expression 4](#):

$$BFI = 1 + \frac{1}{D - 1} \times \frac{B_t}{B_c} \tag{5}$$

[Expression 5](#) is equivalent to:

$$\frac{B_t}{B_c} = (BFI - 1) \times (D - 1) \tag{6}$$

Given [expression 3](#), we can compute PP by solving for [expression 6](#):

$$PP = 1 - ((BFI - 1) \times (D - 1)) \tag{7}$$

Studies included in the analysis

WHOPES supervised EHTs (N = 10) involving 13 ITNs (12 long-lasting factory-impregnated nets and one long-lasting treatment kit (LLT) for manual impregnation) with raw data (daily

collections) available for subsequent statistical analyses were included in the analysis. These studies were carried out between 2006 and 2011 in two sites (Malanville, Northern Benin and the Kou Valley, Western Burkina Faso) according to the WHO guidelines [18]. A brief description of these trials is presented in the Table 1 and a summary of the WHOPES phase II experimental hut trial protocol is provided as a supplementary material (S1 Text). Among the 13 products tested, all were tested after 20 washes and 12 were tested unwashed [19–24].

The malaria vector population in the Malanville site (North of Benin) was composed at 95% by *An. coluzzii* (former M form) with a Kdr frequency (L1014F target-site mutation) that increased from 16% in 2008 to 50% in 2010 [25]. WHO cone bioassays indicated 85% and 93% mortality in 2008 [25] to the deltamethrin and permethrin insecticides, respectively. In 2010, mortality to deltamethrin decreased to 40% [25]. In the Kou Valley (North-West of Burkina Faso), the malaria vector population was composed at 85% by *An. gambiae* s.s. (former S form) and the Kdr frequency was 90% and the mortality rate induced by deltamethrin was 23% [8].

Statistical analysis

In order to assess the deterrence, we analyzed the daily numbers of malaria vectors entering the huts using a negative binomial mixed-effect model with all the treatment arms from the 10 EHTs and the study site (Malanville or Kou Valley) as fixed effects and with the trial and the day in the trial (to deal with daily variations of the mosquito density) as nested random effects (random intercepts). The model was written as follow:

$$\log(\mu_{dt}) = \beta_a^{Arm} + \beta_s^{Site} + a_t + a_{dt}$$

With μ_{dt} the number of anopheles entered a particular hut on day d of trial t . β_a^{Arm} is the effect on $\log(\mu_{dt})$ of classification in category a of the treatment arm and β_s^{Site} the effect of classification in category s (Malanville or Kou Valley) of the trial site. a_t is a random intercept for trial t and a_{dt} the random intercept for day d of trial t .

Using the same modelling approach, we assessed the personal protection by analyzing the number of blood-fed mosquitoes collected daily in the huts. We used the ‘R’ software [26] and the additional ‘glmmADMB’ [27] package for the analysis. Rates ratios (RRs) and 95% confidence intervals were computed.

Results

When compared to the untreated net, the number of *Anopheles* that entered the hut was lower for six of the 12 unwashed ITNs indicating a significant deterrent effect against malaria vectors (Fig 2A). For the 6 other ITNs, we were not able to detect any difference in the number of mosquito collected when compared to an UNTN. When washed 20 times, only three ITNs (Interceptor, Permanet 2.0 and Permanet 3.0) had significant deterrent effect (RRs < 1; $p < 0.05$; Fig 2B) and three others (Icon Maxx LLT, RR = 1.59 [1.15–2.19], $p = 0.0048$; Icon Maxx-Net, RR = 1.57 [1.14–2.16], $p = 0.0059$; and OlysetNet, RR = 1.7 [1.18–2.47], $p = 0.0046$) showed an apparent “attractiveness”. The 7 remaining ITNs did not show any difference with the untreated net ($p < 0.05$).

When compared to the untreated net, all unwashed ITNs showed lower number of blood-fed *Anopheles* indicating a significant personal protection (RR < 1, $p < 0.05$, Fig 2C). However when washed 20 times, the three ITNs that were found to be attractive did not significantly reduced human-vector contact when compared to an untreated net ($p > 0.05$; Fig 2D).

Table 1. Summary of the experimental hut trials included in the analysis.

Trial No.	WHOPES phase II trials				ITNs tested during the trials (0 and 20 washes)					Other arms of the trial (except the control untreated net)*	Control untreated net			
	Product evaluated	Site	Year	WHOPES reference	Duration (days)	Name	Type of ITN*	Fabric	Insecticide			Conc. (mg/m ²)	Impregnation method	2016 WHO recommendation
1	DawaPlus 1.0	Malanville	2006	[20]	60	DawaPlus 1.0	LN	polyester \$	Deltamethrin	40	coated	no (failed in phase II)	CTN 75 den 25 mg/m ² (0, 20 and until exh. washes), CTN 100 den (40 mg/m ² 20 washes and 25 mg/m ² washed until exh.), DawaPlus 1.0 100 den (0 and 20 washes)	untreated polyester net
2	DawaPlus 2.0	Malanville	2008	[22]	72	DawaPlus 1.0 # DawaPlus 2.0	LN	polyester \$	Deltamethrin	80	coated	Interim	CTN 25 mg/m ² (0 and until exh. washes)	untreated polyester net
3	DuraNet	Kou Valley	2007	[20]	25	DuraNet	LN	polyethylene	Alpha-cypermethrin	261	incorporated	Full	CTN 40 mg/m ² (0 and until exh. washes)	untreated net (same fabric and mesh size as DuraNet)
4	Icon Maxx and Icon Maxx-Net	Kou Valley	2007	[20,21]	35	Icon Maxx Icon Maxx-Net	LLT LN	polyester	Lambda-cyhalothrin Lambda-cyhalothrin	50 50	manual treatment kit coated	Full no (failed in phase II)	CTN 15 mg/m ² (0 and until exh. washes)	untreated polyester net
5	Interceptor	Malanville	2006	[19]	66	Interceptor	LN	polyester	Alpha-cypermethrin	200	coated	Full	CTN 40 mg/m ² (0 washes), CTN 200 mg/m ² (20 and until exh. washes)	untreated net (same fabric and mesh size as Interceptor)
6	Lifenet	Malaville	2011	[23]	72	Lifenet	LN	polypropylene	Deltamethrin	340	incorporated	Interim	CTN 25 mg/m ² (20 and until exh. washes), LifeNet washed 30 times	untreated polypropylene net.
7	NetProtect	Kou Valley	2007	[20]	25	NetProtect	LN	polyethylene	Deltamethrin	63	incorporated	no (interim until 2013; failed in phase II)	CTN 25 mg/m ² (0 and until exh. washes)	untreated net (polyethylene)
8	Olyset Plus	Malanville	2011	[24]	72	Olyset Net Olyset Plus	LN LN	polyethylene polyethylene	Permethrin Permethrin + PBO	1000 800	incorporated incorporated	Full Interim	CTN 25 mg/m ² washed until exh.	untreated polyester net.
9	Permanet 2.5 and Permanet 3.0	Malanville	2008	[21]	72	Permanet 2.0 Permanet 2.5 # Permanet 3.0	LN LN LN	polyester polyester polyester + polyethylene	Deltamethrin Deltamethrin Deltamethrin + PBO	55 115 115	coated coated incorporated + coated	Full no (interim until 2013) Interim	CTN 25 mg/m ² washed until exh.	Untreated net (same fabric and same design as of Permanet 3.0)

(Continued)

Table 1. (Continued)

Trial No.	WHOPES phase II trials				ITNs tested during the trials (0 and 20 washes)					Other arms of the trial (except the control untreated net)*	Control untreated net	
	Product evaluated	Site	Year	WHOPES reference	Duration (days)	Name	Type of ITN [§]	Fabric	Insecticide Conc. (mg/m ²)			Impregnation method
10	Permanet 3.0	Kou Valley	2008	[21]	36	Permanet 2.0 Permanet 3.0			see above see above			CTN 25 mg/m ² washed until exh. Untreated net (same fabric and same design as of Permanet 3.0)

[§] LN: Long-Lasting insecticidal net; LLT: "dip-it-yourself" treatment kit for converting mosquito nets into long-lasting insecticide-treated nets.

^{\$} linear mass density of 75 denier, the product exists in a 100 denier version

[#] this product was tested only when washed 20 times

* CTN: conventionnaly treated net, treated with the same insecticide than the evaluated product (concentration expressed in mg/m²); den = denier; exh: exhaustion;

doi:10.1371/journal.pone.0170732.t001




Fig 2. Deterrence (A,B) and personal protection (C,D) of unwashed (A,C) and washed (B,D) insecticidal treated nets evaluated through experimental hut trials in West Africa. Squares indicate Rate Ratios (with the control untreated net as reference) as obtained with Negative Binomial mixed effect models of daily counts of *Anopheles* entered the huts (A,B) and counts of blood-fed *Anopheles* (C,D). Error bars represent 95% confidence intervals of the rate ratios. LL: Long Lasting treatment for manual impregnation of the net.

doi:10.1371/journal.pone.0170732.g002

Discussion

This first analysis of the deterrence effect on personal protection of ITNs in experimental huts suggests that most, but not all of the WHO ITN recommended product tested are expected to provide personal protection against malaria transmission after 20 washes. Due to a negative deterrence effect, three ITN products did not show any significant personal protection against pyrethroid resistant malaria vectors after 20 consecutive washes. The three ITNs cause however greater killing effect on mosquito vectors than untreated nets [20,21,24].

Whatever the direction of the mosquito movement in presence of ITNs (deterrence versus attractiveness), this movement indicates that malaria vectors are able to detect the ITN at distance, before entering the hut. Deterrence of ITNs has been widely described in the last decades [28,29] because it allows reducing the vector density inside the dwellings fitted with ITN and therefore reducing the human-vector contact whether or not under the ITN. However, it is still unknown which volatiles are detected by mosquitoes. These volatiles could be the insecticide itself, additives, degradation products of these later, or products of the interaction among the insecticide, additives, CO₂ and human odors. Despite a low vapor pressure (i.e. a low volatility), pyrethroids have been found in the air around a treated net [30] at concentrations (0.000021–0.000038 mg/m³) that are considered negligible in terms of toxicity for

humans [31,32]. However, given the extraordinary sensitivity of the insects' olfactory system [33–35], we can reasonably suspect that such concentrations might be detected by mosquitoes. This field of investigation (i.e. chemical and behavioral ecology in a context of widespread vector control tool implementation) has been neglected for decades and there is a need for more behavioral and physiological studies.

In this study, 3 ITNs of 13 that used the permethrin or the lambda-cyhalothrin insecticides were found to be attractive for malaria vectors in pyrethroid resistance areas after 20 washes. We were not able to find the same trend with corresponding unwashed ITNs indicating a significant impact of washing on ITN deterrence. The performances of an ITN can be altered by washing. After 20 washes, the mortality is strongly reduced whatever the type of ITN [19–24] indicating a reduction of the concentration of available insecticide on the net [36]. The attraction of washed ITNs might therefore indicate an insecticide dose-dependent reversal effect of orientation behavior as it has been observed for *Anopheles gambiae* with human-derived putative repellents [37] and for *Aedes albopictus* with several carboxylic acids [38,39]. Because ITNs are rarely washed 20 times in their lifetime [40–42], the kinetic of active ingredients on the fiber in relation with behavioral responses of mosquitoes are urgently needed to understand better the effect of consecutive washing on ITNs deterrence.

It should be noted that the untreated (control) net used in trial 8 (Table 1) was a polyester net, a different fabric and mesh size than the evaluated Olyset Net. To our knowledge, there is no study that address the role of net fabrics and mesh sizes of nets on human odor and CO₂ dispersion. However, we cannot exclude that wide mesh ITNs (as Olyset Net [24]) allowed a better dispersion of human odor and CO₂ than nets having smaller mesh size. The role of mesh size in the dispersion of odors and volatile substances would merit further investigations.

The impact of the physiological resistance to insecticide in the host-seeking behavior has been overlooked for decades. Recent findings from our team [43] showed that a lab strain of *An. gambiae* homozygous for the *kdr-w* mutation (L1014F) was significantly attracted by an animal host + permethrin treated net odor plume. Studies are ongoing to investigate the impact of other mutations and metabolic mechanisms conferring resistance to public health insecticides. Both the *An. gambiae* populations from Malanville and Kou Valley carried the *kdr-w* mutation [8,25] among other resistance mechanisms. We suspect that resistance mechanisms might modulate the host-seeking behavior by leading to the attraction of some Anopheles vectors when permethrin treated ITNs are drastically washed. Studies are ongoing to investigate the impact of other mutations and metabolic mechanisms on the behavior of mosquitoes in presence of both human host and ITNs.

We showed that three ITNs having a significant attractive effect did not provide a better personal protection than UTNs. In this particular condition, individual benefits of using these ITNs instead of an UTN (provided the UTN is maintained in good condition and is sufficiently large so that the sleeper do not make contact with it) would appeared to be null [28,44]. However, as shown in Fig 1, attractiveness would induce null or negative personal protection only for nets exhibiting a BFI rate lower than 50%.

The effect of attraction on community protection cannot be assessed precisely with EHTs data. Indeed, washed ITNs that we found to be attractive were efficient to kill an important number of mosquitoes [20,21,24] contributing to the reduction of the adult density and the lifespan of the local population of vectors. However theoretically [45], the community protection provided by an intra-domiciliary vector control tool is highly dependent on the coverage of the intervention (i.e. the proportion of people that use it) that cannot be simulated in EHTs. It is therefore impossible to conclude that the attractive property might have an effect (either positive or negative) on the community protection based on EHT outputs.

The best way to evaluate the community effect of ITNs against transmission should be to monitor and compare EIRs, malaria prevalence and incidence through a phase III Randomized Control Trial (RCT) with a negative control arm (untreated net). Nevertheless, since ITNs are now the baseline intervention for most of NMCP, the use of untreated nets as a negative control raises ethical issues. Alternatively, as compliance with ITNs is never 100%, a parasitological and clinical follow-up of non-users after the distribution of LLIN should help to measure the community effect. Community level trials are costly and time consuming and therefore the use of mathematical models of transmission using EHT data showed useful to predict community protection induced by LLIN. Such models exist and have been used to compare the potential efficacy of insecticide products having or not a repellent effect [46]. The authors of the later study found that purely toxic products with no deterrence are predicted to generally provide superior protection to non-users and even users, even if that product confers no personal protection. By extrapolation of Killeen and colleagues' results [46], we could expect that attractive products might induce superior community protection than deterrent ones. However, according to Okumu *et al.* [47] who adapted this model to be used with EHT data, the model does not allow to deal with negative deterrence. Therefore, as a first step before community-level trials, simulations using mathematical models of transmission adapted to allow for attractive product (for example, an adaptation of the modelling approach recently published by Churcher *et al.* [48] that used EHT data to predict the impact of insecticide resistance on malaria infection) should be run to evaluate the effect of an attractive ITN at the community level. If the community effect might be confirmed, it is important to note that products which confer low or no personal protection will require adapted awareness campaign that emphasize the communal nature of protection [46].

Conclusion

Current WHO efficacy criteria do not take into account the deterrence and the deterrence variability is neither analyzed nor discussed in the majority of the reports of experimental hut studies as illustrated in a recent literature review [6]. Consequently, there is an important gap of knowledge with unknown consequences in terms of public health. Our study points the long range effect (repellent or attractive) of ITNs, the personal protection and above all, the community protection out to be major criteria for the evaluation of ITNs.

Supporting Information

S1 Text. Summary of the WHOPES Experimental Hut Trial protocol.
(DOCX)

Acknowledgments

These Phase II trials and the present study have been run in the frame of ABC network.

Author Contributions

Conceptualization: NM CP.

Data curation: NM CP.

Formal analysis: NM.

Investigation: NM CP FC VC JMH.

Methodology: NM CP.

Project administration: NM CP.

Visualization: NM.

Writing – original draft: NM CP.

Writing – review & editing: NM CP FC VC JMH.

References

1. World Health Organization. World malaria report 2015 [Internet]. Geneva: World Health Organization; 2015 [cited 2016 Sep 6] p. 243. <http://www.who.int/iris/handle/10665/200018>
2. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature*. 2015 Sep 16;
3. World Health Organization. WHO recommended long-lasting insecticidal nets [Internet]. 2016 [cited 2016 Sep 8]. http://www.who.int/whopes/Long-lasting_insecticidal_nets_April_2016.pdf
4. WHO. Guidelines for laboratory and field-testing of long-lasting insecticidal nets. World Health Organization; 2013 p. 89. Report No.: WHO/HTM/NTD/WHOPES/2013.3.
5. WHO. Guidelines for monitoring the durability of long-lasting insecticidal mosquito nets under operational conditions. World Health Organization; 2011. Report No.: WHO/HTM/NTD/WHOPES/2011.5.
6. Strode C, Donegan S, Garner P, Enayati AA, Hemingway J. The Impact of Pyrethroid Resistance on the Efficacy of Insecticide-Treated Bed Nets against African Anopheline Mosquitoes: Systematic Review and Meta-Analysis. *PLoS Med*. 2014 Mar; 11(3):e1001619. doi: [10.1371/journal.pmed.1001619](https://doi.org/10.1371/journal.pmed.1001619) PMID: [24642791](https://pubmed.ncbi.nlm.nih.gov/24642791/)
7. Chandre F, Darriet F, Duchon S, Finot L, Manguin S, Carnevale P, et al. Modifications of pyrethroid effects associated with *kdr* mutation in *Anopheles gambiae*. *Med Vet Entomol*. 2000; 14(1):81–8. PMID: [10759316](https://pubmed.ncbi.nlm.nih.gov/10759316/)
8. Corbel V, Chabi J, Dabire RK, Etang J, Nwane P, Pigeon O, et al. Field efficacy of a new mosaic long-lasting mosquito net (PermaNet 3.0) against pyrethroid-resistant malaria vectors: a multi centre study in Western and Central Africa. *Malar J*. 2010; 9:113. doi: [10.1186/1475-2875-9-113](https://doi.org/10.1186/1475-2875-9-113) PMID: [20423479](https://pubmed.ncbi.nlm.nih.gov/20423479/)
9. Fanello C, Kolaczinski JH, Conway DJ, Carnevale P, Curtis CF. The *kdr* pyrethroid resistance gene in *Anopheles gambiae*: tests of non-pyrethroid insecticides and a new detection method for the gene. *Parassitologia*. 1999 Sep; 41(1–3):323–6. PMID: [10697877](https://pubmed.ncbi.nlm.nih.gov/10697877/)
10. Okumu FO, Mbeyela E, Lingamba G, Moore J, Ntamatungiro AJ, Kavishe DR, et al. Comparative field evaluation of combinations of long-lasting insecticide treated nets and indoor residual spraying, relative to either method alone, for malaria prevention in an area where the main vector is *Anopheles arabiensis*. *Parasit Vectors*. 2013; 6:46. doi: [10.1186/1756-3305-6-46](https://doi.org/10.1186/1756-3305-6-46) PMID: [23433393](https://pubmed.ncbi.nlm.nih.gov/23433393/)
11. Ngufor C, N'Guessan R, Boko P, Odjo A, Vigninou E, Asidi A, et al. Combining indoor residual spraying with chlorfenapyr and long-lasting insecticidal bed nets for improved control of pyrethroid-resistant *Anopheles gambiae*: an experimental hut trial in Benin. *Malar J*. 2011; 10:343. doi: [10.1186/1475-2875-10-343](https://doi.org/10.1186/1475-2875-10-343) PMID: [22087506](https://pubmed.ncbi.nlm.nih.gov/22087506/)
12. Malima RC, Oxborough RM, Tungu PK, Maxwell C, Lyimo I, Mwingira V, et al. Behavioural and insecticidal effects of organophosphate-, carbamate- and pyrethroid-treated mosquito nets against African malaria vectors. *Med Vet Entomol*. 2009 Dec; 23(4):317–25. doi: [10.1111/j.1365-2915.2009.00837.x](https://doi.org/10.1111/j.1365-2915.2009.00837.x) PMID: [19941597](https://pubmed.ncbi.nlm.nih.gov/19941597/)
13. Malima RC, Magesa SM, Tungu PK, Mwingira V, Magogo FS, Sudi W, et al. An experimental hut evaluation of Olyset nets against anopheline mosquitoes after seven years use in Tanzanian villages. *Malar J*. 2008; 7:38. doi: [10.1186/1475-2875-7-38](https://doi.org/10.1186/1475-2875-7-38) PMID: [18307802](https://pubmed.ncbi.nlm.nih.gov/18307802/)
14. Ngufor C, N'Guessan R, Fagbohoun J, Todjinou D, Odjo A, Malone D, et al. Efficacy of the Olyset Duo net against insecticide-resistant mosquito vectors of malaria. *Sci Transl Med*. 2016 Sep 14; 8(356):356ra121. doi: [10.1126/scitranslmed.aad3270](https://doi.org/10.1126/scitranslmed.aad3270) PMID: [27629488](https://pubmed.ncbi.nlm.nih.gov/27629488/)
15. WHO. Guidance note on the control of residual malaria parasite transmission [Internet]. 2014. Report No.: WHO/HTM/GMP/MPAC/2014.5. <http://www.who.int/entity/malaria/publications/atoz/technical-note-control-of-residual-malaria-parasite-transmission-sep14.pdf>
16. Durnez L, Coosemans M. Residual Transmission of Malaria: An Old Issue for New Approaches. In: Manguin S, editor. *Anopheles mosquitoes—New insights into malaria vectors* [Internet]. InTech; 2013 [cited 2015 Aug 4]. <http://www.intechopen.com/books/anopheles-mosquitoes-new-insights-into-malaria-vectors/residual-transmission-of-malaria-an-old-issue-for-new-approaches>
17. Killeen GF. Characterizing, controlling and eliminating residual malaria transmission. *Malar J*. 2014; 13(1):330.

18. WHO. Guidelines for laboratory and field-testing of long-lasting insecticidal mosquito nets. 2005. Report No.: WHO/CDS/WHOPES/GCDPP/2005.11.
19. WHO Pesticide Evaluation Scheme. Report of the tenth WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2007 [cited 2016 Aug 19]. <http://www.who.int/iris/handle/10665/69465>
20. WHO Pesticide Evaluation Scheme. Report of the eleventh WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2008 [cited 2016 Aug 19]. Report No.: WHO/HTM/NTD/WHOPES/2008.1. <http://www.who.int/iris/handle/10665/69732>
21. WHO Pesticide Evaluation Scheme. Report of the twelfth WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2009 [cited 2016 Aug 19]. <http://www.who.int/iris/handle/10665/69986>
22. WHO Pesticide Evaluation Scheme. Report of the thirteenth [13th] WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2009 [cited 2016 Aug 19]. <http://www.who.int/iris/handle/10665/44212>
23. WHO Pesticide Evaluation Scheme. Report of the fourteenth [14th] WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2011 [cited 2016 Aug 19]. Report No.: WHO/HTM/NTD/WHOPES/2011.7. <http://www.who.int/iris/handle/10665/44669>
24. WHO Pesticide Evaluation Scheme. Report of the fifteenth WHOPES working group meeting [Internet]. Geneva: World Health Organization; 2012 [cited 2016 Aug 19]. <http://www.who.int/iris/handle/10665/75304>
25. Djegbe I, Boussari O, Sidick A, Martin T, Ranson H, Chandre F, et al. Dynamics of insecticide resistance in malaria vectors in Benin: first evidence of the presence of L1014S kdr mutation in *Anopheles gambiae* from West Africa. *Malar J*. 2011; 10(1):261.
26. R Development Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2010. <http://www.R-project.org>
27. Skaug H, Fournier D, Nielsen A, Magnusson A, Bolker B. glmmADMB: Generalized Linear Mixed Models Using AD Model Builder [Internet]. 2012. <http://R-Forge.R-project.org/projects/glmmadmb/>
28. Takken W. Do insecticide-treated bednets have an effect on malaria vectors? *Trop Med Int Health*. 2002; 7(12):1022–30. PMID: 12460393
29. Lindsay SW, Adiamah JH, Miller JE, Armstrong JR. Pyrethroid-treated bednet effects on mosquitoes of the *Anopheles gambiae* complex in The Gambia. *Med Vet Entomol*. 1991 Oct; 5(4):477–83. PMID: 1773125
30. Bomann W. How safe are pyrethroid-treated mosquito nets? An evaluation based on the example of Solfac EW 050. *Public Health*. 1995; 12:30–5.
31. Barlow SM, Sullivan FM, Lines J. Risk assessment of the use of deltamethrin on bednets for the prevention of malaria. *Food Chem Toxicol*. 2001 mai; 39(5):407–22. PMID: 11313107
32. World Health Organization. A generic risk assessment model for insecticide-treated nets: revised edition. 2012 [cited 2016 Aug 19]; <http://www.who.int/iris/handle/10665/44862>
33. Angioy AM, Desogus A, Barbarossa IT, Anderson P, Hansson BS. Extreme Sensitivity in an Olfactory System. *Chem Senses*. 2003 May 1; 28(4):279–84. PMID: 12771014
34. Baker TC, Roelofs WL. Initiation and Termination of Oriental Fruit Moth Male Response to Pheromone Concentrations in the Field. *Environ Entomol*. 1981 avril; 10(2):211.
35. Clements AN. The Biology of Mosquitoes. Volume 2: Sensory, Reception, and Behaviour. First edition. Wallingford u.a.: CABI; 1999. 752 p.
36. Pennetier C, Bouraima A, Chandre F, Piamou M, Etang J, Rossignol M, et al. Efficacy of Olyset® Plus, a New Long-Lasting Insecticidal Net Incorporating Permethrin and Piperoniol-Butoxide against Multi-Resistant Malaria Vectors. *PLoS One*. 2013; 8(10):e75134. doi: 10.1371/journal.pone.0075134 PMID: 24116029
37. Logan JG, Stanczyk NM, Hassanali A, Kemei J, Santana AE, Ribeiro KA, et al. Arm-in-cage testing of natural human-derived mosquito repellents. *Malar J*. 2010; 9:239. doi: 10.1186/1475-2875-9-239 PMID: 20727149
38. Guha L, Seenivasagan T, Iqbal ST, Agrawal OP, Parashar BD. Behavioral and electrophysiological responses of *Aedes albopictus* to certain acids and alcohols present in human skin emanations. *Parasitol Res*. 2014 Jul 23; 113(10):3781–7. doi: 10.1007/s00436-014-4044-0 PMID: 25049052
39. Seenivasagan T, Guha L, Parashar BD, Agrawal OP, Sukumaran D. Olfaction in Asian tiger mosquito *Aedes albopictus*: flight orientation response to certain saturated carboxylic acids in human skin emanations. *Parasitol Res*. 2014 Mar 12; 113(5):1927–32. doi: 10.1007/s00436-014-3840-x PMID: 24619069

40. Mutuku FM, Khambira M, Bisanzio D, Mungai P, Mwanzo I, Muchiri EM, et al. Physical condition and maintenance of mosquito bed nets in Kwale County, coastal Kenya. *Malar J.* 2013; 12(1):46.
41. Erlanger TE, Enayati AA, Hemingway J, Mshinda H, Tami A, Lengeler C. Field issues related to effectiveness of insecticide-treated nets in Tanzania. *Med Vet Entomol.* 2004 Jun; 18(2):153–60. doi: [10.1111/j.0269-283X.2004.00491.x](https://doi.org/10.1111/j.0269-283X.2004.00491.x) PMID: [15189240](https://pubmed.ncbi.nlm.nih.gov/15189240/)
42. Lindblade KA, Dotson E, Hawley WA, Bayoh N, Williamson J, Mount D, et al. Evaluation of long-lasting insecticidal nets after 2 years of household use. *Trop Med Int Health.* 2005 Nov; 10(11):1141–50. doi: [10.1111/j.1365-3156.2005.01501.x](https://doi.org/10.1111/j.1365-3156.2005.01501.x) PMID: [16262739](https://pubmed.ncbi.nlm.nih.gov/16262739/)
43. Porciani A, Diop MM, Moiroux N, Kadoke-Lambi T, Cohuet A, Chandre F, et al. A pyrethroid-treated bed net increases host attractiveness for *Anopheles gambiae* s.s carrying the *kdr* allele in a dual choice olfactometer. *bioRxiv [Internet].* 2016 Sep 26;
44. Guyatt HL, Snow RW. The cost of not treating bednets. *Trends Parasitol.* 2002 Jan; 18(1):12–6. PMID: [11850008](https://pubmed.ncbi.nlm.nih.gov/11850008/)
45. Killeen GF, Smith TA, Ferguson HM, Mshinda H, Abdulla S, Lengeler C, et al. Preventing childhood malaria in Africa by protecting adults from mosquitoes with insecticide-treated nets. *PLoS Med.* 2007 Jul; 4(7):e229. doi: [10.1371/journal.pmed.0040229](https://doi.org/10.1371/journal.pmed.0040229) PMID: [17608562](https://pubmed.ncbi.nlm.nih.gov/17608562/)
46. Killeen GF, Chitnis N, Moore SJ, Okumu FO. Target product profile choices for intra-domiciliary malaria vector control pesticide products: repel or kill? *Malar J.* 2011; 10:207. doi: [10.1186/1475-2875-10-207](https://doi.org/10.1186/1475-2875-10-207) PMID: [21798023](https://pubmed.ncbi.nlm.nih.gov/21798023/)
47. Okumu FO, Kiware SS, Moore SJ, Killeen GF. Mathematical evaluation of community level impact of combining bed nets and indoor residual spraying upon malaria transmission in areas where the main vectors are *Anopheles arabiensis* mosquitoes. *Parasit Vectors.* 2013; 6:17. doi: [10.1186/1756-3305-6-17](https://doi.org/10.1186/1756-3305-6-17) PMID: [23324456](https://pubmed.ncbi.nlm.nih.gov/23324456/)
48. Churcher TS, Lissenden N, Griffin JT, Worrall E, Ranson H. The impact of pyrethroid resistance on the efficacy and effectiveness of bednets for malaria control in Africa. *eLife.* 2016 Aug 22; 5.