



HAL
open science

Coagulation and Mental Disorders

Silvia Hoirisch-Clapauch, Antonio Egidio Nardi, Jean-Christophe Gris,
Benjamin Brenner

► **To cite this version:**

Silvia Hoirisch-Clapauch, Antonio Egidio Nardi, Jean-Christophe Gris, Benjamin Brenner. Coagulation and Mental Disorders. Rambam Maimonides Medical Journal, 2014, 5 (4), pp.e0036. 10.5041/RMMJ.10170 . hal-01940134

HAL Id: hal-01940134

<https://hal.umontpellier.fr/hal-01940134v1>

Submitted on 13 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Special Issue on Advances in Hematology

Guest Editor: Benjamin Brenner, M.D.

Coagulation and Mental Disorders

Silvia Hoirisch-Clapauch, M.D.^{1*}, Antonio Egidio Nardi, M.D., Ph.D.², Jean-Christophe Gris, M.D., Ph.D.³, and Benjamin Brenner, M.D.⁴

¹Department of Hematology, Hospital Federal dos Servidores do Estado, Ministry of Health, Rio de Janeiro, Brazil; ²Institute of Psychiatry, Federal University of Rio de Janeiro, National Institute for Translational Medicine (INCT-TM), Brazil; ³Department of Haematology, Centre Hospitalier Universitaire, Nîmes, France; ⁴Department of Hematology and Bone Marrow Transplantation, Rambam Health Care Campus and Technion Institute of Technology, Haifa, Israel

ABSTRACT

The neurovascular unit is a key player in brain development, homeostasis, and pathology. Mental stress affects coagulation, while severe mental illnesses, such as recurrent depression and schizophrenia, are associated with an increased thrombotic risk and cardiovascular morbidity. Evidence indicates that the hemostatic system is involved to some extent in the pathogenesis, morbidity, and prognosis of a wide variety of psychiatric disorders. The current review focuses on emerging data linking coagulation and some psychiatric disorders.

KEY WORDS: Anxiety, cardiovascular diseases, depression, mental disorders, schizophrenia, thrombosis

INTRODUCTION

The human brain critically relies on an elaborate vascular network, containing up to 100 billion vessels, a vessel for each neuron. Apart from the oxygen supply function, endothelial cells also produce instructive signals for neural development, support normal functioning, ensure maintenance, and promote reparative regeneration of the central nervous system. The brain vasculature is increasingly recognized as a key player that actively influences and directs brain development, homeostasis, and

Abbreviations: APLA, antiphospholipid antibodies; PAI, plasminogen activator inhibitor.

Citation: Hoirisch-Clapauch S, Nardi AE, Gris JC, Brenner B. Coagulation and Mental Disorders. Rambam Maimonides Med J 2014;5 (4):e0036. doi:10.5041/RMMJ.10170

Copyright: © 2014 Hoirisch-Clapauch et al. This is an open-access article. All its content, *except where otherwise noted*, is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Conflict of interest: No potential conflict of interest relevant to this article was reported.

* To whom correspondence should be addressed. **E-mail:** sclapauch@ig.com.br

pathology.¹ One of the major challenges of the century is to improve the understanding and treatment of psychiatric diseases. Among the physiologic systems which may be influenced by, and/or may influence, the neurovascular unit is the hemostatic system. Elucidating the cross-talk between hemostasis, thrombosis, and the fibrinolytic system and the nervous system is crucial to understand better the molecular and cellular basis of how the brain operates, but also to develop novel diagnostic and therapeutic strategies for mental illnesses. We herein review the emerging knowledge linking hemostasis and some psychiatric disorders.

ANXIETY DISORDERS

The same stress responses that were essential in ancient times to protect the body against fatal bleeding can be elicited by emotional stimuli or professional and social stress. Fear and anxiety are important responses to stress, mediated by several neurotransmitters and hormones, including epinephrine, norepinephrine, serotonin, acetylcholine, dopamine, prolactin, and cortisol. While fear is the emotional response to real or perceived imminent threat, anxiety is anticipation of future threat.²

Catecholamines have a prominent role in the fight-or-flight reaction. In response to catecholamine and also to serotonin stimulus, platelets manifest a series of functional responses, such as adhesion, secretion, and aggregation.³ Catecholamines, particularly epinephrine, bind to the β_2 -adrenoreceptor, resulting in release of factor VIII, von Willebrand factor, and tissue plasminogen activator. Importantly, tissue plasminogen activator is expressed in chromaffin cells, and co-released with catecholamines by chromaffin cell stimulation.⁴ The simultaneous activation of coagulation and fibrinolysis during acute stress can be demonstrated by increased levels of D-dimer.³

Through α -adrenoreceptor activation, catecholamine induces vasoconstriction. During vasoconstriction plasma, but not large plasma proteins, passes easily into interstitial spaces through minute openings or pores in the capillaries. The presence of large molecules creates a large colloidal osmotic pressure gradient⁵ that increases viscosity, which contributes to a hypercoagulable state. Factor VIII activity remains elevated, even after saline reconstitution, suggesting that the intrinsic pathway is genuinely activated during acute stress.³

Anxiety disorders differ from transient fear or anxiety by an excessive response to stress, with a typically chronic course.² In severe post-traumatic stress disorder, for example, although D-dimer levels might be normal, von Willebrand antigen and factor VIII activity are usually elevated, and platelets exhibit exaggerated reactivity to *in vitro* epinephrine/ADP stimulation mediated by the α_2 -adrenergic receptor.^{3,6} Opposed to acute stress, chronic stress seems to impair fibrinolytic activity.³ Abnormal fibrinolysis is reflected by increased synthesis of plasminogen activator inhibitor (PAI)-1, a major inhibitor of tissue plasminogen activator, probably as a result of serotonin and cortisol secretion.^{7,8} Angiotensin potentiates the production of PAI-1 stimulated by steroids,⁹ which provides a reasonable explanation for the increased thrombotic tendency and cardiovascular risk seen in patients who are both anxious and hypertensive.^{3,10}

DEPRESSIVE DISORDERS

Different from sadness, which is a normal response to loss, depressive disorders are disabling illnesses that may present with psychotic symptoms, such as delusions or hallucinations, and frequently co-occur with lifetime anxiety.¹¹ They include major depressive disorder and many other subtypes, such as dysthymia and premenstrual dysphoric disorder. Major depressive disorder may present with melancholic or with atypical features.² Melancholia is characterized by diurnal variation of depressive symptoms, usually worse in the morning, by insomnia and anorexia, and by a hyperactive stress response. Patients with melancholia are usually anxious, less responsive to the environment, and they dread the future.

Major depression accompanied by atypical features is defined as mood improvement in response to positive events, plus at least two of the following four symptoms: significant weight gain or hyperphagia, hypersomnia, severe lethargy, also described as heavy, leaden feelings in arms or legs, and pathological rejection sensitivity. Whereas patients with melancholia usually have increased levels of corticotrophin-releasing hormone and diminished activities of the growth hormone and reproductive axes, in atypical depression hypothalamic-pituitary-adrenal is typically down-regulated and corticotrophin-releasing hormone is deficient.¹²

Serotonin seems to play a major role in the pathophysiology of depression. More than 99% of the circulating serotonin is found in platelet-dense granules,¹³ and serotonin itself is a platelet agonist. In patients with recurrent depressive disorders, platelet studies have shown increased serotonin response, increased serotonin receptor density, decreased serotonin transporter binding, blunting of adenosine response, increased thrombin response, increased expression of glycoprotein 1b, P-selectin, β thromboglobulin, platelet factor 4, fibrinogen, factor V, and anionic phospholipids, as well as decreased platelet content of brain-derived neurotrophic factor and serotonin.^{14,15} Clot firmness and procoagulant activity of platelet-associated tissue factor may also be significantly elevated. Additionally, studies with circulating blood revealed increased fibrin formation in early diagnosed patients.¹⁵

Abnormalities in the fibrinolytic pathway can be also observed in depressed subjects. Because PAI-1 promoter has cortisol and insulin response, depressed individuals may have elevated levels of PAI-1 activity, even after adjusting for confounders, such as coronary heart disease, smoking, hypertension, triglyceride concentration, and body mass index.¹⁶ Also, serotonin has been shown to increase PAI-1 levels in endothelial cells.¹⁷

SCHIZOPHRENIA

Compared to controls, first-episode psychosis patients display higher levels of markers of thrombogenesis, such as D-dimer and factor VIII, and markers of platelet activation, such as soluble P-selectin.¹⁸ Additionally, first-episode drug-naïve schizophrenia patients may exhibit higher cortisol and insulin levels than healthy subjects, as well as a significantly higher insulin resistance score.¹⁹ The PAI-1 promoter has insulin and cortisol response, and it is possible that elevated PAI-1 levels could contribute to the increased cardiovascular and thrombotic risk seen in schizophrenia patients.

A high prevalence of free-protein S deficiency has been demonstrated in a sample of schizophrenia patients. Fifteen out of 70 (22%) and none of 98 controls showed low levels of the natural anticoagulant. Except for one patient with periodontitis, no patient had clinical or laboratory evidence of inflammation. Even though it was not determined in this study if deficiency was inherited or acquired, low levels of free-protein S increased the chances of having a first-degree relative with schizophrenia by

145 times compared to controls, which suggests that, at least in some of the patients, protein S deficiency could have been inherited.²⁰ Aside from having anti-coagulant activity, free-protein S accelerates neutralization of PAI-1, and therefore increases fibrin clot lysis. Moreover, it also affords neuroprotection.²¹

REPERCUSSION OF MENTAL DISORDERS ON CARDIOVASCULAR RISK

Generalized anxiety disorder, post-traumatic stress disorder, and panic disorder are strongly associated with increased risk for cardiovascular morbidity and mortality.^{22–25} A bidirectional relationship exists between depression and cardiovascular disease. Depression increases the risk for cardiovascular disorders such as coronary artery disease and myocardial infarction 1.5–2-fold and for stroke 1.8-fold.²⁶ Likewise, in patients with coronary heart disease, depression is independently associated with cardiovascular morbidity and mortality.²⁷ Besides susceptibility to blood clotting, various pathophysiological mechanisms may underlie the risk of cardiovascular disease in patients with depression, including oxidative stress, hypothyroidism, hyperactivity of the sympatho-adrenomedullary system and the hypothalamic-pituitary-adrenal axis, and reduction in arterial repair processes.²⁶ Different studies suggest an increased risk for coronary artery disease not only in patients with major depression but also in those with depressive symptoms and dysphoria. In these studies, the magnitude of risk seems to follow a graded effect, with increasing risk for coronary artery disease among patients with more severe depression.²⁸

Patients with bipolar disorders are known to be at high risk of premature death. A systematic review of recently published articles suggested that a multisystemic inflammatory involvement is present from the early stages, and a dysregulated inflammatory background seems to be a common factor underlying cardiovascular and bipolar disorders, with hemostasis activation paying tribute to immuno-inflammation.²⁹

The increased risk for cardiovascular disease observed in schizophrenia patients has been related to inactivity, dietary habits, hyperhomocysteinemia, the high rate of substance abuse, including tobacco and illicit drug use, and alcoholism. Other risk factors may contribute to the increased risk for cardiovascular events. For example, insulin

resistance, a common psychotropic side effect, may cause obesity. Additionally periodontitis is highly prevalent in schizophrenia patients due to both heavy tobacco smoking and inadequate oral hygiene. Obesity and periodontitis may cause systemic inflammation, which in turn may reduce protein S activity and increase PAI-1 levels, thus increasing thrombotic and cardiovascular risk. Increased cardiovascular risk is already observed in first-episode schizophrenia.^{30,31}

REPERCUSSION OF MENTAL DISORDERS ON THROMBOTIC RISK

Patients with pulmonary hypertension have more panic disorders than controls.³² Based on the assumption that recurrent showers of small emboli, which may give rise to pulmonary hypertension, are commonly undetected clinically, it is possible that some episodes of panic attacks might accompany undetected episodes of pulmonary thromboembolism.

Both thromboembolism and depression are common complications of neoplastic disorders. In patients with recurrent depressive disorders without neoplastic disorders, however, despite platelet activation and abnormal fibrinolysis,³³ there is no evidence that depression per se increases the risk of thromboembolism. There are three possibilities. First, there have been no specific studies assessing the incidence of venous thrombosis in depressed patients. Second, serotonergic antidepressants may impair platelet function and decrease PAI-1 levels, therefore reducing the thrombotic tendency.³⁴ Another explanation is that even though plasma levels of interleukin-6 and tumor necrosis factor- α may be increased in patients with depression,³⁵ these patients often display features of immunosuppression, such as accelerated apoptosis of both T lymphocytes and neutrophils.³⁶

Schizophrenia patients may also be at increased risk of thromboembolic events. Thrombotic tendency has been usually associated with psychotropic medication and with immobility, as in restraint or catatonia. In a study conducted in restrained psychotic patients, the incidence of deep vein thrombosis was 12% in spite of prophylaxis with graduated compression stockings and subcutaneous injection of unfractionated heparin.³⁷ The finding of high levels of thrombogenesis markers and platelet activation in first-episode psychosis patients suggests that mechanisms involved in the

pathogenesis of psychosis might also contribute to the thrombotic tendency.¹⁸

ASSOCIATION OF THROMBOTIC RISK MARKERS WITH MENTAL DISORDERS

Among all classical thrombophilias, i.e. abnormalities of blood coagulation that enhance the thrombotic risk, only an acquired one, namely antiphospholipid antibodies (APLA), has been substantially described in patients with severe mental disorders.³⁸

Associations with hyperactivity and behavioral abnormalities were reported in mice exposed to human APLA, as well as in mice immunized with a major APLA cofactor, β 2 glycoprotein 1. Human APLA can directly permeabilize and depolarize brain synaptosomes, with histological examination showing antibody binding to both neuronal and white matter. Immunogold electron microscopy revealed APLA reacting with myelin, mainly the major dense line formed by the cytoplasmic apposition of the oligodendrocyte plasma membrane. The effects of APLA on brain endothelial cells may influence inflammatory and hypercoagulation effects on the brain tissues. Antiphospholipid antibody-mediated endothelial cell activation is likely to disrupt the blood-brain barrier, facilitating the diffusion of pathogenic antibodies, coagulation factors, and inflammatory factors into the brain tissues. Direct effects of autoantibodies on various brain cellular components may induce the behavioral changes observed in the mice passively or actively developing the antiphospholipid syndrome.

Case reports of bipolar disorders, of schizoaffective disorder, and of schizophrenia have been convincingly argued in APLA-positive patients. In some cases, the acute onset of psychiatric symptoms even revealed a significantly sustained positivity for APLA. Some reports have also suggested that APLA may emerge as a response to antipsychotic treatment, since a high prevalence of the antibodies in antipsychotic users has been observed. A recent cross-sectional study determined the presence and a subtype of APLA in 333 psychiatric inpatients: APLA were frequently found positive, and prevalence of the antibodies was not significantly different in patients who were receiving, or not receiving, antipsychotics.³⁹ Long-term prospective epidemiological cohort studies are still needed to analyze possible pathogenic implications of persistent APLA in psychiatric patients, such as worsening of the clinical symptomatology.

TREATMENT OF MENTAL DISORDERS MAY AFFECT COAGULATION

In patients with anxiety and depressive disorders, factor VII and plasminogen activator inhibitor levels normalize after psychotherapy, indicating that improvement of psychiatric symptoms somehow reverses the procoagulant effect of these psychological states.⁴⁰ The same may occur with pharmacological treatment. Of note, compared to non-serotonergic antidepressants, serotonergic antidepressants may decrease the risk of arterial occlusive events, such as myocardial infarction, but may also increase the risk for abnormal bleeding, which includes perioperative, gastrointestinal, and brain hemorrhage.³⁴ The increased bleeding risk observed in some patients on serotonergic antidepressants has been related to platelet and fibrinolytic abnormalities. Platelets of serotonergic antidepressant-medicated patients display a lower serotonin content and lower ADP, collagen, or epinephrine-induced aggregation.⁴⁰ Furthermore, patients on serotonergic antidepressants appear to have fibrinogen and PAI-1 plasma levels that are similar to those of healthy controls, but lower than in depressed patients receiving non-serotonergic antidepressants.⁴¹

In a group of psychotic patients, plasma levels of soluble P-selectin varied significantly in the course of 1-year antipsychotic treatment, mainly between 3 and 6 months after therapy was started, but plasma levels of D-dimer and factor VIII remained elevated.¹⁸ Antipsychotics, especially clozapine and olanzapine, may promote weight accrual and increase the levels of insulin and triglycerides.⁴² Fat tissue stroma synthesizes PAI-1, and both insulin and triglycerides provide stimulus for PAI-1 synthesis, which may increase cardiovascular risk.

Schizophrenia patients with hyperhomocysteinemia might benefit from B-vitamin supplementation: when these patients were treated with folic acid, B₁₂, and pyridoxine, clinical symptoms as measured by the Positive and Negative Syndrome Scale declined significantly.⁴³ Considering that hyperhomocysteinemia is an independent risk factor for both cardiovascular disease and thromboembolism, it remains to be defined if vitamin supplementation reduces morbidity and mortality in this group of patients.

CONCLUSION

Preliminary emerging data suggest that the hemostatic system may play either a direct or an indirect role in the pathogenesis, morbidity, and prognosis of mental disorders. Interdisciplinary efforts are mandatory to elucidate the numerous uncertainties related to the associations of thrombosis, cardiovascular disorders, and psychiatric presentations. This could potentially pave the way for refined diagnosis and improved management.

REFERENCES

1. Quaegebeur A, Lange C, Carmeliet P. The neurovascular link in health and disease: molecular mechanisms and therapeutic implications. *Neuron* 2011;71:406–24. [Full Text](#)
2. American Psychiatric Association. *The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition: DSM-5*. Arlington, VA: American Psychiatric Association; 2013.
3. Austin AW, Wissmann T, von Känel R. Stress and hemostasis: an update. *Semin Thromb Hemost* 2013;39:902–12. [Full Text](#)
4. Parmer RJ, Mahata M, Mahata S, Sebald MT, O'Connor DT, Miles LA. Tissue plasminogen activator (t-PA) is targeted to the regulated secretory pathway catecholamine storage vesicles as a reservoir for the rapid release of t-PA. *J Biol Chem* 1997;272:1976–82. [Full Text](#)
5. Austin AW, Patterson SM, von Känel R. Hemocentration and hemostasis during acute stress: interacting and independent effects. *Ann Behav Med* 2011;42:153–73. [Full Text](#)
6. Robicsek O, Makhoul B, Klein E, et al. Hypercoagulation in chronic post-traumatic stress disorder. *Isr Med Assoc J* 2011;13:548–52.
7. Raikonen K, Lassila R, Keltikangas-Jarvinen L, Hautanen A. Association of chronic stress with plasminogen activator inhibitor-1 in healthy middle-aged men. *Arterioscler Thromb Vasc Biol* 1996;16:363–7. [Full Text](#)
8. Halleux CM, Declercq PJ, Tran SL, et al. Hormonal control of plasminogen activator inhibitor-1 gene expression and production in human adipose tissue: stimulation by glucocorticoids and inhibition by catecholamines. *J Clin Endocrinol Metab* 1999;84:4097–105.

9. Brown NJ, Kim KS, Chen YQ, et al. Synergistic effect of adrenal steroids and angiotensin II on plasminogen activator inhibitor-1 production. *J Clin Endocrinol Metab* 2000;85:336–44.
10. Geiser F, Gessler K, Conrad R, et al. Can activation of coagulation and impairment of fibrinolysis in patients with anxiety and depression be reversed after improvement of psychiatric symptoms? Results of a pilot study. *J Nerv Ment Dis* 2012;200:721–3. [Full Text](#)
11. Varghese D, Scott J, Welham J, et al. Psychotic-like experiences in major depression and anxiety disorders: a population-based survey in young adults. *Schizophr Bull* 2011;37:389–93. [Full Text](#)
12. Gold PW, Chrousos GP. Organization of the stress system and its dysregulation in melancholic and atypical depression: high vs low CRH/NE states. *Mol Psychiatry* 2002;7:254–75. [Full Text](#)
13. Lesch KP, WOLOZIN BL, Murphy DL, Reiderer P. Primary structure of the human platelet serotonin uptake site: identity with the brain serotonin transporter. *J Neurochem* 1993;60:2319–22. [Full Text](#)
14. Williams MS. Platelets and depression in cardiovascular disease: a brief review of the current literature. *World J Psychiatry* 2012;2:114–23. [Full Text](#)
15. Lopez-Vilchez I, Serra-Millas M, Navarro V, et al. Prothrombotic platelet phenotype in major depression: downregulation by antidepressant treatment. *J Affect Disord* 2014;159:39–45. [Full Text](#)
16. Lahlou-Laforet K, Alhenc-Gelas M, Pornin M, et al. Relation of depressive mood to plasminogen activator inhibitor, tissue plasminogen activator, and fibrinogen levels in patients with versus without coronary heart disease. *Am J Cardiol* 2006;97:1287–91. [Full Text](#)
17. Kawano H, Tsuji H, Nishimura H, et al. Serotonin induces the expression of tissue factor and plasminogen activator inhibitor-1 in cultured rat aortic endothelial cells. *Blood* 2001;97:1697–702. [Full Text](#)
18. Masopust J, Maly R, Andrys C, et al. The dynamics of haemostatic parameters in acute psychotic patients: a one-year prospective study. *Psychiatr Danub* 2013;25:142–8.
19. Venkatasubramanian G, Chittiprol S, Neelakantachar N, et al. Insulin and insulin-like growth factor-1 abnormalities in antipsychotic-naïve schizophrenia. *Am J Psychiatry* 2007;164:1557–60. [Full Text](#)
20. Hoirisch-Clapauch S, Nardi AE. Markers of low activity of tissue plasminogen activator/plasmin are prevalent in schizophrenia patients. *Schizophr Res* 2014;159:118–23. [Full Text](#)
21. Rezende SM, Simmonds RE, Lane DA. Coagulation, inflammation, and apoptosis: different roles for protein S and the protein S–C4b binding protein complex. *Blood* 2004;103:1192–201. [Full Text](#)
22. Soares-Filho GL, Machado S, Arias-Carrion O, et al. Myocardial perfusion imaging study of CO₂-induced panic attack. *Am J Cardiol* 2014;113:384–8. [Full Text](#)
23. Batelaan NM, ten Have M, van Balkom AJ, et al. Anxiety disorders and onset of cardiovascular disease: the differential impact of panic, phobias and worry. *J Anxiety Disord* 2014;28:252–8. [Full Text](#)
24. Buckley TC, Kaloupek DG. A meta-analytic examination of basal cardiovascular activity in posttraumatic stress disorder. *Psychosom Med* 2001;63:585–94. [Full Text](#)
25. Coryell W, Noyes R Jr, House JD. Mortality among outpatients with anxiety disorders. *Am J Psychiatry* 1986;143:508–10.
26. Lang UE, Borgwardt S. Molecular mechanisms of depression: perspectives on new treatment strategies. *Cell Physiol Biochem* 2013;31:761–77. [Full Text](#)
27. Lichtman JH, Bigger JT Jr, Blumenthal JA, et al. Depression and coronary heart disease: recommendations for screening, referral, and treatment: a science advisory from the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing, Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Interdisciplinary Council on Quality of Care and Outcomes Research: Endorsed by the American Psychiatric Association. *Circulation* 2008;118:1768–75. [Full Text](#)
28. Januzzi JL Jr, Stern TA, Pasternak RC, DeSanctis RW. The influence of anxiety and depression on outcomes of patients with coronary artery disease. *Arch Intern Med* 2000;160:1913–21. [Full Text](#)
29. Leboyer M, Soreca I, Scott J, et al. Can bipolar disorder be viewed as a multi-system inflammatory disease? *J Affect Disord* 2012;141:1–10. [Full Text](#)
30. Hoirisch-Clapauch S, Nardi AE. Multiple roles of tissue plasminogen activator in schizophrenia pathophysiology. *Semin Thromb Hemost* 2013;39:950–4. [Full Text](#)
31. Beary M, Hodgson R, Wildgust HJ. A critical review of major mortality risk factors for all-cause mortality in first-episode schizophrenia: clinical and research implications. *J Psychopharmacol* 2012;26(5 Suppl): 52–61. [Full Text](#)
32. Matura LA, McDonough A, Carroll DL. Cluster analysis of symptoms in pulmonary arterial hypertension: a pilot study. *Eur J Cardiovasc Nurs* 2012;11:51–61. [Full Text](#)

33. Geiser F, Meier C, Wegener I, et al. Association between anxiety and factors of coagulation and fibrinolysis. *Psychother Psychosom* 2008;77:377–83. [Full Text](#)
34. Hoirisch-Clapauch S, Nardi AE, Gris JC, Brenner B. Are the antiplatelet and profibrinolytic properties of selective serotonin-reuptake inhibitors relevant to their brain effects? *Thromb Res* 2014;134:11–16. [Full Text](#)
35. Maes M. Depression is an inflammatory disease, but cell-mediated immune activation is the key component of depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2011;35:664–75. [Full Text](#)
36. Szuster-Ciesielska A, Slotwinska M, Stachura A, et al. Accelerated apoptosis of blood leukocytes and oxidative stress in blood of patients with major depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2008;32:686–94. [Full Text](#)
37. Ishida T, Katagiri T, Uchida H, et al. Incidence of deep vein thrombosis in restrained psychiatric patients. *Psychosomatics* 2014;55:69–75. [Full Text](#)
38. Gris JC, Brenner B. Antiphospholipid antibodies: neuropsychiatric presentations. *Semin Thromb Hemost* 2013;39:935–42. [Full Text](#)
39. Delluc A, Rousseau A, Le Galudec M, et al. Prevalence of antiphospholipid antibodies in psychiatric patients users and non-users of antipsychotics. *Br J Haematol* 2014;164:272–9. [Full Text](#)
40. Bismuth-Evenzal Y, Gonopolsky Y, Gurwitz D, et al. Decreased serotonin content and reduced agonist-induced aggregation in platelets of patients chronically medicated with SSRI drugs. *J Affect Disord* 2012;136:99–103. [Full Text](#)
41. Geiser F, Conrad R, Imbierowicz K, et al. Coagulation activation and fibrinolysis impairment are reduced in patients with anxiety and depression when medicated with serotonergic antidepressants. *Psychiatry Clin Neurosci* 2011;65:518–25. [Full Text](#)
42. Wu RR, Zhao JP, Liu ZN, et al. Effects of typical and atypical antipsychotics on glucose-insulin homeostasis and lipid metabolism in first-episode schizophrenia. *Psychopharmacology (Berl)* 2006;186:572–8. [Full Text](#)
43. Levine J, Stahl Z, Sela BA, et al. Homocysteine-reducing strategies improve symptoms in chronic schizophrenic patients with hyperhomocysteinemia. *Biol Psychiatry* 2006;60:265–9. [Full Text](#)