

### Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017

Hélène M de Nys, Placide Mbala Kingebeni, Alpha Keita, Christelle Butel, Guillaume Thaurignac, Christian Julián Villabona-Arenas, Thomas Lemarcis, Mare Geraerts, Nicole Vidal, Amandine Esteban, et al.

### ▶ To cite this version:

Hélène M de Nys, Placide Mbala Kingebeni, Alpha Keita, Christelle Butel, Guillaume Thaurignac, et al.. Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017. Emerging Infectious Diseases, 2018, 24 (12), pp.2228-2240. 10.3201/eid2412.180740. hal-01941405

### HAL Id: hal-01941405 https://hal.umontpellier.fr/hal-01941405

Submitted on 26 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017

Helene M. De Nys,<sup>1</sup> Placide Mbala Kingebeni,<sup>1</sup> Alpha K. Keita,<sup>1</sup> Christelle Butel, Guillaume Thaurignac, Christian-Julian Villabona-Arenas, Thomas Lemarcis, Mare Geraerts, Nicole Vidal, Amandine Esteban, Mathieu Bourgarel, François Roger, Fabian Leendertz, Ramadan Diallo, Simon-Pierre Ndimbo-Kumugo, Justus Nsio-Mbeta, Nikki Tagg, Lamine Koivogui, Abdoulaye Toure, Eric Delaporte, Steve Ahuka-Mundeke, Jean-Jacques Muyembe Tamfum, Eitel Mpoudi-Ngole, Ahidjo Ayouba,<sup>2</sup> Martine Peeters<sup>2</sup>

To clarify the role of bats in the ecology of Ebola viruses, we assessed the prevalence of Ebola virus antibodies in a large-scale sample of bats collected during 2015–2017 from countries in Africa that have had previous Ebola outbreaks (Guinea, the Democratic Republic of the Congo) or are at high risk for outbreaks (Cameroon). We analyzed 4,022 blood samples of bats from ≥12 frugivorous and

Author affiliations: TransVIHMI of Institut de Recherche pour le Développement, Institut National de la Santé et de la Recherche Médicale and University of Montpellier, Montpellier, France (H.M. De Nys, P. Mbala Kingebeni, A.K. Keita, C. Butel, G. Thaurignac, C.-J. Villabona-Arenas, T. Lemarcis, M. Geraerts, N. Vidal, A. Esteban, A. Toure, E. Delaporte, A. Ayouba, M. Peeters); National Institute of Biomedical Research, Kinshasa, Democratic Republic of the Congo (P. Mbala Kingebeni, S.-P. Ndimbo-Kumugo, S. Ahuka-Mundeke, J.-J. Muyembe Tamfum); Cliniques Universitaires de Kinshasa, Kinshasa (P. Mbala Kingebeni, S. Ahuka-Mundeke, J.-J. Muyembe Tamfum); Centre de Recherche et de Formation en Infectiologie de Guinée, Conakry, Guinea (A.K. Keita, A. Toure); ASTRE of Centre de coopération internationale en recherche agronomique pour le développement, Institut national de la Recherche Agronomique and Univerisity of Montpellier, Montpellier (M. Bourgarel, F. Roger); Robert Koch-Institute, Berlin, Germany (F. Leendertz); Ministère de l'Elevage et des Productions Animales, Conakry (R. Diallo); Direction de Lutte contre la Maladie, Kinshasa (J. Nsio-Mbeta); Royal Zoological Society of Antwerp, Antwerp, Belgium (N. Tagg); Université de Conakry, Conakry (L. Koivogui); Institut National de Sante Publique, Conakry (A. Toure); Institut de Recherches Médicales et d'Études des Plantes Médicinales, Yaoundé, Cameroon (E. Mpoudi-Ngole); Cameroon Institut de Recherche pout le Développement, Yaoundé (E. Mpoudi-Ngole)

27 insectivorous species; 2-37 (0.05%–0.92%) bats were seropositive for Zaire and 0–30 (0%–0.75%) bats for Sudan Ebola viruses. We observed Ebola virus antibodies in 1 insectivorous bat genus and 6 frugivorous bat species. Certain bat species widespread across Africa had serologic evidence of Zaire and Sudan Ebola viruses. No viral RNA was detected in the subset of samples tested (n = 665). Ongoing surveillance of bats and other potential animal reservoirs are required to predict and prepare for future outbreaks.

Since the first outbreak of Ebola virus disease (EVD) in 1976 in the northern part of the Democratic Republic of the Congo (DRC), 26 recognized outbreaks have occurred in humans across Africa; fatality rates of outbreaks have been 25%-90% (1-4). Each EVD outbreak most likely resulted from independent zoonotic events.

Bats are believed to play a role in the ecology of Ebola viruses as a reservoir species (5). Bats might infect humans directly or via intermediate amplifying hosts, like nonhuman primates or duikers (6, 7). Bats might serve as a source of infection in certain areas where bats are hunted and eaten as bushmeat, but infection could also occur after consumption of fruits contaminated with saliva, urine, or feces from Ebola virus–infected bats (8, 9). Ebola virus emergence through exposure to bats was suspected for at least 2 outbreaks: Luebo (the DRC) in 2007 and West Africa in 2013 (10,11).

Relatively few data are available to support the role of bats in the ecology of Ebola viruses. During the EVD outbreaks of 2003 in Gabon and the Congo, Zaire Ebola

<sup>&</sup>lt;sup>1</sup>These first authors contributed equally to this article. <sup>2</sup>These senior authors contributed equally to this article.

virus RNA and antibodies were detected in live-caught specimens from 3 fruit bat species (*Epomops franqueti*, *Hypsignathus monstrosus*, *Myonycteris torquata*); virus sequences were found in the livers or spleens of a few bats (6). In subsequent studies in Gabon, the Congo, Ghana, and Zambia, antibodies were detected in additional frugivorous bat species (*Eidolon helvum, Epomophorus gambianus, Rousettus aegyptiacus, Micropteropus pusillus*) and 1 insectivorous species (*Mops condylurus*) (12–16). The amplification and sequencing of viral RNA of other filoviruses in bats, such as Marburg virus in bats from Africa (17–20), Lloviu virus in bats from Europe (21), and new filoviruses in bats from China (22), has provided additional evidence for a possible role of bats in Ebola virus ecology.

In general, EVD outbreaks have been limited in terms of their geographic spread and chains of human-to-human transmission (1). However, during the 2013–2016 outbreak, virus spread to the urban areas of 3 countries, infecting  $\approx$ 30,000 persons in Guinea, Sierra Leone, and Liberia, and  $\approx$ 11,000 deaths were recorded (23). This outbreak illustrated the potential for epidemic spread from a single zoonotic transmission, with severe public health and so-cioeconomic impact (24). Additional studies are urgently needed to identify the animal reservoir, predict EVD outbreak risks, and improve our capacity to control epidemics.

In previous modeling studies, areas were defined as at risk for EVD outbreaks on the basis of data collected from a limited number of wildlife bat species from a few geographic regions (5,25). Also, a wide variety of serologic assays and interpretation criteria have been used, making comparison of results challenging (12-16,26,27). For this study, we performed a large serosurvey with a highly specific and sensitive high-throughput assay to assess Ebola virus prevalence in bats from Africa (28). We studied bats from Guinea and the DRC, countries with previous EVD outbreaks, and Cameroon, a country considered at high risk for future EVD outbreaks (5,25).

#### **Materials and Methods**

#### **Study Sites and Sample Collection**

During November 2015–August 2017, we collected samples from free-ranging frugivorous and insectivorous bats in Guinea, Cameroon, and the DRC. We captured bats at night using ground mist nets or harp traps in roosting and foraging sites. We set up ground mist nets  $(12 \times 3.2 \text{ m})$  of 30-mm and 60-mm mesh sizes at different heights (1-7 m) to maximize capture of different species. We opened nets or harp traps just before sunset and checked for bats every 1–2 hours. Captured bats were released the same night immediately after sampling. Using bat whole blood taken by venipuncture of the propatagial or brachial vein, we dropped blood samples directly onto Whatman 903

filter paper (GE Healthcare, Feasterville-Trevose, PA, USA). We air-dried and preserved samples individually in plastic bags containing silica desiccant and stored them in hermetic boxes; 2-3 weeks later, we transferred dried blood spots to -20°C until needed for analysis. Data recorded in the field included information on capture site (global positioning system coordinates, ecologic environment), capture method, morphology (body measurements, weight, color), sex, age class (adult, juvenile), and species (identified visually). We collected negative control samples (n = 145) from a captive-born insectivorous bat species (103 Carollia perspicillata bats) hosted at the Parc Zoologique de Montpellier (Montpellier, France) and 2 frugivorous bat species (19 Pteropus giganteus bats, 23 R. aegyptiacus bats) hosted at Wilhelma Zoo and Botanical Garden (Stuttgart, Germany). We collected and preserved samples the same way we did for free-ranging bats.

#### **Screening for Ebola Virus Antibodies**

We tested dried blood spots with a Luminex-based serologic assay adapted for bats (28) (online Technical Appenhttps://wwwnc.cdc.gov/EID/article/24/12/18-0740dix. Techapp1.pdf). The assay included recombinant Ebola virus proteins glycoprotein, nucleoprotein, or viral protein 40 for different lineages: Zaire, Sudan, Bundibugyo, and Reston. We reconstituted plasma from dried blood spots as previously described (28) and incubated 100 µL of sample (final plasma dilution 1:2,000) with 50 µL of recombinant protein-coated beads (2  $\mu$ g protein/1.25  $\times$  10<sup>6</sup> beads) in 96-well flat-bottom filter plates (Millipore, Tullagreen, Ireland) on a plate shaker at 300 rpm for 16 h at 4°C in the dark. After washing, we added 0.1 µg/mL of goat antibat biotin-labeled IgG (Euromedex, Souffelweyersheim, France) per well and incubated for 30 min at 300 rpm. After another round of washing, we added 50  $\mu$ L of 4  $\mu$ g/mL streptavidin-R-phycoerythrin (Fisher Scientific, Illkirch, France) per well and incubated for 10 min at 300 rpm. Reactions were read with BioPlex-200 (BioRad, Marnes-la-Coquette, France). We expressed results as median fluorescence intensity (MFI) per 100 beads. We included 3 samples on every plate to validate interassay repeatability.

#### **Determination of Cutoffs**

In the absence of positive control samples, we used 4 different statistical methods to determine the MFI cutoff value for each antigen (29,30) (online Technical Appendix Table 1). First, we used a general formula that involved the MFI of the 145 negative control samples, and we assigned the cutoff as mean plus 4 times the SD (mean + 4×SD). Second, we used a change point analysis (31) to identify the value at which statistical properties of the underlying probability distribution changed. This value was used to identify outliers and classify them as reactive. We used the R package

Country, site	No. samples
Democratic Republic of the Congo	
Boma	156
Kimpese	202
Zongo	472
Subtotal	830
Cameroon	
Yaoundé	126
Libellengoi Sud	44
Mbalmayo	48
Bipindi	479
Campo M'an	344
North Dja	295
Ekom	122
Djoum	56
Mambele	348
Mbam Djerem	156
Subtotal	2,018
Guinea	
Conakry	107
Kindia	323
Kankan	378
Koundara	90
Mamou	147
Gueckedou	49
Macenta	9
Nzerekore	71
Subtotal	1,174
Total	4,022

 
 Table 1. Bat samples collected for Ebola virus serology by study site, Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017

changepoint (32) to calculate a single shift in the arithmetic mean with the at-most-1-change method (33). Third, we fitted univariate distributions to our data and defined the cutoff as a 0.001 risk for error, as was used in other virus serology studies (13,34). We reduced the set of candidate distributions following a bootstrapped skewness-kurtosis analysis (35). We performed fitting by maximum-likelihood estimation and selected the best-fit distribution on the basis of the Akaike information criteria with the R library fitdistrplus (36). A negative binomial distribution best-fit the data; however, we also used the negative exponential distribution as in Pourrut et al. and Laing et al. (13,34). For every antigen, we computed bootstrap values using 10,000 replicates and averaged. We performed analyses with R version 3.3.2 software (https://www.r-project.org/). We considered a blood sample reactive if the MFI of the reaction was above the cutoff. We defined Ebola virus antibody positivity as reactivity to glycoprotein and nucleoprotein of the same lineage, as was done in our previous study (28).

## Nucleic Acid Extraction and PCR Screening for Ebola Virus RNA

We extracted total DNA and RNA from dried blood spots as previously described using Nuclisens (bioMerieux, Marcyl'Etoile, France) or m2000sp methods (Abbott Molecular Inc., Des Plaines, IL, USA), which are known for a high performance recovering nucleic acids from dried blood spots (37,38). For bat species from Cameroon and Guinea, we screened for Zaire Ebola virus RNA by seminested reverse transcription PCR (RT-PCR) targeting the nucleoprotein region of the virus genome. We amplified a 126-bp fragment of Zaire Ebola virus using primers NP1F1 (forward, 5'-CGGACACACAAAAAGAAWGAA-3') and NP1R-ZR (reverse, 5'-CTCTATCTTKGTGATRTGGCTCTGA-3') in the first round of PCR and NP1F2 (forward, 5'- TTGT-GTGCGARTAACTAYGAGGAAG-3') plus NP1R-ZR in the second round. For species from the DRC, we performed seminested RT-PCR targeting the viral protein 35 region of the genome using the protocol of He et al. with modifications (41). In the first round, we amplified a 217bp fragment with primers VP35-F (5'-ATYATGTATGAT-CACYTVCCWGG-3') and VP35-R (52-AGCGRATGTG-GATSACRGGT-32) and, in the second round, a 184-bp product with primers VP35-R and VP35-in-F (5'-GCTT-TYCAYCAAYTAGTRCAAG-3').

#### **Molecular Confirmation of Bat Species**

We confirmed bat species identification recorded in the field on a subset of samples by using molecular tests. We amplified an  $\approx$ 800-bp fragment of mitochondrial cytochrome b using primers cytb-L14724 (forward) and cytb-H15506 (reverse) (11,39,40). We substituted the cytb-L14724 primer with cytb-L140217 (5'-ATGACCAACATCC-GAAAATCNCAC-3') to improve PCR performance for certain species. We purified PCR products through agarose gel (1%) and directly sequenced on an ABI 3500 sequencer (Applied Biosystems, Courtaboeuf, France). We performed BLAST analyses (https://blast.ncbi.nlm.nih.gov/Blast.cgi) to identify the most similar bat species. For samples with no or low similarity (<97%) hits with species in GenBank, we performed phylogenetic analyses with newly obtained sequences and reference sequences for different bat species using maximum-likelihood methods implemented with PhyML (http://www.atgc-montpellier.fr/phyml/) to determine genus.

#### Results

#### **Bat Species and Sampling**

We analyzed blood samples from 4,022 wild bats from 21 different regions in Cameroon (n = 10), Guinea (n = 8), and the DRC (n = 3) (Figure 1; Table 1). To increase species diversity, we captured bats in multiple ecologic settings: forests (49%), open fields (10%), villages (29%), plantations (7%), and urban areas (5%). For 1,470 (36.5%) samples, species identification in the field was confirmed by sequence analysis. At each site,  $\geq$ 1 sample was confirmed per sampling date, capture method, and morphologic description. For the remaining samples, species identification was extrapolated by combining molecular and morphologic data, including photographs whenever available. For some insectivorous bat



Figure 1. Study sites for bat blood sample collection for Ebola virus serology, Guinea, Cameroon, and the Democratic Republic of the Congo, 2015-2017. Yellow dots indicate sampling sites for bats in our study, and green dots indicate sampling sites in previously published studies. Dark red shading indicates highest and light yellow lowest risk for Ebola virus spillover events. Study sites are numbered: 1, Koundara; 2, Conakry; 3, Kindia; 4, Mamou; 5, Kankan; 6, Gueckedou; 7, Macenta; 8, Nzerekore; 9, Mbam Djerem; 10, Libellengoi Sud; 11, Yaoundé; 12, Ekom; 13, North Dja; 14, Bipindi; 15, Mbalmayo; 16, Djoum; 17, Mambele; 18, Campo M'an: 19. Boma: 20. Kimpese; 21, Zongo. Countries with reported index Ebola cases and countries without such cases but deemed at risk are indicated. Map of Africa adapted from Pigott et al. (5) (https://creativecommons.org/ licenses/by/4.0/) by adding locations of collection sites.

families (Miniopteridae, Molossidae, Nycteridae, Rhinolo*phidae*), identification was possible only at the genus level; for some *Molossidae* bats, we could not distinguish between Mops and Chaerephon genera because of the lack of sequences in GenBank (Table 2). For 87 (2.16%) samples, species identification was not possible because incomplete data were recorded in the field, and available biologic materials were insufficient for molecular confirmation. We collected samples from 1,736 (43.2%) frugivorous bats (family Pteropodidae) of 12 species and 2,199 (54.7%) insectivorous bats (7 families) of >27 species. The insectivorous bat families sampled, in order of decreasing frequency, were Hipposideridae (31.9%), Molossidae (13.4%), Miniopteridae (5.8%), Rhinolophidae (2.1%), Vespertilionidae (0.8%), Nycteridae (0.5%), and Emballonuridae (0.12%). Overall, 54.7% of bats were female and 43.8% were male; for 1.5% (n = 60) of bats, sex was unknown. Most (77.9%) bats were adults, and 9.6% were juveniles; for 12.5% (n = 502) of bats, age could not be determined or was not recorded.

**Bats Antibodies against Different Ebola Virus Antigens** We tested all samples for Ebola virus antibodies. The number of samples reacting with >1 antigen was 734 (18.2%)

by the mean + 4×SD method, 274 (6.8%) for the changepoint method, 175 (4.4%) for the binomial method, and 457 (11.4%) for the exponential method. Blood samples frequently reacted with glycoprotein antigens; samples reacted most with Zaire and Sudan Ebola virus antigens and least with Reston (Table 3). Simultaneous reactivity to >1 antigen (i.e., glycoprotein, nucleoprotein, viral protein 40) from the same virus lineage was rare. Simultaneous reactivity to the same antigen from different virus lineages was frequent; 32.3%-76.7% of blood samples were reactive to glycoprotein from >2 Ebola virus species, 18.4%-34.0% to viral protein 40, and 1.5%-4.4% to nucleoprotein (online Technical Appendix Table 2). When using the criterion simultaneous presence of antibodies to nucleoprotein and glycoprotein, the antibody positivity for Zaire or Sudan Ebola virus antibodies was generally <1% for all bats tested, regardless of cutoff method, and was lower among insectivorous than frugivorous bats: 0.05%-0.27% (insectivorous) and 0.06%-1.79% (frugivorous) for Zaire Ebola virus versus 0%-0.09% (insectivorous) and 0%-1.61% (frugivorous) for Sudan Ebola virus (Table 3; Figure 2). Three samples were positive for Zaire and Sudan Ebola viruses, but only by less stringent cutoff methods (i.e., mean  $+ 4 \times SD$ ).

Family	Species	DRC, no.	Cameroon, no.	Guinea, no.	Total, no.
Emballonuridae	Coleura afra	0	5	0	5
Hipposideridae	Hipposideros abae	0	0	37	37
	H. beatus	0	4	0	4
	H. cyclops	0	14	0	14
	H. fuliginosus	0	2	0	2
	H. gigas	2	9	2	13
	H. jonesi	0	1	12	13
	H. ruber/caffer	127	807	237	1,171
	Hipposideros sp.	28	0	0	28
Subtotal		157	837	288	1,282
Miniopteridae	<i>Miniopterus</i> sp.	205	0	27	232
Molossidae	Chaerephon sp.	0	0	44	44
	Mops condylurus	0	0	110	110
	Mops sp.	0	256	0	256
	Mops/Chaerephon sp.	0	8	120	128
Subtotal		0	264	274	538
Nycteridae	<i>Nycteris</i> sp.	0	7	15	22
Rhinolophidae	Rhinolophus alcyone	0	16	0	16
	R. darlingii	3	0	0	3
	R. fumigatus	0	0	19	19
	R. landeri	0	0	6	6
	Rhinolophus sp.	3	38	1	42
Subtotal		6	54	26	86
Vespertilionidae	Glauconycteris variegata	0	3	0	3
	Kerivoula sp.	0	1	0	1
	Myotis bocagii	0	3	0	3
	Neoromicia sp.	0	5	0	5
	Scotophilus leucogaster	0	0	15	15
	S. nigrita	0	0	1	1
	S. nux	0	6	0	6
Subtotal		0	18	16	34
Pteropodidae	Eidolon helvum	305	158	17	480
	Epomophorus gambianus	0	0	191	191
	Epomophorus wahlbergi	0	16	0	16
	Epomops buettikoferi	0	0	4	4
	Epomops franqueti	20	256	0	276
	Hypsignathus monstrosus	1	176	8	185
	Lissonycteris angolensis	22	30	32	84
	Megaloglossus woermanni	1	19	0	20
	Micropteropus pusillus	44	2	18	64
	Myonycteris torquata	35	21	0	56
	Rousettus aegyptiacus	0	131	228	359
	Scotonycteris zenkeri	0	1	0	1
Subtotal		428	810	498	1,736
Inderminate species		34	23	30	87
Total		830	2,018	1,174	4,022
*DBC the Democratic Bor	while of the Congo		· · ·	· · ·	

Table 2. Bat species sam	pled for Ebola virus serology	/, Guinea, Cameroon	i, and the DRC,	2015-2017*

#### Zaire and Sudan Ebola Virus Reactivity of Different Bat Species

We estimated specific reactivity to Zaire and Sudan Ebola viruses by bat species. We did not include Bundibugyo and Reston because recombinant nucleoproteins were not available. Among insectivorous bats, only blood samples from *Mops* sp. bats (1–6/494) were positive for Zaire or Sudan Ebola virus antibodies (Table 4). Among frugivorous bats, samples from *E. helvum*, *H. monstrosus*, and *R. aegyptiacus* bats had the highest reactivity. We observed Zaire and Sudan Ebola virus seropositivity in these 3 species with almost all cutoff methods: 0.2%– 3.3% for Zaire Ebola virus and 1.0%–2.9% for Sudan Ebola virus in *E. helvum* bat samples, 0.5%–1.6% for Zaire Ebola virus and 1.1%–4.3% for Sudan Ebola virus in *H. monstrosus* bat samples, and 0.6%–2.5% for Zaire Ebola virus and 0.8%–1.4% for Sudan Ebola virus in *R. aegyptiacus* bat samples. We observed 2.4% Zaire Ebola virus–seropositive samples for *Lissonycteris angolensis* bats and 0.5% for *Epomophorus* sp. bats, but only by less stringent cutoff methods. One sample from *M. pusillus* bats was seropositive for Sudan Ebola virus. No samples from *E. franqueti* or *M. torquata* bats were reactive with any Ebola virus antigens. Samples from the 1 *Scotonycteris zenkeri* bat and 20 *Megaloglossus woermanni* bats were seronegative. Overall, Zaire or Sudan Ebola virus antibodies were observed in 7 (1 insectivorous and 6 frugivorous) bat species.

		Statistical method, no. (%)					
Ebola virus species, antigen	Bat type	Mean + 4×SD	Change point	Binomial	Exponential	range, %	
Zaire							
NP	Frugivorous	57 (3.28)	8 (0.46)	24 (1.38)	51 (2.94)	0.46-3.28	
NP	Insectivorous	15 (0.68)	1 (0.05)	6 (0.27)	15 (0.68)	0.05-0.68	
NP	Total	72 (1.79)	9 (0.22)	30 (0.75)	66 (1.64)	0.23-1.79	
GP-K	Frugivorous	365 (21.03)	141 (8.12)	20 (1.15)	113 (6.51)	1.15–21.03	
GP-K	Insectivorous	73 (3.32)	18 (0.82)	2 (0.09)	12 (0.55)	0.09-3.32	
GP-K	Total	440 (10.94)	160 (3.98)	22 (0.55)	125 (3.11)	0.55–10.94	
GP-M	Frugivorous	226 (13.02)	128 (7.37)	16 (0.92)	103 (5.93)	0.92-13.02	
GP-M	Insectivorous	31 (1.41)	14 (0.64)	2 (0.09)	12 (0.55)	0.09-1.41	
GP-M	Total	259 (6.44)	143 (3.56)	18 (0.45)	115 (2.86)	0.45-6.44	
VP	Frugivorous	55 (3.17)	8 (0.46)	24 (1.38)	44 (2.53)	0.46-3.17	
VP	Insectivorous	19 (0.86)	5 (0.23)	6 (0.27)	14 (0.64)	0.23-0.86	
VP	Total	75 (1.86)	14 (0.35)	30 (0.75)	59 (1.47)	0.35-1.86	
NP + GP	Frugivorous	31 (1.79)	31 (1.79)	1 (0.06)	7 (0.40)	0.06-1.79	
NP + GP	Insectivorous	6 (0.27)	6 (0.27)	1 (0.05)	1 (0.05)	0.05-0.27	
NP + GP	Total	37 (0.92)	37 (0.92)	2 (0.05)	8 (0.20)	0.05–0.92	
Sudan							
NP	Frugivorous	71 (4.09)	15 (0.86)	34 (1.96)	77 (4.44)	0.86-4.44	
NP	Insectivorous	12 (0.55)	1 (0.05)	5 (0.23)	18 (0.82)	0.05-0.82	
NP	Total	84 (2.09)	17 (0.42)	39 (0.97)	96 (2.39)	0.42-2.39	
GP	Frugivorous	459 (26.44)	147 (8.47)	17 (0.98)	121 (6.97)	0.98–26.44	
GP	Insectivorous	49 (2.23)	6 (0.27)	1 (0.05)	1 (0.05)	0.05-2.23	
GP	Total	509 (12.66)	154 (3.83)	18 (0.45)	125 (3.11)	0.45-12.66	
VP	Frugivorous	102 (5.88)	20 (1.15)	28 (1.61)	61 (3.51)	1.15–5.88	
VP	Insectivorous	19 (0.86)	4 (0.18)	6 (0.27)	18 (0.82)	0.18–0.86	
VP	Total	121 (3.01)	24 (0.60)	34 (0.85)	80 (1.99)	0.60-3.01	
NP + GP	Frugivorous	28 (1.61)	28 (1.61)	0	10 (0.58)	0–1.61	
NP + GP	Insectivorous	2 (0.09)	2 (0.09)	0	0	0-0.09	
NP + GP	Total	30 (0.75)	30 (0.75)	0	10 (0.25)	0-0.75	
Bundibugyo							
GP	Frugivorous	301 (17.34)	59 (3.40)	0	93 (5.36)	0–17.34	
GP	Insectivorous	58 (2.64)	8 (0.36)	5 (0.23)	13 (0.59)	0.23–2.64	
GP	Total	361 (8.98)	68 (1.69)	22 (0.55)	107 (2.66)	0.55-8.98	
VP	Frugivorous	9 (0.52)	7 (0.40)	12 (0.69)	37 (2.13)	0.40–2.14	
VP	Insectivorous	0	1 (0.05)	8 (0.36)	20 (0.91)	0-0.91	
VP	Total	9 (0.22)	8 (0.20)	20 (0.5)	57 (1.42)	0.20-1.42	
Reston							
GP	Frugivorous	17 (0.98)	28 (1.61)	26 (1.50)	61 (3.51)	0.98–3.51	
GP	Insectivorous	3 (0.14)	10 (0.45)	10 (0.45)	29 (1.32)	0.14–1.32	
GP	Total	20 (0.50)	38 (0.94)	36 (0.90)	90 (2.24)	0.50-2.24	
*VP refers to viral protein 40 of Ebola virus. Rest	ilts are presented f	or frugivorous $(n =$	1,736), insectivorou	s (n = 2,199), a	and total (n = $\overline{4,022}$	2) bats. GP,	

Table 3. Blood samples from bats reactive with Ebola virus antigens in Luminex assay, by antigen, bat type, and statistical method used to determine cutoff, Guinea, Cameroon, and the Democratic Republic of the Congo, 2015-2017\*

glycoprotein; K, Kissoudougou strain; M, Mayinga strain; NP, nucleoprotein; VP, viral protein.

#### Comparison of Zaire Ebola Virus Seroprevalence in **Bats from Africa across Studies**

For comparison, we compiled data regarding Zaire Ebola virus serology in bats of known species from previous studies (n = 4,493) and this study (n = 3,935; 46.7%) (Tables 5, 6). Data were available for 3,023 insectivorous bats of  $\approx 30$ species from 7 different families; 2,199 (72.7%) were from this study (Table 5). Insectivorous bat samples originated from Guinea, Cameroon, the DRC, and Gabon. Zaire Ebola virus reactivity has been observed only in M. condylurus bat samples from Gabon and Mops sp. bat samples from Cameroon. Data were available for 5,405 frugivorous bats of 17 species from 12 genera from West (Guinea, Ghana), West Central (Cameroon, Gabon, the Congo, the DRC), and East (Zambia) Africa (Table 6). No Zaire Ebola virus reactivity has been seen in blood samples from bat species

Casinycteris, Megaloglossus, Nanonycteris, and Scotonyc*teris*, but only a limited number of samples (n = 152) have been tested. Overall, blood samples from 8 frugivorous bat species have been found reactive with Zaire Ebola virus antigens. Blood samples from E. helvum, H. monstrosus, and R. aegyptiacus bats from several countries across Africa have been reported to be seropositive. Reactivity has been observed with samples from E. gambianus bats in Ghana (10.8%) and Guinea. Reactivity was observed with large sample sets from E. franqueti bats derived from Gabon and the Congo and a small sample set from Ghana but not Guinea, Cameroon, or the DRC. M. pusillus and M. torquata bats tested positive for Zaire Ebola virus antibodies in studies in which large sample sets were collected. Among L. angolensis bat samples, only those from Cameroon have tested positive for antibodies.



Figure 2. Bat blood samples reactive to Ebola virus antigens, by statistical method used to determine cutoff, Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017. Samples from frugivorous bats (n = 1,736) and insectivorous bats (n = 2,199) were tested by Luminex assay with GP, NP, and VP of the Zaire and Sudan lineages; GP and VP of the Bundibugyo lineage; and GP of the Reston lineage. GP, glycoprotein; K, Kissoudougou strain; M, Mayinga strain; NP, nucleoprotein; VP, viral protein 40.

#### **RT-PCR Screening for Zaire Ebola Virus RNA**

We screened 665 samples from the DRC (n = 193), Cameroon (n = 399), and Guinea (n = 73) by RT-PCR for the presence of Zaire Ebola virus RNA. Of the 294 samples originating from bats previously documented to carry Zaire Ebola virus RNA (6) (i.e., *H. monstrosus* [132 from Cameroon, 1 from the DRC], *M. torquata* [20 from Cameroon, 25 from the DRC], and *E. franqueti* [116 from Cameroon]), all were negative for Zaire Ebola virus RNA. Of the 371 samples from bat species *E. helvum* (58 from Cameroon, 165 from the DRC, 3 from Guinea), *L. angolensis* (8 from Cameroon, 4 from Guinea), *M. pusillus* (2 from the DRC, 1 from Guinea), *R. aegyptiacus* (45 from Cameroon, 40 from Guinea), *E. gambianus* (25 from Guinea), and *Mops* sp. (20 from Cameroon), all were negative for Zaire Ebola virus RNA.

#### Discussion

To clarify the role of bats in Ebola virus ecology and identify where the virus circulates between outbreaks, we tested >4,000 bats, almost doubling the total number of samples tested in all previous studies in Africa (5-7,42). We provided data on bats from Cameroon, added to the existing data on bats from Guinea and the DRC, and substantially increased the data available on insectivorous bats. We tested samples with the same assay, enabling comparison across species and countries. We used different statistical methods to determine positive sample numbers and expressed the proportion of reactive samples as a range on the basis of the different cutoff values proposed by those methods. As has been done in studies of human Zaire Ebola virus survivors (28,43), we defined Zaire and Sudan Ebola virus positivity as the presence of antibodies to both nucleoprotein and glycoprotein. As such, we estimated that 2-37 (0.05%-0.92%) bats were seropositive for Zaire Ebola virus and 0-30 (0%-0.75%) bats were seropositive for Sudan Ebola virus (Table 3). Among

insectivorous bats, we observed Zaire and Sudan Ebola virus antibodies only in Mops sp. bats, an observation that has previously been observed (13). We provided information on insectivorous Miniopterus and Rhinolophus bats and extended knowledge on Mops and Hipposideros bats; all 1,200 Hipposideros samples were seronegative. We confirmed the presence of Zaire Ebola virus antibodies in only 1 of 3 frugivorous species in which Zaire Ebola virus RNA has been reported, that is, in H. monstrosus but not E. franqueti or M. torquata bats (6). However, this result might have been influenced by sample size, test used, and interpretation criteria. We confirmed antibodies in E. helvum bats and showed that Zaire Ebola virus antibodies are widespread among this species across Africa: Ghana and Zambia, and with our data, also Cameroon, Guinea, and the DRC (13,14,16). We confirmed antibodies in R. aegyptiacus bats from Cameroon and Guinea, in agreement with previous findings in these bats from the Congo and Gabon (13). For E. gambianus bats from Ghana, we also observed Zaire Ebola virus reactivity of samples from this species in Guinea (15). In contrast with a previous study, we observed Sudan Ebola virus antibodies (not Zaire Ebola virus antibodies) in M. pusillus bats (13). We also identified Zaire Ebola virus antibodies in L. angolensis bats from Cameroon, although only when using less stringent cutoff calculations.

When combining data from previous Zaire Ebola virus seroprevalence studies in bats with data from our study, only 1 insectivorous bat species (*Mops* sp.) and 8 frugivorous bat species (*E. helvum*, *E. gambianus*, *E. franqueti*, *H. monstrosus*, *L. angolensis*, *M. pusillus*, *M. torquata*, *R. aegyptiacus*) exhibited Zaire Ebola virus antibodies (13–16). As seen in bat samples from Zambia, we observed in this study Sudan Ebola virus antibodies in *E. helvum* bats from Guinea, Cameroon, and the DRC, suggesting that Zaire and Sudan Ebola viruses co-circulate and could be widespread among this species. However, only 1 other study has tested

		Fbola	Statistical method							
	No.	virus	M	lean + 4×SD	С	hange-point		Binomial	E	Exponential
Bat family, genus	tested	species	No.	% (95% CI)	No.	% (95% CI)	No.	% (95% CI)	No.	% (95% CI)
Hipposideridae										
Hipposideros sp.	1.282	Zaire	0	0 (0-0.3)	0	0 (0-0.3)	0	0 (0-0.3)	0	0 (0-0.3)
	1.282	Sudan	0	0 (0-0.3)	0	0 (0-0.3)	0	0 (0-0.3)	0	0 (0-0.3)
Miniopteridae	, -		_		-		-	- ( /	-	
Miniopterus sp.	232	Zaire	0	0 (0–1.6)	0	0 (0-1.6)	0	0 (0-1.6)	0	0 (0-1.6)
-,	232	Sudan	0	0 (0–1.6)	0	0 (0–1.6)	0	0 (0–1.6)	0	0 (0–1.6)
Molossidae	-		_		-		-	- \ /	-	
Chaerephon sp.	44	Zaire	0	0 (0-8.0)	0	0 (0-8.0)	0	0 (0-8.0)	0	0 (0-8.0)
, 1	44	Sudan	0	0 (0-8.0)	0	0 (0-8.0)	0	0 (0-8.0)	0	0 (0-8.0)
Mops sp.	494	Zaire	6	1.2 (0.6–2.6)	6	1.2 (0.6–2.6)	1	0.2 (0.03-1.1)	1	0.2 (0.03–1.1)
, ,	494	Sudan	2	0.4 (0.1–1.5)	2	0.4 (0.1–1.5)	0	0 (0–0.8)	0	0`(0–0.8)
Nycteridae										
<i>Nycteris</i> sp	22	Zaire	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)
2	22	Sudan	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)
Rhinolophidae				<b>, , ,</b>		<b>x</b>		, , , , , , , , , , , , , , , , , , ,		<b>x x</b>
Rhinolophus sp.	86	Zaire	0	0 (0-4.3)	0	0 (0-4.3)	0	0 (0-4.3)	0	0 (0-4.3)
, ,	86	Sudan	0	0 (0–4.3)	0	0 (0–4.3)	0	0 (0–4.3)	0	0 (0–4.3)
Vespertilionidae				х <i>Г</i>				<b>y y</b>		, i
Glauconycteris sp.*	3	Zaire	0		0		0		0	
2	3	Sudan	0		0		0		0	
<i>Kerivoula</i> sp.*	1	Zaire	0		0		0		0	
	1	Sudan	0		0		0		0	
Myotis bocagii*	3	Zaire	0		0		0		0	
	3	Sudan	0		0		0		0	
<i>Neoromicia</i> sp.*	5	Zaire	0		0		0		0	
	5	Sudan	0		0		0		0	
Scotophilus sp.	22	Zaire	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)	0	0 (0–14.9)
	22	Sudan	0	0 (0-14.9)	0	0 (0–14.9)	0	0 (0-14.9)	0	0 (0–14.9)
Pteropodidae										
Eidolon helvum	480	Zaire	16	3.3 (2.1–5.4)	16	3.3 (2.1–5.4)	1	0.2 (0–1.2)	4	0.8 (0.3–2.1)
	480	Sudan	14	2.9 (1.7–4.8)	14	2.9 (1.7–4.8)	0	0 (0–0.8)	5	1.0 (0.4–2.4)
Epomophorus sp.	207	Zaire	1	0.5 (0.08–2.7)	1	0.5 (0.08–2.7)	0	0 (0–1.4)	0	0 (0–1.8)
	207	Sudan	0	0 (0–1.8)	0	0 (0–1.8)	0	0 (0–1.8)	0	0 (0–1.8)
Epomops sp.	280	Zaire	0	0 (0–1.4)	0	0 (0–1.4)	0	0 (0–1.4)	0	0 (0–1.4)
	280	Sudan	0	0 (0–1.4)	0	0 (0–1.4)	0	0 (0–1.4)	0	0 (0–1.4)
Hypsignathus	185	Zaire	3	1.6 (0.6–4.7)	3	1.6 (0.6–4.7)	0	0 (0–2.0)	1	0.5 (0.05–3.0)
monstrosus	185	Sudan	8	4.3 (2.2–8.3)	8	4.3 (2.2–8.3)	3	1.6 (0.6–4.7)	2	1.1(0.3–3.9)
Lissonycteris	84	Zaire	2	2.4 (0.7–8.3)	2	2.4 (0.7–8.3)	0	0 (0–4.4)	0	0 (0–4.4)
angolensis	84	Sudan	0	0 (0–4.4)	0	0 (0–4.4)	0	0 (0–4.4)	0	0 (0–4.4)
Megaloglossus	20	Zaire	0	0 (0–16.1)	0	0 (0–16.1)	0	0 (0–16.1)	0	0 (0–16.1)
woermanni	20	Sudan	0	0 (0–16.1)	0	0 (0–16.1)	0	0 (0–16.1)	0	0 (0–16.1)
Micropteropus	64	Zaire	0	0 (0–5.7)	0	0 (0–5.7)	0	0 (0–5.7)	0	0 (0–5.7)
pusillus	64	Sudan	1	1.6 (0.3–8.3)	1	1.6 (0.3–8.3)	0	0 (0–5.7)	0	0 (0–5.7)
Myonycteris	56	Zaire	0	0 (0–6.4)	0	0 (0–6.4)	0	0 (0–6.4)	0	0 (0–6.4)
torquata	56	Sudan	0	0 (0–6.4)	0	0 (0–6.4)	0	0 (0–6.4)	0	0 (0–6.4)
Rousettus	359	Zaire	9	2.5 (1.3–4.7)	9	2.5 (1.3–4.7)	0	0 (0–1.1)	2	0.6 (0.2–2.0)
aegyptiacus	359	Sudan	5	1.4 (0.6–3.2)	5	1.4 (0.6–3.2)	0	0 (0–1.1)	3	0.8 (0.3–2.4)
Scotonycteris	1	Zaire	0		0		0		0	
zenkeri*	1	Sudan	0		0		0		0	
*Percentages were not cal	culated be	cause the r	number	of samples collected	was to	oo low.				

 Table 4. Blood samples from bats reactive with both nucleoprotein and glycoprotein of Zaire or Sudan Ebola virus, by statistical method used to determine cutoff, Guinea, Cameroon, and the Democratic Republic of the Congo, 2015–2017

for Ebola viruses other than Zaire Ebola virus in *E. helvum* bats (*16*). In our study, we also observed Sudan Ebola virus antibodies in *Mops* sp., *H. monstrosus*, and *R. aegyptiacus* bats in Cameroon. Almost all samples were positive for either Zaire or Sudan Ebola virus but not for both.

Despite the presence of Ebola virus antibodies, the role of bats as reservoir species remains unclear because viral RNA detection is rare. In only 1 study Zaire Ebola virus RNA was amplified in a few bats (6). Thus, antibodies

might reflect previous acute infection with viral clearance. Unlike inoculations with Marburg virus (44-46), experimental inoculation of *R. aegyptiacus* bats with Zaire Ebola virus leads to antibody development but infrequent or absent detection of viral RNA or shedding (44,47). *R. aegyptiacus* bats are therefore able to clear Zaire Ebola virus after a short infectious period without viral shedding and with little or no transmission. No antibodies or viral RNA were detected in noninoculated

Family         Species         Country         No.	published studies	5						
Family         Species         Country         (reference)         Test         tested         positive/feeted (%)†           Embalanundae         Coleura afra         Cameroon         2015-2017         Luminex         5         0-0 (0-0)         0/14 (0)           Hipposiderideros sp.         DRC         2015-2017         Luminex         837         0-0 (0-0)         0/1,395 (0)           Hipposiderideros sp.         Guinea         2015-2017         Luminex         837         0-0 (0-0)         0/1,395 (0)           Miniopterideros sp.         Guinea         2014 (17)         ELISA         0         0(0)           Miniopterus sp.         Guinea         2014 (2017)         Luminex         288         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         1995 (27)         ELISA         2         0 (0)         0/001 (0)           C. aminus         Guinea         2015-2017         Luminex         24         0 (0)         0         0/01 (0)         0           C. aminus         Guinea         2014 (17)         ELISA         1         0 (0)         0         0         0         0         0         0         0         0         0         0         0         0         0 </td <td></td> <td>_</td> <td>-</td> <td>Year of study</td> <td>_</td> <td>No.</td> <td>No. (%)</td> <td>Total, no.</td>		_	-	Year of study	_	No.	No. (%)	Total, no.
Emballorunidae         Coleura afra         Cameroon         2015-2017‡         Luminex         5         0-0 (0-0)         0/14 (0)           Hipposiderios sp.         DRC         2015-2017‡         Luminex         157         0-0 (0-0)         0/1,395 (0)           Hipposideros sp.         Cameroon         2015-2017‡         Luminex         837         0-0 (0-0)         0/1,395 (0)           Miniopteridae         Miniopteros sp.         Guinea         2015-2017‡         Luminex         288         0-0 (0-0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         270         0-0 (0-0)         0/234 (0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         205         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         1985 (27)         ELISA         2         0 (0)         0           Molossidae         Chaerephon sp.         Guinea         2015-2017‡         Luminex         264         1-6 (0-2)         0/401 (0)           C. amajor         DRC         1995 (27)         ELISA         1         0 (0)         0           C. amajor         DRC         1995 (27)         ELISA         2         0 (0)         0         0	Family	Species	Country	(reference)	Test	tested	positive†	positive/tested (%)†
Saccolarinus peli         DRC         1970-1980 (26)         IFA         9         0 (0)           Hipposideros sp.         DRC         2015-2017‡         Luminex         837         0-0 (0-0)         0/1,395 (0)           Hipposideros sp.         DRC         1979-1980 (26)         IFA         69         0 (0)           Hipposideros sp.         Guinea         2015-2017‡         Luminex         288         0-0 (0-0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         27         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         2015-2017‡         Luminex         27         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         1995 (27)         ELISA         1         0 (0)           C. ansorgei         DRC         1979-1980 (26)         IFA         26         0 (0)           C. ansorgei         DRC         1979-1980 (26)         IFA         264         1-6 (0.4-2.3)           Mops sp.         Guinea         2015-2017‡         Luminex         254         1-6 (0.4-2.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980	Emballonuridae	Coleura afra	Cameroon	2015–2017‡	Luminex	5	0-0 (0-0)	0/14 (0)
Hipposideridae         Hipposideros sp.         DRC         2015–2017‡         Luminex         157         0-0 (0-0)         0/1,395 (0)           Hipposideros sp.         DRC         1979-1980 (26)         IFA         69         0 (0)           Miniopteridaes         Sp.         Guinea         2015-2017‡         Luminex         288         0-0 (0-0)           Miniopteridaes         Miniopteridas sp.         Guinea         2015-2017‡         Luminex         205         0-0 (0-0)         0/234 (0)           Molossidae         Charerophon sp.         DRC         1985 (27)         ELISA         4         0-0 (0-0)         0/401 (0)           C. amarori         DRC         1995 (27)         ELISA         1         0 (0)         0         0         0           C. amarori         DRC         1995 (27)         ELISA         1         0 (0)         0		Saccolaimus peli	DRC	1979–1980 (26)	IFA	9	0 (0)	
Hipposideros sp.         Cameroon         2015–2017‡         Luminex         83         0–0 (0–0)           Hipposideros sp.         Guinea         2015–2017‡         Luminex         288         0–0 (0–0)           Miniopterus sp.         Guinea         2015–2017‡         Luminex         228         0–0 (0–0)           Miniopterus sp.         DRC         2015–2017‡         Luminex         227         0–0 (0–0)           Miniopterus sp.         DRC         1995 (27)         ELISA         2         0 (0)           Molossidae         Chaerephon sp.         Guinea         2015–2017‡         Luminex         40         0 (0)           C. major         DRC         1995 (27)         ELISA         1         0 (0)         0           C. nagor         DRC         1995 (27)         ELISA         1         0 (0)         0           Mops sp.         Cameroon         2015–2017‡         Luminex         230         0–0 (0–0)         4–9/705 (0.6–1.3)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops sp.         DRC         1995 (27) <t< td=""><td>Hipposideridae</td><td>Hipposideros sp.</td><td>DRC</td><td>2015–2017‡</td><td>Luminex</td><td>157</td><td>0-0 (0-0)</td><td>0/1,395 (0)</td></t<>	Hipposideridae	Hipposideros sp.	DRC	2015–2017‡	Luminex	157	0-0 (0-0)	0/1,395 (0)
Hipposideros sp.         DRC         1979-1980 (26)         IFA         69         0 (0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         28         0-0 (0-0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         205         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         1995 (27)         ELISA         2         0 (0)           Molossidae         Chaerephon sp.         Guinea         2015-2017‡         Luminex         20         0(0)           Molossidae         Chaerephon sp.         Guinea         2015-2017‡         Luminex         44         0-0 (0-0)         0/401 (0)           C. amorgei         DRC         1995 (27)         ELISA         120         0 (0)         0           C. amorgei         DRC         1995 (27)         Elisa         210         0 (0)         0           Mops sp.         Cameroon         2015-2017‡         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)         0           Mops sp.         DRC         1995 (27)         ELISA         24         3 (12.5)		Hipposideros sp.	Cameroon	2015-2017	Luminex	837	0-0 (0-0)	
Hipposideros sp.         Guinea         2015-2017‡         Luminex         248         0-0 (0-0)           Miniopterus sp.         Guinea         2015-2017‡         Luminex         27         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         2015-2017‡         Luminex         27         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         1955 (27)         ELISA         2         0 (0)           Molossidae         Chaerephon sp.         Guinea         2015-2017‡         Luminex         44         0-0 (0-0)         0/401 (0)           C. pumilus         Guinea         2015-2017‡         Luminex         44         0-0 (0-0)         0/401 (0)           C. pumilus         DRC         1995 (27)         ELISA         120         0 (0)           Mops sp.         Guinea         2015-2017‡         Luminex         264         1-6 (0-4-2.3)           Mops sp.         DRC         1995 (27)         ELISA         2         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         2         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         2         0 (0)           Mops sp.         Guinea		Hipposideros sp.	DRC	1979–1980 (26)	IFA	69	0 (0)	
Hipposideros sp.         Guinea         2014 (11)         ELISA         44         0 (0)           Miniopterus sp.         DRC         2015-2017‡         Luminex         20         0-0 (0-0)         0/234 (0)           Miniopterus sp.         DRC         2015-2017‡         Luminex         205         0-0 (0-0)         0/401 (0)           Molossidae         Carsorgei         DRC         1985 (27)         ELISA         2         0 (0)           C. ansorgei         DRC         1995 (27)         ELISA         1         0 (0)           C. ansorgei         DRC         1995 (27)         ELISA         10 (0)         0           Mops sp.         Guinea         2015-2017‡         Luminex         266         0 (0)           Mops sp.         Cameroon         2015-2017‡         Luminex         264         1-6 (0.4-2.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         15         0-0 (0-0)         0/43 (0)           Myoteris sp.         Guinea         2015-2017‡		Hipposideros sp.	Guinea	2015–2017‡	Luminex	288	0-0 (0-0)	
Miniopterulae         Miniopterus sp. Miniopterus sp. Miniopterus sp.         Guinea DRC         2015-2017‡ 1995 (27)         Luminex Luminex         27         0-0 (0-0)         0/234 (0)           Molossidae         Chaerephon sp. C. pumilus         Guinea         2015-2017‡         Luminex         20         0 (0)           Molossidae         Chaerephon sp. C. ansorgei         DRC         1995 (27)         ELISA         1         0 (0)           C. major         DRC         1995 (27)         ELISA         1         0 (0)           C. major         DRC         1995 (27)         ELISA         0 (0)           Mops sp.         Guinea         2015-2017‡         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1995 (27)         ELISA         210         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         2         0 (0)           Mops sp.         Guinea         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Mycteris sp.         Guinea         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Mycteris sp.         Guinea         2015-2017‡         Luminex         7<		Hipposideros sp.	Guinea	2014 (11)	ELISA	44	0 (0)	
Miniopterus sp. M. minor         DRC         2015–2017‡         Luminex         205         0–0 (0–0)           Molossidae         Chaerephon sp. C. pumilus         Guinea         2014–2017‡         Luminex         44         0–0 (0–0)         0/401 (0)           C. ansorgei         DRC         1995 (27)         ELISA         1         0 (0)           C. ansorgei         DRC         1995 (27)         Elisa         210         0 (0)           C. major         DRC         1995 (27)         Elisa         210         0 (0)           Mops sp.         Caimea         2015–2017‡         Luminex         230         0–0 (0–0)         4–9/705 (0.6–1.3)           Mops sp.         Cameroon         2015–2017‡         Luminex         236         160 (0         -           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)         -           Mops sp.         DRC         1995 (27)         ELISA         24         3 (12.5)         -           Mops sp.         DRC         1995 (27)         ELISA         2         0 (0)         -           Myopterus whitleyi         DRC         1995 (27)         ELISA         1         0 (0)         - <td< td=""><td>Miniopteridae</td><td>Miniopterus sp.</td><td>Guinea</td><td>2015-2017‡</td><td>Luminex</td><td>27</td><td>0-0 (0-0)</td><td>0/234 (0)</td></td<>	Miniopteridae	Miniopterus sp.	Guinea	2015-2017‡	Luminex	27	0-0 (0-0)	0/234 (0)
M. minor         DRC         1995 (27)         ELISA         2         0 (0)           Molossidae         Chaerephon sp. C. pumilus         Guinea         2015-2017‡         Luminex         44         0-0 (0-0)         0/401 (0)           C. ansorgei         DRC         1995 (27)         ELISA         120         0 (0)           C. ansorgei         DRC         1995 (27)         ELISA         210         0 (0)           C. major         DRC         1995 (27)         Elisa         210         0 (0)           Mops sp.         Cameroon         2015-2017‡         Luminex         284         1-6 (0.4-2.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         10 (0)         0/043 (0)           Myoteris sp.         Guinea         2014 (11)         ELISA         2         0 (0)           Myoteris sp.         Guinea         2015-2017‡         Luminex         15         0-0 (0-0)         0/43 (0)           Myoteris sp.         Guinea         2015-2017‡         Luminex		Miniopterus sp.	DRC	2015–2017±	Luminex	205	0–0 (0–0)	
Molossidae         Cheerephon sp. C. pumilus         Guinea         2015-2017 2014         Luminex         44         0-0 (0-0)         0/401 (0)           C. ansorgei         DRC         1995 (27)         ELISA         1         0 (0)           C. major         DRC         1995 (27)         ELISA         20         0 (0)           C. major         DRC         1995 (27)         Elisa         210         0 (0)           Mops sp.         Cameroon         2015-2017‡         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         24         3 (12.5)           M. condylurus         Gabon         2003-2008 (13)         ELISA         1         0 (0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         6         0-0 (0-0)         0/66 (0)           Nycteris sp.         Cameroo		M. minor	DRC	1995 (27)	ELISA	2	0 (0)	
C. pumilus         Guinea         2014 (11)         ELISA         1         0 (0)         0 (0)           C. ansorgei         DRC         1995 (27)         ELISA         120         0 (0)           C. major         DRC         1995 (27)         ELISA         120         0 (0)           Mops sp.         Guinea         2015-2017         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         Cameroon         2015-2017         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops sondylurus         Guinea         2014 (11)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2015-2017         Luminex         15         0-0 (0-0)           Mycteris sp.         Guinea         2014 (11)         ELISA         6         0 (0)           Nycteris sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1	Molossidae	Chaerephon sp.	Guinea	2015–2017±	Luminex	44	0-0 (0-0)	0/401 (0)
C. ansorgei         DRC         1995 (27)         ELISA         120         0 (0)           C. maijor         DRC         1979–1980 (26)         IFA         26         0 (0)           Mops sp.         Guinea         2015–2017‡         Luminex         230         0–0 (0–0)         4–9/705 (0.6–1.3)           Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2014 (11)         ELISA         2         0 (0)           Mycteris sp.         Guinea         2014 (11)         ELISA         0         0 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Rhinolophus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)           Rhinolophus s		C. pumilus	Guinea	2014 (11)	ELISA	1	0 (0)	
C. major         DRC         1979–1980 (26)         IFA         26         0 (0)           C. purnilus         DRC         1995 (27)         Elia         210         0 (0)           Mops sp.         Guinea         2015–2017‡         Luminex         230         0-0 (0-0)         4–9/705 (0.6–1.3)           Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops condylurus         Gabon         2003–2008 (13)         ELISA         2         0 (0)           Myoterix swhitleyi         DRC         1995 (27)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         15         0-0 (0-0)         0/43 (0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         1         0 (0)         0           Nycteris sp.         Cameroon         2015–2017‡         Luminex         6         0-0 (0-0)         0/86 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Lumin		C. ansorgei	DRC	1995 (27)	ELISA	120	0 (0)	
C. pumilius         DRC         1995 (27)         Elisa         210         0 (0)           Mops sp.         Guinea         2015-2017‡         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops condylurus         Gabon         2003-2008 (13)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2014 (11)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Nycteris sp.         Cameroon         2015-2017‡         Luminex         6         0 (0)         0           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)         0           Nycteris sp.         Cameroon         2015-2017‡         Luminex         6         0-0 (0-0)         0/43 (0)           Rhinolophus sp.         Cameroon         2015-2017‡         Luminex         6         0-0 (0-0)         0/143 (0)         Eptesicus sp.         DRC <td></td> <td>C. maior</td> <td>DRC</td> <td>1979–1980 (26)</td> <td>IFA</td> <td>26</td> <td>0 (0)</td> <td></td>		C. maior	DRC	1979–1980 (26)	IFA	26	0 (0)	
Mops sp.         Guinea         2015-2017‡         Luminex         230         0-0 (0-0)         4-9/705 (0.6-1.3)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1979-1980 (26)         IFA         158         0 (0)           Mops condylurus         Gabon         2003-2008 (13)         ELISA         28         0 (0)           Mops condylurus         Gabon         2003-2008 (13)         ELISA         2         0 (0)           Mops condylurus         Gabinea         2015-2017‡         Luminex         15         0-0 (0-0)           Mycteris sp.         Guinea         2014 (11)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2014 (27)         ELISA         0         0           Nycteris sp.         Cameroon         2014 (11)         ELISA         0         0           Nycteris sp.         Cameroon         2015-2017‡         Luminex         7         0-0 (0-0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         6         0-0 (0-0)           Rhinolophus sp.         Gueroon         2015-2017‡         Luminex         3         0-0 (		C. pumilus	DRC	1995 (27)	Elisa	210	0 (0)	
Mops sp.         Cameroon         2015–2017‡         Luminex         264         1-6 (0.4–2.3)           Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops condylurus         Gabon         2003–2008 (13)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2014 (11)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2015–2017‡         Luminex         15         0-0 (0-0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         7         0-0 (0-0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         7         0-0 (0-0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         6         0-0 (0-0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         6         0-0 (0-0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         3         0-0 (0-0)         0/143 (0)           Vespertillonidae         Glauconycteris variegata         C		Mops sp.	Guinea	2015–2017±	Luminex	230	0-0 (0-0)	4-9/705 (0.6-1.3)
Mops sp.         DRC         1979–1980 (26)         IFA         158         0 (0)           Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops condylurus         Gabon         2003–2008 (13)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2014 (11)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)         0/43 (0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         7         0–0 (0–0)         0/43 (0)           Nycteris sp.         Cameroon         2015–2017‡         Luminex         7         0–0 (0–0)         0/43 (0)           Nycteris hispida         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         6         0–0 (0–0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         54         0–0 (0–0)           Vespertilionidae         Chalinolobus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)           Eptesicus sp.         DRC		Mops sp.	Cameroon	2015–2017±	Luminex	264	1-6(0.4-2.3)	
Mops sp.         DRC         1995 (27)         ELISA         28         0 (0)           Mops condylurus         Gabon         2003-2008 (13)         ELISA         24         3 (12.5)           M. condylurus         Guinea         2014 (11)         ELISA         2         0 (0)           Myopterus whitleyi         DRC         1995 (27)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         15         0-0 (0-0)         0/43 (0)           Nycteris sp.         Guinea         2014 (11)         ELISA         6         0 (0)         0/43 (0)           Nycteris sp.         Cameroon         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         Cameroon         2015-2017‡         Luminex         6         0-0 (0-0)         0/86 (0)           Eptesicus renuipinnis         DRC         1979-1980 (26)         IFA         15         0 (0)           Eptesicus tenuipinnis         DRC <td< td=""><td></td><td>Mops sp</td><td>DRC</td><td>1979–1980 (26)</td><td>IFA</td><td>158</td><td>0 (0)</td><td></td></td<>		Mops sp	DRC	1979–1980 (26)	IFA	158	0 (0)	
Mops condylurus         Gabon         2003–2008 (13)         ELISA         24         3 (12,5)           M. condylurus         Guinea         2014 (11)         ELISA         1         0 (0)           Myoteris sp.         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)         0/43 (0)           Nycteris sp.         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)         0/43 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         2         0 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         6         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)         143 (0)         143 (0)         143 (0)         143 (0)         143 (0)         143 (0)         144 (11)         15         144 (11		Mops sp	DRC	1995 (27)	FLISA	28	0(0)	
M. condylurus         Guinea         2014 (11)         ELISA         1         0 (0)           Myopterus whitleyi         DRC         1995 (27)         ELISA         2         0 (0)           Nycteris sp.         Guinea         2015-2017‡         Luminex         15         0-0 (0-0)         0/43 (0)           Nycteris sp.         Cameroon         2015-2017‡         Luminex         7         0-0 (0-0)         0/43 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)           Nycteris sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophidae         Rhinolophus sp.         Guinea         2015-2017‡         Luminex         6         0-0 (0-0)           Rhinolophus sp.         DRC         2015-2017‡         Luminex         6         0-0 (0-0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015-2017‡         Luminex         3         0-0 (0-0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015-2017‡         Luminex         3         0-0 (0-0)           Kerivoula sp.         DRC         1979-1980 (26)         IFA         1         0 (0)         <		Mops condvlurus	Gabon	2003 - 2008(13)	FLISA	24	3 (12 5)	
Myopterus whitleyi         DRC         1995 (27)         ELISA         2         0 (0)           Nycteris p.         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)         0/43 (0)           Nycteris sp.         Guinea         2014 (11)         ELISA         6         0 (0)           Nycteris sp.         Cameroon         2015–2017‡         Luminex         7         0–0 (0–0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         6         0–0 (0–0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         6         0–0 (0–0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Vespertilionidae sp.         DRC         1979–1980 (26)         IFA         15         0 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         1         0 (0)           Kerivoula sp.         Cameroon         2015–201		M condvlurus	Guinea	2014 (11)	FLISA	1	0(0)	
Nycteridae         Nycteris sp. Nycteris sp.         Guinea Guinea         2015–2017 2017 Luminex         Luminex         15         0–0 (0–0)         0/43 (0)           Nycteris sp. Nycteris sp.         Cameroon         2014(11)         ELISA         6         0 (0)           Nycteris sp. Nycteris sp.         DRC         1995(27)         ELISA         1         0 (0)           Rhinolophus sp. Rhinolophus sp.         Guinea         2015–2017 2015–2017 Luminex         1         0 (0)           Rhinolophus sp. Rhinolophus sp.         Cameroon         2015–2017 2015–2017 Luminex         1         0 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017 2015–2017 Luminex         1         0 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017 2015–2017 Luminex         3         0–0 (0–0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017 2015–2017 Luminex         1         0 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)           Kerivoula sp.         Guinea         2014(11)         ELISA         1         0 (0)           Myotis bocagii         DRC         1995 (27)         ELISA		Myopterus whitlevi	DRC	1995 (27)	FLISA	2	0(0)	
Nycteris sp.         Guinea         2014 (11)         ELISA         6         0 (0)         Dist (0)           Nycteris sp.         Cameroon         2015–2017‡         Luminex         7         0–0 (0–0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Rhinolophus sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         DRC         2015–2017‡         Luminex         26         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         DRC         2015–2017‡         Luminex         54         0–0 (0–0)         0/86 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         1         0 (0)           Kerivoula sp.         Guinea         2014 (17)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Myotis b	Nvcteridae	Nvcteris sp.	Guinea	2015–2017±	Luminex	15	0-0 (0-0)	0/43 (0)
Nycteris sp.         Cameroon         2015–2017‡         Luminex         7         0–0 (0–0)           Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Rhinolophidae         Rhinolophus sp.         Guinea         2015–2017‡         Luminex         26         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         26         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)         Eptesicus sp.         DRC         1995 (27)         ELISA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)         0         0           Myctis bocagii         DRC         1995 (27)         ELISA         2		Nycteris sp	Guinea	2014 (11)	FLISA	6	0(0)	0, 10 (0)
Nycteris sp.         DRC         1979–1980 (26)         IFA         14         0 (0)           Nycteris hispida         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophus sp.         Guinea         2015–2017‡         Luminex         26         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         DRC         2015–2017‡         Luminex         6         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         54         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Eptesicus senuipinnis         DRC         1979–1980 (26)         IFA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Myotis bocagii         DRC         1995 (27)         <		Nycteris sp	Cameroon	2015-2017+	Luminex	7	0-0(0-0)	
Nycteris hispida         DRC         1995 (27)         ELISA         1         0 (0)           Rhinolophidae         Rhinolophus sp.         Guinea         2015–2017‡         Luminex         26         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         DRC         2015–2017‡         Luminex         6         0–0 (0–0)         0/86 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         54         0–0 (0–0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         12         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         1         0–0 (0–0)           Myctis bocagii         DRC         1995 (27)         ELISA         2         0 (0)           Meoromicia sp.         Cameroo		Nycteris sp	DRC	1979-1980 (26)	IFA	14		
Rhinolophidae         Rhinolophus sp.         Guinea         2015–2017‡         Luminex         26         0–0         0–0         0/86 (0)           Rhinolophus sp.         DRC         2015–2017‡         Luminex         6         0–0 (0–0)         0/86 (0)           Rhinolophus sp.         Cameroon         2015–2017‡         Luminex         54         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           M. bocagii         DRC         1979–1980 (26)         IFA         17         0		Nycteris hispida	DRC	1995 (27)	FLISA	1	0(0)	
Rhinolophus sp.       DRC       2015–2017‡       Luminex       6       0-0 (0-0)         Vespertilionidae       Glauconycteris variegata       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0)         Vespertilionidae       Glauconycteris variegata       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0)         Vespertilionidae       Glauconycteris variegata       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0)         Vespertilionidae       Glauconycteris variegata       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0) <i>Eptesicus</i> sp.       DRC       1979–1980 (26)       IFA       15       0 (0)         Eptesicus tenuipinnis       DRC       1995 (27)       ELISA       1       0 (0)         Kerivoula sp.       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0)         Myotis bocagii       Cameroon       2015–2017‡       Luminex       3       0-0 (0-0)         Myotis bocagii       DRC       1995 (27)       ELISA       22       0 (0)         Mocagii       DRC       1995 (27)       ELISA       20       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex </td <td>Rhinolophidae</td> <td>Rhinolophus sp</td> <td>Guinea</td> <td>2015-2017+</td> <td></td> <td>26</td> <td>0-0(0-0)</td> <td>0/86 (0)</td>	Rhinolophidae	Rhinolophus sp	Guinea	2015-2017+		26	0-0(0-0)	0/86 (0)
Rhinolophus sp.         Cameroon         2015 2017‡         Luminex         54         0-0 (0-0)           Vespertilionidae         Glauconycteris variegata Chalinolobus sp.         Cameroon         2015-2017‡         Luminex         3         0-0 (0-0)         0/143 (0)           Vespertilionidae         Glauconycteris variegata Chalinolobus sp.         DRC         1979-1980 (26)         IFA         15         0 (0)           Eptesicus sp.         DRC         1979-1980 (26)         IFA         22         0 (0)           Eptesicus tenuipinnis         DRC         1995 (27)         ELISA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015-2017‡         Luminex         3         0-0 (0-0)           Myotis bocagii         Cameroon         2015-2017‡         Luminex         3         0-0 (0-0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           M. bocagii         DRC         1979-1980 (26)         IFA         17         0 (0)           Scotophilus nanus         DRC         1995 (27)         ELISA         2         0 (0)           Scoto	1 annoiopinaao	Rhinolophus sp	DRC	2015-2017+	Luminex	6	0-0(0-0)	0,00 (0)
Vespertilionidae         Glauconycteris variegata Chalinolobus sp.         Cameroon DRC         1979–1980 (26)         IFA         15         0 (0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         15         0 (0)         0/143 (0)           Eptesicus sp.         DRC         1979–1980 (26)         IFA         22         0 (0)           Eptesicus tenuipinnis         DRC         1995 (27)         ELISA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           Myotis bocagii         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           M. bocagii         DRC         1979–1980 (26)         IFA         17         0 (0)           Neoromicia sp.         Cameroon         2015–2017‡         Luminex         5         0–0 (0–0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         15         0–0 (0–0)           Scotophilus leucogaster		Rhinolophus sp	Cameroon	2015-2017+	Luminex	54	0-0(0-0)	
Chalinolobus sp.       DRC       1979–1980 (26)       IFA       15       0 (0)         Eptesicus sp.       DRC       1979–1980 (26)       IFA       22       0 (0)         Eptesicus tenuipinnis       DRC       1995 (27)       ELISA       1       0 (0)         Kerivoula sp.       Guinea       2014 (11)       ELISA       1       0 (0)         Kerivoula sp.       Cameroon       2015–2017‡       Luminex       1       0–0 (0–0)         Myotis bocagii       Cameroon       2015–2017‡       Luminex       3       0–0 (0–0)         M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex       5       0–0 (0–0)         Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19 <td>Vespertilionidae</td> <td>Glauconvcteris variegata</td> <td>Cameroon</td> <td>2015-2017+</td> <td>Luminex</td> <td>3</td> <td>0-0(0-0)</td> <td>0/143 (0)</td>	Vespertilionidae	Glauconvcteris variegata	Cameroon	2015-2017+	Luminex	3	0-0(0-0)	0/143 (0)
Eptesicus sp.       DRC       1979–1980 (26)       IFA       22       0 (0)         Eptesicus tenuipinnis       DRC       1995 (27)       ELISA       1       0 (0)         Kerivoula sp.       Guinea       2014 (11)       ELISA       1       0 (0)         Kerivoula sp.       Cameroon       2015–2017‡       Luminex       1       0–0 (0–0)         Myotis bocagii       Cameroon       2015–2017‡       Luminex       3       0–0 (0–0)         M. bocagii       DRC       1995 (27)       ELISA       22       0 (0)         M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex       5       0–0 (0–0)         Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       1	reeperanornaae	Chalinolobus sp	DRC	1979-1980 (26)	IFA	15		0,110 (0)
Epicological op.         Drive         1010 for 000 (20)         In r         22         0 (0)           Eptesicus tenuipinnis         DRC         1995 (27)         ELISA         1         0 (0)           Kerivoula sp.         Guinea         2014 (11)         ELISA         1         0 (0)           Kerivoula sp.         Cameroon         2015–2017‡         Luminex         1         0–0 (0–0)           Myotis bocagii         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           Neoromicia sp.         Cameroon         2015–2017‡         Luminex         5         0–0 (0–0)           Pipistrellus nanus         DRC         1995 (27)         ELISA         2         0 (0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         1         0–0 (0–0)		Entesicus sp	DRC	1979_1980 (26)	IFA	22	0(0)	
Lick         Lick <thlick< th="">         Lick         Lick         <thl< td=""><td></td><td>Entesicus tenuininnis</td><td>DRC</td><td>1995 (27)</td><td>FLISA</td><td>1</td><td>0(0)</td><td></td></thl<></thlick<>		Entesicus tenuininnis	DRC	1995 (27)	FLISA	1	0(0)	
Kerrivoula sp.       Cameroon       2015–2017‡       Luminex       1       0–0 (0–0)         Myotis bocagii       Cameroon       2015–2017‡       Luminex       3       0–0 (0–0)         M. bocagii       DRC       1995 (27)       ELISA       22       0 (0)         M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex       5       0–0 (0–0)         Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       15       0–0 (0–0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       15       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19       0 (0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19       0 (0)         Scotophilus sp.       DRC       1979–1980 (26)       IFA<		Kerivoula sp	Guinea	2014(11)	FLISA	1	0(0)	
Myotis bocagii         Cameroon         2015–2017‡         Luminex         3         0–0 (0–0)           M. bocagii         DRC         1995 (27)         ELISA         22         0 (0)           M. bocagii         DRC         1979–1980 (26)         IFA         17         0 (0)           M. bocagii         DRC         1979–1980 (26)         IFA         17         0 (0)           Neoromicia sp.         Cameroon         2015–2017‡         Luminex         5         0–0 (0–0)           Pipistrellus nanus         DRC         1995 (27)         ELISA         2         0 (0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         15         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus dinganii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus dinganii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)		Kerivoula sp	Cameroon	2015-2017+	Luminex	1	0-0(0-0)	
M. bocagii       DRC       1995 (27)       ELISA       22       0 (0)         M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex       5       0–0 (0–0)         Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus leucogaster       Guinea       2015–2017‡       Luminex       15       0–0 (0–0)         Scotophilus leucogaster       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19       0 (0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19       0 (0)         Scotophilus sp.       DRC       1979–1980 (26)       IFA       10       0 (0)         Total       4–9/3 023 (0 13–0 30)       4–9/3 023 (0 13–0 30)		Myotis bocagii	Cameroon	2015-2017+	Luminex	3	0-0(0-0)	
M. bocagii       DRC       1979–1980 (26)       IFA       17       0 (0)         Neoromicia sp.       Cameroon       2015–2017‡       Luminex       5       0–0 (0–0)         Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus leucogaster       Guinea       2015–2017‡       Luminex       15       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus dinganii       DRC       1995 (27)       ELISA       19       0 (0)         Scotophilus dinganii       DRC       1979–1980 (26)       IFA       10       0 (0)         Scotophilus sp.       DRC       1979–1980 (26)       IFA       10       0 (0)		M bocagii	DRC	1995 (27)	FLISA	22	0(0)	
Neoromicia sp.         Cameroon         2015–2017‡         Luminex         5         0–0 (0–0)           Pipistrellus nanus         DRC         1995 (27)         ELISA         2         0 (0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)           Scotophilus nux         Cameroon         2015–2017‡         Luminex         15         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus singrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus singanii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)		M bocagii	DRC	1979 - 1980(26)	IFA	17	0(0)	
Pipistrellus nanus       DRC       1995 (27)       ELISA       2       0 (0)         Scotophilus nux       Cameroon       2015–2017‡       Luminex       6       0–0 (0–0)         Scotophilus leucogaster       Guinea       2015–2017‡       Luminex       15       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus nigrita       Guinea       2015–2017‡       Luminex       1       0–0 (0–0)         Scotophilus sigrita       BRC       1995 (27)       ELISA       19       0 (0)         Scotophilus sp.       DRC       1979–1980 (26)       IFA       10       0 (0)		Neoromicia sp	Cameroon	2015-2017+	Luminex	5	0-0(0-0)	
Scotophilus nux         Cameroon         2015–2017‡         Luminex         6         0–0 (0–0)           Scotophilus leucogaster         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus sigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus sigrita         BRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)		Pipistrellus nanus	DRC	1995 (27)	FLISA	2	0(0)	
Scotophilus leucogaster         Guinea         2015–2017‡         Luminex         15         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus nigrita         Guinea         2015–2017‡         Luminex         1         0–0 (0–0)           Scotophilus dinganii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)           Total         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)		Scotophilus nux	Cameroon	2015-2017+	Luminex	6	0 - 0 (0 - 0)	
Scotophilus nigrita         Guinea         2015–2017‡         Luminex         10         0-0 (0-0)           Scotophilus dinganii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)           Total         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)		Scotophilus leucogaster	Guinea	2015_2017+	Luminev	15	$0_{-0}(0_{-0})$	
Scotophilus dinganii         DRC         1995 (27)         ELISA         19         0 (0)           Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)           Total         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)         4–9/3 023 (0 13–0 30)		Scotophilus niarita	Guinea	2015-2017+	Luminev	1	0-0(0-0)	
Scotophilus sp.         DRC         1979–1980 (26)         IFA         10         0 (0)           Total         4–9/3 023 (0 13–0 30) <t< td=""><td></td><td>Scotophilus dingenii</td><td>DRC</td><td>1995 (27)</td><td>FLISA</td><td>19</td><td></td><td></td></t<>		Scotophilus dingenii	DRC	1995 (27)	FLISA	19		
Total 4–9/3 023 (0 13–0 30)		Scotophilus sp	DRC	1979–1980 (26)	IFA	10	0(0)	
	Total		2.10				0 (0)	4-9/3 023 (0 13-0 30)

Table 5. Zaire Ebola virus antibodies in insectivorous bats from our research, Guinea, Cameroon, and the DRC, 2015–2017, and other nublished studies

Total

\*DRC, the Democratic Republic of the Congo; IFA, immunofluorescence assay.

+For data from cited studies, the number of positive samples reported in the original study is indicated. For our results, we show the range in the number of samples simultaneously reactive with glycoprotein and nucleoprotein of Zaire Ebola virus on the basis of 4 different statistical methods used to determine cutoff values.

<u>‡This study</u>

bats housed with experimentally Zaire Ebola virusinfected bats (44). Whether this low level of infectiousness also occurs for other bat species that carry Ebola virus antibodies remains to be determined. Zaire Ebola virus was experimentally inoculated in other bat species (M. condylurus, Chaerephon pumilus, and Epomophorus wahlbergi) in only 1 study; virus replication was seen in all species, and fecal shedding was seen in E. wahlbergi bats (48). R. aegyptiacus bats experimentally infected with Marburg virus were shown to develop antibodies that protect against reinfection (49). Long-term survival with Zaire Ebola virus antibodies has been reported with E. helvum bats from Ghana but without information on protection (14). Among insectivorous bats, the presence of Ebola virus antibodies in only Mops sp. is striking, suggesting higher exposure or susceptibility compared with other insectivorous bats.

In conclusion, we demonstrated higher rates of Ebola virus antibodies in frugivorous than in insectivorous

, ,		Year of study		No.	No. (%)	
Species	Country	(reference)	Test	tested	positivé	Total, no. positive/tested (%)
Casinycteris ophiodon	Guinea	2014 (11)	ELISA	1	0	0/20
Casinycteris argynnis	Gabon, Congo	2003-2008 (13)	ELISA	18	0	
C. argynnis	DRC	1995 (27)	ELISA	1	0	
Eidolon helvum†	Guinea	2014 (11)	ELISA	6	0	21-36/1,551 (1.4-2.3)
-	Guinea	2015–2017‡	Luminex	17	0–3 (0–17.6)	. ,
	Ghana	2008 (14)	IFA	262	1 (0.39)	
	Cameroon	2015–2017‡	Luminex	158	1–9 (0.6–5.7)	
	Gabon, Congo	2003-2008 (13)	ELISA	49	0	
	DRC	1979–1980 (26)	IFA	6	0	
	DRC	2015–2017‡	Luminex	305	0-4 (0-1.3)	
	Zambia	2006–2013 (16)	ELISA	748	19 (2.55)	
Epomophorus gambianus	Guinea	2015–2017‡	Luminex	191	0–1 (0–0.5)	4-5/244 (1.6-2.0)
	Ghana	2007 (15)	ELISA	37	4 (10.82)	
Epomophorus wahlbergi	Cameroon	2015–2017‡	Luminex	16	0-0 (0-0)	
Epomops buettikoferi	Guinea	2014 (11)	ELISA	17	0	47/1,269 (3.7)
	Guinea	2015–2017‡	Luminex	4	0-0 (0-0)	
Epomops franqueti	Ghana	2007 (15)	ELISA	27	3 (11.2)	
	Cameroon	2015–2017‡	Luminex	256	0-0 (0-0)	
	Gabon, Congo	2001-2005 (6)	ELISA	117	8 (6.8)	
	Gabon, Congo	2003-2008 (13)	ELISA	805	36 (4.5)	
	DRC	2015–2017‡	Luminex	20	0-0 (0-0)	
	DRC	1979–1980 (26)	IFA	21	0	
	DRC	1995 (27)	ELISA	2	0	
Hypsygnathus monstrosus	Guinea	2015-2017‡	Luminex	8	0-0 (0-0)	15-18/347 (4.3-5.2)
	Guinea	2014 (13)	ELISA	1	0	
	Ghana	2008 (14)	IFA	3	0	
	Ghana	2007 (15)	ELISA	16	2 (12.5)	
	Cameroon	2015–2017‡	Luminex	176	0–3 (0–1.7)	
	Gabon, Congo	2001–2005 (6)	ELISA	17	4 (23.5)	
	Gabon, Congo	2003–2008 (13)	ELISA	125	9 (7.2)	
	DRC	2015–2017‡	Luminex	1	0-0 (0-0)	
Lissonycteris angolensis	Guinea	2014 (11)	ELISA	45	0	0-2/129 (0-1.6)
	Guinea	2015-2017‡	Luminex	32	0-0 (0-0)	
	DRC	2015–2017‡	Luminex	22	0-0 (0-0)	
	Cameroon	2015–2017‡	Luminex	30	0-2 (0-6.7)	
Megaloglossus azagnyi	Guinea	2014 (11)	ELISA	3	0	0/110
Megaloglossus woermanni	Cameroon	2015–2017‡	Luminex	19	0-0 (0-0)	
	Gabon, Congo	2003–2008 (13)	ELISA	49	0	
	DRC	2015–2017‡	Luminex	1	0-0 (0-0)	
	DRC	1995 (27)	ELISA	38	0	
Micropteropus pusillus	Guinea	2015–2017‡	Luminex	18	0-0 (0-0)	4/339 (1.2)
	Cameroon	2015–2017‡	Luminex	2	0-0 (0-0)	
	Gabon, Congo	2003–2008 (13)	ELISA	197	4 (2.04)	
	DRC	2015–2017‡	Luminex	44	0-0 (0-0)	
	DRC	1995 (27)	ELISA	78	0	
Myonycteris leptodon	Guinea	2014 (11)	ELISA	21	0	23-27/708 (3.2-3.8)
Myonycteris torquata	Cameroon	2015–2017‡	Luminex	21	0-0 (0-0)	
	Gabon, Congo	2001–2005 (6)	ELISA	58	4 (6.9)	
	Gabon, Congo	2003–2008 (13)	ELISA	573	19 (3.32)	
	DRC	2015–2017‡	Luminex	35	0-0 (0-0)	
Nanonycteris veldkampii	Guinea	2014 (11)	ELISA	17	0	0/21
	Ghana	2007 (15)	ELISA	4	0	
Rousettus aegyptiacus	Guinea	2015–2017‡	Luminex	228	0–1 (0–0.4)	24-33/666 (3.6-5.0)
	Cameroon	2015–2017‡	Luminex	131	0-8 (0-6.1)	
	Gabon, Congo	2003–2008 (13)	ELISA	307	24 (7.8)	
Scotonycteris zenkeri	Cameroon	2015–2017‡	Luminex	1	0-0 (0-0)	0-0/1 (0-0)
Total						138–172/5,405 (2.55–3.18)

Table 6. Zaire Ebola virus antibodies in frugivorous (Pteropodidae family) bats from our research, Guinea, Cameroon, and the DRC, 2015-2017, and published studies\*

\*DRC, the Democratic Republic of the Congo; IFA, immunofluorescence assay. +For cited studies, the number of positive samples reported in the original study is indicated. For our results, we show the range in the number of samples simultaneously reactive with glycoprotein and nucleoprotein of Zaire Ebola virus on the basis of 4 different statistical methods used to determine cutoff values.

‡This study.

bats. The total number of frugivorous species shown to be Zaire Ebola virus seropositive has increased to 8, and 1 insectivorous bat species (Mops sp.) was confirmed to be seropositive. Zaire and Sudan Ebola viruses circulate in different species across Africa, with potential co-circulation of both viruses in some species. Although we have data on >8,000 bats from >40 species, this sample size is small, given the high numbers of bats that constitute colonies. This study illustrates the complexity of tracking the animal reservoir of Ebola viruses, not only because sampling of wild bats without performing euthanasia is difficult and time-consuming but also because of the absence of a reference standard for serologic tests. To clarify the significance of Ebola virus antibodies, documenting the extent to which viral RNA and shedding can be detected in species with antibodies is crucial for predicting and controlling the risk for new outbreaks. Efforts must continue not only to sample bats but also other animals to elucidate where the virus circulates in wildlife.

#### Acknowledgments

We thank the staffs from the Ministry of Health and Ministry of Environment and the national ethics committees from the DRC, Cameroon, and Guinea for permission to perform this study. We thank all the field staff from the DRC (Guy Midingi and Servet Kimbonza); Guinea (Souana Goumou, Mamadou Kalif Diallo, Pépé Justin Beavogui, Philippe Kolié, Michel Guilavogui); and Cameroon (Innocent Ndong Bass, Aime Mebenga, Joseph Moudindo, Thomas Atemkem) for the collection of bat samples. We thank the staffs of the National Institute of Biomedical Research (Kinshasa, the DRC), the Kongo Central Provincial Government (Matadi, the DRC), and Projet PRESICA; Donald Mbohli from Project Grand Singes; and the staff of the Institut National de Santé Publique (Conakry, Guinea) for logistical support in the field. We thank Seny Mane for his involvement and support in the implementation of this project and Daouda Bangoura for his constant support for this project and the facilitation of field missions. We thank the veterinary staff from the Parc Zoologique de Montpellier and Wilhelma Zoo and Botanical Garden for providing control samples.

This work was supported in part by grants from Institut National de la Santé et de la Recherche Médicale, the Ebola Task Force, REACTing, EBO-SURSY project funded by the European Union, Institut de Recherche pour le Développement (IRD), and Christophe Mérieux Prize 2015 (to J-.J.M.T.). A.K.K. was supported by a fellowship from the IRD and the University of Montpellier (MUSE, ANR-16-IDEX-0006). C.-J.V.-A. was supported by a fellowship from IRD, Labex EpiGenMed via the National Research Agency, Programme for Future Investment (ANR-10-LABX-12-01), and the University of Montpellier.

#### About the Author

Drs. Mbala Kingebeni and Keita are researchers from ASTRE of Centre de coopération internationale en recherche agronomique pour le développement, Institut national de la recherche agronomique and University of Montpellier, Montpellier, France. Their research interests include characterization of hosts that harbor zoonotic pathogens.

#### References

- Mylne A, Brady OJ, Huang Z, Pigott DM, Golding N, Kraemer MU, et al. A comprehensive database of the geographic spread of past human Ebola outbreaks. Sci Data. 2014;1:140042. http://dx.doi.org/10.1038/sdata.2014.42
- Baize S, Pannetier D, Oestereich L, Rieger T, Koivogui L, Magassouba N, et al. Emergence of Zaire Ebola virus disease in Guinea. N Engl J Med. 2014;371:1418–25. http://dx.doi.org/ 10.1056/NEJMoa1404505
- Maganga GD, Kapetshi J, Berthet N, Kebela Ilunga B, Kabange F, Mbala Kingebeni P, et al. Ebola virus disease in the Democratic Republic of Congo. N Engl J Med. 2014;371:2083–91. http://dx.doi.org/10.1056/NEJMoa1411099
- World Health Organization. Ebola outbreak Democratic Republic of the Congo 2017. 2017 Jul 2 [cited 2018 May 3]. http://www.who.int/emergencies/ebola-DRC-2017/en/
- Pigott DM, Millear AI, Earl L, Morozoff C, Han BA, Shearer FM, et al. Updates to the zoonotic niche map of Ebola virus disease in Africa. eLife. 2016;5:e16412. http://dx.doi.org/10.7554/eLife.16412
- Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, et al. Fruit bats as reservoirs of Ebola virus. Nature. 2005;438:575–6. http://dx.doi.org/10.1038/438575a
- Leendertz SA, Gogarten JF, Düx A, Calvignac-Spencer S, Leendertz FH. Assessing the evidence supporting fruit bats as the primary reservoirs for Ebola viruses. EcoHealth. 2016;13:18–25. http://dx.doi.org/10.1007/s10393-015-1053-0
- Kamins AO, Rowcliffe JM, Ntiamoa-Baidu Y, Cunningham AA, Wood JL, Restif O. Characteristics and risk perceptions of Ghanaians potentially exposed to bat-borne zoonoses through bushmeat. EcoHealth. 2015;12:104–20. http://dx.doi.org/10.1007/ s10393-014-0977-0
- Leroy E, Gonzalez JP, Pourrut X. Ebolavirus and other filoviruses. Curr Top Microbiol Immunol. 2007;315:363–87. http://dx.doi.org/ 10.1007/978-3-540-70962-6\_15
- Leroy EM, Epelboin A, Mondonge V, Pourrut X, Gonzalez JP, Muyembe-Tamfum JJ, et al. Human Ebola outbreak resulting from direct exposure to fruit bats in Luebo, Democratic Republic of Congo, 2007. Vector Borne Zoonotic Dis. 2009;9:723–8. http://dx.doi.org/10.1089/vbz.2008.0167
- Marí Saéz A, Weiss S, Nowak K, Lapeyre V, Zimmermann F, Düx A, et al. Investigating the zoonotic origin of the West African Ebola epidemic. EMBO Mol Med. 2015;7:17–23. http://dx.doi.org/ 10.15252/emmm.201404792
- Pourrut X, Délicat A, Rollin PE, Ksiazek TG, Gonzalez JP, Leroy EM. Spatial and temporal patterns of Zaire ebolavirus antibody prevalence in the possible reservoir bat species. J Infect Dis. 2007;196(Suppl 2):S176–83. http://dx.doi.org/ 10.1086/520541
- Pourrut X, Souris M, Towner JS, Rollin PE, Nichol ST, Gonzalez JP, et al. Large serological survey showing cocirculation of Ebola and Marburg viruses in Gabonese bat populations, and a high seroprevalence of both viruses in *Rousettus aegyptiacus*. BMC Infect Dis. 2009;9:159. http://dx.doi.org/10.1186/1471-2334-9-159
- 14. Hayman DT, Emmerich P, Yu M, Wang LF, Suu-Ire R, Fooks AR, et al. Long-term survival of an urban fruit bat seropositive for

Ebola and Lagos bat viruses. PLoS One. 2010;5:e11978. http://dx.doi.org/10.1371/journal.pone.0011978

- Hayman DT, Yu M, Crameri G, Wang LF, Suu-Ire R, Wood JL, et al. Ebola virus antibodies in fruit bats, Ghana, West Africa. Emerg Infect Dis. 2012;18:1207–9. http://dx.doi.org/10.3201/ eid1807.111654
- Ogawa H, Miyamoto H, Nakayama E, Yoshida R, Nakamura I, Sawa H, et al. Seroepidemiological prevalence of multiple species of filoviruses in fruit bats (*Eidolon helvum*) migrating in Africa. J Infect Dis. 2015;212(Suppl 2):S101–8. http://dx.doi.org/10.1093/ infdis/jiv063
- Towner JS, Pourrut X, Albariño CG, Nkogue CN, Bird BH, Grard G, et al. Marburg virus infection detected in a common African bat. PLoS One. 2007;2:e764. http://dx.doi.org/10.1371/ journal.pone.0000764
- Towner JS, Amman BR, Sealy TK, Carroll SA, Comer JA, Kemp A, et al. Isolation of genetically diverse Marburg viruses from Egyptian fruit bats. PLoS Pathog. 2009;5:e1000536. http://dx.doi.org/10.1371/journal.ppat.1000536
- Kuzmin IV, Niezgoda M, Franka R, Agwanda B, Markotter W, Breiman RF, et al. Marburg virus in fruit bat, Kenya. Emerg Infect Dis. 2010;16:352–4. http://dx.doi.org/10.3201/eid1602.091269
- Swanepoel R, Smit SB, Rollin PE, Formenty P, Leman PA, Kemp A, et al.; International Scientific and Technical Committee for Marburg Hemorrhagic Fever Control in the Democratic Republic of the Congo. Studies of reservoir hosts for Marburg virus. Emerg Infect Dis. 2007;13:1847–51. http://dx.doi.org/ 10.3201/eid1312.071115
- Negredo A, Palacios G, Vázquez-Morón S, González F, Dopazo H, Molero F, et al. Discovery of an ebolavirus-like filovirus in europe. PLoS Pathog. 2011;7:e1002304. http://dx.doi.org/10.1371/ journal.ppat.1002304
- Yang XL, Zhang YZ, Jiang RD, Guo H, Zhang W, Li B, et al. Genetically diverse filoviruses in *Rousettus* and *Eonycteris* spp. bats, China, 2009 and 2015. Emerg Infect Dis. 2017;23:482–6. http://dx.doi.org/10.3201/eid2303.161119
- World Health Organization. Ebola situation report 30 March 2016. 2016 [cited 2018 May 3]. http://apps.who.int./ebola/ current-situation/ebola-situation-report-30-march-2016
- Dudas G, Carvalho LM, Bedford T, Tatem AJ, Baele G, Faria NR, et al. Virus genomes reveal factors that spread and sustained the Ebola epidemic. Nature. 2017;544:309–15. http://dx.doi.org/ 10.1038/nature22040
- Pigott DM, Deshpande A, Letourneau I, Morozoff C, Reiner RC Jr, Kraemer MUG, et al. Local, national, and regional viral haemorrhagic fever pandemic potential in Africa: a multistage analysis. Lancet. 2017;390:2662–72. http://dx.doi.org/10.1016/ S0140-6736(17)32092-5
- Breman JG, Johnson KM, van der Groen G, Robbins CB, Szczeniowski MV, Ruti K, et al.; Ebola Virus Study Teams. A search for Ebola virus in animals in the Democratic Republic of the Congo and Cameroon: ecologic, virologic, and serologic surveys, 1979–1980. J Infect Dis. 1999;179(Suppl 1):S139–47. http://dx.doi.org/10.1086/514278
- Leirs H, Mills JN, Krebs JW, Childs JE, Akaibe D, Woollen N, et al. Search for the Ebola virus reservoir in Kikwit, Democratic Republic of the Congo: reflections on a vertebrate collection. J Infect Dis. 1999;179(Suppl 1):S155–63. http://dx.doi.org/ 10.1086/514299
- Ayouba A, Touré A, Butel C, Keita AK, Binetruy F, Sow MS, et al. Development of a sensitive and specific serological assay based on luminex technology for detection of antibodies to Zaire Ebola virus. J Clin Microbiol. 2017;55:165–76. http://dx.doi.org/ 10.1128/JCM.01979-16
- 29. Peel AJ, McKinley TJ, Baker KS, Barr JA, Crameri G, Hayman DT, et al. Use of cross-reactive serological assays for detecting novel

pathogens in wildlife: assessing an appropriate cutoff for henipavirus assays in African bats. J Virol Methods. 2013;193:295– 303. http://dx.doi.org/10.1016/j.jviromet.2013.06.030

- 30. Gilbert AT, Fooks AR, Hayman DT, Horton DL, Müller T, Plowright R, et al. Deciphering serology to understand the ecology of infectious diseases in wildlife. EcoHealth. 2013;10:298–313. http://dx.doi.org/10.1007/s10393-013-0856-0
- Lardeux F, Torrico G, Aliaga C. Calculation of the ELISA's cut-off based on the change-point analysis method for detection of *Trypanosoma cruzi* infection in Bolivian dogs in the absence of controls. Mem Inst Oswaldo Cruz. 2016;111:501–4. http://dx.doi.org/10.1590/0074-02760160119
- Killick R, Eckley IA. changepoint: an R package for changepoint analysis. J Stat Softw. 2014;58:1–19. http://dx.doi.org/10.18637/ jss.v058.i03
- Hinkley DV. Inference about the change-point in a sequence of random variables. Biometrika. 1970;57:1–17. http://dx.doi.org/ 10.1093/biomet/57.1.1
- Laing ED, Mendenhall IH, Linster M, Low DHW, Chen Y, Yan L, et al. Serologic evidence of fruit bat exposure to filoviruses, Singapore, 2011–2016. Emerg Infect Dis. 2018;24:114–7. http://dx.doi.org/10.3201/eid2401.170401
- Cullen AC, Frey HC. Probabilistic techniques in exposure assessment. New York: Plenum Press; 1999. p. 81–159.
- Delignette-Muller ML, Dutang C. fitdistrplus: an R package for fitting distributions. J Stat Softw. 2015;64:1–34. http://dx.doi.org/ 10.18637/jss.v064.i04
- 37. Monleau M, Montavon C, Laurent C, Segondy M, Montes B, Delaporte E, et al. Evaluation of different RNA extraction methods and storage conditions of dried plasma or blood spots for human immunodeficiency virus type 1 RNA quantification and PCR amplification for drug resistance testing. J Clin Microbiol. 2009;47:1107–18. http://dx.doi.org/10.1128/ JCM.02255-08
- Guichet E, Serrano L, Laurent C, Eymard-Duvernay S, Kuaban C, Vidal L, et al. Comparison of different nucleic acid preparation methods to improve specific HIV-1 RNA isolation for viral load testing on dried blood spots. J Virol Methods. 2018;251:75–9. http://dx.doi.org/10.1016/j.jviromet.2017.10.014
- Irwin DM, Kocher TD, Wilson AC. Evolution of the cytochrome b gene of mammals. J Mol Evol. 1991;32:128–44. http://dx.doi.org/ 10.1007/BF02515385
- 40. Kocher TD, Thomas WK, Meyer A, Edwards SV, Pääbo S, Villablanca FX, et al. Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. Proc Natl Acad Sci U S A. 1989;86:6196–200. http://dx.doi.org/10.1073/pnas.86.16.6196
- He B, Feng Y, Zhang H, Xu L, Yang W, Zhang Y, et al. Filovirus RNA in fruit bats, China. Emerg Infect Dis. 2015;21:1675–7. http://dx.doi.org/10.3201/eid2109.150260
- Han BA, Schmidt JP, Alexander LW, Bowden SE, Hayman DT, Drake JM. Undiscovered bat hosts of filoviruses. PLoS Negl Trop Dis. 2016;10:e0004815. http://dx.doi.org/10.1371/ journal.pntd.0004815
- Rimoin AW, Lu K, Bramble MS, Steffen I, Doshi RH, Hoff NA, et al. Ebola virus neutralizing antibodies detectable in survivors of theYambuku, Zaire outbreak 40 years after infection. J Infect Dis. 2018;217:223–31. http://dx.doi.org/10.1093/infdis/jix584
- Paweska JT, Storm N, Grobbelaar AA, Markotter W, Kemp A, Jansen van Vuren P. Experimental inoculation of Egyptian fruit bats (*Rousettus aegyptiacus*) with Ebola virus. Viruses. 2016;8:29. http://dx.doi.org/10.3390/v8020029
- Amman BR, Jones ME, Sealy TK, Uebelhoer LS, Schuh AJ, Bird BH, et al. Oral shedding of Marburg virus in experimentally infected Egyptian fruit bats (*Rousettus aegyptiacus*). J Wildl Dis. 2015;51:113–24. http://dx.doi.org/10.7589/2014-08-198

- 46. Schuh AJ, Amman BR, Jones ME, Sealy TK, Uebelhoer LS, Spengler JR, et al. Modelling filovirus maintenance in nature by experimental transmission of Marburg virus between Egyptian rousette bats. Nat Commun. 2017;8:14446. http://dx.doi.org/10.1038/ncomms14446
- 47. Jones ME, Schuh AJ, Amman BR, Sealy TK, Zaki SR, Nichol ST, et al. Experimental inoculation of Egyptian rousette bats (*Rousettus aegyptiacus*) with viruses of the *Ebolavirus* and *Marburgvirus* genera. Viruses. 2015;7:3420–42. http://dx.doi.org/10.3390/v7072779
- 48. Swanepoel R, Leman PA, Burt FJ, Zachariades NA, Braack LE, Ksiazek TG, et al. Experimental inoculation of plants and

animals with Ebola virus. Emerg Infect Dis. 1996;2:321–5. http://dx.doi.org/10.3201/eid0204.960407

49. Storm N, Jansen Van Vuren P, Markotter W, Paweska JT. Antibody responses to Marburg virus in Egyptian rousette bats and their role in protection against infection. Viruses. 2018;10:73. http://dx.doi.org/10.3390/v10020073

Adress for correspondence: Martine Peeters, TransVIHMI, Institut de Recherche pour le Développement, 911 Ave Agropolis, BP34394 Montpellier CEDEX 1, France; e-mail: martine.peeters@ird.fr



### EMERGING INFECTIOUS DISEASES

### February 2016

# Ebola

- Ebola and Its Control in Liberia, 2014–2015
- Epidemiology of Epidemic Ebola Virus Disease in Conakry and Surrounding Prefectures, Guinea, 2014–2015
- Hospital Preparations for Viral Hemorrhagic Fever Patients and Experience Gained from the Admission of an Ebola Patient
- Trematode Fluke *Procerovum varium* as Cause of Ocular Inflammation in Children, South India
- Association between Landscape Factors and Spatial Patterns of *Plasmodium knowlesi* Infections in Sabah, Malaysia
- Feasibility of Xpert Ebola Assay in Médecins Sans Frontières Ebola Program, Guinea
- Prognostic Indicators for Ebola Patient Survival
- Invasive Group A *Streptococcus* Infection among Children, Rural Kenya
- Randomized Controlled Trial of Hospital-Based Hygiene and Water Treatment Intervention (CHoBI7) to Reduce Cholera
- Sustained Transmission of Pertussis in Vaccinated, 1–5-Year-Old Children in a Preschool, Florida, USA
- Molecular Characterization of Invasive Streptococcus dysgalactiae subsp. equisimilis, Japan
- Epidemiology of Serotype 1 Invasive Pneumococcal Disease, South Africa, 2003–2013
- Hemorrhagic Fever with Renal Syndrome, Zibo City, China, 2006–2014

- Dogs and Opossums Positive for Vaccinia Virus during Outbreak Affecting Cattle and Humans, São Paulo State, Brazil
- African Buffalo Movement and Zoonotic Disease Risk across Transfrontier Conservation Areas, Southern Africa
- Anaplasmataceae–Specific PCR for Diagnosis and Therapeutic Guidance for Symptomatic Neoehrlichiosis in Immunocompetent Host
- Candidatus Coxiella massiliensis Infection
- Ebola Virus Persistence in Semen Ex Vivo
- Ebola Virus RNA Stability in Human Blood and Urine in West Africa's Environmental Conditions
- Uveitis and Systemic Inflammatory Markers in Convalescent Phase of Ebola Virus Disease
- Louseborne Relapsing Fever among East African Refugees, Italy, 2015
- Mediterranean Fin Whales (*Balaenoptera physalus*) Threatened by Dolphin Morbillivirus
- *Blastomyces gilchristii* as Cause of Fatal Acute Respiratory Distress Syndrome
- Effectiveness of Meningococcal B Vaccine against Endemic Hypervirulent *Neisseria meningitidis* W Strain, England
- Frequency and Distribution of Rickettsiae, Borreliae, and Ehrlichiae Detected in Human-Parasitizing Ticks, Texas, USA
- High Prevalence of *Borrelia miyamotoi* among Adult Blacklegged Ticks from White-Tailed Deer

#### To revisit the February 2016 issue, go to: https://wwwnc.cdc.gov/eid/articles/issue/22/2/table-of-contents