



HAL
open science

Consequences of insecticide resistance on malaria transmission

Haoues Alout, Benjamin Roche, Roch Kounbobr Dabiré, Anna Cohuet

► **To cite this version:**

Haoues Alout, Benjamin Roche, Roch Kounbobr Dabiré, Anna Cohuet. Consequences of insecticide resistance on malaria transmission. *PLoS Pathogens*, 2017, 13 (9), pp.e1006499. <10.1371/journal.ppat.1006499>. <hal-01942723>

HAL Id: hal-01942723

<https://hal.umontpellier.fr/hal-01942723v1>

Submitted on 3 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization


PEARLS

Consequences of insecticide resistance on malaria transmission

Haoues Alout^{1*}, Benjamin Roche^{2,3}, Roch Kounbobr Dabiré⁴, Anna Cohuet²

1 Institut des Sciences de l'Evolution de Montpellier, CNRS, IRD, University of Montpellier, Montpellier, France, **2** Institut de recherche pour le développement (IRD), Maladies Infectieuses et Vecteurs, Ecologie, Génétique, Evolution et Contrôle (MIVEGEC), Montpellier, France, **3** UMI IRD/UPMC 209, Unité de Modélisation Mathématique et Informatique des Systèmes Complexes (UMMISCO lab), Bondy, France, **4** Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso

* haoues.alout@gmail.com



Malaria burden and control

Human malaria is caused by protozoan parasites of the genus *Plasmodium* (*Plasmodium falciparum*, *P. vivax*, *P. malariae*, *P. ovale*, and *P. knowlesi*). In 2015, 212 million malaria cases were reported worldwide with approximately 429,000 deaths, 92% of which occurred in Africa. However, several countries within sub-Saharan Africa have reported decreased numbers of malaria cases over the last several decades (<http://www.who.int/malaria/publications/world-malaria-report-2016/en/>). *Plasmodium* parasites are dependent on completing a complex life cycle in mosquitoes of the genus *Anopheles*, specifically, for transmission to occur. Thus, reducing the mosquito abundance or interfering with its ability to support the parasite cycle (i.e., the vector competence) can interrupt malaria transmission. However, not every species of *Anopheles* is a vector, and not all species of *Plasmodium* occur in all regions. In natural populations, the combination of *Plasmodium*–*Anopheles* species is very specific and leads to distinct epidemiology and transmission patterns. For example, the transmission of malaria parasites between humans is due primarily to the *A. gambiae s.l.*–*P. falciparum* combination, while in Southeast Asia, *A. dirus* and *A. minimus*–*P. vivax* are the main vectorial systems. This transmission potential between human hosts of a specific *Anopheles* population is characterized by the “vectorial capacity” and depends on the abundance, the feeding rate, the longevity, and the vector competence of female mosquitoes as well as the extrinsic incubation period of the parasite, which is the time for the pathogen to reach the transmissible stage from the midgut to the salivary glands of its vector. Between 2000 and 2015, the number of malaria infection deaths have been halved, with 79% of this reduction being attributed to insecticide-based vector control (insecticide-treated nets [ITNs] and indoor residual spraying [IRS]) [1]. Therefore, insecticide-based vector control still represents the most important and affordable method to alleviate the malaria burden by interrupting the transmission cycle, although insecticide-resistance frequency and intensity have increased dramatically in malaria vector populations. In an attempt to explain this discrepancy, we present 5 facts about insecticide resistance in regards to malaria control when the epidemiological and ecological contexts are considered.

The threat of insecticide resistance

Only a few insecticides have been approved for public health, and, among them, only the pyrethroids are allowed for ITNs. Nowadays, resistance against this insecticide family is widespread among malaria vectors in Africa (see IR mapper [www.irmapper.com] for the latest update) and is expected to jeopardize the success of malaria control. According to the World Health

OPEN ACCESS

Citation: Alout H, Roche B, Dabiré RK, Cohuet A (2017) Consequences of insecticide resistance on malaria transmission. PLoS Pathog 13(9): e1006499. <https://doi.org/10.1371/journal.ppat.1006499>

Editor: Laura J. Knoll, University of Wisconsin Medical School, UNITED STATES

Published: September 7, 2017

Copyright: © 2017 Alout et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: HA is supported by Marie Skłodowska-Curie under the grant agreement number 749897, BR is supported by the ANR project PANIC, and AC is supported by ANSES under the grant number PNR EST 2016, by Languedoc-Roussillon “Chercheur(se)s d’avenir 2015” and Expertise France: Initiative 5% under the grant number 15SANIN2013. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Organization (<http://www.who.int/malaria/publications/atoz/gpirm/en/>), resistance to insecticides is defined as “an ability to tolerate doses of toxicants, which would prove lethal to the majority of individuals in a normal population of the same insect species”. Resistance arises from the selection of individuals able to survive and reproduce in an insecticide-treated environment or after being in contact with insecticides. Mechanisms that decrease the insecticide toxicity can rely on modifications in one or several genes of the mosquito, and, as a result, resistance is a heritable trait. In the presence of insecticides, the frequency of alleles responsible for insecticide resistance increases in the population because they confer a strong advantage, which is expected to hinder the effect of insecticides on vector survival and abundance.

The impact of insecticide resistance on vector control efficacy is usually measured by comparing survival rates of mosquito strains or natural populations exposed to various insecticides, both in the laboratory (e.g., WHO cones or tube assays) and in natural settings (e.g., experimental huts). The loss of insecticide efficacy may thus lead to partial operational failure of malaria control and probably to increased disease transmission. A meta-analysis of the published results synthesized the ITN-associated mortality of pyrethroid-resistant vectors and predicted a strong negative epidemiological impact of insecticide resistance [2].

Discrepancy between entomological and epidemiological studies on insecticide-resistance impact

In contrast to these expectations based on entomological observations, implementation of vector control tools (ITNs and/or IRS) has significantly decreased malaria incidence and parasite infection prevalence in children in multiple endemic countries across Africa (Equatorial Guinea, Burundi, Ivory Coast, Malawi, and Kenya), despite moderate-to-high pyrethroid resistance observed in local malaria vectors [3–6]. These results show reductions in malaria indices following ITN distribution similar to those in areas with pyrethroid-susceptible vectors [4,7,8]. Only 1 study suggested that the selection of insecticide resistance has led to a rebound in malaria incidence in South Africa, but this could also be attributed to climatic factors, reestablishment of other malaria vector species, and resistance to antimalarial drugs [9]. There is thus no clear evidence that insecticide resistance would lead to malaria control failure; however, these studies were not specifically designed to test its epidemiological impact and thus lack the power to assess it correctly. A trial in 5 countries (Benin, Cameroon, Kenya, Sudan, and India) comparing malaria incidence in areas of insecticide resistant versus susceptible vector populations is ongoing, and preliminary results did not show any evidence of an association between malaria burden and pyrethroid resistance (<http://www.who.int/malaria/publications/atoz/insecticide-resistance-implications/en/>), further supporting the previous observations that entomological efficacy of vector control does not directly correlate with epidemiological efficacy. In order to assess the actual threat of insecticide resistance on malaria control, it is fundamental to explain this contrast between the entomological and epidemiological outcomes of ITNs in insecticide-resistant vector populations.

The phenotypic expression of insecticide resistance in malaria vectors

The detection of insecticide resistance in the field relies on standard protocols provided by WHO that use 1 insecticide dose, 1 exposure time, and the observation of mortality 24 h post-exposure. However, the absence of mortality resulting from such methods does not imply a complete absence of mortality due to insecticides. For instance, the WHO protocols fail at capturing insecticide-induced delayed mortality, which is mortality occurring more than 24 h later [10]. In addition, the phenotypic expression of resistance and the resistance level are

highly dependent upon environmental variables like temperature [11], food quality/quantity [12], multiple blood meals [13] and preexisting pesticide exposure [10], all variations that are not captured in standardized bioassays. Therefore, the standard protocols, while useful to monitor insecticide susceptibility during vector control campaigns, may not provide evidence for significant failure of disease control.

Similarly, the level of resistance has been shown to decline with age (reviewed in [14]), so that young mosquitoes found resistant could still die from exposure to insecticides when they get older. The fact that mosquitoes exposed to insecticide tend to die early-on in spite of their resistance alleles or the fact that they become more susceptible to insecticides when aging can have tremendous implications in terms of epidemiology: most of the infectious mosquitoes are indeed old females (*Plasmodium* requires about 10 days to become infectious once ingested by *Anopheles* vectors, which have a daily survivorship of approximately 0.8/day [15]), so insecticides may still be efficient at reducing their proportion and, thus, at controlling malaria transmission.

The fitness cost of insecticide resistance and its impact on malaria transmission

The mutations responsible for resistance are often associated with modification of physiological processes or resource availability, which often leads to decreased performance and fitness disadvantage (i.e., cost). The deleterious pleiotropic effects (i.e., the negative influence on unrelated phenotypic traits) of insecticide-resistance alleles could affect a wide range of vector life-history traits, such as longevity, biting behavior, and vector competence [16], which are important factors of the vectorial capacity to transmit the pathogen. For instance, insecticide-resistance mutations have been shown to increase the vector competence of *A. gambiae* for malaria parasites [17], which may potentiate malaria transmission. The cost of insecticide resistance also depends on other biotic (i.e., parasite infection) and abiotic (i.e., environmental) factors, which may increase or decrease mosquitoes' ability to transmit parasites. Parasite infection itself may impose an additional cost to vectors carrying insecticide resistance alleles, thereby further influencing the mosquito life-history traits involved in vectorial capacity and also the phenotypic expression of insecticide resistance. Consistently, infection by *Plasmodium* parasites was shown to alter the survival of resistant mosquitoes in the absence of insecticides [18] and to partially restore the susceptibility to insecticide in *A. gambiae* carrying the *kdr* mutation [19]. Moreover, exposure to pyrethroid insecticides reduced the prevalence of infection by *P. falciparum* in *A. gambiae* s.s. harboring the *kdr* mutation [20]. Altogether, these observations appear as assets for vector control. Therefore, the interactions between insecticide resistance, exposure, and infection induce contrasted effects on mosquito traits, so anticipating the “net” outcome of insecticide resistance on malaria epidemiology remains challenging.

Conclusion

Considering only the direct effect of insecticide on mosquito survival often leads to the simplistic conclusion that insecticide resistance annihilates the efforts to control malaria incidence. There are at least 5 facts about the effect of insecticide resistance on malaria control: (1) the discrepancies between the entomological and epidemiological studies of malaria vector control efficacy in relation to insecticide resistance, (2) the overestimated phenotype of resistance in standard protocols compared to natural context (i.e., increased insecticide toxicity due to delayed insecticide effects or to age), (3) the fitness cost associated with insecticide resistance (on mosquito density, biting behavior, vector competence, and survival), (4) the increased parasite-induced mortality in insecticide-resistant mosquitoes (interactive cost between infection

and resistance), and (5) the impact of insecticides on vector–parasite interactions (i.e., increased toxicity on infected/infectious vectors, reduced parasite development, and reduced transmission). In addition to these, other reasons for successful malaria control despite the selection of insecticide resistance exist, such as bed–net physical barrier and housing improvement (such as window screens and eave tubes). The local epidemiological, ecological, and entomological context (population structure, seasonality, multiple mechanisms of insecticide resistance, etc. . .) are thus crucial to consider to reach a more precise estimate of insecticide-resistance impact on vectorial capacity. We should also consider that continuous insecticide selective pressure, as well as parasitism, may select for enhanced resistance phenotypes and/or for reduced fitness cost. The epidemiological consequences of insecticide resistance are thus expected to be continuously moving. This emphasizes the necessity to precisely and continuously monitor the consequences of vector control strategies, considering the vectorial system in all its ecological interactions.

Acknowledgments

We are indebted to the anonymous reviewers for their excellent comments that led to improvement of the manuscript. Contribution number 2017–125 of the Institut des Sciences de l’Evolution de Montpellier, UMR 5554 CNRS-UM-IRD-EPHE.

References

1. Cibulskis RE, Alonso P, Aponte J, Aregawi M, Barrette A, Bergeron L, et al. Malaria: Global progress 2000–2015 and future challenges. *Infect Dis Poverty*. Infectious Diseases of Poverty; 2016; 5: 61. <https://doi.org/10.1186/s40249-016-0151-8> PMID: 27282148
2. Churcher TS, Lissenden N, Griffin JT, Worrall E, Ranson H. The impact of pyrethroid resistance on the efficacy and effectiveness of bednets for malaria control in Africa. *Elife*. 2016; 5: e16090. <https://doi.org/10.7554/eLife.16090> PMID: 27547988
3. Kleinschmidt I, Sharp B, Benavente LE, Schwabe C, Torrez M, Kuklinski J, et al. Reduction in infection with *Plasmodium falciparum* one year after the introduction of malaria control interventions on Bioko island, Equatorial Guinea. *Am J Trop Med Hyg*. 2006; 74: 972–978. 74/6/972 [pii] PMID: 16760506
4. Henry MC, Asisi SB, Rogier C, Dossou-Yovo J, Chandre F, Guillet P, et al. Protective efficacy of lambda-cyhalothrin treated nets in *Anopheles gambiae* pyrethroid resistance areas of Côte d’Ivoire. *Am J Trop Med Hyg*. 2005; 73: 859–864. PMID: 16282294
5. Lindblade KA, Mwandama D, Mzilahowa T, Steinhart L, Gimnig J, Shah M, et al. A cohort study of the effectiveness of insecticide-treated bed nets to prevent malaria in an area of moderate pyrethroid resistance, Malawi. *Malar J*. 2015; 14: 31. <https://doi.org/10.1186/s12936-015-0554-1> PMID: 25627987
6. Okoyo C, Mwandawiro C, Kihara J, Simiyu E, Gitonga CW, Noor AM, et al. Comparing insecticide-treated bed net use to *Plasmodium falciparum* infection among schoolchildren living near Lake Victoria, Kenya. *Malar J*. 2015; 14: 515. <https://doi.org/10.1186/s12936-015-1031-6> PMID: 26696416
7. Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database of Systematic Reviews*. 2004. p. CD000363. <https://doi.org/10.1002/14651858.CD000363.pub2> PMID: 15106149
8. Alout H, Roche B, Dabiré RK, Cohuet A. Malaria vector control still matters despite insecticide resistance. *Trends Parasitol*. 2017. In press. <https://doi.org/10.1016/j.pt.2017.04.006> PMID: 28499699
9. Maharaj R, Mthembu DJ, Sharp BL. Impact of DDT re-introduction on malaria transmission in KwaZulu-Natal. *South African Med J*. 2005; 95: 871–874.
10. Viana M, Hughes A, Matthiopoulos J, Ranson H, Ferguson HM. Delayed mortality effects cut the malaria transmission potential of insecticide-resistant mosquitoes. *Proc Natl Acad Sci*. 2016; 113: 8975–8980. <https://doi.org/10.1073/pnas.1603431113> PMID: 27402740
11. Glunt KD, Paaijmans KP, Read AF, Thomas MB. Environmental temperatures significantly change the impact of insecticides measured using WHOPES protocols. *Malar J*. 2014; 13: 350. <https://doi.org/10.1186/1475-2875-13-350> PMID: 25187231
12. Kulma K, Saddler A, Koella JC. Effects of age and larval nutrition on phenotypic expression of insecticide-resistance in *Anopheles* mosquitoes. *PLoS One*. 2013; 8: e58322. <https://doi.org/10.1371/journal.pone.0058322> PMID: 23484017

13. Oliver S V, Brooke BD. The effect of multiple blood-feeding on the longevity and insecticide resistant phenotype in the major malaria vector *Anopheles arabiensis* (Diptera: Culicidae). *Parasites and Vectors*. 2014; 7: 390. <https://doi.org/10.1186/1756-3305-7-390> PMID: 25150975
14. Saddler A, Koella JC. Modelling the impact of declining insecticide resistance with mosquito age on malaria transmission. *Malar World J*. 2015; 6: 1–7.
15. Beier JC. Malaria parasite development in mosquitoes. *Annu Rev Entomol*. 1998; 43: 519–43. <https://doi.org/10.1146/annurev.ento.43.1.519> PMID: 9444756
16. Rivero A, Vézilier J, Weill M, Read AF, Gandon S. Insecticide control of vector-borne diseases: When is insecticide resistance a problem? *PLoS Pathog*. 2010; 6: 5–6. <https://doi.org/10.1371/journal.ppat.1001000> PMID: 20700451
17. Alout H, Krajacich BJ, Meyers JI, Grubaugh ND, Brackney DE, Kobylinski KC, et al. Evaluation of ivermectin mass drug administration for malaria transmission control across different West African environments. *Malar J*. 2014; 13: 417. <https://doi.org/10.1186/1475-2875-13-417> PMID: 25363349
18. Alout H, Dabiré RK, Djogbénou LS, Abate L, Corbel V, Chandre F, et al. Interactive cost of *Plasmodium* infection and insecticide resistance in the malaria vector *Anopheles gambiae*. *Sci Rep*. 2016; 6: 29755. <https://doi.org/10.1038/srep29755> PMID: 27432257
19. Alout H, Yameogo B, Djogbenou LS, Chandre F, Dabire RK, Corbel V, et al. Interplay between *Plasmodium* infection and resistance to insecticides in vector mosquitoes. *J Infect Dis*. 2014; 210: 1464–1470. <https://doi.org/10.1093/infdis/jiu276> PMID: 24829465
20. Alout H, Djegbe I, Chandre F, Djogbenou LS, Dabire RK, Corbel V, et al. Insecticide exposure impacts vector-parasite interactions in insecticide-resistant malaria vectors. *Proc R Soc B Biol Sci*. 2014; 281: 20140389–20140389. <https://doi.org/10.1098/rspb.2014.0389> PMID: 24850924