



HAL
open science

Meeting the challenge of tick-borne disease control: A proposal for 1000 Ixodes genomes

Maria Murgia, Lesley Bell-Sakyi, José de La Fuente, Timothy Kurtti, Benjamin Makepeace, Ben Mans, Karen Mccoy, Ulrike Munderloh, Olivier Plantard, Claude Rispe, et al.

► To cite this version:

Maria Murgia, Lesley Bell-Sakyi, José de La Fuente, Timothy Kurtti, Benjamin Makepeace, et al.. Meeting the challenge of tick-borne disease control: A proposal for 1000 Ixodes genomes. *Ticks and Tick-borne Diseases*, 2019, 10 (1), pp.213-218. 10.1016/j.ttbdis.2018.08.009 . hal-01998059

HAL Id: hal-01998059

<https://hal.umontpellier.fr/hal-01998059>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Meeting the challenge of tick-borne disease control: A proposal for 1000 *Ixodes* genomes

Author: Maria V. Murgia

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Lesley Bell-Sakyi

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: José de la Fuente

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Timothy J. Kurtti

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Benjamin L. Makepeace

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Ben Mans

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Karen D. McCoy

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Uli Munderloh

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Authors: Olivier Plantard, Claude Rispe

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Authors: Manuel Rodriguez Valle, Ala Tabor

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091


To appear in:

Author: Saravanan Thangamani

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Jyothi Thimmapuram



PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Author: Catherine A. Hill

PII: S1877-959X(18)30074-8
DOI: <https://doi.org/10.1016/j.ttbdis.2018.08.009>
Reference: TTBDIS 1091

To appear in:

Received date: 16-2-2018

Please cite this article as: Murgia MV, Bell-Sakyi L, de la Fuente J, Kurtti TJ, Makepeace BL, Mans B, McCoy KD, Munderloh U, Plantard O, Risper C, Valle MR, Tabor A, Thangamani S, Thimmapuram J, Hill CA, Meeting the challenge of tick-borne disease control: A proposal for 1000 *Ixodes* genomes, *Ticks and Tick-borne Diseases* (2018), <https://doi.org/10.1016/j.ttbdis.2018.08.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

"Meeting the challenge of tick-borne disease control: A proposal for 1000 *Ixodes* genomes"

Ixodes 1000 Genomes Consortium, Leadership Team

Maria V. Murgia^a

^a*Department of Entomology, Purdue University, West Lafayette, IN, United States*

Lesley Bell-Sakyi^b

^b*Department of Infection Biology, Institute of Infection and Global Health, University of Liverpool, Liverpool, United Kingdom*

José de la Fuente^{c,d}

^c*SaBio Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13005 Ciudad Real, Spain.*

^d*Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK 74078, USA.*

Timothy J. Kurtti^e

^e*Department of Entomology, University of Minnesota, Saint Paul, MN, USA*

Benjamin L. Makepeace^b

^b*Department of Infection Biology, Institute of Infection and Global Health, University of Liverpool, Liverpool, United Kingdom*

Ben Mans^{f,g}

^f*Epidemiology, Parasites and Vectors, Onderstepoort Veterinary Research, Agricultural Research Council, Onderstepoort, South Africa*

^g*Department of Life and Consumer Sciences, University of South Africa, South Africa*

Karen D. McCoy^h

^h*MIVEGEC, UMR CNRS – IRD, University of Montpellier, Centre IRD, Montpellier, France*

Uli Munderloh^e

^e*Department of Entomology, University of Minnesota, St. Paul, MN, USA*

Olivier Plantardⁱ

ⁱ*BIOEPAR, INRA, Oniris, Université Bretagne Loire, Nantes, France*

Claude Rispeⁱ

ⁱ*BIOEPAR, INRA, Oniris, Université Bretagne Loire, Nantes, France*

Manuel Rodriguez Valle^j

^j*Queensland Alliance for Agriculture & Food Innovation, The University of Queensland, St. Lucia, Queensland, Australia*

Ala Tabor^j

^j*Queensland Alliance for Agriculture & Food Innovation, The University of Queensland, St. Lucia, Queensland, Australia*

Saravanan Thangamani^k

^k*Department of Pathology, Institute for Human Infections and Immunity, University of Texas Medical Branch, Galveston, Texas, USA*

Jyothi Thimmapuram^l

^l*Bioinformatics Core, Purdue University, West Lafayette, IN, United States*

Catherine A. Hill^{a,m,*}

^a*Department of Entomology, Purdue University, West Lafayette, IN, United States*

^m*Purdue Institute for Inflammation, Immunology and Infectious Diseases, Purdue University, West Lafayette, IN, United States*

*Corresponding author. Tel.: +1 765 496 6157

Email address: hillca@purdue.edu (C.A. Hill)

Keywords: *Ixodes*; Ixodida; Genomics; Tick; Tick-borne diseases; Ultra-long read

ACCEPTED MANUSCRIPT

Introduction

At the ‘One Health’ 9th Tick and Tick-borne Pathogen Conference and 1st Asia Pacific Rickettsia Conference (TTP9-APRC1; <http://www.ttp9-aprc1.com>), 27 August-1 September 2017 in Cairns, Australia, members of the tick and tick-borne disease (TBD) research communities assembled to discuss a high priority research agenda. Diseases transmitted by hard ticks (subphylum Chelicerata; subclass Acari; family Ixodidae) have substantial impacts on public health and are on the rise globally due to human population growth and change in geographic ranges of tick vectors (de la Fuente et al., 2016). The genus *Ixodes* is a global menace. Members of the genus impact human and animal health directly via host parasitism, and indirectly via transmission of multiple viral, bacterial and protozoan diseases. The first tick genome assembly was completed in 2016 for *Ixodes scapularis* (blacklegged tick), the North American vector of Lyme disease (LD), human babesiosis, human anaplasmosis and Powassan virus (Gulia-Nuss et al., 2016). The assembly provided insight into the genome biology of hard (ixodid) ticks and supported molecular studies for many species of Acari (ticks and mites). Draft genome assemblies are available for *Ixodes ricinus* (castor bean tick; Cramaro et al., 2015; 2017) and *Rhipicephalus (Boophilus) microplus* (southern cattle tick; Barrero et al., 2017). However, high quality reference assemblies to rival those produced for the mosquito vectors *Anopheles gambiae* (Neafsey et al., 2015) and *Aedes aegypti* (Mathews et al., 2017), have not been produced for a tick vector. Conference attendees identified the need to expand genomic resources for tick research, beginning with the genus *Ixodes* – one of the most important phyletic groups affecting human and animal health worldwide. Inspired by the 1000 *Anopheles* genomes effort (Malaria Genomic Epidemiology Network, Ag1000G), an ambitious goal to sequence and assemble the genomes of 1000 *Ixodes* ticks was proposed. The *Ixodes* 1000 genomes project (Ix1000G) outlines a “hub and spoke” model to sequence both laboratory reference strains and natural populations of *Ixodes*. The project is aligned with other ambitious genome initiatives such as the i5K effort that proposes to sequence 5000 arthropod species (Evans et al., 2013) and the Earth BioGenomes project, a moonshot to sequence and catalog all of Earth’s biodiversity (Lewin et al., 2018). The *Ixodes* 1000 Genomes Consortium (IGC) represents an international scientific collaboration formed to launch and guide the initiative. This Letter to the Editor defines the strategic vision of the Ix1000G and serves as a call to the broader scientific community for engagement.

The need for tick genome resources

Tick-borne diseases affect human and animal health globally. Lyme disease (LD) is one of the most important TBDs in North America and Europe. It is estimated that the approximately 30,000 human cases reported to the U.S. Centers for Disease Control and Prevention (CDC) each year represent only about 10% of actual cases (CDC; Hinckley et al., 2014; Nelson et al., 2015). In Europe, roughly 85,000 LD cases are reported annually, although actual case numbers are unknown (European Centre for Disease Prevention and Control (ECDC), 2012). Recent studies are also shedding light on the transmission of human and animal pathogens by Australian ticks and the role of *Ixodes holocyclus*, as a vector (reviewed in Graves and Stenos, 2017; Greay et al., 2018). Options to control hard ticks and the pathogens they transmit are limited. Human vaccines are not available, except against the tick-borne encephalitis virus (Heinz and Stiasny, 2017). Diagnosis and treatment of many TBDs is a challenge because reliable, comprehensive molecular and immunological diagnostics are lacking. Control is also complicated by environmental impacts of acaricides and development of resistant tick populations. Human-tick encounters and pathogen transmission are increasing as geographic ranges of ticks expand and humans exploit new habitats. The need for new strategies to control ticks has never been greater. This need is reflected in recent funding announcements from the U.S. National Institutes of Health (NIH), the Department of Defense (DoD) Congressionally Directed Medical Research Programs (CDMRP), FY17 “Tick-Borne Disease Research Program (TBDRP)”, and the U.K. Biotechnology and Biological Sciences Research Council (BBSRC) and Medical Research Council (MRC) “Networks in Vector Borne Disease Research”. Pressing questions remain in TBD research: how do pathogen complements vary in tick populations at local and regional scales? How do tick microbiomes (including symbionts) and vector genetics affect pathogen transmission and disease outcome? How can scientists best develop comprehensive, region-specific diagnostics, risk matrices and treatment regimes? Fortunately, advances in sequencing and the rapidly-evolving field of data science offer the tools to address many of these questions. The *I. scapularis* genome sequence (Gulia-Nuss et al., 2016) and other molecular and genetic resources for a number of *Ixodes* species such as transcriptome, sialome and mialome datasets (reviewed in Chmelař et al., 2016; Rodriguez Valle et al., 2018), and the establishment of the DNA and tissue repository via the Global Genome Biodiversity Network (Gonzales et al, 2017), are an important beginning. However, incremental advances in science will not be adequate to meet pressing public health needs; ambitious goals must be proposed to effect a rapid paradigm shift in TBD research.

Data-driven solutions for tick and tick-borne disease control

Multiple species of ticks and mites of medical significance have been recognized as high priority targets for sequencing by the NIH (Hill, 2010; Gulia-Nuss et al., 2016). These projects were not advanced due to the limitations of sequencing platforms available at the time. Recently, ultra-long read, single-molecule and genome scaffolding technologies have enabled production of genome assemblies for multiple eukaryotes, including mosquitoes (Jiao et al., 2017; Zimin et al., 2017; Ag1000G, 2017; Mathews et al., 2017). The genomes of 765 field-collected *Anopheles gambiae* and *Anopheles coluzzii* from 15 locations were sequenced by the Ag1000G project (Ag1000G, 2017). The *Ae. aegypti* L5 genome assembly produced by Matthews et al. (2017) comprises chromosome-length scaffolds produced via a hybrid PacBio (Pacific Biosciences) and HiC (Dovetail Genomics) approach. These landmark achievements suggest that modern sequencing technologies will have utility for assembling the large, repeat-rich genomes of ixodid ticks.

The broad objectives of the *Ix*1000G are to: (1) build multiple reference assemblies to support research on *Ixodes* vectors in North America, Europe, Africa and the Asia-Pacific, (2) catalogue natural genetic variation among wild populations of *Ixodes* and their microbial assemblies, (3) describe the genetic history and population structure of *Ixodes* species, and (4) combine genetic variation with biological data to understand tick ecology, epidemiology and disease transmission. The *Ix*1000G will generate fundamental resources for identification and analysis of tick genes and other genomic features, and will support three critical areas needed to advance research on tick and TBD control:

The systems biology of tick-host-pathogen interactions. Every tick bite represents a unique, complex molecular interaction between the tick, the pathogen and the immunological landscape of the vertebrate host. Potentially hundreds of tick salivary factors and microbes are delivered to the host dermis during blood feeding, which lasts for days to over a week. Furthermore, molecular interactions between ticks and commensal or pathogenic microorganisms within tick microbiota may also shape pathogen infection and transmission. Genomics offers opportunities to unravel the systems biology of tick-host-pathogen interactions across multiple *Ixodes* species.

The link between tick genetics, vector competence and disease transmission. To understand this link, there is need to evaluate the correlation between the genetic structure of tick populations and their microbial complement, including pathogens and symbionts. Molecular studies have revealed multiple genetic clades, signatures of north-south population structure (Gulia-Nuss et al., 2016), and evidence for gene flow between populations of *I. scapularis* (Van Zee et al., 2015), but implications for pathogen transmission remain unclear. As sequences of tick-associated microbes are typically recovered with tick genome data, results from the *Ix*1000G will provide fundamental insight into this tripartite relationship.

*Genetic frameworks for *Ixodes* control.* Population genomic studies can directly inform pest management strategies for *Ixodes* and other tick pests. Applications may involve the development of acaricides with species-selective activity and innovative approaches through microbiome manipulation to disrupt pathogen transmission similar to strategies being explored for triatomine bugs and tsetse flies (Hurwitz et al., 2012; Medlock et al., 2013). Population genetics data can also support the development of resistance management and genetic control strategies. For example, genome-wide analyses revealed high levels of heterogeneity between hundreds of *Anopheles* in Africa, suggesting potential for introduction of resistance alleles and evolution of insecticide resistance (Ag1000G, 2017). This indicates the need for genetic control programs tailored to the genetic background of local mosquito populations, and molecular analyses of resistance alleles suggest the same phenomenon will apply to tick populations.

The *Ix*1000G roadmap

The *Ix*1000G strategy calls for whole genome deep sequencing to generate reference assemblies for multiple *Ixodes* species (Priority Area 1) as well as a high-resolution view of genetic variation and microbiome structure in natural populations across the genus (Priority Area 2). Global project partners will source materials and work with teams skilled in data analysis to execute the project in phases, with oversight provided by the IGC. Importantly, the IGC will define a set of standards for genome assembly and annotation aligned with those recognized by sequencing institutes and the Genomic Standards Consortium (Chain et al., 2009; <http://gensc.org/>).

*Priority area 1: High quality reference assemblies for *Ixodes* node species*

The Consortium proposes sequencing and assembly of up to six node species via a combination of PacBio ultra-long-read technology and Hi-C scaffolding of assembled regions. This approach has been used to

generate high quality reference assemblies for *An. gambiae* and *Ae. aegypti* (Ag1000G, 2017; Matthews et al., 2017). The *Ixodes* species listed in Figure 1 have been identified as high priority sequencing targets on the basis of medical significance and potential to anchor additional genome studies across the genus *Ixodes*. This includes population genomic studies as described under Priority Area 2 below. Sequencing will generate a high-quality reference assembly for *I. scapularis*, building on the existing draft genome. Efforts to generate an improved assembly for *I. ricinus* are already underway via the GenIric project (<http://www.angers-nantes.inra.fr/en/All-the-news/GenIric>). Reference genomes will support production of assemblies for other species of *Ixodes*, as well as alignment and analysis of genomic data from field-caught specimens. Species such as *Ixodes affinis*, *Ixodes uriae* and *Ixodes cookei* that are thought to play a minor role in disease transmission, or that do not primarily parasitize humans, are recommended for comparative analyses. The engagement of laboratories with the capability to source ticks and prepare genetic material for sequencing will be critical to these efforts (see Table 1). To obtain sufficient DNA for long read libraries (PacBio, 10X or Nanopore) and reduce heterogeneity in sequenced samples, the IGC proposes to sequence one individual of each node species (or a minimal number). In-bred reference strains should be established via sib-matings for each node species in the case of species listed in Table 1 that have not already been colonized. This would provide source material of similar genetic background for Chicago and Hi-C libraries to support genome scaffolding, and complementary transcriptomic and proteomic studies to validate predicted coding sequences and analyses of gene expression.

The IGC also identified the generation of reference assemblies for tick cell lines in regular use by the scientific community as a priority (Oliver et al., 2015; Miller et al., 2018; Tick Cell Biobank: <https://www.liverpool.ac.uk/infection-and-global-health/research/tick-cell-biobank/>; Table 1). Forward genetic approaches can be used to identify genes associated with vector competence and acaricide resistance. Inbred strains of ticks with quantifiable traits will facilitate genome-wide association studies (GWAS) and mapping of quantitative trait loci (QTL) that must be developed.

Priority area 2: Genomics of natural Ixodes populations and microbiome composition/structure

Under this priority area, the Consortium recommends deep sequencing of several hundred individual ticks belonging to the six node species via an Illumina approach. These should be collected from 10 or more geographic locations across North America, Europe, Africa and Asia. Dual digest Restriction Site Associated DNA sequencing (ddRADseq) may offer a complementary and more financially-viable strategy to generate population genomic data. These studies will also reveal sequences of symbionts and pathogens, and provide insight into the microbiome structure of *Ixodes* at macro-geographic scales. The initial Ix1000G goal proposes a target of 1000 genomes, but the effort is expected to encourage generation of genome data from many thousands of individual ticks in subsequent phases.

Conclusions and Broader Impacts

The Ix1000G will provide a critical framework to support genome studies of other ticks, and mites, of medical importance, including members of the metastriate genera *Amblyomma*, *Rhipicephalus* and *Dermacentor* that include multiple vector species. Moreover, genomes for several mite species of medical importance are already available (Chan et al, 2015; Mofiz et al., 2016; Rider et al., 2017), including a recent genomic analysis of the scrub typhus vector *Leptotrombidium deliense* (Dong et al., 2018). These members of the Acari, with smaller nuclear genomes compared to hard tick species, will support genomic analyses while the more challenging reference assemblies and population genomic studies of wild *Ixodes* populations are in development. Given the worldwide medical and veterinary importance and smaller nuclear genome size of *Ixodes* species as compared to metastriate ticks, coupled with the availability of supporting genomic resources, the genus *Ixodes* represents a logical starting point for generation of high quality reference assemblies and population genomic data. This Letter to the Editor is an official call for project partners and strategic investment to advance the Ix1000G initiative. The broader, long-term vision is an assemblage of data from many thousands of *Ixodes* ticks and a detailed understanding of genetic diversity among *Ixodes* populations worldwide. Data will be used to understand the transmission of TBDs in the context of tick genetics and microbiome communities. The Ix1000G will operate across geographic boundaries to provide the first global genomic resource for tick research and help meet the Grand Challenge of tick and TBD control on a global scale.

Acknowledgements:

The authors thank Ben Matthews, The Rockefeller University, New York, United States and Martin Donnelly, Liverpool School of Tropical Medicine, Liverpool, United Kingdom for helpful discussions.

Declaration of interest: The authors declare no competing interests.

ACCEPTED MANUSCRIPT

References

- 9th Tick and Tick-borne Pathogen Conference/1st Asia Pacific Rickettsia Conference, 2017. <http://www.ttp9-aprc1.com>
- Ag1000g, 2017. Genetic diversity of the African malaria vector *Anopheles gambiae*. *Nature* **552**, 96–100. doi:10.1038/nature24995
- Barrero, R.A., Guerrero, F.D., Black, M., McCooke, J., Chapman, B., Schilkey, F., Pérez de León, A.A., Miller, R.J., Bruns, S., Dobry, J., Mikhaylenko, G., Stormo, K., Bell, C., Tao, Q., Bogden, R., Moolhuijzen, P.M., Hunter, A., Bellgard, M.I., 2017. Gene-enriched draft genome of the cattle tick *Rhipicephalus microplus*: assembly by the hybrid Pacific Bioscience/Illumina approach enabled analysis of the highly repetitive genome. *Int. J. Parasitol.* **47**, 569-583
- Centers for Disease Control and Prevention. Lyme Disease Frequently Asked Questions. How many people get Lyme disease? <https://www.cdc.gov/lyme/faq/index.html> (accessed 05 February 2018)
- Chan, T.F., Ji, K.M., Yim, A.K., Liu, X.Y., Zhou, J.W., Li, R.Q., Yang, K.Y., Li, J., Li, M., Law, P.T., Wu, Y.L., Cai, Z.L., Qin, H., Bao, Y., Leung, R.K., Ng, P.K., Zou, J., Zhong, X.J., Ran, P.X., Zhong, N.S., Liu, Z.G., Tsui, S.K., 2015. The draft genome, transcriptome, and microbiome of *Dermatophagoides farinae* reveal a broad spectrum of dust mite allergens. *J. Allergy Clin. Immunol.* **135**, 539-548. doi:10.1016/j.jaci.2014.09.031
- Chain, P.S., Grafham, D.V., Fulton, R.S., Fitzgerald, M.G., Hostetler, J., Muzny, D., Ali, J., Birren, B., Bruce, D.C., Buhay, C., Cole, J.R., Ding, Y., Dugan, S., Field, D., Garrity, G.M., Gibbs, R., Graves, T., Han, C.S., Harrison, S.H., Highlander, S., Hugenholtz, P., Khouri, H.M., Kodira, C.D., Kolker, E., Kyrpides, N.C., Lang, D., Lapidus, A., Malfatti, S.A., Markowitz, V., Metha, T., Nelson, K.E., Parkhill, J., Pitluck, S., Qin, X., Read, T.D., Schmutz, J., Sozhamannan, S., Sterk, P., Strausberg, R.L., Sutton, G., Thomson, N.R., Tiedje, J.M., Weinstock, G., Wollam, A.; Genomic Standards Consortium Human Microbiome Project Jumpstart Consortium, Detter, J.C. 2009. Genomics. Genome project standards in a new era of sequencing. *Science*. **326**:236-7. doi: 10.1126/science.1180614.
- Chmelař, J., Kotál, J., Karim, S., Kopacek, P., Francischetti, I.M.B., Pedra, J.H.F., Kotsyfakis, M., 2016. Sialomes and mialomes: a systems biology view of tick tissues and tick-host interactions. *Trends Parasitol.* **32**, 242-254. doi:10.1016/j.pt.2015.10.002
- Cramaro, W.J., Revets, D., Hunewald, O.E., Sinner, R., Reye, A.L., Muller, C.P., 2015. Integration of *Ixodes ricinus* genome sequencing with transcriptome and proteome annotation of the naïve midgut. *BMC Genomics* **16**, 871
- Cramaro, W.J., Hunewald, O.E., Bell-Sakyi, L., Muller, C.P., 2017. Genome scaffolding and annotation for the pathogen vector *Ixodes ricinus* by ultra-long single molecule sequencing. *Parasites & Vectors* **10**, 71
- de la Fuente, J., Waterhouse, R.M., Sonenshine, D.E., Roe, R.M., Ribeiro, J.M., Sattelle, D.B. Hill, C.A., 2016. Tick Genome Assembled: New Opportunities for Research on Tick-Host-Pathogen Interactions. *Front. Cell. Infect. Microbiol.* **6**, 103. doi:10.3389/fcimb.2016.00103
- Dong, X., Chaisiri, K., Xia, D., Armstrong, S.D., Fang, Y., Donnelly, M.J., Kadowaki, T., McGarry, J.W., Darby, A.C., Makepeace, B.L., 2018. Genomes of trombidid mites reveal novel predicted allergens and laterally-transferred genes associated with secondary metabolism. *bioRxiv* preprint first posted online Feb.2, 2018. doi: <https://doi.org/10.1101/259044>
- European Centre for Disease Prevention and Control (ECDC), 2012. Second expert consultation on tick-borne diseases with emphasis on Lyme Borreliosis and tick-borne encephalitis. Meeting Report, Stockholm, Sweden, 22-23 November, 2011. <https://ecdc.europa.eu/sites/portal/files/media/en/publications/Publications/Tick-borne-diseases-meeting-report.pdf>
- Evans, J.D., Brown, S.J., Hackett, K.J., Robinson, G., Richards, S., Lawson, D., Elsik, C., Coddington, J., Edwards, O., Emrich, S., Gabaldon, T., Goldsmith, M., Hanes, G., Misof, B., Muñoz-Torres, M., Niehuis, O., Papanicolaou, A., Pfrender, M., Poelchau, M., Purcell-Miramontes, M., Robertson, H.M., Ryder, O., Tagu, D., Torres, T., Zdobnov, E., Zhang, G., Zhou, X., 2013. The i5K Initiative: advancing

arthropod genomics for knowledge, human health, agriculture, and the environment. *J. Hered.* 104, 595-600. doi: 10.1093/jhered/est050

- Heinz, F. X., Stiasny, K., 2012. Flaviviruses and flavivirus vaccines. *Vaccine* 30, 4301-4306. doi:10.1016/j.vaccine.2011.09.114
- González, V.L., Devine, A.M., Trizna, M., Mulcahy, D.G., Barker, K.B., Coddington, J.A., 2017. Open access genomic resources for terrestrial arthropods. *Curr. Opin. Insect Sci.* 25, 91-98. doi: 10.1016/j.cois.2017.12.003.
- Graves, S.R., Stenos, J., 2017. Tick-borne infectious diseases in Australia. *Med. J. Aust.* 206, 320-324. doi: 10.5694/mja17.00090
- Gulia-Nuss, M., Nuss, A.B., Meyer, J.M., Sonenshine, D.E., Roe, R.M., Waterhouse, R.M., Sattelle, D.B., de la Fuente, J., Ribeiro, J.M., Megy, K., Thimmapuram, J., Miller, J.R., Walenz, B.P., Koren, S., Hostetler, J.B., Thiagarajan, M., Joardar, V.S., Hannick, L.I., Bidwell, S., Hammond, M.P., Young, S., Zeng, Q., Abrudan, J.L., Almeida, F.C., Ayllón, N., Bhide, K., Bissinger, B.W., Bonzon-Kulichenko, E., Buckingham, S.D., Caffrey, D.R., Caimano, M.J., Croset, V., Driscoll, T., Gilbert, D., Gillespie, J.J., Giraldo-Calderón, G.I., Grabowski, J.M., Jiang, D., Khalil, S.M., Kim, D., Kocan, K.M., Koči, J., Kuhn, R.J., Kurtti, T.J., Lees, K., Lang, E.G., Kennedy, R.C., Kwon, H., Perera, R., Qi, Y., Radolf, J.D., Sakamoto, J.M., Sánchez-Gracia, A., Severo, M.S., Silverman, N., Šimo, L., Tojo, M., Tornador, C., Van Zee, J.P., Vázquez, J., Vieira, F.G., Villar, M., Wespiser, A.R., Yang, Y., Zhu, J., Arensbürger, P., Pietrantonio, P.V., Barker, S.C., Shao, R., Zdobnov, E.M., Hauser, F., Grimmelikhuijzen, C.J., Park, Y., Rozas, J., Benton, R., Pedra, J.H., Nelson, D.R., Unger, M.F., Tubio, J.M., Tu, Z., Robertson, H.M., Shumway, M., Sutton, G., Wortman, J.R., Lawson, D., Wikel, S.K., Nene, V.M., Fraser, C.M., Collins, F.H., Birren, B., Nelson, K.E., Caler, E., Hill, C.A., 2016. Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. *Nat. Commun.* 7, 10507
- Hill, C.A., 2010. Genome analysis of major tick and mite vectors of human pathogens. Tick and Mite Genomes Consortium. <https://www.vectorbase.org/projects/genome-analysis-major-tick-and-mite-vectors-human-pathogens> (accessed 4 February 2018)
- Hinckley, A.F., Connally, N.P., Meek, J.I., Johnson, B.J., Kemperman, M.M., Feldman, K.A., White, J.L., Mead, P.S., 2014. Lyme disease testing by large commercial laboratories in the United States. *Clin. Infect. Dis.* 59, 676-681.
- Hurwitz, I., Fieck, A., Klein, N., Jose, C., Kang, A., Durvasula, R., 2012. A Paratransgenic Strategy for the Control of Chagas Disease. *Psyche* 2012, Article ID 178930
- Jiao, Y., Peluso, P., Shi, J., Liang, T., Stitzer, M.C., Wang, B., Campbell, M.S., Stein, J.C., Wei, X., Chin, C.S., Guill, K., Regulski, M., Kumari, S., Olson, A., Gent, J., Schneider, K.L., Wolfgruber, T.K., May, M.R., Springer, N.M., Antoniou, E., McCombie, W.R., Presting, G.G., McMullen, M., Ross-Ibarra, J., Dawe, R.K., Hastie, A., Rank, D.R., Ware, D., 2017. Improved maize reference genome with single-molecule technologies. *Nature* 546, 524–527. doi:10.1038/nature22971
- Lewin, H.A., Robinson, G.E., Kress, W.J., Baker, W.J., Coddington, J., Crandall, K.A., Durbin, R., Edwards, S.V., Forest, F., Gilbert, M.T.P., Goldstein, M.M., Grigoriev, I.V., Hackett, K.J., Haussler, D., Jarvis, E.D., Johnson, W.E., Patrinos, A., Richards, S., Castilla-Rubio, J.C., van Sluys, M.A., Soltis, P.S., Xu, X., Yang, H., Zhang, G., 2018. Earth BioGenome Project: Sequencing life for the future of life. *Proc. Natl. Acad. Sci. U S A.* 115, 4325-4333. doi: 10.1073/pnas.1720115115.
- Malaria Genomic Epidemiology Network. Ag1000G. <https://www.malariagen.net/projects/ag1000g>. (Accessed on 02-04-2018)
- Matthews, B.J., Dudchenko, O., Kingan, S., Koren, S., Antoshechkin, I., Crawford, J.E., Glassford, W.J., Herre, M., Redmond, S.N., Rose, N.H., Weedall, G.D., Wu, Y., Batra, S.S., Brito-Sierra, C.A., Buckingham, S.D., Campbell, C.L., Chan, S., Cox, E., Evans, B. R., Fansiri, T., Filipović, I., Fontaine, A., Gloria-Soria, A., Hall, R., Joardar, V. S., Jones, A.K., Kay, R.G.G., Kodali, V.K., Lee, J., Lycett, G.J., Mitchell, S.N., Muehling, J., Murphy, M.R., Omer, A.D., Partridge, F.A., Peluso, P., Aiden, A.P., Ramasamy, V., Rašić, G., Roy, S., Saavedra-Rodriguez, K., Sharan, S., Sharma, A., Smith, M.L., Turner, J., Weakley, A.M., Zhao, Z., Akbari, O.S., Black IV, W.C., Cao, H., Darby, A.C., Hill, C., Johnston, J. S., Murphy, T.D., Raikhel, A.S., Sattelle, D.B., Sharakhov, I.V., White, B.J., Zhao, L., Aiden, E.L., Mann, R.S., Lambrechts, L., Powell, J.R., Sharakhova, M.V., Tu, Z., Robertson, H.M., McBride, C.S., Hastie, A.R.,

- Korlach, J., Neafsey, D.E., Phlipp, A.M., Vosshall, L.B., 2017. Improved *Aedes aegypti* mosquito reference genome assembly enables biological discovery and vector control. doi:<https://doi.org/10.1101/240747>
- Medlock, J., Atkins, K.E., Thomas, D.N., Aksoy, S., Galvani, A.P., 2013. Evaluating Paratransgenesis as a Potential Control Strategy for African Trypanosomiasis. *PLoS Negl. Trop. Dis.* 7, e2374. <https://doi.org/10.1371/journal.pntd.0002374>
- Miller, J.R., Koren, S., Dilley, K.A., Harkins, D.M., Stockwell, T.B., Shabman, R.S., Sutton, G.G. 2018. A draft genome sequence for the *Ixodes scapularis* cell line, ISE6.. *F1000 Research* 7, 297
- Mofiz, E., Holt, D.C., Seemann, T., Currie, B.J., Fischer, K., Papenfuss, A.T., 2016. Genomic resources and draft assemblies of the human and porcine varieties of scabies mites, *Sarcoptes scabiei* var. *hominis* and var. *suus*. *Gigascience* 5, 23. doi:10.1186/s13742-016-0129-2
- Neafsey, D.E., Waterhouse, R.M., Abai, M.R., Aganezov, S.S., Alekseyev, M.A., Allen, J.E., Amon, J., Arcà, B., Arensburger, P., Artemov, G., Assour, L.A., Basseri, H., Berlin, A., Birren, B.W., Blandin, S.A., Brockman, A.I., Burkot, T.R., Burt, A., Chan, C.S., Chauve, C., Chiu, J.C., Christensen, M., Costantini, C., Davidson, V.L., Deligianni, E., Dottorini, T., Dritsou, V., Gabriel, S.B., Guelbeogo, W.M., Hall, A.B., Han, M.V., Hlaing, T., Hughes, D.S., Jenkins, A.M., Jiang, X., Jungreis, I., Kakani, E.G., Kamali, M., Kempainen, P., Kennedy, R.C., Kirmizoglou, I.K., Koekemoer, L.L., Laban, N., Langridge, N., Lawniczak, M.K., Lirakis, M., Lobo, N.F., Lowy, E., MacCallum, R.M., Mao, C., Maslen, G., Mbogo, C., McCarthy, J., Michel, K., Mitchell, S.N., Moore, W., Murphy, K.A., Naumenko, A.N., Nolan, T., Novoa, E.M., O'Loughlin, S., Oranganje, C., Oshaghi, M.A., Pakpour, N., Papathanos, P.A., Peery, A.N., Povelones, M., Prakash, A., Price, D.P., Rajaraman, A., Reimer, L.J., Rinker, D.C., Rokas, A., Russell, T.L., Sagnon, N., Sharakhova, M.V., Shea, T., Simão, F.A., Simard, F., Slotman, M.A., Somboon, P., Stegny, V., Struchiner, C.J., Thomas, G.W., Tojo, M., Topalis, P., Tubio, J.M., Unger, M.F., Vontas, J., Walton, C., Wilding, C.S., Willis, J.H., Wu, Y.C., Yan, G., Zdobnov, E.M., Zhou, X., Catteruccia, F., Christophides, G.K., Collins, F.H., Cornman, R.S., Crisanti, A., Donnelly, M.J., Emrich, S.J., Fontaine, M.C., Gelbart, W., Hahn, M.W., Hansen, I.A., Howell, P.I., Kafatos, F.C., Kellis, M., Lawson, D., Louis, C., Luckhart, S., Muskavitch, M.A., Ribeiro, J.M., Riehle, M.A., Sharakhov IV, Tu, Z., Zwiebel, L.J., Besansky, N.J., 2015. Mosquito genomics. Highly evolvable malaria vectors: the genomes of 16 *Anopheles* mosquitoes. *Science* 347, 1258522. doi:10.1126/science.1258522.
- Nelson, C.A., Saha, S., Kugeler, K.J., Delorey, M.J., Shankar, M.B., Hinckley, A.F., Mead, P.S., 2015. Incidence of clinician-diagnosed Lyme disease, United States, 2005-2010. *Emerg. Infect. Dis.* 21, 1625-1631.
- Chain, P.S., Grafham, D.V., Fulton, R.S., Fitzgerald, M.G., Hostetler, J., Muzny, D., Ali, J., Birren, B., Bruce, D.C., Buhay, C., Cole, J.R., Ding, Y., Dugan, S., Field, D., Garrity, G.M., Gibbs, R., Graves, T., Han, C.S., Harrison, S.H., Highlander, S., Hugenholtz, P., Khouri, H.M., Kodira, C.D., Kolker, E., Kyrpides, N.C., Lang, D., Lapidus, A., Malfatti, S.A., Markowitz, V., Metha, T., Nelson, K.E., Parkhill, J., Pitluck, S., Qin, X., Read, T.D., Schmutz, J., Sozhamannan, S., Sterk, P., Strausberg, R.L., Sutton, G., Thomson, N.R., Tiedje, J.M., Weinstock, G., Wollam, A.; Genomic Standards Consortium Human Microbiome Project Jumpstart Consortium, Detter, J.C. 2009. Genomics. Genome project standards in a new era of sequencing. *Science*. 326:236-7. doi: 10.1126/science.1180614.
- Oliver, J.D., Chávez, A.S., Felsheim, R.F., Kurtti, T.J., Munderloh, U.G., 2015. An *Ixodes scapularis* cell line with a predominantly neuron-like phenotype. *Exp. Appl. Acarol.* 2015, 66,427-442. doi:10.1007/s10493-015-9908-1.
- Rider, S.D. Jr, Morgan, M.S., Arlian, L.G., 2017. Allergen homologs in the *Euroglyphus maynei* draft genome. *PLoS One* 12, e0183535. doi: 10.1371/journal.pone.0183535.
- Rodriguez-Valle, M., Moolhuijzen, P., Barrero, R.A., Ong, C.T., Busch, G., Karbanowicz, T., Booth, M., Clark, R., Koehbach, J., Ijaz, H., Broady, K., Agnew, K., Knowles, A.G., Bellgard, M.I., Tabor, A.E., 2018. Transcriptome and toxin family analysis of the paralysis tick, *Ixodes holocyclus*. *Int. J. Parasitol.* 48, 71-82. doi: 0.1016/j.ijpara.2017.07.007.
- The Tick Cell Biobank: <https://www.liverpool.ac.uk/infection-and-global-health/research/tick-cell-biobank/>
- Van Zee, J., Piesman, J.F., Hojgaard, A., Black, W.C. IV., 2015. Nuclear markers reveal predominantly North to South gene flow in *Ixodes scapularis*, the tick vector of the Lyme Disease spirochete. *PLoS One.* 10, e0139630. doi:10.1371/journal.pone.0139630

Zimin, A.V., Puiu, D., Hall, R., Kingan, S., Clavijo, B.J., Salzberg, S.L., 2017. The first near-complete assembly of the hexaploid bread wheat genome, *Triticum aestivum*. *GigaScience* 6, 1–7.
<https://doi.org/10.1093/gigascience/gix097>

ACCEPTED MANUSCRIPT

Figure Legends:

Figure 1. Schematic showing six “node” species of *Ixodes* recommended as targets for deep genome sequencing. The species *Ixodes scapularis*, *Ixodes pacificus*, *Ixodes ricinus*, *Ixodes persulcatus*, *Ixodes rubicundus* and *Ixodes holocyclus* were selected based on status as disease vectors and their potential to anchor population genomic studies across the genus *Ixodes*. Segments of the pie chart show diseases transmitted or caused by ticks: BAC, Bacterial Artificial Chromosome library; LD, Lyme disease; HGA, Human granulocytic anaplasmosis; TBE, Tick-borne encephalitis; SFGR, Spotted fever group rickettsiosis.

ACCEPTED MANUSCRIPT

Table 1. *Ixodes* species proposed for genome sequencing (priority area 1) and availability of colonies and cell lines for sequencing.

Tick Species		Pathogen Transmitted	Disease	Genome Project Resources
Scientific Name	Common Name			
North America				
<i>Ixodes scapularis</i>	Blacklegged tick	<i>Borrelia burgdorferi</i> <i>Babesia microti</i> <i>Anaplasma phagocytophilum</i> Powassan virus <i>Borrelia miyamotoi</i>	LD Babesiosis HGA Powassan encephalitis <i>Borrelia miyamotoi</i> disease	Colonies; UTMB; BEI Resources ^a ; field specimens
<i>Ixodes pacificus</i>	Western blacklegged tick	<i>Borrelia burgdorferi</i> <i>Anaplasma phagocytophilum</i> <i>Borrelia miyamotoi</i>	LD HGA <i>Borrelia miyamotoi</i> disease	Colonies; BEI Resources
Europe				
<i>Ixodes ricinus</i>	Castor bean tick	<i>Borrelia</i> spp <i>Borrelia miyamotoi</i> TBE virus <i>Candidatus</i> Neoehrlichia mikurensis <i>Anaplasma phagocytophilum</i> <i>Francisella tularensis</i> Louping ill virus <i>Babesia divergens</i> , <i>Babesia microti</i> , other <i>Babesia</i> spp. <i>Rickettsia helvetica</i>	LD <i>Borrelia miyamotoi</i> disease TBE Neoehrlichiosis Tick-borne fever of ruminants, HGA Tularemia Louping ill of sheep and grouse Babesiosis Human SFGR	Colonies; UTMB
Asia-Pacific				
<i>Ixodes holocyclus</i>	Australian paralysis tick	<i>Rickettsia australis</i> <i>Coxiella burnetti</i>	Tick paralysis	Field specimens;
<i>Ixodes persulcatus</i>	Taiga tick	<i>Borrelia</i> spp. <i>Borrelia miyamotoi</i> TBE virus <i>Anaplasma phagocytophilum</i> , <i>Ehrlichia</i> spp.	LD <i>Borrelia miyamotoi</i> disease TBE HGA Ehrlichiosis	-
Africa				
<i>Ixodes rubicundus</i>	South African Karoo paralysis tick	No known etiological agents	Tick paralysis	Laboratory colonies; Univ. South Africa
Minor species				
<i>Ixodes affinis</i>		<i>Borrelia burgdorferi</i>	LD	-
<i>Ixodes uriae</i>	Seabird tick	<i>Borrelia garinii</i>	LD	-

<i>Ixodes cookei</i>	Powassan virus	Powassan encephalitis	-
----------------------	----------------	-----------------------	---

***Ixodes* cell lines**

Species	Name of Cell Line	Origin of Cell Line
<i>Ixodes scapularis</i>	ISE6, ISE18, IDE8	United States
<i>Ixodes ricinus</i>	IRE/CTVM19, IRE/CTVM20	United Kingdom
<i>Ixodes ricinus</i>	IRE11	Germany

HGA, Human granulocytic anaplasmosis; LD, Lyme disease; SFGR, Spotted fever group rickettsiosis; TBE, Tick-borne encephalitis; UTMB, University of Texas Medical Branch, Galveston, Texas, USA

^aNIH, NIAID BEI Resources Repository: <https://www.niaid.nih.gov/research/bei-resources-repository>

ACCEPTED MANUSCRIPT

Fig-1

