



HAL
open science

Phlebotomus (Legeromyia) multihamatus subg. nov., sp. nov. from Gabon (Diptera: Psychodidae)

Nil Rahola, Jérôme Dépaquit, Boris Kevin Makanga, Christophe Paupy

► To cite this version:

Nil Rahola, Jérôme Dépaquit, Boris Kevin Makanga, Christophe Paupy. *Phlebotomus (Legeromyia) multihamatus subg. nov., sp. nov. from Gabon (Diptera: Psychodidae)*. *Memórias do Instituto Oswaldo Cruz*, 2013, 108 (7), pp.845-849. 10.1590/0074-0276130172 . hal-02065049

HAL Id: hal-02065049

<https://hal.umontpellier.fr/hal-02065049v1>

Submitted on 21 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Phlebotomus (Legeromyia) multihamatus subg. nov., sp. nov. from Gabon (Diptera: Psychodidae)

Nil Rahola^{1,2}, Jérôme Depaquit³, Boris Kevin Makanga^{2,4}, Christophe Paupy^{1,2/+}

¹Laboratoire Maladies Infectieuses et Vecteurs: Écologie, Génétique, Évolution et Contrôle, Unité Mixte de Recherche 224-5290, Centre National de la Recherche Scientifique, Institut de Recherche pour le Développement, Universités Montpellier 182, Montpellier, Hérault, Languedoc Rousillon, France ²Centre International de Recherches Médicales de Franceville, Franceville, Mpassa, Haut-Ogooué, Gabon ³Transmission Vectorielle et Épidémiologie de Maladies Parasitaires EA 4688-USC, Agence Nationale de la Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail, Université de Reims Champagne-Ardenne, Reims, Marne, Champagne-Ardenne, France ⁴Institut de Recherche en Écologie Tropicale, Libreville, Libreville, Estuaire, Gabon

During a research project aimed at the study of the Culicinae fauna of Gabon and carried out in the National Park of La Lopé, we captured an unknown sandfly male specimen (genus *Phlebotomus*) by CDC miniature light trap belonging to a new species for Science. Furthermore, the originality of his genitalia does not allow us to include this species in one of the existing subgenus, thus in this paper we propose the creation of a new subgenus, as *Phlebotomus (Legeromyia) multihamatus* sp. nov., subg. nov. described from the National Park of La Lopé, through one male captured with CDC miniature light trap. A new species and a new subgenus of sandfly is characterised by a short style with three spines, a paramere wearing a basal hook as well as a basal pouch and the absence of basal lobe on the coxite. The originality of the genitalia of the male gives way to discussion about potential primary homologies between *P. multihamatus* sp. nov. and *Phlebotomus (Abonnencius) fortunatarum*, *Phlebotomus (Anaphlebotomus) stantoni* and *Phlebotomus (Euphlebotomus) argentipes*, which should be verified for future studies. The discovery of this new species in Gabon must encourage the study of sandflies in this country.

Key words: sandfly - *Phlebotomus* - new species - Gabon

Despite the former mention of a native case of visceral leishmaniasis (Tournier 1920), Gabon does not actually constitute an endemic country for leishmaniasis (Alvar et al. 2012). As a result, the sandfly fauna in this country remains poorly documented and prospected. Unlike in the neighbouring countries where no less than 23 species in Cameroon, 30 in the Central African Republic, 28 in the Democratic Republic of the Congo and 42 in the Republic of the Congo were previously found (Niang et al. 2004); an exhaustive review of literature reports only four species in Gabon (Galliard & Nitzulescu 1931, Abonnenc 1972): *Sergentomyia (Parrotomyia) africana africana* (Newstead, 1912), *Sergentomyia (Sergentomyia) antennata* (Newstead, 1912), *Sergentomyia (Sergentomyia) schwetzi* (Adler, Theodor & Parrot, 1929) and *Spelaeophlebotomus gigas* (Parrot & Schwetz, 1937).

MATERIALS AND METHODS

The sandfly described here was collected in a secondary equatorial rainforest inside the National Park of La Lopé, Gabon at approximately 10 km away of the village of Mikongo (0°18'41.34"S 11°42'23.58"E) at an altitude of 303 m above mean sea level (Fig. 1). The insect collections and the access to the Park were authorised by the

Ministry of Scientific Research and Technological Development of Gabon (AR0006/12/MENERSI/CENAR-EST/CG/CST/CSAR) and the National Agency of Natural Parks. The capture was performed in 2012 during the night of the 22 October, using a CDC miniature light trap placed 2 m above the ground level in an open forest undergrowth allowing a good visual attraction.

The sandfly was preserved in 100% ethanol then mounted whole in Euparal after successive different baths: 2 h in potassium hydroxide 10%, 2 h in distilled water, 10 h in a Marc-André solution (Abonnenc 1972), 10 h in distilled water, 20 min in ethanol 70%, 20 min in ethanol 90%, 20 min in ethanol 100% and 10 h in a beech wood solution.

Specimen was observed using a BX 50 microscope (Olympus, Japan). Measurements were collected using the Perfect Image software (ARIES Company, Chatillon, France) and a video camera connected to the microscope.

RESULTS

Phlebotomus multihamatus Rahola, Depaquit
and Paupy, sp. nov.
(Figs 2-5)

Male - Holotype - Sandfly of small size (~1.5 mm) and rather light colour.

Head (Figs 2A-C, 3A): orientated laterally. Interocular suture, hardly visible with this orientation, seems to be complete. Cibarium with 13 discreet teeth arranged on a concave arc directed backwards and with rare denticles not well defined. Pigmented plate absent. Pharynx fine and short slightly shrunk backwards. Pharyngeal armature not well developed constituted with some wrinkles. Palpal

doi: 10.1590/0074-0276130172

Financial support: CIRMF, IRD (PPR FT-AC)

+ Corresponding author: christophe.paupy@ird.fr

Received 27 March 2013

Accepted 10 June 2013

formula: 1-2-4-3-5, third palpal article with five club-like Newstead spines. Ascoid formula: 2/III-IX, articles X-XVI are missing on both antennae. Ascoids long going over the next articulation. Sensillae present on article III-V. A III = 155 μm , A IV = 104 μm , A V = 104 μm , A III < A IV + A V; epipharynx = 143 μm . A III/E = 1.08.

Thorax (Figs 2D, 3B): presence of three antero-inferior mesanepisternal setae. Wing length = 1,205 μm , width = 412 μm , length/width = 2.9°, α = 241 μm° , β = 140 μm , δ = 35, \square = 270 μm , wing width/ \square = 1.5° ad π = 59 μm .

Genitalia (Figs 3C, 4A-C): coxite length: 148 μm bearing a small number of scattered setae in its internal face. Style extremely short (60 μm), globular, pear-shaped, with three spines: one terminal and two median. Paramere is very complex measuring 134 μm in its whole. The superior lobe is measuring 84 μm [measurement *sensu* (Leng & Lewis 1987)]. It bears a very high number of setae. The inferior lobe is subdivided in two parts: (i) a short pointed hook without any setae curved towards the top and pointing backwards and (ii) a small rounded bump bearing three to four setae. At the base of the paramere, near the implantation of the penis, a remarkable process fits in a huge pigmented hook measuring 78 μm long, directed backwards and supplied with a little dewclaw in subapical position. The set composed by this hook and the penis is covered on its external face by a membrane constituting a basal pouch. Surstyle: 168 μm . Penis: 51 μm , wide at its base and contracting regularly to its apex. Apex translucent in its inferior part. Genital pump heavily sclerotised and well developed measuring 114 μm .

Female - Unknown.

Type locality - National Park of La Lopé, Gabon.

Type data and depository - Holotype: male: Gabon, National Park of La Lopé 0°18'41.34"S 11°42'23.58"E,




Fig. 1: location of the collection site in Gabon.

22 October, 2012, CDC miniature light trap, N Rahola, J Depaquit and C Paupy leg. Deposited in the Institut de Recherche pour le Développement, Montpellier, France.

Etymology - The subgenus *Legeromyia* is dedicated to our colleague Nicole Léger which has greatly contributed to the systematics of sandflies. The species name *multihamatus* refers to the paramere's hooks that characterise this species.

DISCUSSION

To date, seven genera of sandflies are recorded in the Old World according Rispaill and Léger (1998b): *Phlebotomus* Rondani, 1843, *Sergentomyia* França & Parrot, 1921, *Spelaephlebotomus* Theodor 1948, *Spelaomyia* Theodor, 1948, *Idiophlebotomus* Quate & Fairchild, 1961, *Australophlebotomus* Theodor, 1948 and *Chinius* Leng, 1987. Moreover, these authors considered the taxa *Parvidens* Theodor & Mesghali, 1964 as incertae sedis. The transmission of *Leishmania* parasites in the Old World is mainly ensured by species belonging to the genus *Phlebotomus* (Killick-Kendrick 1990). However, it was recently found that species from other genera could also play a role in leishmaniasis transmission in




Fig. 2: drawing of palp (A), third segment of palp showing five club-like Newstead spines (B), antennal segment III and IV + V with ascoids (C) and wing (D) of *Phlebotomus multihamatus* sp. nov.

the Old World. It is the case of the genus *Sergentomyia* (*Sergentomyia darlingi*) involved in the transmission of *Leishmania major* in Mali (Berdjane-Brouk et al. 2012). The genus *Phlebotomus* is currently subdivided in nine subgenera according Rispaill and Léger (1998b): (*Phlebotomus*) Rondani, 1843, (*Paraphlebotomus*) Theodor, 1948, (*Synphlebotomus*) Theodor, 1948, (*Larroussius*) Theodor, 1948, (*Adlerius*) Nitzulescu, 1931, (*Transphlebotomus*) Artemiev, 1984, (*Euphlebotomus*) Theodor, 1948, (*Anaphlebotomus*) Theodor, 1948, and (*Kasaulius*) Lewis, 1982. However, Rispaill and Léger (1998a) thought the subgenus (*Anaphlebotomus*) is artificial, because it includes two lineages. New species from Madagascar belonging to this group were recently described (Depaquit et al. 2002, 2004, Randrianambinintsoa et al. 2013), suggesting this subgenus needs to be revised. Until it, we think that the subgenus (*Abonnencius*) Ontiveros et al. (1982), considered as a non-valid taxa (Lane & Alexander 1988), has to be retained.

The main species involved in the *Leishmania* transmission are: *Phlebotomus* (*Phlebotomus*) *papatasi* (Scopoli, 1786), *Phlebotomus* (*Paraphlebotomus*) *sergenti* Parrot, 1917, *Phlebotomus* (*Euphlebotomus*) *argentipes* Annandale & Brunetti, 1908, as well as many species of the subgenus *Larroussius* for the transmission of *L.*

major, *Leishmania tropica*, *Leishmania donovani* and *Leishmania infantum*, respectively. The creation of all these subgenera was gradually made initially without phylogenetic arguments, but has become customary among specialists. *P. multihamatus* sp. nov. possesses antero-inferior mesanepisternal setae. Therefore its inclusion in the genus *Phlebotomus* is obvious. In addition, its geographical localisation is also in adequacy with this systematic position. However, in sight of the morphology of its genitalia, this species cannot be included in any of the existing subgenus of *Phlebotomus*. The presence of three spines on the style is a criteria shared with genus *Australophlebotomus* confined to the Australian region. However, in *P. multihamatus* sp. nov., the paramere is complex and the genital filaments are long whereas the paramere is simple and the genital filaments are short and even nonexistent in *Australophlebotomus* (Lewis & Dyce 1982, Léger & Pesson 1993).

We cannot include it either in the subgenera with simple paramere such as (*Larroussius*), (*Adlerius*), (*Transphlebotomus*), (*Paraphlebotomus*), (*Synphlebotomus*) and (*Kasaulius*). The absence of a basal lobe on the coxite prevents its inclusion in the subgenus (*Phlebotomus*). Possibly, it could be related to the subgenus *Euphlebotomus*, *Anaphlebotomus* and *Abonnencius*, the




Fig. 3: drawing of pharynx and cibarium (A), mesothoracic part showing mesanepisternal setae (B) and full view of male genitalia (C) of *Phlebotomus multihamatus* sp. nov.




Fig. 4: drawing of style (A), paramere with penis and genital filaments (B) and paramere showing pouch surrounding penis (C) of *Phlebotomus multihamatus* sp. nov.

validity of the latter being obvious to us despite the curious position without real scientific foundation adopted by Lane and Alexander (1988).

The systematics of these groups deserves a revision based on phylogenetic arguments. Meanwhile, we note down the existence of points of convergence in the structure of the male genitalia of these three subgenera with that of *P. multihamatus* sp. nov.: genital pump well developed and strongly pigmented, complex parameres and short styles. We also note a point of convergence between the hook implanted at the base of the paramere in *P. multihamatus* sp. nov. and in *Phlebotomus fortunatarum*.

Moreover, we also observe a certain degree of homology between these structures and the accessory sticks observed in some (*Anaphlebotomus*) and (*Euphlebotomus*) like their respective type-species: *Phlebotomus stantoni* and *P. argentipes* (Fig. 5). The hypothetical establishment of primary homologies between these two structures will certainly act upon an attempt of cladistics morphological analysis in these groups. On the other hand, the membrane forming a pouch on the paramere seems very original to us among sandflies.

On the basis of this argumentation, we propose the creation of a new subgenus: (*Legeromyia*) subg. nov. defined as follows: style with three spines, paramere wearing a basal hook as well as a basal pouch, absence of basal lobe on the coxite. Type-species of the subgenus (*Legeromyia*) subg. nov.: *P. multihamatus* sp. nov.

The discovery of this new species of sandfly must encourage prospecting in this direction in Gabon. Even if only one case of leishmaniasis has been reported in this country (Tournier 1920), the role of sandflies in the transmission of such parasites or other infectious agents, such as virus or bacteria (Depaquit et al. 2010) to wild-life and domestic animals or to human and their trophic preferences, should be evaluated.




Fig. 5: parameres including organs and accessory sticks from penis of *Phlebotomus multihamatus* sp. nov. (A), *Phlebotomus argentipes* (B), *Phlebotomus stantoni* (C) and *Phlebotomus fortunatarum* (D).

ACKNOWLEDGEMENTS

To Donald Ndongo and Patrice Mikelet, for their technical help during the field work, to the Station d'Etude des Gorilles et Chimpanzés, for the logistic support, to the Agence Nationale des Parcs Naturels and the Centre National de la Recherche Scientifique et Technologique of Gabon, that authorised this study and facilitated the access to the National Park of La Lopé.

REFERENCES

- Abonnenc E 1972. *Les phlébotomes de la Région Ethiopienne (Diptera, Psychodidae)*, Mémoires ORSTOM 55, Paris, 289 pp.
- Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, den Boer M, WHO Leishmaniasis Control Team 2012. Leishmaniasis worldwide and global estimates of its incidence. *PLoS ONE* 7: e35671.
- Berdjane-Brouk Z, Koné AK, Djimdé AA, Charrel RN, Ravel C, De-launay P, del Giudice P, Diarra AZ, Doumbo S, Goita S, Thera MA, Depaquit J, Marty P, Doumbo OK, Izri A 2012. First detection of *Leishmania major* DNA in *Sergentomyia (Spelaemyia) darlingi* from cutaneous leishmaniasis foci in Mali. *PLoS ONE* 7: e28266.
- Depaquit J, Grandadam M, Fouque F, Andry PE, Peyrefitte C 2010. Arthropod-borne viruses transmitted by phlebotomine sand flies in Europe: a review. *Euro Surveill* 15: 19507.
- Depaquit J, Léger N, Robert V 2002. Première mention de *Phlebotomus* à Madagascar (Diptera: Psychodidae). Description de *Phlebotomus (Anaphlebotomus) fertei* n. sp. et de *Phlebotomus (Anaphlebotomus) huberti* n. sp. *Parasite* 9: 325-331.
- Depaquit J, Léger N, Robert V 2004. Les phlébotomes de Madagascar. III - Description de *Phlebotomus (Anaphlebotomus) fontenillei* n. sp. *Parasite* 11: 261-265.
- Galliard H, Nitzulescu V 1931. Contribution à l'étude des phlébotomes du Gabon. *Phlebotomus sanneri* n. sp. *Ann Parasitol Hum Comp* 9: 233-246.
- ICZN - International Commission on Zoological Nomenclature 2000. *International Code of Zoological Nomenclature*, 4th ed., The International Trust for Zoological Nomenclature 1999/The Natural History Museum, London, 124 pp.
- Killick-Kendrick R 1990. Phlebotomine vectors of the leishmaniasis: a review. *Med Vet Entomol* 4: 1-24.
- Lane RP, Alexander B 1988. Sand flies (Diptera: Phlebotominae) of the Canary Islands. *J Nat Hist* 22: 313-319.
- Léger N, Pesson B 1993. Première mention de phlébotomes en Nouvelle-Calédonie et description de deux espèces nouvelles (Diptera, Psychodidae, Phlebotominae). *Mémoires du Muséum National d'Histoire Naturelle* 157: 123-129.
- Leng YJ, Lewis DJ 1987. The subgenus *Euphlebotomus* (Diptera: Psychodidae) in China, with description of a new species, *Phlebotomus yunshengensis*. *Ann Trop Med Parasitol* 81: 305-309.
- Lewis D, Dyce A 1982. The subgenus *Australophlebotomus* Theodor of *Phlebotomus* Rondani and Berté (Diptera: Psychodidae). *Aust J Entomol Soc* 21: 37-54.
- Niang AA, Geoffroy B, Davidson I, Depaquit J, Hervy J-P, Boussès P, Leger N, Trouillet J, Killick-Kendrick R, Killick-Kendrick M, Rioux J-A, Brunhes J 2004. *Sand flies of the Afrotropical Region: identification and training program*, CD Rom, Collection Didactiques, IRD Editions, Paris.
- Ontiveros JMU, Marquez FM, Benitez DCG, Roman RL, Barrios CC 1982. Flebotomos de las Islas Canarias (España). *Rev Iber Parasitol* 42: 197-206.

- Randrianambinintsoa FJ, Léger N, Robert V, Depaquit J 2013. Phlebotomine sand flies from Madagascar (Diptera: Psychodidae). VII. An identification key for *Phlebotomus* with the description of *Phlebotomus (Anaphlebotomus) vaomalalae* n. sp. *Parasite* 20: 6.
- Rispail P, Léger N 1998a. Numerical taxonomy of Old World Phlebotominae (Diptera: Psychodidae). 1. Considerations of morphological characters in the genus *Phlebotomus* Rondani & Berté 1840. *Mem Inst Oswaldo Cruz* 93: 773-785.
- Rispail P, Léger N 1998b. Numerical taxonomy of Old World Phlebotominae (Diptera: Psychodidae). 2. Restatement of classification upon subgeneric morphological characters. *Mem Inst Oswaldo Cruz* 93: 787-793.
- Theodor O 1948. Classification of the Old World species of the subfamily Phlebotominae (Diptera: Psychodidae). *Bull Entomol Res* 39: 85-118.
- Tournier E 1920. Note sur un cas de Kala-azar infantile observé au Gabon. *Bull Soc Pathol Exot* 13: 175-176.