Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950–2005

Moussa Sidibe, Bastien Dieppois, Yu Fisjak, Jean-Emmanuel Paturel, Ernest Amoussou, Babatunde Anifowose, Damian Lawler

► To cite this version:

Moussa Sidibe, Bastien Dieppois, Yu Fisjak, Jean-Emmanuel Paturel, Ernest Amoussou, et al.. Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950–2005. *Journal of Hydrology*, 2018, 561, pp.478-493. 10.1016/j.jhydrol.2018.04.024 . hal-02106650

HAL Id: hal-02106650

<https://hal.umontpellier.fr/hal-02106650>

Submitted on 23 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950 – 2005

Sidibe, M, Dieppois, B, Mahé, G, Paturel, J-E, Amoussou, E, Anifowose, B & Lawler, D

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Sidibe, M, Dieppois, B, Mahé, G, Paturel, J-E, Amoussou, E, Anifowose, B & Lawler, D 2018, 'Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950 – 2005', *Journal of Hydrology*, vol. 561, pp. 478-493.

<https://dx.doi.org/10.1016/j.jhydrol.2018.04.024>

DOI 10.1016/j.jhydrol.2018.04.024

ISSN 0022-1694

ESSN 1879-2707

Publisher: Elsevier

NOTICE: this is the author's version of a work that was accepted for publication in *Journal of Hydrology*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Journal of Hydrology*, (561), (2018) DOI: [10.1016/j.jhydrol.2018.04.024](https://doi.org/10.1016/j.jhydrol.2018.04.024)

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version

may remain and you are advised to consult the published version if you wish to cite from it.

Accepted Manuscript

Research papers

Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950 – 2005

Moussa Sidibe, Bastien Dieppois, Gil Mahé, Jean-Emmanuel Paturel, Ernest Amoussou, Babatunde Anifowose, Damian Lawler

PII: S0022-1694(18)30277-4
DOI: <https://doi.org/10.1016/j.jhydrol.2018.04.024>
Reference: HYDROL 22727

To appear in: *Journal of Hydrology*

Received Date: 26 October 2017
Revised Date: 30 March 2018
Accepted Date: 7 April 2018

Please cite this article as: Sidibe, M., Dieppois, B., Mahé, G., Paturel, J-E., Amoussou, E., Anifowose, B., Lawler, D., Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950 – 2005, *Journal of Hydrology* (2018), doi: <https://doi.org/10.1016/j.jhydrol.2018.04.024>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Trend and variability in a new, reconstructed streamflow dataset for West and Central Africa, and climatic interactions, 1950 – 2005

Moussa Sidibe¹, Bastien Dieppo^{1,2,3}, Gil Mahé⁴, Jean-Emmanuel Paturel⁴,
Ernest Amoussou⁵, Babatunde Anifowose⁶, Damian Lawler¹

¹Centre for Agroecology, Water and Resilience (CAWR), Coventry University, Coventry, UK

²Department of Oceanography, University of Cape Town, Cape Town, RSA

³School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

⁴HydroSciences (HSM), IRD, Montpellier, France

⁵Département de Géographie et Aménagement du Territoire (DGAT), Université de Parakou, Parakou, Benin

⁶School of Energy, Construction & Environment, Coventry University, UK

Abstract

Over recent decades, regions of West and Central Africa have experienced different and significant changes in climatic patterns, which have significantly impacted hydrological regimes. Such impacts, however, are not fully understood at the regional scale, largely because of scarce hydroclimatic data. Therefore, the aim of this study is to (a) assemble a new, robust, reconstructed streamflow dataset of 152 gauging stations; (b) quantify changes in streamflow over 1950 – 2005 period, using these newly reconstructed datasets; (c) significantly reveal trends and variability in streamflow over West and Central Africa based on new reconstructions; and (d) assess the robustness of this dataset by comparing the results with those identified in key climatic drivers (*e.g.* precipitation and temperature) over the region. Gap filling methods applied to monthly time series (1950-2005) yielded robust results (median Kling-Gupta Efficiency >0.75). The study underlines a good agreement between precipitation and streamflow trends and reveals contrasts between western Africa (negative trends) and Central Africa (positive trends) in the 1950s and 1960s. Homogenous dry conditions of the 1970s and 1980s, characterized by reduced significant negative trends resulting from quasi-decadal modulations of the trend, are replaced by wetter conditions in the recent period (1993-2005). The effect of this rainfall recovery (which extends to West and Central Africa) on increased river flows are further amplified by land use change in some Sahelian basins. This is partially offset, however, by higher potential evapotranspiration rates over parts of Niger and Nigeria. Crucially, the new reconstructed

streamflow datasets presented here will be available for both the scientific community and water resource managers.

Keywords: *West and Central Africa, streamflow trend and variability, hydroclimate variability, multi-temporal trend identification, gap filling methods.*

1. INTRODUCTION

Rainfall in Africa during the 20th century was characterized by decreasing annual trends over the continent except regions in Cameroon, Sierra Leone and southern equatorial Africa (Hulme *et al.*, 2001). Since 1950, most of the extreme climatic conditions occurred in the Sahel region, which has experienced several drought events from the end of the 1960s to the 1990s (Dai *et al.*, 2004; Lebel, 2003; Nicholson, 2013). For the 1968–1997 period, rainfall in August in the West African Sahel showed a decrease of up to 37% compared to the 1931–1960 period (Nicholson *et al.*, 2000). Rainfall patterns in the post-1990 period are, however, less well documented, given data scarcity: this has led to controversial findings regarding the end of Sahel drought. For example, Ozer *et al.* (2003) claimed that the Sahel drought ended in the 1990s, whereas L'Hôte *et al.* (2002) suggested that the drought continued beyond the 1990s. These contradictions partly reflect the significant changes in the spatial distribution of precipitation, which make findings highly dependent on the specific region, and the years and even months considered. This underlines the need for studies covering larger spatial scales. However, there is agreement on an increase in annual rainfall over the West African Sahel since the 1990s (*e.g.* Ali and Lebel, 2009; Jury, 2013; Lebel and Ali, 2009; Mahé and Paturel, 2009). See also Maidment *et al.* (2015), who described rainfall trends over Africa during the period 1983–2010, using different observational datasets and simulations from the current state-of-the-art global climate models.

Interestingly, while rainfall variability has been investigated at the continental scale in Africa, its effects on runoff regimes have mostly been investigated at catchment scales, using different statistical approaches and hydrological models (*e.g.* Ibrahim *et al.*, 2015). This is mainly due to restricted data, and several factors

(*e.g.* Gyau-Boakye and Schultz, 1994) resulted in missing values in streamflow records. Such data restrictions have limited attempts to systematically assess streamflow trend, variability and changes at the regional scales. Descroix *et al.* (2009) reported a negative trend (more than 15% decrease) in streamflow for the 1960-2010 period in Sudanian regions (receiving approximately 700 - 1300 mm yr⁻¹ annual rainfall) as a response to changes in rainfall regimes. Also, Mahé *et al.* (2013) found that a decrease in annual rainfall of around 20% since 1970 has resulted in a streamflow decrease of up to 60% for some rivers in West Africa (*e.g.* Niger and Senegal rivers). Amogu *et al.* (2010), in their attempt to regionalize runoff evolution over western Africa (1950-2010), found a clear difference between the Sahelian zone (where, curiously, runoff increases despite reduction in rainfall) and Sudanian and Guinean areas (where runoff decreases logically with rainfall). While major rivers of West Africa show a distinct runoff decrease since 1970, river flows in Central Africa do not show any long-term trend (Mahé *et al.*, 2013). However, these results are restricted to a few long and gap-free time series, making it difficult to describe regional streamflow variability.

Changes in the observational networks (*e.g.* station locations, gauge maintenance and data collection methods) have limited attempts to study streamflow trends and variability at regional scales. Different gap filling methods have been used (*e.g.* regression analysis, time series analysis, artificial neural network and interpolation). Multiple imputations approaches, such as proposed by Rubin (1987), were recently implemented to construct a complete rainfall-runoff database in Iran (Kaltch and Hjorth, 2009). More complex methods such as artificial neural networks (Kim and Pachepsky, 2010) and random forest-based approaches (Stekhoven and Bühlmann, 2012) have also been implemented for gap filling problems with satisfactory results. Despite many hydrological data gap filling studies, few African examples exist. Most African studies focus on gap-free stations (*e.g.* Nka *et al.*, 2015) or reconstructions using linear interpolation techniques. A decision support system based on length of data-gaps, seasons, climatic zones, model performances and data availability has been provided by Gyau-Boakye and Schultz (1994), but such a system would be difficult to implement at the regional scale due to substantial input data requirement and it

may result in: i) spatially non-homogenous reconstructions, and ii) non-statistically independent reconstructions from climate variables. The development of regional climate models (RCM) also open new prospects for hydrological data reconstruction. For instance, Moalafhi *et al.* (2017), testing such approaches over the Limpopo basin, in southern Africa, found that dynamical downscaling of reanalysis products can be very useful for semi-arid, data-scarce environments. However, important biases in RCM physics combined with uncertainties in hydrological modeling could substantially impact the quality of streamflow estimates. The present study aims at (1) providing a new, robust reconstructed streamflow dataset, using only streamflow records as predictands, over West and Central Africa, and (2) using the new dataset, together with gridded climatic data, to examine and assess flow changes and variability over the region between 1950 and 2005. This paper is organized as follows. After introducing the study area and the different datasets in section 2, we present the methods in section 3. In section 4.1, two gap filling methods are used to generate a robust and complete streamflow dataset for West and Central Africa. Then, we examine changes (abrupt and gradual) and variability in streamflow, and we compare these results to those observed in climatic variables from section 4.2 to section 4.4. Results are interpreted, and their wider implications discussed in Section 5.

2. STUDY AREA AND DATASETS

2.1 Research Area

The study area covers West and Central Africa, from -10°N to 25°N and -20°E to 30°E . The selection of the research area was motivated by the key importance of climate change and variability in this region, which also has a dense streamflow gauging network (Figure 1). Some records were short or incomplete, mainly due to equipment failure, instrumentation maintenance issues and sometimes political instability. Most hydrological studies in the region primarily refer to four eco-climatic zones, which are based on both annual rainfall amounts and agricultural properties (FAO, 2004; Roudier *et al.*, 2014): (a) the Sahelian zone (mean annual precipitation ranges between 250 and 500 mm yr⁻¹), (b) the Sudano-sahelian zone (mean

annual precipitation ranges between 500 to 900 mm yr⁻¹), (c) the Sudanian zone (mean annual precipitation ranges from 900 to 1100 mm yr⁻¹) and (d) the Guinean zone (mean annual precipitation exceeds 1100 mm yr⁻¹). However, more complex classifications based on seasonal to interannual variability of rainfall can be found (Badr *et al.*, 2016; Hermann and Mohr, 2011; Janicot, 1992; L'Hôte *et al.*, 1996; Mahé *et al.*, 2001). West African rainfall is primarily related to the displacement of the Intertropical Convergence Zone (ITCZ), which results in two major seasonal cycles. Regions with less than ~1100 mm yr⁻¹ annual rainfall are characterized by a single rainy season with a maximum in August, while, further south, the rainfall seasonal cycle is characterized by two rainy seasons (September-November and March-July) (*e.g.* L'Hôte *et al.*, 1996; Roudier *et al.*, 2014). The boundary between these two zones is however not very clear, with areas experiencing both cycles from year to year because of high rainfall variability (*e.g.* Le Barbé *et al.*, 2002). These different rainfall patterns result in different streamflow regimes. If the characteristics of the flow hydrographs coincide with the rainfall seasonal cycle, aspects such as the timing of the peak and the shape of hydrographs are mainly related to the size and physical properties of drainage basins (Roudier *et al.*, 2014). For example, headwater catchments in the Niger river basin (*e.g.* Mopti, Koulikoro), are characterized by hydrographs with shorter lag times compared to their downstream counterparts (*e.g.* Niamey, Malanville).

Figure 1: Study area with locations of the main catchments (grey shaded), the river network (blue) and streamflow gauges collected from the SIEREM database (light blue dots).

Furthermore, water related issues have led to the construction of several hydraulic structures, which can have significant impacts on hydrological regimes in some basins. According to the Global Reservoir and Dam database (GRanD; Lehner *et al.*, 2011), large dams (capacity >10⁶.m³), as defined by the International Commission on Large Dams (ICOLD; <http://www.icold-cigb.net/GB/Dictionary/dictionary.asp>), are primarily located in the Volta basin (53.5%) and in the Niger River basin (35.2%; Figure 2). The other large dams are distributed within the Lake Chad basin (9.4%), the Senegal River basin (1.2%) and the

127 Congo basin (<1%; Figure 2). This corroborates the study by Adeaga *et al.* (2012) who found that the Volta
128 River and the lower Niger River are the most impacted rivers in western Africa. A summary of the key
129 characteristics of the existing major water resource schemes (hydropower, irrigation) in the Volta basin is
130 provided by McCartney *et al.* (2012).

131 **Figure 2: Large dams (Capacity > 10⁶ .m³) in the study area and their start of operation (purple: 1920-1950; blue: 1950-
132 1970; green:1970-1990; red: 1990-2006). Data source: Global Reservoir and Dam database (GRanD; Lehner *et al.*, 2011).**

133 2.2 Data

134 2.2.1 Streamflow data

135 Mean daily streamflow data were collected from the SIEREM (“Système d’Informations
136 Environnementales sur les Ressources en Eaux et leur Modélisation”) database, which initially consisted of
137 data collected by the “Institut de Recherche pour le Développement” (IRD). Further developments include
138 data quality assessment and a coupling to gridded environmental data over West and Central Africa (Boyer
139 *et al.*, 2006). Station metadata and GIS format files (basin contours, hydrological network, soil water
140 holding capacity, vegetation, and geology) can freely be retrieved from
141 <http://www.hydrosciences.org/sierem>. Additional streamflow data for the Niger River (Idah, Lokoja,
142 Makurdi and Onitsha) were collected from the National Inland Waterways Authority of Nigeria.

143 Over the study area, 863 daily streamflow datasets were collected, and monthly time series were generated
144 but only for the complete months. The percentage of missing data was then calculated for the entire region
145 (Figure 3A), and only stations with less than 50% missing records were selected for analysis. This approach
146 covered most of the study area, and rigorously minimized reconstruction errors. (see Appendix A for the
147 list of reconstructed stations). Figure 3B shows that most gaps are in the 1950s and 2000s for the 152
148 selected stations. This is due, in some countries, to the absence of gauging stations (*e.g.* Burkina Faso) and
149 lack of updated records for the recent period (*e.g.* Central African Republic).

150 **Figure 3: A) River network (blue lines) and spatial distribution of stream gauges over the study area, and with their
151 percentage of missing data (purple=0-25%, blue=25-50%, green=50-75% and red= >75%. Major catchments are**

152 displayed in grey shaded. B) Time-evolution of missing values for the 152 selected stream gauges. Missing values are in
 153 red, while observations are in grey. Stations are ordered by country (BF: Burkina Faso, BJ: Benin, CF: Central African
 154 Republic, CG: Democratic Republic of Congo, CI: Cote d'Ivoire, CM: Cameroon, GA: Gabon, GH: Ghana, GN: Guinea
 155 Conakry, ML: Mali, NG: Nigeria, SN: Senegal, TD: Chad, TO: Togo). The black line represents the number of records
 156 per month over the study area for the 1950–2005 period.

157 2.2.2 Gridded climate data

158 To investigate climate variability and its impact on streamflow regimes over West and Central Africa,
 159 gridded monthly climatic datasets (P, T, PET) from the Climatic Research Unit (Mitchel and Jones, 2005)
 160 were used. The dataset consists of monthly climatic data for the entire world (generated with more than
 161 4000 weather stations at the global scale) at half a degree resolution. The development of this database
 162 required seven data sources, the most important being: the Global Historical Climatology Network (GHCN;
 163 Peterson and Vose, 1997), Jones Surface Temperature Anomaly dataset (Jones, 1994; Jones and Moberg,
 164 2003) and Hulme Historical Monthly Precipitation (Hulme *et al.*, 1998). The latest release (CRU TS
 165 v.4.00) was preferred as it was built using a new gridding technique (Angular Distance Weighting), which
 166 provides more robust results due to a better selection of observation stations for gridding (Harris and Jones,
 167 2017). Unlike precipitation and temperature, which are observed variables, potential evapotranspiration
 168 was derived based on a variant of the Penman-Monteith formula, i.e. the FAO (Food and Agricultural
 169 Organization) grass reference evapotranspiration equation (Ekström *et al.*, 2007), which assumes a
 170 homogenous grass surface (0.12 m height) with no moisture stress, surface albedo of 0.23 and bulk surface
 171 resistance of 0.70 s/m. All climate variables are measured at 2m AGL (Above Ground Level), except for
 172 wind speed (commonly recorded at 10m AGL) which has been reduced to 2m AGL using a conversion
 173 coefficient. Absolute values of the different variables were computed using the baseline values (*i.e.* 1961–
 174 1990 long-term average) (see Harris *et al.*, 2014; Appendix 1).

175 Even though the high spatial resolution of the dataset makes it very convenient for investigating local
 176 processes, limited number of operational stations over West and Central Africa before 1940 could have
 177 resulted in inconsistencies in the CRU dataset (Mitchell and Jones, 2005). Therefore, analyses in this study

will be performed from 1950 to 2005. In addition, Harris *et al.* (2014) compared the CRU dataset to datasets developed by the University of Delaware (UDEL) and the Global Precipitation Climatology Centre (GPCC), which both used different observation datasets, interpolation and quality control techniques than the CRU dataset and found good agreement. For instance, for the period considered in this study, mean annual precipitation values from the CRU dataset and the GPCC dataset have a correlation coefficient of 0.9885 significant at $p \leq 0.1$.

3. METHODOLOGY

Methods have been implemented using R, a free software environment for statistical computing and graphics (<https://www.R-project.org/>).

3.1 Gap filling Methods

Although parametric gap-filling methods are more commonly used (*e.g.* Gyau-Boakye and Schultz, 1994; Kalteh and Hjorth, 2009), non-parametric tests are more suitable for hydroclimate variables, as there is no assumption regarding the distribution of the data. Both parametric and non-parametric gap filling methods are therefore tested in this study to generate robust streamflow reconstructions.

3.1.1 Multiple Imputation by Chained Equations (MICE)

Based on a set of imputation models defined for individual variables with missing values, Multiple Imputation by Chained Equations (MICE; Van Buuren and Oudshoorn, 1999) is a practical approach for handling missing data. The method has been successfully tested for both continuous and categorical variables in hydrology (*e.g.* Kalteh and Hjorth, 2009).

In this study, for each incomplete streamflow record, the first step consists of imputing missing values by randomly sampling with replacement from the distribution of observed values. The observed values of each streamflow station are then regressed to other streamflow stations, and missing values are completed by simulated draws from the corresponding posterior predictive distribution of the considered variable (*e.g.* observed values of x_1 are regressed on all other variables $x_2 \dots x_k$, and the missing values of x_1 are sampled

from its posterior predictive distribution). Several simulations are required to generate a stable single reconstructed streamflow dataset, and the process is repeated several times to generate multiple complete streamflow datasets. In most applications, linear regression models are used for imputing normally distributed continuous variables. The different steps are summarized below:

Considering an incomplete variable z (with n_{obs} observed values) to be reconstructed using other complete variables $X = (x_1 \dots x_k)$ the following linear model is used:

$$z|x; \beta \sim N(\beta x, \sigma^2) \quad (\text{eq. 1})$$

Let $\hat{\beta}$ be a row vector of length k , a realization of the estimated parameters from fitting the model with the observed z . V represents the covariance matrix of $\hat{\beta}$, and $\hat{\sigma}$ the estimated root mean-squared error.

Imputation parameters σ^* and β^* are drawn from the exact joint distribution of σ, β such that:

$$\sigma^* = \hat{\sigma} \sqrt{(n_{obs} - k)/g} \quad (\text{eq. 2})$$

$$\beta^* = \hat{\beta} + \frac{\sigma^*}{\hat{\sigma}} u_1 V^{1/2} \quad (\text{eq. 3})$$

with g , a random draw from a χ^2 distribution on $n_{obs} - k$ degrees of freedom, u_1 a row vector of k independent random draws from a standard Normal distribution and $V^{1/2}$ the Cholesky decomposition of V .

For each missing observation z_i estimates are calculated:

$$z_i^* = \beta^* x_i + u_{2i} \sigma^* \quad (\text{eq. 4})$$

where u_{2i} is a random draw from a standard normal distribution.

As the normal assumption is not often valid for streamflow data (e.g. Kundzewicz and Radziejewski, 2006), missing values were estimated using the Predictive Mean Matching (PMM) approach, which samples estimates from the observed values of the variable z . Instead of estimating missing values of z as in eq. 4, PMM identifies α elements with the smallest error $|\hat{\beta} x_h - \beta^* x_i|$ ($h=1, \dots, n_{obs}$). One of these elements is randomly selected and the imputed value of z_i is z_i^* . The method was implemented using 50 iterations and 100 multiple imputations, which produce a standard deviation only 0.25% wider than a case

of infinite multiple imputations according to Rubin (1987). The median was taken as the better estimate to combine the multiple reconstructed datasets.

3.1.2 Random forest-based reconstruction

The method is based on the random forest (RF) technique (Breiman, 2001), and involves iteratively training a RF on observed values for predicting the missing values. This method was chosen for its ability to perform under high dimensions, complex interactions and non-linearity (Stekhoven and Bühlmann, 2012). Furthermore, compared to other gap filling methods (*e.g.* KNNimpute: Troyanskaya *et al.*, 2001; MICE: Van Buuren and Oudshoorn, 1999), it does not require tuning parameters and prior knowledge of the data, making it computationally attractive. The main limitation, however, is the lack of understanding around the construction of the different trees. The different steps are presented below:

Assuming $X = (X_1, X_2, \dots, X_p)$ a $n \times p$ -dimensional dataset (in our case n observations and p streamflow gauges), the missing values are estimated based on a RF trained on the observed parts of the dataset. For a given gauging station X_s with missing values at the indices $i_{mis}^{(s)}$, the dataset is separated in four parts:

- The observed streamflow values at the station X_s , denoted by $Y_{obs}^{(s)}$;
- The missing values at the station X_s , denoted by $Y_{mis}^{(s)}$;
- The other gauging stations with streamflow records at the indices $i_{obs}^{(s)} = \{1, \dots, n\} \setminus i_{mis}^{(s)}$ denoted $X_{obs}^{(s)}$;
- The other gauging stations with streamflow records at $i_{mis}^{(s)}$ denoted by $X_{mis}^{(s)}$.

The initial step consists of an initial guess of missing values using mean values. The data frame is then sorted and gauging stations are placed in increasing order, based on the proportion of missing data. For each gauging station X_s , the missing data is imputed by first fitting a RF taking $Y_{obs}^{(s)}$ as response variable and $X_{obs}^{(s)}$ as predictors; then estimating missing values $Y_{mis}^{(s)}$ by applying the trained RF to $X_{mis}^{(s)}$. The procedure is repeated until the difference between the newly filled data matrix and the previous one increases for the first time. The stopping criteria is defined as follows:

$$\Delta = \frac{\sum_{j \in N} (X_{new}^{imp} - X_{old}^{imp})^2}{\sum_{j \in N} (X_{new}^{imp})^2} \quad (\text{eq. 5})$$

249 The simulations were performed using 1000 trees with the maximum number of iterations set to 100.

250 **3.1.3 Validation of gap filling methods**

251 The validation method used to assess the performance of the implemented reconstruction techniques
 252 involves generating artificial gaps in the time series, performing the reconstructions on the new dataset and
 253 estimating agreements between predictions and observations. Over the study area, the assumption of data
 254 missing completely at random was considered. First, observed values (12, 24, 36, 48, 60 and 120 months)
 255 over the entire period, 1950–2005, were randomly removed in each of the stations and later compared to
 256 the predictions. Secondly, we randomly removed entire segments of observed data to assess the ability of
 257 the gap filling methods to reconstruct contiguous missing data. The modified Kling-Gupta Efficiency
 258 (KGE) was used as an indicator of agreement between observations and predictions. This efficiency
 259 criterion ensures that the temporal dynamics (measured by the correlation coefficient) as well as the
 260 distribution of flows (measured by both the bias and variability ratio) are well represented (Kling *et al.*,
 261 2012).

262 **3.2 Step change detection and trend analysis**

263 Changes (natural or artificial) in hydro-climatic time series can occur abruptly (step change) or gradually
 264 (trend) or in more complex forms (Machiwal and Jha, 2006). Step-like changes, induced by reservoir
 265 construction and changes of gauging structures, for example, might also result from gradual changes since
 266 nonlinear system dynamics may show cumulative effects and thresholds (Kundzewicz and Radziejewski,
 267 2006). In this study, step-like changes in the mean are investigated in reconstructed mean annual
 268 streamflow time series using a multiple change-points detection analysis (Killick and Eckley, 2014). This
 269 technique, which is similar to the method proposed by Hubert *et al.* (1989), is based on the segment
 270 neighborhood algorithm (Auger and Lawrence, 1989). The non-parametric cumulative sum test statistic
 271 (Page, 1954) is used to assess the optimal position of change-points.

272 Linear trends are then investigated for periods defined based on the results of the multiple change-points
 273 analysis at the regional scale. The significance of the Mann-Kendall (MK) test (Kendall, 1975; Mann,

1945) is highly sensitive to positive serial correlation (Von Storch, 1995), so its variant (Yue *et al.*, 2002) was preferred for linear trend detection here. The Yue *et al.* (2002) method assumes trends are linear; datasets are first detrended before extracting the lag-1 serial correlation from the detrended series (i.e. a trend-free pre-whitening procedure (TFPW) which should generate independent residuals series). The detected trend and the residuals are combined, before the MK test for significance is applied. The Theil Sen Approach (TSA) is used to estimate the slope of the trend in a dataset. This approach is less sensitive to outliers and therefore provides a better estimate of slope for skewed data, compared to regression methods. In addition, as trend values are highly dependent on start and end dates, a multitemporal trend analysis approach has been implemented here (Liebmann *et al.*, 2010; McCabe and Wolock, 2002). Trends here are calculated for all possible segments (minimal length of 5 years) from 1950 to 2005 to explore and define the time series internal variability. For each time series, the multitemporal trend analysis generates a diagram in which each possible pair of start and end dates is associated with a trend value. To investigate the spatial extent and zonal coherence of the different variability patterns in precipitation and streamflow, the multi-temporal trend analysis results were grouped using hierarchical clustering, using the Euclidean distance as the metric of dissimilarity. Different approaches exist to determine the optimal number of clusters (Charrad *et al.*, 2014), but we used the multiscale bootstrapping approach of Suzuki and Shimodaira (2006), which allows uncertainty estimation for each cluster. This is achieved through thousands of bootstraps resampling and used to estimate the probability that a cluster appears in the replicates.

4. RESULTS AND DISCUSSIONS

4.1 Reconstruction outputs

Two reconstruction methods were applied to the subset of streamflow stations with less than 50% missing data (*i.e.* 152 streamflow gauges here). All 152 stations were reconstructed with satisfactory results as illustrated in Figures 4 and 5.

298 The validation shows that gap filling methods perform well for both cases of randomly removed
 299 observations and contiguous missing segments. Figure 4 shows that the median of the KGE is always
 300 greater than 0.75, which indicates that for half of the stations, the worst component (*e.g.* correlation, bias
 301 ratio or variability ratio) is higher or equal to 0.75: this suggests good reconstruction performance. Very
 302 similar results were achieved using the Nash-Sutcliffe Efficiency and the normalized Root Mean Squared
 303 Error (not shown). Also, both methods are reasonably stable when artificially increasing the number of
 304 missing observations and when artificially increasing the length of missing segments, despite an artefact
 305 suggesting better performances with increasing missing data, which is in fact caused by the sensitivity of
 306 efficiency criteria to sample size (*e.g.* Schönbrodt and Perugini, 2013). However, MICE seem to perform
 307 better than RF when increasing the number and the length of missing data (Figure 4).

308 **Figure 4: Validation of gap filling methods: boxplot of validation efficiencies for all the reconstructed stations; upper**
 309 **panels for randomly removed values and lower ones for cases of randomly missing data segments. A red line is drawn at**
 310 **KGE=0.75. Outliers are represented in blue dots.**

311 To compare both gap filling methods, results from five stations from different climatic zones and
 312 hydrological regimes are presented in Figure 5. While both methods show similar results overall,
 313 significant dissimilarity appears in some cases, such as in the Niger River at Niamey, where MICE show an
 314 abrupt increase in minimum flow, and decrease in peak flow from 1999 (Figure 5). This pattern, which is
 315 similar to those induced by large dams (higher low flows and lower peak flows in downstream reaches), is
 316 not consistent with recent studies in the region (*e.g.* Amogu *et al.*, 2010; Mahé *et al.*, 2013), highlighting
 317 increased runoff coefficients at Niamey from the 1990s. MICE generate estimates of missing values by
 318 sampling from the observed values and might therefore fail at reconstructing flows beyond observed
 319 ranges. Thus, even though MICE often seem to provide better estimates than the RF based method, the
 320 latter appears to be more appropriate in the context of changing hydrological regimes, because of its ability
 321 to capture complex nonlinear relations between predictors and predictands.

Figure 5: Reconstructed time series for five streamflow stations representative of different climatic conditions: Wayen (Sahelian), Bonou (Tropical humid), Mbasso (Tropical humid), Niamey (Tropical humid, Sudanian and Sahelian), Bangui (Tropical humid). Blue lines represent observations; black dotted lines represent MICE estimates and Red dotted lines represent Random Forest estimates. Red Boxes highlight time windows of interest.

4.1. Streamflow changes between 1950 and 2005

With the assumption that two major break points occurred in the streamflow time series, the step change analysis detected changes in 147 stations over the study area. Both reconstruction methods presented similar results and only those of random-forest based reconstructions are presented. At the regional scale, the first discontinuity in mean annual streamflow occurred in 1970 (Figure 6), with a marked negative shift in the mean (up to -60%). Similar results were found by Hubert *et al.* (1989), for the Niger and Senegal rivers. The second discontinuity at the regional scale occurred around 1993 and is characterized by a positive shift for more than 70% of the stations (with an average increase of about +23%, Figure 6). Despite this positive shift in mean streamflow, recent conditions are still below the 1950s and 1960s wet periods.

Some sub-regional differences, however, emerge along the Gulf of Guinea and regions in Central Africa, where a discontinuity in the mean annual streamflow occurred in the 1950s and early 1960s, with an average positive shift of around 18% (Figure 6). These results are consistent with the findings of Mahé *et al.* (2001), underlining differences in rainfall variability between West and Central Africa from 1951 to 1989. Also, some discontinuities are revealed before the 1990s in some stations (Figure 6), probably induced by the wet episodes observed at the end of the 1980s. Based on the data collected from the Global Reservoir and Dam database (GRanD; Lehner *et al.*, 2011), presented in Figure 2, regional scale discontinuities in streamflow were more likely induced by climate variability and land use change rather than reservoirs as only 4% of the large dams in the region were completed between 1968 and 1970 and 14% between 1985 and 1993.

Figure 6: Locations of step changes in random-forest based streamflow reconstructions: positive shift in the mean (blue), negative shift in the mean (red). Stations are ordered by country (BF: Burkina Faso, BJ: Benin, CF: Central African

348 Republic, CG: Democratic Republic of Congo, CI: Cote d'Ivoire, CM: Cameroon, GA: Gabon, GH: Ghana, GN: Guinea
 349 Conakry, ML: Mali, NG: Nigeria, SN: Senegal, TD: Chad, TO: Togo). The black curve on top presents the temporal
 350 distribution of change-points over the study area.

351 Gradual changes (trends) are investigated in mean annual reconstructed streamflow time series (MICE and
 352 RF) over the periods defined by the change-points analysis, which highlights two major discontinuities at
 353 the regional scale (1970 and 1993): 1950-1970 (wet conditions), 1970-1993 (drought conditions), 1993-
 354 2005 (partial recovery). Figure 7 presents the correlation between the results from both reconstruction
 355 methods for the different time intervals. Both reconstruction methods show similar streamflow trends at the
 356 regional scale (Figure 7). However, although the results from both methods remain significantly correlated
 357 ($p \leq 0.1$), trends differ slightly in the post-1990 period, mainly due to the limited ability of MICE to
 358 extrapolate beyond the range of observed values, highlighting that hydrological regimes may have changed
 359 in the 1993–2005 period.

360 **Figure 7: Spatial correlation between normalized trends calculated using both reconstructed datasets, for the three**
 361 **periods of investigation: 1950-1970 (red), 1970-1993 (green) and 1993-2005 (blue).**

362 Trend analysis over the three different time intervals revealed that, during the 1950–1970 period, even
 363 though mean annual streamflow values are at the highest, streamflow trends are significantly negative (up
 364 to -4% per year) over the Sahelian and Sudanian regions of West Africa (Figure 8a-b): this suggests that the
 365 step change observed around 1970 in this region was mainly induced by a gradual aridification pattern.
 366 During the same period, significant positive trends are identified over Central Africa (up to +2.5% per year)
 367 (Figure 7a-b). At the regional scale, 35% and 30% of trends are significant for MICE and RF respectively.
 368 Among those significant trends, 52% and 40% are positive mainly in Sudanian and coastal regions (Figure
 369 8a-b) for MICE and RF respectively. Most of the significant negative trends are in the Sahelian region,
 370 driven by dryer conditions in the end of the 1960s compared to the 1950s (Figure 8a-b).

371 These negative streamflow trends along the Sahelian band spread toward the Gulf of Guinea and over
 372 Central Africa during the well-known drought period of the 1970s and 1980s (Dai *et al.*, 2004; Lebel, 2003;
 373 Nicholson, 2013; Figure 8c-d), marking a stronger spatial coherence. During this dry period, mean annual

streamflow values decrease by up to 69% compared to the 1950s and 1960s. Also, more than 90% of all significant trends (40% and 38% using MICE and RF, respectively) are negative (up to -5% per year), as a result of intensified dry conditions from the end of the 1960s (Figure 8c-d).

The last period (1993-2005) is characterized by a reduction in significant trends [MICE (26%) and RF (8%)] and contrasting patterns mainly due to the limited ability of MICE to fully capture complex streamflow interactions (Figure 8e-f). Compared to the previous period (1970-1993) mean annual streamflow values mark an increase of at least 15% over more than half of the study area and a decrease of around 7% in some regions (Figure 8c-f). Significant positive trends on the Niger River, as shown using RF, would however be consistent with the “Sahelian paradox” (Descroix *et al.*, 2013; Mahé *et al.*, 2005), with a higher flow contribution from the Sahelian tributaries. Despite positive rainfall trends in some Sudanian areas (Northern Ghana and Ivory Coast), which are detected using both MICE and RF, streamflow trends remain negative (Figure 8e-f). This might have resulted from severe groundwater depletion during the dry periods 1970s and 1980s (Mahé *et al.*, 2005), but this needs further research.

Figure 8: Streamflow trends estimated for both reconstructed datasets, upward triangles for positive trends and downward triangles for negative trends, filling highlights the significance of trend at 10% (negative trends in red and positive trends in blue). River basins are greyed and the river network in blue.

4.2 Observed climatic trends between 1950 and 2005

4.2.1 Trends in annual precipitations

Annual rainfall trends for the 1950–1970 period decline by ~10 mm yr⁻¹ (significant for around 34% of the study area) along the entire Sahelian band, but locally increase in parts of the Central African Republic and Democratic Republic of Congo (Figure 9a). This suggests that the drying trends might have started earlier than hitherto recognized. The negative trends observed along the Sahelian band then spread towards the Gulf of Guinea during the 1970–1993 period (Figure 9b), similarly to the pattern observed in streamflow (Figure 8 c-d).

398 However, although this period is widely recognized to be extremely dry from comparisons of mean values,
 399 we find here that only 11.5% of the study area show significantly negative precipitation trends.
 400 Interestingly, however, significant positive trends are identified in the Congo River basin (Figure 9b). This
 401 highlights a potential hiatus in the regional drying trend during the 1970s and 1980s, supporting earlier
 402 studies (Le Barbé and Lebel, 1997; D'Amato and Lebel, 1998). These could result from increasing quasi-
 403 decadal rainfall variability as suggested in Dieppois *et al.* (2013, 2015). In the post-1993 period, we note an
 404 increase of annual precipitation compared to the previous period (trends significant for 11% of the study
 405 area), corroborating previous findings (Biasutti, 2013; Lebel and Ali, 2009; Nicholson *et al.*, 2000). This
 406 potential annual rainfall recovery ($\sim +11.5$ mm yr⁻¹) is particularly pronounced in western and eastern
 407 Sahel and Liberia (Figure 9c), which agrees with the findings of Ogungbenro and Morakinyo (2014) in
 408 northern Nigeria. At the same time, regions in northern Cameroon and in the Democratic Republic of
 409 Congo, are characterized by significant negative trends (~ -7 mm yr⁻¹, to ~ -30 mm yr⁻¹), in agreement
 410 with the recent study of central African rainfall by Maidment *et al.* (2015).
 411 The same analysis, conducted using the GPCC V7 datasets, show similar patterns. The relationships are,
 412 however, slightly more significant over the study area for the three periods (35%, 11.43%, and 14.65% for
 413 the 1950-1970, 1970-1993 and post-1993 periods, respectively; not shown). In addition, during the post-
 414 1993 period, the GPCC V7 dataset underlines a significant decreasing trend in Guinea (which,
 415 interestingly, does not appear in the CRU dataset) and a wider spatial extent of negative trends in
 416 Cameroon and Central African Republic. Despite these slight differences probably resulting from the
 417 greater number of observation stations used to generate the GPCC V7 dataset, agreement between
 418 precipitation and streamflow trends remains strong.
 419 Overall, there is a good agreement between annual streamflow and precipitation trends over the entire study
 420 area highlighting the importance of precipitation in driving hydrological systems. However, quantifying
 421 runoff response to increasing precipitation is likely to be a complex task since rising temperatures and
 422 potential evapotranspiration could offset increasing precipitation. This issue is addressed in the following

section by investigating trends in temperatures and potential evapotranspiration and their impact on runoff responses.

4.2.2 Trends in mean annual minimum and maximum temperatures, and potential evapotranspiration

As widely accepted, temperatures over the African continent have been increasing during the 20th century (since 1950), and this is primarily associated with anthropogenic causes (*e.g.* IPCC, 2014; Stott *et al.*, 2010). However, here we aim to discuss temperature trends in term of impact on water resources, through its impact on evapotranspiration and effective rainfall.

Trends in annual minimum and maximum temperatures over the study area show different amplitude and spatial extents. For instance, in West and Central Africa, the 1950–1970 period is characterized by positive trends (+0.5 to +1.5°C) in minimum annual temperatures (significant for 32.5% of the study area). However, weaker and spatially less coherent trends are detected for annual maximum temperatures (~ +0.5°C; significant for 9.5% of the study area). Maximum values are reported only in the western Sahel (Figure 9d, g). The rest of the study area shows few significant trends, apart from some significant negative trends in both minimum and maximum annual temperatures (Figure 9d, g). According to the CRU potential evapotranspiration estimates, the patterns in both minimum and maximum temperatures could have resulted in significant positive evapotranspiration trends (~ +2.5 mm yr⁻¹) in western and central Sahel, and significant decreasing trends (~ -3.75 mm yr⁻¹) over the Gulf of Guinea and Central Africa regions (Figure 9j).

The 1970–1993 period is marked by a homogeneous increase in annual minimum temperatures, which is significant over 63% of the study area (including regions in the Congo River basin, where significant cooling is identified; Figure 9e). These trends contrast with annual maximum trends, which are negative in the Sahelian region (~ -1°C), but positive in the Gulf of Guinea coastal regions and Central Africa (Figure 9h). This configuration is, however, consistent with a weaker meridional thermal gradient, which characterizes a southward shift of the ITCZ and dry conditions in the Sahel (Chiang and Friedman, 2012;

Webster *et al.*, 1998). The fluctuation of temperature range during this period has driven a uniform decrease in potential evapotranspiration over the Sahelian band but increased significant positive trends in the Gulf of Guinea and Central Africa (Figure 9k).

Since 1993, greater spatial coherence emerges, with increasing trends of both annual minimum temperatures (significant for 65% of the study area) and maximum temperatures (significant for 85% of the study area; Figure 9f, i). Trends in annual maximum temperatures, however, are more pronounced ($\sim 0.1^{\circ}\text{C}$ higher in average) than in annual minimal temperature (Figure 9i). This could be an artefact of the baseline period used in our study, as this result contrasts with those revealed in some other studies (*e.g.* Funk *et al.*, 2012; Ringard *et al.*, 2016), which suggested that minimum temperatures have risen faster compared to maximum temperatures in the post-1990 period. Nonetheless, temperature trends are consistent with trends in potential evapotranspiration (Figure 9l), which highlight a uniformly significant (for around 46.8% of the study area) and positive trend ($\sim < +3.8 \text{ mm yr}^{-1}$) over almost the entire eastern part of the study region. Regions in western, eastern Sahel and part of the Gulf of Guinea, however, show non-significant negative trends (Figure 9l), which may result from the spurious trends (above) in minimum temperatures and errors resulting from the use of the same monthly wind speed values (1961-1990) for each year.

Trends in effective rainfall, approximated here as the difference between rainfall totals and potential evapotranspiration are presented in Figure 9m-o. Over the two first periods (1950-1970 and 1970-1993), these trends are similar to precipitation trends: this suggests the limited effect of potential evapotranspiration on the relationship between rainfall and streamflow (Figure 9m-n). However, from 1993, the situation is reversed, mainly in the eastern part of the Sahel (eastern Niger, Chad and northern Nigeria), where high potential evapotranspiration rates significantly subdue the potential impact of the rainfall recovery (Figure 9o) on streamflow. This might help explain, at least partially, why the rainfall recovery over these regions is not associated with significant positive streamflow trends (Figure 8c-d). Over Central Africa (areas in the Congo basin), it also appears from trends in effective rainfall that during

recent decades, the decrease in rainfall is exacerbated by increased evapotranspiration (Figure 9c, i, o). This suggests enhanced climatic stress on Central African streamflow in relation to warming temperatures.

Figure 9: Hydroclimatic trends over the study area for three different time intervals (1950-1970, 1970-1993 and 1993-2005) according to the CRU.TS. V4.00 dataset: a-c) Annual precipitation trends d-f) Minimum annual temperature trends g-i) Maximum annual temperature trends j-l) Annual potential evapotranspiration trends m-o) Annual effective rainfall trends. Sen's slope values are displayed through a red-white-blue color scale. Solid red lines highlight trend significance at $p \leq 0.1$ according to a modified MK trend test accounting for serial correlation in the time series.

4.3 Precipitation and streamflow variability

Standard trend analysis methods assess the slope of the considered variables over the period of investigation. The value of the slope is, however, highly dependent on the selected time window and changes significantly for different start and end dates, mainly because of internal variability. Such limitations are tackled in the multitemporal trend analysis method (Liebmann *et al.*, 2010; McCabe and Wolock, 2002). We used the Liebmann *et al.* (2010) approach, to calculate precipitation and streamflow trends for all possible segments of 5 to 56 years between 1950 and 2005. The results are stored in two-dimensional diagrams (*e.g.* Figure 11), which have been analyzed using multiscale bootstrapped agglomerative hierarchical clustering.

Clustering streamflow variability diagrams resulted in three main clusters, which are highly significant ($p \leq 0.1$) based on the multiscale bootstrapping test, with associated spatial distributions presented in Figure 10, identified using hierarchical clustering.

Figure 10: Spatial distribution of streamflow variability (1950–2005) clusters based on multi-temporal trend analysis superimposed on the river network (blue) and major river basins (grey shaded). All the clusters are highly significant at $p \leq 0.1$ according to the multiscale bootstrapping test. Different colours displayed the location of the different clusters.

Overall, these three clusters show decreasing flow trends over the entire period (1950–2005), but we also identify decadal periods of alternating positive and negative trends with different amplitudes, modulating the general trend, according to the three clusters (Figure 11). For instance, a pronounced positive trend in the mid-1970s during the drought period emerges in cluster 1 (Congo Basin at Brazzaville), which

progressively disappears in cluster 2 (lower Niger River, Benue and stations in the upper Congo basin) and cluster 3 (all the other stations; Figure 11). This emphasizes the importance of decadal variability in modulating streamflow trends (which has hitherto been little studied) and provides a new picture of the behaviour of hydrological systems in West and Central Africa.

These differences in the contribution of interannual to decadal variability could, however, arise from differences in the large-scale climate drivers. According to Mahé *et al.* (2013), Cluster 1, which is located at the outlet of the Congo Basin at Brazzaville, could be more sensitive to changes in the thermal gradient between the Atlantic and Indian oceans resulting in a unique runoff variability. Such decadal fluctuations have also been reported for eastern Sahel rainfall in Dieppois *et al.* (2013, 2015), suggesting that differences between clusters should at least partly be related to different interactions with catchment properties (*e.g.* reduction in soil water holding capacity) and water management. In addition, while trend amplitude is a distinctive element between clusters, the sign and temporal scale during the humid period (1950-1960) and the recovery period (post-1990) are also very important. For instance, stations in clusters 1 and 2 are characterized by wet conditions in the 1950s-1960s, whereas most of the stations in cluster 3 show decreasing trends during the same period (Figure 11). Furthermore, cluster 3 highlights less intense dry conditions in the 1980s and a more pronounced recovery in the recent years compared to the first two clusters (Figure 11). A further classification of the stations in cluster 3 is provided as supplementary materials. The significant negative trend (observed in the 1980s) in stations of cluster 2, for instance might have been partly accentuated by large dams in Nigeria (*e.g.* the Dadin Kowa Dam and the Kiri dam, on a main tributary of the Benue river).

Figure 11: Multi-temporal diagrams of the different cluster centroids: trends in m^3/s are presented in red (negative) – white (null) – blue (positive) colour scale, contours lines represent trend significance at $p \leq 0.1$.

Applying the same clustering method to gridded annual rainfall, variability diagrams resulted in 12 major clusters ($p \leq 0.1$) and few grid points with lower probabilities ($p \leq 0.2$) and therefore unclassified (Figure 12).

Figure 12: Clusters of rainfall variability generated using CRU TS V4.00 (2.5°x2.5°) on the period 1950-2005: colours and numbers from 1 to 12 refer to the grid points within the 12 initial clusters at $p \leq 0.1$. Red boxes represent grid points which did not fall within the clusters. All the clusters are highly significant at $p \leq 0.1$ according to the multiscale bootstrapping test.

West African regions predominantly fall within clusters 11, 8, 2 and 1 (Figures 12, 13) which are mainly characterized by persistent dry conditions from the end of the 1960s, and positive trends starting earlier in clusters 2 and 8 (1970s) compared to clusters 1 and 11 (end of 1980s). Comparing, for instance, patterns observed in streamflow cluster 3 and rainfall cluster 11, it appears that the significant negative rainfall trend in the 1980s is attenuated in the streamflow signal and, furthermore, the observed streamflow recovery is more widespread compared to the recovery observed in rainfall. This suggests a combination of drivers which might have enhanced the runoff response, described by some authors as the “Sahelian paradox” (Descroix *et al.*, 2013; Mahé *et al.*, 2005) which refers to a counterintuitive increase in runoff coefficient despite decreasing rainfall. In fact, parts of this region are known to have experienced drastic changes in land cover resulting from several coupled interactions between increasing cultivated areas (Cappelaere *et al.*, 2009), and natural vegetation changes after the 1970s and 1980s major drought periods (Gal *et al.*, 2017).

The clustering underlines a high variability in rainfall over the western part of West Africa, where some grid points are left outside the clusters. Some parts of this region are characterized by the pattern observed in cluster 9 (Figures 12-13). After the humid period of the 1960s, rainfall is characterized by decreasing trends until the 1990s (Figure 13). From the end of the 1990s rainfalls largely returned to their level of the 1960s as a result of a recovery which started in the 1980s (Figure 13). From these different clusters, it appears that most regions over western Africa have experienced improved streamflow conditions because of the recent rainfall recovery even though long-term trends remain negative.

Over Central Africa, rainfall shows high decadal variability (succession of wet and dry periods) with no clear long-term trends (clusters 4, 5, 6, 10 and 12; Figure 13). This region is characterized by a humid

period from the mid-1950s to the 1970s, with dry episodes around 1980 (Figure 13). In cluster 4, for instance, recent conditions (1990s-2000s) are almost as wet as the humid period, which is not the case for cluster 6 where recent conditions remain relatively drier (Figure 13). The streamflow variability displayed in cluster 1 (Congo basin at Brazzaville) appears to have resulted from a combination of rainfall clusters 6, 10 and 12, highlighting the diverse climatic influences in this basin (Figure 11-13). Rainfall-runoff relations over this region suggest that rainfall is the main driving factor, with no, or limited, effect from other moderating factors (*e.g.* land use change, intensification of agriculture, deforestation, and warming temperatures). The change in seasonal rainfall distribution may likely be the major factor related to climatic change in this area to have an impact on discharges' seasonal regimes. This can be observed at the scale of small basins like the Kienke at Kribi in the South coastal Cameroon, where the small dry season disappeared during the last decades (Lienou *et al.*, 2008), and at the larger scale for the Ogooue river in Gabon, where the Spring flood lost 30% of discharge value after the rainfall regimes slightly changed over past decades (Mahé *et al.*, 2013), the same being observed to a lesser extent for the whole Congo basin (Alsdorf *et al.*, 2016; Tshimanga *et al.*, 2016).

Figure 13: Multi-temporal diagrams of the 12 rainfall variability clusters derived from the multi-scale bootstrap clustering: trends (mm) are presented in red (negative) – white (null) - blue (positive) color scale, contours lines represent trend significance at $p \leq 0.1$.

5. CONCLUSION

Using parametric (MICE) and non-parametric (RF) gap filling methods, a new and complete streamflow dataset, spatially distributed over West and Central Africa and encompassing different climatic zones and hydrological regimes has been generated. Gap filling results highlighted that both methods performed well, though, in general, MICE was slightly outperforming RF. However, due to its parametric nature, MICE analyses, in some cases, failed to capture changes in streamflow conditions (case of Niamey on the Niger River). The complete streamflow dataset (RF method) was then used to investigate streamflow changes and variability and their interactions with key climatic variables (P, T, PET) over West and Central Africa

573 between 1950 and 2005.

574 Majority of streamflow stations over the study area present a step change in 1970 mainly induced by a
 575 gradual aridification pattern. In the 1990s a positive shift in mean discharge is observed, but it is difficult to
 576 conclude whether this change is led by positive rainfall trends or single wet episodes amplified by land use
 577 change, warming temperature and evapotranspiration reduction. In general, there is a good agreement
 578 between streamflow and precipitation trends, with an offsetting effect of potential evapotranspiration
 579 observed in some regions. Over the study area, the period 1950–1970 was characterized by negative
 580 streamflow trends in Sahelian and Sudanian regions of West Africa, which seems counterintuitive
 581 considering that this period was the wettest on record. The opposite is observed over Central Africa where
 582 significant positive streamflow trends emerge. The following period (1970–1993), is marked mostly by
 583 negative trends due to dryer conditions. This pattern is reversed during the 13-year period 1993–2005, with
 584 mainly positive trends resulting from increased rainfall and changes in land use in Sahelian regions. Annual
 585 streamflow trends reflect annual precipitation trends which decrease from the 1950s to 1980s and increase
 586 from the 1990s. More importantly, the study showed that, even though the 1950s and 1960s were the
 587 wettest decades in terms of total rainfall amounts, decreasing annual rainfall trends were more prominent,
 588 suggesting an earlier start of the drought. The drought peaked during the 1970s/80s, over most of western
 589 Africa, but the reduced negative trends in precipitation suggest a hiatus, which have resulted from quasi-
 590 decadal variability.

591 Furthermore, over most of the study area, hydrological regimes during the recent period have been
 592 impacted by the rainfall recovery which is not limited to the west African Sahel. Even though other
 593 climatic variables such as wind speed and vapor pressure deficit might have played an important role,
 594 temperature trends appeared to be highly related to trends in potential evapotranspiration, which seem to
 595 have hampered the effect of the rainfall recovery on hydrological regimes in some areas over the eastern
 596 Sahel (eastern Niger, Chad and northern Nigeria).

597 Building significantly on previous studies, which generally provide trend estimates over a certain period,

we have provided novel information and analyses of the impact of internal variability using the multitemporal trend analysis method. The results highlight strong interannual to decadal signals which clearly modulate streamflow and precipitation trends. In West Africa, for instance, the 1970-1989 period is characterized by two main dry events (1972-1973 and 1983-1984) separated by a wet period (Nicholson *et al.*, 2000; Dai *et al.*, 2004). This probably resulted in increased runoff coefficients in Sahelian catchments, as observed by Albergel (1987) in Burkina Faso over the period (1969-1984) and later in larger Sahelian catchments (Descroix *et al.*, 2013; Mahé *et al.*, 2005). Such a rainfall-runoff response (referred to as the Sahelian paradox) indeed seems paradoxical when considering long-term trends but becomes less counterintuitive when investigating variability in precipitation and streamflow time series. Therefore, rather than describing the “Sahelian paradox” as an increase in runoff despite reduced rainfall since 1970, it should be considered as enhancing runoff response to positive rainfall anomalies, as a result of changes in land-surface properties.

If flow trends can be largely explained by decadal variability in rainfall (Dieppois *et al.*, 2013), influence of other driving factors should also be considered at the catchment level (such as geology, soils, agricultural land use change, water consumption and urbanization). For instance, large dams constructed in the 1980s in Nigeria (*e.g.* the Dadin Kowa Dam and the Kiri dam, on a main tributary of the Benue river), might have affected to some extent the variability of the lower Niger river, but this is beyond the scope of the present paper.

This study has shed light on hydroclimatic variability and its associated impact on streamflow regimes over large, key parts of West and Central Africa over recent decades, and also provides water practitioners with reconstructed streamflow time series which can be used as input for water balance models to develop sound water and agricultural management policies. These useful time series here can form the basis of future developments, to include updating of the streamflow datasets through national water offices. This should further improve the quality of the reconstructions and open up investigations of more recent conditions. In addition, future in-depth studies are required of climate processes (*e.g.* sea-surface temperature,

atmospheric circulation), catchment land use properties, and water management policies, all of which can drive streamflow variability at interannual to decadal timescales. As these potentially modulate the climate signal, such work is required to further improve our understanding of hydrological variability in West and Central Africa, and our ability to model hydrological changes in this region.

Acknowledgements

We acknowledge the HydroSciences Montpellier Laboratory, particularly Nathalie Rouché for providing us with streamflow datasets used in this study. Moussa gratefully acknowledges the funding received towards his PhD studies from Coventry University, UK, and resources provided by the Centre for Agroecology, Water and Resilience (CAWR). We thank the anonymous reviewers for their insightful comments and suggestions.

Appendix A: List of reconstructed streamflow time series

ID	Basin	Station name	latitude	Longitude
BFQ0010	LERABA	YENDERE au pont	10.1667	-5.0683
BFQ0060	VOLTA	WAYEN	12.3789	-1.08
BFQ0064	VOLTA	BOROMO	11.7833	-2.9167
BFQ0065	VOLTA	DAPOLA	10.5667	-2.9167
BFQ0072	VOLTA	NWOKUY	12.5278	-3.55
BFQ0074	VOLTA	SAMANDENI	11.4667	-4.4667
BJQ0009	SOTA	COUBERI	11.74	3.3333
BJQ0022	COUFFO	LANHOUNTA - LANTA	7.1	1.8833
BJQ0028	MONO	ATHIEME	6.9167	1.6667
BJQ0033	OUEME	BONOU	6.9	2.45
BJQ0036	OUEME	HETIN SOTA	6.6	2.5
BJQ0047	OKPARA	KABOUA	8.25	2.7167
BJQ0050	SOTA	RTE KANDI-SEGBANA AMONT	10.9833	3.25
BJQ0075	WE-WE	WE-WE	9.1667	2.1083
BJQ1000	PENDJARI	PORGA	10.99401	0.9773
BJQ2000	NIGER	MALANVILLE	11.888	3.383
BJQ2004	OUEME	PONT DE BETEROU	9.199179	2.267582
BJQ2005	OUEME	PONT DE SAVE	8	2.4167
BJQ2006	ZOU	ATCHERIGBE	7.5333	2.0333
CFQ0005	OUHAM	BOSSANGOA	6.4667	17.45
CFQ0025	OUBANGUI	ZINGA TRANSIT	3.713833	18.58716
CFQ0027	MBOMOU	ZEMIO	5.028726	25.1471
CFQ0028	BANGUI-KETTE	ALINDAO	5.04457	21.20172

CFQ0034	LOBAYE	M'BATA	3.666296	21.98114
CFQ0040	M'POKO	BOSSELE-BALI	4.530737	18.46878
CFQ0057	SANGHA	SALO	3.181621	16.11362
CFQ2000	OUBANGUI	BANGUI	4.364275	18.59487
CGQ0003	ALIMA	TCHIKAPIKA	-1.26385	16.16937
CGQ0013	LEFINI	BWEMBE	-2.9167	15.6308
CGQ0014	LIKOUALA	ETOUMBI	0.0167	14.95
CGQ0015	LIKOUALA	MAKOUA	0.00167	15.633
CGQ0017	N'KENI	GAMBOMA	-1.9	15.85
CGQ0020	KOUYOU	LINNEGUE	-0.5	15.9333
CGQ0026	LIKOUALA	BOTOUALI	-0.55	17.45
CGQ2000	CONGO	BEACH - V.N. Brazzaville	-4.27285	15.29392
CGQ2001	SANGHA	OUESSO	1.6167	16.05
CIQ0007	BANDAMA	MBRIMBO	6.0125	-4.425
CIQ0013	BANDAMA	KIMOUKRO BALISE 10201	6.5056	-5.3053
CIQ0032	MARAOUE	RTE BEOUMI-SEQUELA - KONGASSO 10145	7.8319	-6.2542
CIQ0033	MARAOUE	BOUAFLE 10147	6.979988	-5.75437
CIQ0058	NZI	BOCANDA	7.0442	-4.52
CIQ0061	NZI	DIMBOKRO 10141	6.6358	-4.71
CIQ0154	KOUROUKELE	IRADOUGOU	9.7069	-7.8028
CIQ0292	KAVI	MBESSE	5.8386	-4.2961
CIQ0312	CAVALLY	FLAMPLEU	7.2833	-8.0583
CIQ0314	CAVALLY	TAI	5.86	-7.45
CIQ0319	NSE	TAI 1 (TAI PONT)	5.875	-7.4583
CIQ4020	BANDAMA	BADA	8.1069	-5.4972
CIQ4022	BANDAMA	TIASSALE 10144	5.8947	-4.8178
CIQ4025	NZI	FETEKRO	7.8106	-4.6875
CIQ4026	NZI	MBAHIAKRO 10133	7.4458	-4.3556
CIQ4027	NZI	NZIENOA 10136	5.9964	-4.8125
CIQ4028	COMOE	ANIASSE PONT 10138	6.6375	-3.7126
CIQ4029	COMOE	MBASSO	6.125	-3.48
CIQ4030	COMOE	SEREBOU	7.9383	-3.9419
CIQ4031	SASSANDRA	SEMIEN 10130	7.7083	-7.0669
CIQ4032	SASSANDRA	SOUBRE	5.7833	-6.6131
CIQ4033	BAFING	BAFINGDALA (BADALA) BIANKOUMA 10162	7.841611	-7.66658
CIQ4034	LOBO	NIBEHIBE (NIBEIGBEU)	6.8003	-6.7
CIQ4035	COMOE	AKAKOMOEKRO 10149	7.447418	-3.5094
CMQ0008	DOUME	DOUME	4.2333	13.45
CMQ0029	SANAGA	NACHTIGAL	4.35	11.6333
CMQ0030	SANAGA	NANGA EBOKO	4.7	12.3833
CMQ0038	MBAM	BAC DE GOURA	4.5667	11.3667
CMQ0071	NYONG	DEHANE	3.5667	10.1167
CMQ5001	VINA NORD	PONT DE BEREM	7.55	13.95
CMQ5005	DJA	SOMALOMO	3.3667	12.7333

CMQ5006	BENOUE	BUFFLE NOIR	8.1167	13.8333
CMQ5007	BENOUE	GAROUA	9.294019	13.4041
CMQ5015	MAPE	AU PONT DE MAGBA AMONT	5.9833	11.2667
CMQ5016	VINA DU SUD	LAHORE	7.25	13.5667
CMQ5018	LOBE	BAC KRIBI-CAMPO	2.8667	9.8833
CMQ5019	LOKOUNDJE	LOLORDORF	3.2333	10.7333
CMQ5038	MUNGO	MUNDAME	4.5667	9.5333
CMQ5040	NTEM	BAC DE NGOAZIK	2.1333	11.3
CMQ5044	LOM	BETARE OYA	5.9167	14.1333
CMQ5047	KIENKE	KRIBI SCIERIE	2.9333	9.9
CMQ5050	KADEI	BATOURI	4.4167	14.3167
GAQ0006	OGOOUE	BOOUE (LMNG)	-0.1025	11.9367
GAQ0015	OGOOUE	NDJOLE OPERATIONNEL	-0.455	10.4025
GAQ0028	IVINDO	MAKOKOU (LMNG)	0.5689	12.8611
GAQ0041	NGOUNIE	FOUGAMOU S H O (LMNG)	-1.2156	10.5908
GAQ0046	NGOUNIE	MOUILA VAL MARIE	-1.8869	11.0558
GHQ0045	NASIA	NASIA	10.15	-0.8
GHQ1005	VOLTA NOIRE	BUI AMONT	8.2833	-2.2333
GNQ0015	NIGER	FARANAH	10.03744	-10.7495
GNQ0016	NIGER	KOUROUSSA	10.65169	-9.87096
GNQ0018	NIGER	TIGUIBERY (Siguiri)	11.3545	-9.16459
GNQ0026	MILO	KANKAN	10.3833	-9.3
GNQ0030	NIANDAN	BARO	10.6167	-9.7
GNQ0034	NIANDAN	KISSIDOUYOU (NIANDAN SCIERIE)	9.25	-10.0167
GNQ0200	BADI	BAC DE BADI	10.2833	-13.4
GNQ0204	KONKOURE	PONT DE LINSAN	10.3	-12.4167
MLQ0009	NIGER	DIRE	16.27595	-3.395
MLQ0012	NIGER	KE MACINA	13.95831	-5.35896
MLQ0019	NIGER	KOULIKORO	12.85727	-7.55811
MLQ0022	NIGER	MOPTI	14.49605	-4.20127
MLQ0036	NIGER	TOSSAYE	16.9333	-0.5833
MLQ0052	DIKA	KARA	14.1667	-5.0167
MLQ0091	BANI	SOFARA	14.01393	-4.2429
MLQ0123	SENEGAL	GALOUGO	13.8333	-11.1333
MLQ0130	SENEGAL	BAFING MAKANA	12.55	-10.2667
MLQ0131	SENEGAL	SOUKOUTALI	13.2	-10.4167
MLQ0134	BAKOYE	OUALIA	13.6	-10.3833
MLQ0135	BAKOYE	TOUKOTO	13.45	-9.8833
MLQ0137	FALEME	FADOUYOU	12.5167	-11.3833
MLQ0145	BAOULE	SIRAMAKANA (Balenda)	13.5833	-9.8833
MLQ2003	NIGER	KENIEROBA	12.1	-8.3167
MLQ2007	SANKARANI	SELINGUE	11.5833	-8.1667
MLQ2008	BANI	DOUNA	13.21385	-5.90311
MLQ2064	SENEGAL	DAKA SAIDOU	11.95	-10.6167

MLQ2066	SENEGAL	DIBIA	13.2333	-10.8
MLQ2069	FALEME	GOURBASSY	13.4	-11.6333
MLQ2070	SENEGAL	KAYES	14.45	-11.45
NEQ2000	NIGER	NIAMEY	13.5016	2.105
NGQ0001	BENUE	MAKURDI	7.75	8.5333
NGQ0002	NIGER	ONITSHA	6.166667	6.75
NGQ2000	NIGER	LOKOJA	7.8	6.7667
NGQ2004	NIGER	IDAH	7.1	6.716667
SNQ2039	GAMBIE	KEDOUGOU	12.55	-12.1833
SNQ2045	GAMBIE	MAKO	12.8667	-12.35
SNQ2055	GAMBIE	SIMENTI	13.0333	-13.3
SNQ2060	GAMBIE	WASSADOU-AMONT	13.35	-13.3667
SNQ2062	GAMBIE	WASSADOU-AVAL	13.35	-13.3833
SNQ2063	SENEGAL	BAKEL	14.9	-12.45
SNQ2064	SENEGAL	DAGANA	16.5167	-15.5
SNQ2065	FALEME	KIDIRA UHEA	14.45466	-12.205
SNQ2066	SENEGAL	MATAM	15.65	-13.25
SNQ2067	DOUE	NGOUI	16.15	-13.9167
SNQ2068	SENEGAL	SALDE	16.16325	-13.8795
TDQ0004	CHARI	SARH (EX.FORT-ARCHAMBAULT)	9.15	18.4167
TDQ0009	CHARI	MAILAO	11.6	15.2833
TDQ0013	BAHR-SARA	MANDA	9.1833	18.2
TDQ0014	BAHR-SARA	MOISSALA	8.3333	17.7667
TDQ0036	LIM	OULI BANGALA	7.8333	15.8333
TDQ0041	PENDE	GORE	7.95	16.6167
TDQ0043	TANDJILE	TCHOA	9.3333	16.0833
TDQ2011	CHARI	BOUSSO	10.5	16.7167
TDQ5004	LOGONE	KATOA	10.8333	15.0833
TDQ5005	LOGONE	LAI (MISSION)	9.4	16.3
TDQ5006	LOGONE	LOGONE-GANA	11.55	15.15
TOQ0006	KARA	LAMA KARA 1	9.5333	1.1833
TOQ0037	SIO	KPEDJI	6.5317	1.0083
TOQ0042	MONO	CORREKOPE	7.8	1.3
TOQ0043	MONO	DOTAIKOPE	7.8167	1.2667
TOQ0046	MONO	TETETOU	7.0167	1.5333
TOQ0048	AMOU	AMOU OBLO	7.4	0.8667
TOQ0053	ANIE	PONT C F T	7.7333	1.2
TOQ0056	KOLOWARE	KOLOWARE	8.9667	1.2833
TOQ0057	NA	PARATAO	8.95	1.1833
TOQ0059	OGOUI	SIRKA	7.9167	1.3667

REFERENCES

Adeaga, O., Mahé, G., Dieulin, C., Elbaz-Poulichet, F., Rouche, N., Seide, J. L., & Servat, E., 2012. Rainfall-Runoff

- Simulation in Part of Lower Niger Basin. *Journal of Environmental Science and Engineering. B*, 1(6B), 812.
- Albergel, J., 1987. Sécheresse, désertification et ressources en eau de surface : application aux petits bassins du Burkina Faso. *The influence of climate change and climatic variability on the hydrologic regime and water resources*, 168, 355-365.
- Ali, A., Lebel, T., 2009. The Sahelian standardized rainfall index revisited. *Int. J. Climatol.* 29, 1705–1714. <https://doi.org/10.1002/joc.1832>
- Alsdorf, D., Beighley, E., Laraque, A., Lee, H., Tshimanga, R., O'Loughlin, F., Mahé, G., Dinga, B., Moukandi, G., Spencer, R.G.M., 2016. Opportunities for hydrologic research in the Congo Basin. *Rev. Geophys.* <https://doi.org/10.1002/2016RG000517>
- Amogu, O., Descroix, L., Yéro, K.S., Le Breton, E., Mamadou, I., Ali, A., Vischel, T., Bader, J.-C., Moussa, I.B., Gautier, E., Boubkraoui, S., Belleudy, P., 2010. Increasing River Flows in the Sahel? *Water* 2, 170–199. <https://doi.org/10.3390/w2020170>
- Auger, I.E., Lawrence, C.E., 1989. Algorithms for the optimal identification of segment neighborhoods. *Bull. Math. Biol.* 51, 39–54. <https://doi.org/10.1007/BF02458835>
- Badr, H.S., Dezfuli, A.K., Zaitchik, B.F., PETers-Lidard, C.D., 2016. Regionalizing Africa: Patterns of precipitation variability in observations and global climate models. *J. Clim.* 29, 9027–9043. <https://doi.org/10.1175/JCLI-D-16-0182.1>
- Biasutti, M., 2013. Forced Sahel rainfall trends in the CMIP5 archive. *J. Geophys. Res. Atmos.* 118, 1613–1623. <https://doi.org/10.1002/jgrd.50206>
- Boyer, J. F., Dieulin, C., Rouche, N., Cres, A., Servat, E., Paturel, J. E., Mahé, G., 2006. SIEREM: An environmental information system for water resources. *Climate Variability and Change—Hydrological Impacts*, S. Demuth et al., Eds., IAHS Publication 308, IAHS/AISH, 19–25
- Breiman, L., 2001. Random Forests. *Mach. Learn.* 45, 5–32. <https://doi.org/10.1023/A:1010933404324>
- Cappelaere, B., Descroix, L., Lebel, T., Boulain, N., Ramier, D., Laurent, J.P., Favreau, G., Boubkraoui, S., Boucher, M., Bouzou Moussa, I., Chaffard, V., Hiernaux, P., Issoufou, H.B.A., Le Breton, E., Mamadou, I., Nazoumou, Y., Oi, M., Ottlé, C., Quantin, G., 2009. The AMMA-CATCH experiment in the cultivated Sahelian area of south-west Niger - Investigating water cycle response to a fluctuating climate and changing environment. *J. Hydrol.* 375, 34–51. <https://doi.org/10.1016/j.jhydrol.2009.06.021>
- Charrad, M., Ghazzali, N., Boiteau, V., Niknafs, A., 2014. **NbClust**: An R Package for Determining the Relevant Number of Clusters in a Dataset. *J. Stat. Softw.* 61. <https://doi.org/10.18637/jss.v061.i06>
- Chiang, J.C.H., Friedman, A.R., 2012. Extratropical Cooling, Interhemispheric Thermal Gradients, and Tropical Climate Change. *Annu. Rev. Earth Planet. Sci.* 40, 383–412. <https://doi.org/10.1146/annurev-earth-042711-105545>
- D'amato, N., Lebel, T., 1998. On the characteristics of the rainfall events in the Sahel with a view to the analysis of climatic variability. *Int. J. Climatol.* 18, 955–974. [https://doi.org/10.1002/\(SICI\)1097-0088\(199807\)18:9<955::AID-JOC236>3.0.CO;2-6](https://doi.org/10.1002/(SICI)1097-0088(199807)18:9<955::AID-JOC236>3.0.CO;2-6)
- Dai, A., Lamb, P.J., Trenberth, K.E., Hulme, M., Jones, P.D., Xie, P., 2004. The recent Sahel drought is real. *Int. J. Climatol.* 24, 1323–1331. <https://doi.org/10.1002/joc.1083>
- Descroix, L., Mahé, G., Lebel, T., Favreau, G., Galle, S., Gautier, E., Olivry, J.C., Albergel, J., Amogu, O., Cappelaere, B., Dessouassi, R., Diedhiou, A., Le Breton, E., Mamadou, I., Sighomnou, D., 2009. Spatio-temporal variability of hydrological regimes around the boundaries between Sahelian and Sudanian areas of West Africa: A synthesis. *J. Hydrol.* 375, 90–102. <https://doi.org/10.1016/j.jhydrol.2008.12.012>
- Descroix, L., Moussa, I. B., Genthon, P., Sighomnou, D., Mahé, G., Mamadou, I., Vandervaere, J.-P., Gautier, E., Maiga, O. F., Rajot, J.-L., Abdou, M. M., Dessay, N., Ingatan, A., Noma, I., Yéro, K.S., Karambiri, H., Fensholt, R., Albergel, J., and Olivry, J.-C.: Impact of Drought and Land – Use Changes on Surface – Wa-ter Quality and Quantity:

- 697 The Sahelian Paradox, *Curr. Perspect. Contam. Hydrol. Water Res. Sustain.*, in: *Current Perspectives in Contaminant*
 698 *Hydrology and Water Resources Sustainability*, chap. 10, edited by: Bradley, P. M., 243–271, doi:10.5772/54536,
 699 2013
- 700
- 701 Dieppois, B., Diedhiou, A., Durand, A., Fournier, M., Massei, N., Sebag, D., Xue, Y., Fontaine, B., 2013. Quasi-
 702 decadal signals of Sahel rainfall and West African monsoon since the mid-twentieth century. *J. Geophys. Res.*
 703 *Atmos.* 118, 12587–12599. <https://doi.org/10.1002/2013JD019681>
- 704
- 705 Dieppois, B., Durand, A., Fournier, M., Diedhiou, A., Fontaine, B., Massei, N., Nouaceur, Z., Sebag, D., 2015. Low-
 706 frequency variability and zonal contrast in Sahel rainfall and Atlantic sea surface temperature teleconnections during
 707 the last century. *Theor. Appl. Climatol.* 121, 139–155. <https://doi.org/10.1007/s00704-014-1229-5>
- 708
- 709 Ekström, M., Jones, P.D., Fowler, H.J., Lenderink, G., Buishand, T. a., Conway, D., 2007. Regional climate model
 710 data used within the SWURVE project – 1: projected changes in seasonal patterns and estimation of PET.
 711 *Hydrol. Earth Syst. Sci.* 11, 1069–1083. <https://doi.org/10.5194/hess-11-1069-2007>
- 712
- 713 FAO: Sahel: situation météorologique et état des cultures, Rapport du 11 juin 2004,
 714 <http://www.fao.org/docrep/006/j2517f/j2517f00.HTM> (last access: October 2016), 2004.
- 715
- 716 Funk, C., Michaelsen, J., Marshall, M., 2012. Mapping recent decadal climate variations in
 717 precipitation and temperature across Eastern Africa and the Sahel. *Drought and Water Crises*,
 718 pp. 331–358. CRC Press.
- 719
- 720 Gal, L., Grippa, M., Hiernaux, P., Pons, L., Kergoat, L., 2017. The paradoxical evolution of runoff in the pastoral
 721 Sahel: Analysis of the hydrological changes over the Agoufou watershed (Mali) using the KINEROS-2 model. *Hydrol.*
 722 *Earth Syst. Sci.* 21, 4591–4613. <https://doi.org/10.5194/hess-21-4591-2017>
- 723
- 724 Gyau-Boakye, P., Schultz, G.A., 1994. Filling gaps in runoff time series in west africa. *Hydrol. Sci. J.* 39, 621–636.
 725 <https://doi.org/10.1080/02626669409492784>
- 726
- 727 Harris, I., Jones, P.D., Osborn, T.J., Lister, D.H., 2014. Updated high-resolution grids of monthly climatic
 728 observations - the CRU TS3.10 Dataset. *Int. J. Climatol.* 34, 623–642. <https://doi.org/10.1002/joc.3711>
- 729
- 730 Herrmann, S.M., Mohr, K.I., 2011. A continental-scale classification of rainfall seasonality regimes in Africa based on
 731 gridded precipitation and land surface temperature products. *J. Appl. Meteorol. Climatol.* 50, 2504–2513.
 732 <https://doi.org/10.1175/JAMC-D-11-024.1>
- 733
- 734 Hubert, P., Carbonnel, J.P., Chaouche, A., 1989. Segmentation des séries hydrométéorologiques - application à des
 735 séries de précipitations et de débits de l’afrique de l’ouest. *J. Hydrol.* 110, 349–367. [https://doi.org/10.1016/0022-1694\(89\)90197-2](https://doi.org/10.1016/0022-1694(89)90197-2)
- 736
- 737
- 738
- 739 Hulme, M., Doherty, R., Ngara, T., New, M., Lister, D., 2001. African climate change: 1900-2100. *Clim. Res.* 17, 145–
 740 168. <https://doi.org/10.3354/cr017145>
- 741
- 742 Hulme, M., Osborn, T.J., Johns, T.C., 1998. Precipitation sensitivity to global warming: Comparison of observations
 743 with HadCM2 simulations. *Geophys. Res. Lett.* 25, 3379. <https://doi.org/10.1029/98GL02562>
- 744
- 745 Ibrahim, B., Karambiri, H., Polcher, J., 2015. Hydrological Impacts of the Changes in Simulated Rainfall Fields on
 746 Nakanbe Basin in Burkina Faso. *Climate* 3, 442–458. <https://doi.org/10.3390/cli3030442>
- 747
- 748 IPCC, 2014. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of*
 749 *Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge
 750 Univ. Press 688. <https://doi.org/10.1017/CBO9781107415324.004>
- 751
- 752 Janicot, S., 1992. Spatiotemporal variability of West African rainfall. Part I: Regionalizations and Typings. *J. Clim.* 5,
 753 489–497. [https://doi.org/10.1175/1520-0442\(1992\)005<0489:SVOWAR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1992)005<0489:SVOWAR>2.0.CO;2)
- 754
- 755 Jones, P.D., 1994. Hemispheric surface air temperature variations: a reanalysis and an update to 1993. *J. Clim.* 7,
 756 1794–1802. [https://doi.org/10.1175/1520-0442\(1994\)007<1794:HSATVA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1994)007<1794:HSATVA>2.0.CO;2)

- 757 Jones, P.D., Moberg, A., 2003. Hemispheric and large-scale surface air temperature variations: An extensive revision
758 and an update to 2001. *J. Clim.* 16, 206–223. [https://doi.org/10.1175/1520-0442\(2003\)016<0206:HALSSA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0206:HALSSA>2.0.CO;2)
759
- 760 Jury, M.R., 2013. A return to wet conditions over Africa: 1995–2010. *Theor. Appl. Climatol.* 111, 471–481.
761 <https://doi.org/10.1007/s00704-012-0677-z>
762
- 763 Kalteh, A.M., Hjorth, P., 2009. Imputation of missing values in a precipitation–runoff process database. *Hydrol. Res.*
764 40, 420. <https://doi.org/10.2166/nh.2009.001>
765
- 766 Kendall, M., 1975. *Multivariate analysis*. Charles Griffin.
767
- 768 Killick, R., Eckley, I., 2014. changepoint: An R package for changepoint analysis. *Journal of statistical*
769 *software*, 58(3), 1–19.
770
- 771 Kim, J.W., Pachepsky, Y.A., 2010. Reconstructing missing daily precipitation data using regression trees and artificial
772 neural networks for SWAT streamflow simulation. *J. Hydrol.* 394, 305–314.
773 <https://doi.org/10.1016/j.jhydrol.2010.09.005>
774
- 775 Kling, H., Fuchs, M., Paulin, M., 2012. Runoff conditions in the upper Danube basin under an ensemble of climate
776 change scenarios. *J. Hydrol.* 424–425, 264–277. <https://doi.org/10.1016/j.jhydrol.2012.01.011>
777
- 778 Kundzewicz Z.W., Radziejewski M., 2006. Methodologies for trend detection. In: *Climate Variability and Change-*
779 *Hydrological Impacts (FRIEND)*. S. Demuth, A. Gustard, E. Planos, F. Scatena & E. Servat (Eds), IAHS Publ. 308,
780 538–550.
781
- 782 L'Hôte, Y., Dubreuil, P., Lerique, J., 1996. Carte des types de climats “en Afrique Noire à l’ouest du Congo”. Rappels
783 et extension aux régimes hydrologiques. In: *L’hydrologie Tropicale: Geoscience et Outil Pour le Développement*
784 (Mélanges à la mémoire de Jean Rodier. Paris, mai 1995), 55–65. *IAHS Publ.* no. 238.
785
- 786 L'Hôte, Y., Mahé, G., Somé, B., Triboulet, J.P., 2002. Analysis of a Sahelian annual rainfall index from 1896 to 2000;
787 the drought continues. *Hydrol. Sci. J.* 47, 563–572. <https://doi.org/10.1080/02626660209492960>
788
- 789 Le Barbé, L., Lebel, T., 1997. Rainfall climatology of the HAPEX-Sahel region during the years 1950–1990. *J. Hydrol.*
790 188–189, 43–73. [https://doi.org/10.1016/S0022-1694\(96\)03154-X](https://doi.org/10.1016/S0022-1694(96)03154-X)
791
- 792 Le Barbé, L., Lebel, T., Tapsoba, D., 2002. Rainfall variability in West Africa during the years 1950–90. *J. Clim.* 15,
793 187–202. [https://doi.org/10.1175/1520-0442\(2002\)015<0187:RVIWAD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0187:RVIWAD>2.0.CO;2)
794
- 795 Lebel, T., 2003. Seasonal cycle and interannual variability of the Sahelian rainfall at hydrological scales. *J. Geophys.*
796 *Res.* 108, 8389. <https://doi.org/10.1029/2001JD001580>
797
- 798 Lebel, T., Ali, A., 2009. Recent trends in the Central and Western Sahel rainfall regime (1990–2007). *J. Hydrol.* 375,
799 52–64. <https://doi.org/10.1016/j.jhydrol.2008.11.030>
800
- 801 Lehner, B., Liermann, C.R., Revenga, C., Vörösmarty, C., Fekete, B., Crouzet, P., Döll, P., Endejan, M., Frenken,
802 K., Magome, J., Nilsson, C., Robertson, J.C., Rödel, R., Sindorf, N., Wisser, D., 2011. High-resolution mapping of the
803 world’s reservoirs and dams for sustainable river-flow management. *Front. Ecol. Environ.*
804 <https://doi.org/10.1890/100125>
805
- 806 Liebmann, B., Dole, R.M., Jones, C., Bladé, I., Allured, D., 2010. Influence of choice of time period on global surface
807 temperature trends estimates. *Bull. Am. Meteorol. Soc.* 1485–1491. <https://doi.org/10.1175/2010BAMS3030.1>
808
- 809 Lienou, G., Mahe, G., Paturel, J.E., Servat, E., Sighomnou, D., Ekodeck, G.E., Dezetter, A., Dieulin, C., 2008.
810 Evolution des régimes hydrologiques en région équatoriale camerounaise: Un impact de la variabilité climatique en
811 Afrique équatoriale? *Hydrol. Sci. J.* 53, 789–801. <https://doi.org/10.1623/hysj.53.4.789>
812
- 813 Machiwal, D., Jha, M.K., 2006. Time series analysis of hydrologic data for water resources planning and
814 management: a review. *J. Hydrol. Hydromechanics* 54, 237–257.
815

- 817 Mahé, G., L'Hôte, Y., Olivry, J.C., Wotling, G., 2001. Trends and discontinuities in regional rainfall of West and
818 Central Africa: 1951–1989. *Hydrol. Sci. J.* 46, 211–226. <https://doi.org/10.1080/02626660109492817>
819
- 820 Mahé, G., Lienou, G., Descroix, L., Bamba, F., Paturel, J.E., Laraque, A., Meddi, M., Habaieb, H., Adeaga, O.,
821 Dieulin, C., Chahnez Kotti, F., Khomsi, K., 2013. The rivers of Africa: Witness of climate change and human impact
822 on the environment. *Hydrol. Process.* 27, 2105–2114. <https://doi.org/10.1002/hyp.9813>
823
- 824 Mahé, G., Paturel, J.E., 2009. 1896-2006 Sahelian annual rainfall variability and runoff increase of Sahelian Rivers.
825 *Comptes Rendus - Geosci.* 341, 538–546. <https://doi.org/10.1016/j.crte.2009.05.002>
826
- 827 Mahé, G., Paturel, J.E., Servat, E., Conway, D., Dezetter, A., 2005. The impact of land use change on soil water
828 holding capacity and river flow modelling in the Nakambe River, Burkina-Faso. *J. Hydrol.*
829 <https://doi.org/10.1016/j.jhydrol.2004.04.028>
830
- 831 Maidment, R.I., Allan, R.P., Black, E., 2015. Recent observed and simulated changes in precipitation over Africa.
832 *Geophys. Res. Lett.* 42, 8155–8164. <https://doi.org/10.1002/2015GL065765>
833
- 834 Mann, H.B., 1945. Nonparametric Tests Against Trend. *Econometrica* 13, 245. <https://doi.org/10.2307/1907187>
835
- 836 McCabe, G.J., Wolock, D.M., 2002. A step increase in streamflow in the conterminous United States. *Geophys. Res.*
837 *Lett.* 29, 38-1-38–4. <https://doi.org/10.1029/2002GL015999>
838
- 839 McCartney, M., Forkuor, G., Sood, A., Amisigo, B., Hattermann, F., Muthuwatta, L., 2012. The water resource
840 implications of changing climate in the Volta River Basin, IWMI Research Report. <https://doi.org/10.5337/2012.219>
841
- 842 Mitchell, T.D., Jones, P.D., 2005. An improved method of constructing a database of monthly climate observations
843 and associated high-resolution grids. *Int. J. Climatol.* 25, 693–712. <https://doi.org/10.1002/joc.1181>
844
- 845 Moalafhi, D.B., Sharma, A., Evans, J.P., 2017. Reconstructing hydro-climatological data using dynamical
846 downscaling of reanalysis products in data-sparse regions – Application to the Limpopo catchment in southern Africa.
847 *J. Hydrol. Reg. Stud.* 12, 378–395. <https://doi.org/10.1016/j.ejrh.2017.07.001>
848
- 849 Nicholson, S.E., 2013. The West African Sahel: A Review of Recent Studies on the Rainfall Regime and Its
850 Interannual Variability. *ISRN Meteorol.* 2013, 1–32. <https://doi.org/10.1155/2013/453521>
851
- 852 Nicholson, S.E., Some, B. and Kone, B. (2000). An Analysis of Recent Rainfall Conditions in West Africa, Including
853 the Rainy Seasons of the 1997 El Niño and the 1998 La Niña Years. *Journal of Climate*, 13, 2628-2640.
854 [https://dx.doi.org/10.1175/1520-0442\(2000\)013<2628:AAORRC>2.0.CO;2](https://dx.doi.org/10.1175/1520-0442(2000)013<2628:AAORRC>2.0.CO;2)
855
- 856 Nka, B.N., Oudin, L., Karambiri, H., Paturel, J.E., Ribstein, P., 2015. Trends in floods in West Africa: Analysis based
857 on 11 catchments in the region. *Hydrol. Earth Syst. Sci.* 19, 4707–4719. <https://doi.org/10.5194/hess-19-4707-2015>
858
- 859 Ogungbenro, S.B., Morakinyo, T.E., 2014. Rainfall distribution and change detection across climatic zones in Nigeria.
860 *Weather Clim. Extrem.* 5, 1–6. <https://doi.org/10.1016/j.wace.2014.10.002>
861
- 862 Ozer, P., Erpicum, M., Demarée, G., Vandiepenbeeck, M., 2003. The Sahelian drought may have ended during the
863 1990s. *Hydrol. Sci. J.* 48, 489–492. <https://doi.org/10.1623/hysj.48.3.489.45285>
864
- 865 Page, E.S., 1954. Continuous Inspection Schemes. *Biometrika* 41, 100. <https://doi.org/10.2307/2333009>
866
- 867 Peterson, T.C., Vose, R.S., 1997. An Overview of the Global Historical Climatology Network Temperature Database.
868 *Bull. Am. Meteorol. Soc.* 78, 2837–2849. [https://doi.org/10.1175/1520-0477\(1997\)078<2837:AOTGH>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<2837:AOTGH>2.0.CO;2)
869
- 870 Ringard, J., Dieppois, B., Rome, S., Diedhiou, A., Pellarin, T., Konaré, A., Diawara, A., Konaté, D., Dje, B.K.,
871 Katiellou, G.L., Seidou Sanda, I., Hassane, B., Vischel, T., Garuma, G.F., Mengistu, G., Camara, M., Diongue, A.,
872 Gaye, A.T., Descroix, L., 2016. The intensification of thermal extremes in west Africa. *Glob. Planet. Change* 139, 66–
873 77. <https://doi.org/10.1016/j.gloplacha.2015.12.009>
874
- 875 Roudier, P., Ducharne, A., Feyen, L., 2014. Climate change impacts on runoff in West Africa: A review. *Hydrol. Earth*
876 *Syst. Sci.* <https://doi.org/10.5194/hess-18-2789-2014>

- Rubin, D.B., 1987. Introduction, in: Multiple Imputation for Nonresponse in Surveys. pp. 1–26. <https://doi.org/10.1002/9780470316696.ch1>
- Schönbrodt, F.D., Perugini, M., 2013. At what sample size do correlations stabilize? J. Res. Pers. 47, 609–612. <https://doi.org/10.1016/j.jrp.2013.05.009>
- Stekhoven, D.J., Bühlmann, P., 2012. Missforest-Non-parametric missing value imputation for mixed-type data. Bioinformatics 28, 112–118. <https://doi.org/10.1093/bioinformatics/btr597>
- Stott, P.A., Gillett, N.P., Hegerl, G.C., Karoly, D.J., Stone, D.A., Zhang, X., Zwiers, F., 2010. Detection and attribution of climate change: A regional perspective. Wiley Interdiscip. Rev. Clim. Chang. 1, 192–211. <https://doi.org/10.1002/wcc.34>
- Suzuki, R., Shimodaira, H., 2006. Pvcust: An R package for assessing the uncertainty in hierarchical clustering. Bioinformatics 22, 1540–1542. <https://doi.org/10.1093/bioinformatics/btl117>
- Troyanskaya, O., Cantor, M., Sherlock, G., Brown, P., Hastie, T., Tibshirani, R., Botstein, D., Altman, R.B., 2001. Missing value estimation methods for DNA microarrays. Bioinformatics 17, 520–525. <https://doi.org/10.1093/bioinformatics/17.6.520>
- Tshimanga, R.M., Tshitenge, J.M., Kabuya, P., Alsdorf, D., Mahe, G., Kibukusa, G., Lukanda, V., 2016. A Regional Perceptive of Flood Forecasting and Disaster Management Systems for the Congo River Basin, in: Flood Forecasting: A Global Perspective. pp. 87–124. <https://doi.org/10.1016/B978-0-12-801884-2.00002-5>
- University of East Anglia Climatic Research Unit; Harris, I.C.; Jones, P.D. (2017): CRU TS4.00: Climatic Research Unit (CRU) Time-Series (TS) version 4.00 of high resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2015). Centre for Environmental Data Analysis, 25 May 2017
- Van Buuren, S., Oudshoorn, K. (1999). Flexible multivariate imputation by MICE. *Leiden, The Netherlands: TNO Prevention Center*.
- Von Storch, H., 1995. Misuses of Statistical Analysis in Climate. Anal. Clim. Var. Appl. Stat. Tech. 11–26. <https://doi.org/10.1007/978-3-662-03744-7>
- Webster, P.J., Magaña, V.O., Palmer, T.N., Shukla, J., Tomas, R.A., Yanai, M., Yasunari, T., 1998. Monsoons: Processes, predictability, and the prospects for prediction. J. Geophys. Res. Ocean. 103, 14451–14510. <https://doi.org/10.1029/97JC02719>
- Yue, S., Pilon, P., Phinney, B., Cavadias, G., 2002. The influence of autocorrelation on the ability to detect trend in hydrological series. Hydrol. Process. 16, 1807–1829. <https://doi.org/10.1002/hyp.1095>

HIGHLIGHTS

- The first imputed streamflow dataset for West and central Africa
- Good agreement between historical trends in streamflow and rainfall
- Partial modulations of post-1990s rainfall recovery by enhancing evapotranspiration
- Decadal modulations of Trends in hydroclimatic trends
- Homogenous zones of streamflow and precipitation variability

STUDY AREA

Large Dams in the Study Area

A) Percentage of missing data

B) Location of missing values

Clusters of streamflow variability

ACCEPTED MANUSCRIPT