



Afropectinariella (Vandeae, Orchidaceae), a new genus of the Angraecum alliance

Murielle Simo-Droissart, Bonaventure Sonké, Vincent Droissart, Tariq Stévart

► To cite this version:

Murielle Simo-Droissart, Bonaventure Sonké, Vincent Droissart, Tariq Stévart. Afropectinariella (Vandeae, Orchidaceae), a new genus of the Angraecum alliance. PhytoKeys, 2018, 96 (96), pp.79-86. 10.3897/phytokeys.96.23933 . hal-02186939

HAL Id: hal-02186939

<https://hal.umontpellier.fr/hal-02186939>

Submitted on 17 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Afropectinariella (Vandeae, Orchidaceae), a new genus of the *Angraecum* alliance

Murielle Simo-Droissart¹, Bonaventure Sonké^{1,2,3},
Vincent Droissart^{1,2,3,4}, Tariq Stévert^{2,3,5}

1 Plant Systematics and Ecology Laboratory, Higher Teachers' Training College, University of Yaoundé I, P.O. Box 047, Yaoundé, Cameroon **2** Missouri Botanical Garden, Africa and Madagascar Department, P.O. Box 299, St. Louis, Missouri 63166-0299, USA **3** Herbarium et Bibliothèque de Botanique africaine, C.P. 265, Université Libre de Bruxelles, Campus de la Plaine, Boulevard du Triomphe 1050, Brussels, Belgium **4** AMAP, IRD, CIRAD, CNRS, INRA, Université de Montpellier, Montpellier, France **5** Botanic Garden Meise, Domein van Bouchout, Nieuwelaan 38, B-1860 Meise, Belgium

Corresponding author: Murielle Simo-Droissart (murielle.simo@gmail.com)

Academic editor: Yaser Mutafchiev | Received 28 January 2018 | Accepted 19 March 2018 | Published 13 April 2018

Citation: Simo-Droissart M, Sonké B, Droissart V, Stévert T (2018) *Afropectinariella* (Vandeae, Orchidaceae), a new genus of the *Angraecum* alliance. *PhytoKeys* 96: 79–86. <https://doi.org/10.3897/phytokeys.96.23933>

Abstract

A recent phylogenetic study showed that species assigned to the newly recognised genus *Pectinariella* Szlach., Mytnik & Grochocka (previously treated as *Angraecum* Bory sect. *Pectinaria* Benth.) are polyphyletic, comprising a clade with species primarily in Madagascar and the Western Indian Ocean islands (including the type) and another non-sister clade whose members occur in continental Africa and the Gulf of Guinea islands. In order to render *Pectinariella* monophyletic, the five continental African species must therefore be removed. A new genus, *Afropectinariella* M.Simo & Stévert, is described and the following combinations are made: *Afropectinariella atlantica* (Stévert & Droissart) M.Simo & Stévert, *Afropectinariella doratophylla* (Summerh.) M.Simo & Stévert, *Afropectinariella gabonensis* (Summerh.) M.Simo & Stévert, *Afropectinariella pungens* (Schltr.) M.Simo & Stévert and *Afropectinariella subulata* (Lindl.) M.Simo & Stévert.

Résumé

Une analyse phylogénétique récente a montré que les espèces attribuées au nouveau genre *Pectinariella* Szlach., Mytnik & Grochocka (anciennement *Angraecum* Bory sect. *Pectinaria* Benth.) forment un groupe polyphylétique, comprenant un clade avec des espèces présentes principalement à Madagascar et dans les îles de l'ouest de l'Océan Indien (incluant le type) et un autre clade qui ne lui est pas apparenté et dont les espèces sont distribuées en Afrique continentale et dans les îles du Golfe de Guinée. Afin de rétablir la monophylie du genre *Pectinariella*, les cinq espèces d'Afrique continentale et des îles du Golfe de Guinée doivent donc en être exclues. Un nouveau genre, *Afropectinariella* M.Simo & Stévert, est décrit et les combinaisons suivantes sont proposées: *Afropectinariella atlantica* (Stévert & Droissart) M.Simo & Stévert, *Afropectinariella doratophylla* (Summerh.) M.Simo & Stévert, *Afropectinariella gabonensis* (Summerh.) M.Simo & Stévert, *Afropectinariella pungens* (Schltr.) M.Simo & Stévert et *Afropectinariella subulata* (Lindl.) M.Simo & Stévert.

Keywords

Angraecoid orchids, continental Africa, Malagasy and Indian Ocean islands, *Pectinaria*, phylogenetics, taxonomy

Introduction

Since its description, the delimitation of *Angraecum* Bory (1804) has been controversial (Garay 1973). Much importance has been ascribed to the structure of the pollinia within *Angraecum*, which has led to the recognition of spurious genera to encompass closely related species (Schlechter 1913). Based on vegetative and floral features, along with geographic distribution, a number of partial or comprehensive revisions of the genus have been proposed over the decades (see Finet 1907; Garay 1973; Perrier de la Bathie 1941; Schlechter 1918, 1925; Senghas 1986; Stewart et al. 2006; Summerhayes 1958). Garay (1973) was the last to provide a full taxonomic treatment for *Angraecum*. He recognised 19 sections, of which nine are endemic to Madagascar and the Western Indian Ocean islands, one to Madagascar, the Western Indian Ocean islands and Sri Lanka, two to continental Africa and the Gulf of Guinea islands and seven have representatives in both Africa and Madagascar.

With the advent of molecular techniques, relationships within *Angraecum* have been explored in more detail over the last 15 years. The studies of Carlsward et al. (2006) on the phylogenetics of leafless angraecoids and of Micheneau et al. (2008) on the biogeography of Mascarene angraecoid orchids have shown the polyphyly of *Angraecum* and of at least five of the 19 recognised sections. In fact, African species of *Angraecum* belong to a group clearly distinct from that of members of the genus in Madagascar and the Indian Ocean islands (Micheneau et al. 2008). While investigating the diversification of the genus in Madagascar, Andriananjamanantsoa et al. (2016) also confirmed the polyphyly of *Angraecum* s.l. and of all *Angraecum* sections with representatives in Madagascar, with the sole exception of section *Hadrangis* Schltr. Agreeing with Micheneau et al. (2008), Szlachetko et al. (2013) indicated that most of the sectional

arrangements within the genus are unnatural. Based on molecular and morphological data, Szlachetko et al. (2013) raised most of the sections, *sensu* Schlechter (1918) and Garay (1973), to the rank of genus, albeit with a different placement for some species. However, their sampling of DNA material lacked more than 3/4 of the species of *Angraecum* (only 53 of the 221 currently recognised species were included in their phylogeny) and several sections were unrepresented. In all, Szlachetko et al. (2013) recognised 18 genera that included species previously placed within *Angraecum*, 12 of which resulted from raising sections to the generic level, five of which were described as new genera and one of which involved resurrecting a genus previously placed in synonymy.

Szlachetko et al. (2013) established the genus *Pectinariella* Szlach., Mytnik & Grochocka to accommodate the species of *Angraecum* sect. *Pectinaria* Benth. *sensu* Garay (1973) since the sectional name was already occupied at the generic level in another family (*Pectinaria* Haworth, Apocynaceae). They circumscribed *Pectinariella* to include all members previously assigned to sect. *Pectinaria* from continental Africa and the Gulf of Guinea islands as well as those from the western Indian Ocean islands. However, their molecular study lacked material of the type species, *A. pectinatum* Thouars and included only five of the ten species of *Angraecum* sect. *Pectinaria*: four from Africa (viz. *A. doratophyllum* Summerh., *A. gabonense* Summerh., *A. pungens* Schltr. and *A. subulatum* Lindl.) and one from Madagascar (*A. dasycarpum* Schltr.).

Cribb (2014) suggested that part of *Angraecum* sect. *Pectinaria* might be separated from *Angraecum* *sensu stricto*, given that the generitype, *A. eburneum* Bory, was placed in the large Malagasy/Mascarene clade identified by Micheneau et al. (2008). The polyphyly of the section was confirmed by Simo-Droissart et al. (2013) based on sequence data from all five currently recognised species from Africa (the four mentioned above plus *A. atlanticum* Stévart & Droissart), along with three species from Madagascar and the Mascarene Islands. Confirming the results of Micheneau et al. (2008), Simo-Droissart et al. (2013) showed that the African members of *A. sect. Pectinaria* formed a well-supported clade that appears to be most closely related to *A. sect. Dolabrilifolia* Pfitzer and is not sister to the Malagasy/Mascarene clade, which includes the type species of the section.

Based on these results, Simo-Droissart et al. (2013) suggested that it would be necessary to remove the African species from *Angraecum* sect. *Pectinaria* in order to maintain its monophyly. The generic name *Pectinariella* proposed by Szlachetko et al. (2013) cannot, however, be applied to these African species since its type, *P. pectinata* (Thouars) Szlach., Mytnik & Grochocka, belongs to the clade comprising Malagasy/Mascarene taxa. Following the treatments of Szlachetko and Romowicz (2007) and Szlachetko et al. (2013), which treat the taxa generally placed in *A. sect. Dolabrilifolia* as a genus, *Dolabrilifolia* (Pfitzer) Szlach. & Romowicz and, since the African members of their polyphyletic genus *Pectinariella* are sister to those of *Dolabrilifolia*, they must also be recognised at the rank of genus.

Here we thus propose a new genus to accommodate the species from continental Africa and the Gulf of Guinea islands of *Pectinariella*.

Taxonomy

Afropectinariella M.Simo & Stévert, gen. nov.

urn:lsid:ipni.org:names:60476170-2

Type. *Afropectinariella doratophylla* (Summerh.) M.Simo & Stévert [= *Angraecum doratophyllum* Summerh.].

Etymology. The name of the genus is based on the geographic distribution of its five species, all of which occur in Africa and the generic name *Pectinariella* in which they were previously placed.

Diagnosis. *Afropectinariella* resembles the related genera *Dolabrifolia* and *Pectinariella* in having a sessile ovary, i.e. without a pedicel and with a very short peduncle that is hardly developed, but differs from *Dolabrifolia* by its elongate leaves that are never compressed laterally (vs. imbricate and laterally compressed) and from *Pectinariella* by its transversely oval lip that is wider than long (vs. the lip longer than wide) and its occurrence in continental Africa and Gulf of Guinea islands (vs. Madagascar and adjacent islands).

Description. Epiphytic herbs. Stem erect to pendent, branched and loosely leafy. Leaves fleshy, alternate and elongate, subulate or linear to oblong-ovate, apex acute to apiculate, petiole twisted. Inflorescences suberect and subsessile, in general 1-flowered, sometimes 2-flowered, borne along the stem or opposite a leaf, sheath brown. Flowers small, nearly sessile in the axils of the leaves, white and often scented. Floral bract one, amplexicaul, broadly ovate. Sepals and petals elliptic to obovate, apex subacute. Lip entire, ovate-triangular, ecallose, apex acute to apiculate. Spur ellipsoid or subcylindric, straight or slightly curved, sometimes hooked, with a wide mouth at the base of the lip, apex acute, often blunt or swollen in the apical half. Peduncle short, hardly developed. Pollinia 2, pyriform, often sessile on one or two viscidia, without a distinct stipe or shortly stipitate, rarely with a well-developed stipe.

Afropectinariella includes five species from continental Africa and the Gulf of Guinea Islands, one of which is endemic to São Tomé and Príncipe Islands. Four of these five species were placed in *Angraecum* sect. *Pectinaria* *sensu* Garay (1973) and, more recently, in *Pectinariella*, as originally circumscribed by Szlachetko et al. (2013). A detailed taxonomic treatment of these five species is presented in Simo-Droissart et al. (2014).

Afropectinariella atlantica (Stévert & Droissart) M.Simo & Stévert, comb. nov.

urn:lsid:ipni.org:names:60476171-2

Basionym. *Angraecum atlanticum* Stévert et al. (2010: 253). Type: Equatorial Guinea (Rio Muni). Monte Alén National Park: Engong inselberg, 5 km NW from Engong village, 01°37'25.8"N, 10°17'49.2"E, 1,100 m alt., 20 July 2001, Stévert 1020 (holotype: BRLU!; isotypes: MO!, K!, WAG!).

***Afropectinariella doratophylla* (Summerh.) M.Simo & Stévert, comb. nov.**
urn:lsid:ipni.org:names:60476172-2

Basionym. *Angraecum doratophyllum* Summerhayes (1937: 465). Type: São Tomé and Príncipe (São Tomé Island) Vanhulst (Macambrará): virgin forest, 1,050–1,200 m alt., 5 November 1932, *Exell* 254 (holotype: BM [BM000539167]! isotype: K [K000306497]!).

Homotypic synonym. *Pectinariella doratophylla* (Summerh.) Szlach., Mytnik & Grochocka

***Afropectinariella gabonensis* (Summerh.) M.Simo & Stévert, comb. nov.**
urn:lsid:ipni.org:names:60476173-2

Basionym. *Angraecum gabonense* Summerhayes (1954: 587). Type: Gabon. Upper Ngounié River, Nimalaba, N. E. of Les Echiras, 15 February 1927, *Le Testu* 6384 (holotype: K [K000306496]! isotypes: BM [BM000539175]! P [P00388248]!).

Homotypic synonym. *Pectinariella gabonense* (Summerh.) Szlach., Mytnik & Grochocka

***Afropectinariella pungens* (Schltr.) M.Simo & Stévert, comb. nov.**
urn:lsid:ipni.org:names:60476174-2

Basionym. *Angraecum pungens* Schlechter (1906: 163). Type: Cameroon. Man O'War Bay, auf Bäumen bei Kriegschiffhafen, 03°57'N, 09°14'E, 27 September 1905, *Schlechter* 15774 (holotype: B, destroyed; lectotype: K [K000107121]!, isolectotypes: BM [BM000539131]!, BR [BR000000642136]!).

Homotypic synonyms. *Angraecopsis pungens* (Schltr.) Rice; *Pectinariella pungens* (Schltr.) Szlach., Mytnik & Grochocka

Heterotypic synonyms. *Mystacydium arthrophyllum* Kraenzl.; *Angraecum arthrophyllum* (Kraenzl.) Schltr.

***Afropectinariella subulata* (Lindl.) M.Simo & Stévert, comb. nov.**
urn:lsid:ipni.org:names:77177891-1

Basionym. *Angraecum subulatum* Lindley (1837: 206). Type: Nigeria. Nun River: s.d., Barter 20125 (lectotype: K!, isolectotype: K!).

Homotypic synonyms. *Epidorkis subulata* (Lindl.) Kuntze; *Listrostachys subulata* (Lindl.) Rchb.f.; *Pectinariella subulata* (Lindl.) Szlach., Mytnik & Grochocka

Heterotypic synonyms. *Angraecum canaliculatum* De Wild.; *Listrostachys canaliculata* De Wild.

Discussion

Although we have removed species from continental Africa and the Gulf of Guinea islands from *Pectinariella*, its delimitation remains open to further discussion. Garay (1973) included four species from Madagascar and the Mascarenes within *Angraecum* section *Pectinaria* (i.e. *A. dasycarpum*, *A. hermannii* (Cordem.) Schltr., *A. humblotianum* Schltr. and *A. pectinatum*). He regarded *A. pterophyllum* H.Perrier as a synonym of *A. hermannii* and placed *A. panicifolium* H.Perrier in *Angraecum* sect. *Conchoglossum* Schltr. In addition to the four species recognised by Garay (1973) as comprising section *Pectinaria*, Stewart et al. (2006) also included *A. panicifolium*. Micheneau et al. (2008) suggested that *A. hermannii* might best be placed in *A. sect. Lemurangis* Garay based on its morphological characters, which was followed by Cribb and Hermans (2009) and Verlynde et al. (2016). Cribb and Hermans (op. cit.) recognised *A. pterophyllum* as a distinct species and thus included five species within the section. As they followed the treatment of Garay (1973), Szlachetko et al. (2013) placed only four species from Madagascar and the Mascarenes within their newly described genus *Pectinariella* (*P. dasycarpa* (Schltr.) Szlach., Mytnik & Grochocka, *P. hermannii* (Cordem.) Szlach., Mytnik & Grochocka, *P. humblotiana* (Schltr.) Szlach., Mytnik & Grochocka and *P. pectinata*). In fact, these authors made no mention of *Angraecum pterophyllum*, but they did propose the combination *Angraecoides panicifolia* (H. Perrier) Szlach., Mytnik & Grochocka. Verlynde et al. (2016) described two new species of *Pectinariella* (*P. edmundi* Bosser ex Verlynde & Ramandimbisoa and *P. scroticalcar* Verlynde & Ramandimbisoa) and made the combination *P. pterophylla* (H.Perrier) Verlynde & Ramandimbisoa, thus recognising six species of *Pectinariella* from Madagascar and the Mascarenes. They did not mention *Angraecum panicifolium* or *Angraecoides panicifolia*, even though Simo-Droissart et al. (2013) placed it close to *Pectinariella pectinata*. *Pectinariella* thus comprises seven species, four of which have been included in published molecular phylogenetic studies. These seven species are found in Madagascar or neighbouring islands, including the type species, which occurs in La Réunion, Mauritius and the Comoros Islands.

In the studies of Simo-Droissart et al. (2013, 2016), species belonging to *Dolabri-folia* and *Afropectinariella* form two distinct, well-supported clades (bootstrap support (BS) = 100% and posterior probability (PP) = 1). The sister relationship of these two groups is also well supported by our Bayesian analysis (PP 1). Species assigned to *Pectinariella* also form a clade well supported in both analyses and are not sister to the clade comprising species of our new genus, *Afropectinariella*. The shared presence of elongate leaves that are never compressed, an ovary lacking a stalk and whose peduncle is scarcely developed appear to represent a convergence in *Pectinariella* and *Afropectinariella*. However, members of *Afropectinariella* possess a transversely oval lip that is wider than long, while those of *Pectinariella* have a lip that is longer than wide. Unpublished molecular data indicate that *P. edmundi* and *P. scroticalcar* are not sister to *P. pectinata*, suggesting that the genus, as circumscribed by Verlynde et al. (2016), may not be monophyletic. The fact that some *Angraecum* species assigned to section *Pseudojumellea* Schltr. are placed with *P. pectinata* in molecular phylogenies (see Micheneau et al. 2008) indicates that further studies will be needed to assess the monophyly of *Pectinariella*.

Acknowledgements

We are grateful to the American Orchid Society for financial support provided for the first author's work in Cameroon. The U.S. National Science Foundation (Grant 1051547, T. Stévert as PI) also supported fieldwork, laboratory studies and herbarium visits. DNA sequencing was supported by the Belgian Fund for Scientific Research (F.R.S.-FNRS) through grants FRFC 2.4.577.10 and MIS 4.519.10. Fieldwork was also funded by DIVEAC, ECOFAC, the Central Africa Regional Program of the Environment (CARPE) and "Sud Expert Plantes" project (project #375) under the French Ministry of Foreign Affairs. We are indebted to the authorities of the University of Yaoundé I and especially the Higher Teachers' Training College who authorised us to build the orchid shadehouse in Yaoundé. We would like to thank Dr. Porter P. Lowry II for commenting and improving the English on an earlier version of the manuscript and the two reviewers for their constructive comments.

References

- Andriananjamanantsoa HN, Engberg S, Louis Jr EE, Brouillet L (2016) Diversification of *Angraecum* (Orchidaceae, Vandeae) in Madagascar: revised phylogeny reveals species accumulation through time rather than rapid radiation. *PloS ONE* 11: e0163194. <https://doi.org/10.1371/journal.pone.0163194>
- Bory de Saint Vincent JBGM (1804) *Voyage dans les quatre principales îles des mers d'Afrique*. F. Buisson, Paris, 409 pp.
- Carlsward BS, Whitten WM, Williams NH, Bytebier B (2006) Molecular phylogenetics of Vandeae (Orchidaceae) and the evolution of leaflessness. *American Journal of Botany* 93: 770–786. <https://doi.org/10.3732/ajb.93.5.770>
- Cribb PJ (2014) *Angraecum*. In: Pridgeon AM, Cribb PJ, Chase MW, Rasmussen FN (Eds) *Genera Orchidacearum Volume 6 Epidendroideae (Part 3)*. Oxford University Press, Oxford, 358–363.
- Cribb PJ, Hermans J (2009) Field guide to the orchids of Madagascar. Royal Botanic Gardens, Kew, 440 pp.
- Finet EA (1907) Classification et énumération des Orchidées africaines de la tribu des Sarcanthées, d'après les collections du Muséum de Paris. *Bulletin de la Société Botanique de France* 54: 1–65. [Pl 1–12] <https://doi.org/10.1080/00378941.1907.10835937>
- Garay LA (1973) Systematics of the genus *Angraecum* (Orchidaceae). *Kew Bulletin* 28: 495–516. <https://doi.org/10.2307/4108894>
- Lindley J (1837) Notes upon some genera and species of Orchidaceae in the collection formed by Mr. Drége, at the Cape of Good Hope. In: Hooker WJ (Ed.) *Companion to the Botanical Magazine*, vol. 2. James Ridgway, London, 200–210.
- Micheneau C, Carlsward BS, Fay MF, Bytebier B, Pailler T, Chase MW (2008) Phylogenetics and biogeography of Mascarene angraecoid orchids (Vandeae, Orchidaceae). *Molecular Phylogenetics and Evolution* 46: 908–922. <https://doi.org/10.1016/j.ympev.2007.12.001>
- Perrier de la Bathie H (1941) *Orchidées*, tome II. In: Humbert H (Ed.) *Flore de Madagascar*, 49 familie. Imprimerie Officielle, Tananarive, 202–321.

- Schlechter R (1906) Orchidaceae africanae IV. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 38: 144–165.
- Schlechter R (1913) Orchidacées de Madagascar. In: Heckel E (Ed.) Annales du musée colonial de Marseille. Musée colonial, Marseille, 148–202.
- Schlechter R (1918) Versuch einer natürlichen Neuordnung der afrikanschen angraekoiden Orchidaceen. Beihefte zum Botanischen Centralblatt 36: 62–181.
- Schlechter R (1925) Orchidaceae Perrieriana. Fedde, Repertorium specierum novarum regni vegetabilis, Beihefte 33: 306–360.
- Senghas K (1986) 15. Tribus Vandaeae, 48. Subtribus Angraecineae, 49. Subtribus Aerangidineae. In: Brieger FG, Maatsch R, Senghas K (Eds) Rudolf Schlechter, Die Orchideen. Paul Parey, Berlin, 975–1044.
- Simo-Droissart M, Micheneau C, Sonké B, Droissart V, Lowry II PP, Plunkett GM, Hardy OJ, Stévert T (2013) Morphometrics and molecular phylogenetics of the continental African species of *Angraecum* section *Pectinaria* (Orchidaceae). Plant Ecology and Evolution 146: 295–309. <https://doi.org/10.5091/plecevo.2013.900>
- Simo-Droissart M, Sonké B, Droissart V, Geerinck D, Micheneau C, Lowry II PP, Plunkett GM, Hardy OJ, Stévert T (2014) Taxonomic revision of the continental African species of *Angraecum* section *Pectinaria* (Orchidaceae). Systematic Botany 39: 725–739. <https://doi.org/10.1600/036364414X682184>
- Simo-Droissart M, Sonké B, Droissart V, Micheneau C, Lowry II PP, Hardy OJ, Plunkett GM, Stévert T (2016) Morphometrics and molecular phylogenetics of *Angraecum* section *Dolabrifolia* (Orchidaceae, Angraecinae). Plant Systematics and Evolution 302: 1027–1045. <https://doi.org/10.1007/s00606-016-1315-5>
- Stévert T, Cawoy V, Damen T, Droissart V (2010) Taxonomy of Atlantic Central African Orchids 1. A new species of *Angraecum* sect. *Pectinaria* (Benth.) Schltr. (Orchidaceae) from Gabon and Equatorial Guinea. Systematic Botany 35: 252–256. <https://doi.org/10.1600/036364410791638298>
- Stewart J, Hermans J, Campbell B (2006) Angraecoid Orchids: Species from the African Region. Timber Press, Portland, Oregon, 431 pp.
- Summerhayes VS (1937) African orchids: IX. Kew Bulletin 9: 457–476. <https://doi.org/10.2307/4107143>
- Summerhayes VS (1954) African orchids: XXII. Kew Bulletin 8: 575–591. <https://doi.org/10.2307/4117387>
- Summerhayes VS (1958) African Orchids: XXVI. Kew Bulletin 13: 257–281. <https://doi.org/10.2307/4109530>
- Szlachetko DL, Romowicz A (2007) *Dolabrifolia*, un nouveau genre d'orchidées de l'alliance *Angraecum*. Richardiana 7: 53–54.
- Szlachetko DL, Tukałło P, Mytnik-Ejsmont J, Grochocka E (2013) Reclassification of the *Angraecum*-alliance (Orchidaceae, Vandoideae) based on molecular and morphological data. Biodiversity Research and Conservation 29: 1–23. <https://doi.org/10.2478/biorc-2013-0004>
- Verlynde S, Ramandimbisoa B, Bosser J, Stévert T (2016) Contribution à l'étude des Orchidaceae de Madagascar. XXXVIII. Deux nouvelles espèces et une nouvelle combinaison pour le genre *Pectinariella* Szlach., Mytnik & Grochocka à Madagascar. Adansonia 38: 219–232. <https://doi.org/10.5252/a2016n2a6>