



HAL
open science

Improving plant allometry by fusing forest models and remote sensing

Fabian Jörg Fischer, Isabelle Maréchaux, Jérôme Chave

► **To cite this version:**

Fabian Jörg Fischer, Isabelle Maréchaux, Jérôme Chave. Improving plant allometry by fusing forest models and remote sensing. *New Phytologist*, 2019, 223 (3), pp.1159-1165. 10.1111/nph.15810 . hal-02191073

HAL Id: hal-02191073

<https://hal.umontpellier.fr/hal-02191073>

Submitted on 22 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tansley insights

Improving plant allometry by fusing forest models and remote sensing

Author for correspondence:

Jérôme Chave

Tel: +33 561 556 760

Email: jerome.chave@univ-tlse3.fr

Received: 29 December 2018

Accepted: 5 March 2019

Fabian Jörg Fischer¹ , Isabelle Maréchaux²  and Jérôme Chave¹ 

¹Laboratoire Evolution et Diversité Biologique, UMR5174, CNRS–Université Paul Sabatier–IRD, Bâtiment 4R1, 118 route de

Narbonne, F-31062 Toulouse Cedex 9, France; ²AMAP, INRA, IRD, CIRAD, CNRS, University of Montpellier, F-34000

Montpellier, France

Contents

Summary	1159	V. Challenges and perspectives	1163
I. Introduction	1159	Acknowledgements	1164
II. Tree allometry and transferability	1160	References	1164
III. Condensing the point cloud: allometry from space	1161		
IV. Bayesian merging of data in IBMs	1162		

New Phytologist (2019) **223**: 1159–1165
doi: 10.1111/nph.15810

Key words: allometry, Approximate Bayesian Computation, data-model fusion, forest model, LiDAR.

Summary

Allometry determines how tree shape and function scale with each other, related through size. Allometric relationships help scale processes from the individual to the global scale and constitute a core component of vegetation models. Allometric relationships have been expected to emerge from optimisation theory, yet this does not suitably predict empirical data. Here we argue that the fusion of high-resolution data, such as those derived from airborne laser scanning, with individual-based forest modelling offers insight into how plant size contributes to large-scale biogeochemical processes. We review the challenges in allometric scaling, how they can be tackled by advances in data-model fusion, and how individual-based models can serve as data integrators for dynamic global vegetation models.

I. Introduction

Forests provide important services to societies globally, sequestering large amounts of carbon, limiting erosion, regulating the water cycle, and providing a habitat for many species. Size, shape and function relationships among plants, or allometries, play a key role in understanding these services. Such relationships encapsulate ontogenetic, ecological and evolutionary constraints (Niklas, 1994) and have been widely used in quantitative tools to aid forest management. How much carbon is stored in the world's forests, for instance, is estimated from forest inventories using allometric models and then scaled up to regional and global scales, based on Earth observation data and modelling (Pan *et al.*, 2011).

Allometries also describe how metabolic functions, such as respiration rates and net primary production, scale with each other.

A theory has been developed to infer allometric scaling from evolutionary optimisation principles (Enquist & Niklas, 2002), but this theory does not account for recent advances in plant physiology (Rogers *et al.*, 2017; Scoffoni *et al.*, 2017), and its predictions do not match empirical data well (Muller-Landau *et al.*, 2006; Poorter *et al.*, 2012). Our ability to simulate the vegetation response to environmental change in Dynamic Global Vegetation Models (DGVMs) is, however, directly dependent on the robustness of these scaling relationships. Because DGVMs adopt a coarse-grained description of forests, allometries are often used to link fluxes and pools, but the results do not always correspond to empirical observations (Wolf *et al.*, 2011).

A great opportunity to bring processes and field information into a consistent modelling framework is offered by individual-based models (IBMs) of forest dynamics (DeAngelis & Grimm, 2014). In

IBMs, the forest ecosystem emerges from a combination of individual tree physiological and demographic processes at a scale that is relevant for forest resource management and ecological data assimilation, as in the FORMIND model (Rödig *et al.*, 2017). This approach can be extended to larger scales, either by informing DGVMs through IBMs (for example ED2; Medvigy *et al.*, 2009; LPJ-GUESS, Smith *et al.*, 2014) or by directly scaling them up (SEIB-DGVM, Sato *et al.*, 2007; FORMIND, Fischer *et al.*, 2016). Like DGVMs, forest IBMs often rely on empirical allometric models to predict tree shape and function but, during model calibration, information can also be gained about the allometric models themselves and the processes that shape them.

Proper calibration and validation of forest IBMs should be based on a variety of independent data sources, ranging from forest inventories to eddy-flux data, as recently exemplified with the TROLL model, a physiology-based and fully spatially explicit forest IBM (Maréchaux & Chave, 2017). A promising additional data source is provided by remote sensing. With its ability to generate detailed information over unprecedented scales and at locations that are otherwise hard to access (for example upper canopy layers, remote ecosystems), remote sensing has already had a transformative effect on vegetation modelling (Shugart *et al.*, 2015).

Here, we examine how a fusion of IBMs and airborne laser scanning (ALS), a remote-sensing technology that provides structural information at landscape scale, can be used to improve allometric relationships and better understand the processes that shape them. We argue that by linking forest IBMs with ALS, we can reduce unexplained variation in allometric estimates and extend these to large spatial scales, as displayed in




Fig. 1. This is an important step towards increased biological knowledge and improved predictions of ecosystem functioning. It is also a test case for the integration of future remote-sensing sources such as hyperspectral imaging or spaceborne laser scanning.

II. Tree allometry and transferability

When tree size, shape and function relate to each other across scales and environmental conditions, then the measurement of a single dimension can already provide a rough estimate of whole-tree attributes. This factor is particularly relevant when one quantity is more easily measured (for example trunk diameter) than the others (for example metabolic rate or biomass). Empirical studies provide a strong support for generalised allometric relationships. Whole-plant autotrophic respiration, for example, scales predictably with biomass across several orders of magnitude and from boreal to tropical forests (Mori *et al.*, 2010), and general patterns of allocation into aboveground vs belowground plant organs exist at individual and stand levels globally (Poorter *et al.*, 2012; Chen *et al.*, 2019). Similarly, allometries that relate trunk diameter to tree height, as shown in Fig. 2, can be found across forest types and have been used to supplement height measurements that are error prone and time consuming without optimised protocols (Sullivan *et al.*, 2018).

The notion that a model developed at one site may be valid elsewhere is called transferability (Wenger & Olden, 2012). An important application is exemplified by the calculation of carbon stocks from forest inventories. The product of wood density, trunk cross-sectional area, and tree height turns out to be a good

Fig. 1 Individual-based models (IBMs) as data assimilators, in interface with dynamic global vegetation models (DGVMs): application to allometric inference. Ground-based censuses and airborne laser scanning (ALS) provide complementary views on trees and forest canopies. Both techniques can be incorporated into the model–data fusion cycle, as formalised by Approximate Bayesian Computation (ABC). Increasingly diverse data can therefore be used to improve model representation and allometric parameter inference. Such improvement can be a benefit to DGVMs, whose simulations typically reach larger extents than IBMs, but which are currently run at coarse resolution, preventing them from making direct comparisons with data provided at finer spatial resolution.




Fig. 2 Empirical allometric relations between tree height and trunk diameter (DBH). Michaelis–Menten type allometric models were fitted with nonlinear least squares and a heteroscedastic error structure at six sites, typical of tropical forests, as follows: (a) Ulu Ulu National Park, Brunei (4.54°N, 115.15°E); (b) Parque Estadual Cristalino, Mato Grosso, Brazil (9.06°S, 55.94°W); (c) Grebo National Forest, southeast Liberia (5.4°N, 7.62°W); (d) Nouragues Ecological Research Station, French Guiana (4.09°N, 52.67°W); (e) Dja Faunal Reserve, Cameroon (1.89°S, 13.22°E); (f) Tambopata National Reserve, Peru (12.84S, 69.29W). Data are from Sullivan *et al.* (2018), and metadata can be accessed on the forestplots.net data portal.

predictor of tree biomass obtained from destructive harvesting (Chave *et al.*, 2014). This holds true across a wide range of values for the predictor variables and broad bioclimatic gradients, from dry forest woodlands to tropical rainforests. Recent work based on an extensive destructive harvest experiment in African tropical forests suggests that relatively simple biomass models are transferable (Fayolle *et al.*, 2018), and could therefore be useful in biomass assessments across the tropics.

However, in most cases, allometries are influenced by environmental factors, both abiotic and biotic, and are not easily transferable. The scaling of tree height with trunk diameter, for example, depends on bioclimatic constraints (Lines *et al.*, 2012; Olson *et al.*, 2018), and tree growth is shaped by interactions with other trees (Coomes *et al.*, 2011; Jucker *et al.*, 2015). Furthermore, allometries typically have a multiplicative error structure. Residual standard deviations for predictions translate into large absolute errors for the biggest individuals and result in inflated uncertainty in the predicted variables.

To quantify variation in scaling of tree shape, remote sensing offers new perspectives. Terrestrial laser scanning (TLS), for example, provides accurate estimates of tree dimensions without requiring destructive harvesting (Momo Takoudjou *et al.*, 2017). It therefore holds great potential for exploring geometric scaling properties in forest trees and their dependence on environmental conditions (Disney, 2019).

III. Condensing the point cloud: allometry from space

Where TLS is a type of remote sensing ‘from the ground’, airborne LiDAR scanning extends the 3D-mapping capacity of forest and tree structure to the landscape scale. The technology and its application to forest scanning have been developed for over 3 decades (Schreier *et al.*, 1985; Nelson *et al.*, 1988), and studies now commonly cover several 1000 hectares of forest at high point densities, that is high resolution. As a result, individual tree shapes can be measured in open woodlands, allowing researchers to monitor the growth and death of individual plants (Levick & Asner, 2013; Duncanson & Dubayah, 2018). Even more impressively, clustering algorithms have been developed to segment ALS point clouds into individual tree crowns in closed-canopy forests (Ferraz *et al.*, 2016). As tree trunk diameter was recently found to be correlated with the product of tree height and crown size, the segmented crowns can then be used to estimate ground-based measurements (Jucker *et al.*, 2017); this technology is being increasingly used in routine forest monitoring programmes.

Tree-delineation from ALS is not without its problems, however. Trees often have irregular crowns, they may partly overlap, and the sharp light attenuation within dense canopy means that understorey trees are sparsely scanned, rendering the direct retrieval of tree dimensions difficult. IBMs such as TROLL (Maréchaux & Chave, 2017) offer an indirect, yet powerful alternative.

The spatially explicit rendering of treefalls and the competition for light resources introduce ecological constraints on the simulated forest structure, limiting tree density and dimensions across size classes. Instead of translating point clouds back into individual tree dimensions, we can create better fits between virtual and empirical canopies by adjusting vital rates and allometric parameters that can therefore be derived from mechanistic principles – even for trees that are difficult to observe directly from ALS. As TROLL’s virtual canopies have a high spatial resolution (m^3), they compare naturally to ALS data and a few statistics are often suffice to link them. For example, Fig. 3 shows the match between top-canopy height obtained by ALS and a TROLL-based reconstruction. In the future it would be critical to extend this approach to other data sources, including TLS and spaceborne missions. Examples are the spaceborne laser scanner GEDI, a LiDAR now on board the International Space Station, and the BIOMASS synthetic aperture radar satellite, scheduled for launch in 2022, that will both provide a radically new view of the world’s forests.

Because vegetation models and remote sensing have long proven mutually informative (Sellers *et al.*, 1997), the available approaches for data-model fusion have been well tested. Possibilities include the derivation of tree-level data from ALS for model parameterisation, the comparison of outputs with observed canopies for model validation (Seidl *et al.*, 2012, Fig. 3), and so-called model inversion, in which models are run with a wide range of parameter combinations and systematically compared with remotely derived metrics (Fig. 1). A hybrid between these approaches – partly inverse modelling, partly initialisation – was developed early on and has recently been applied to derive biomass maps across Amazonia using spaceborne LiDAR (Hurt *et al.*, 2004; Rödig *et al.*, 2017). Moreover, when models




Fig. 3 Comparison of canopies sensed by airborne laser scanning (ALS) and simulated by individual-based models (IBMs). Shown are canopy-height models (height in m across 12 ha, 1-m² resolution; (a, b), and the corresponding height distributions (c, d). (a, c) Canopy-height model derived from an ALS campaign at the Nouragues field station, French Guiana, in 2012. (b, d) Canopy-height model as constructed by TROLL, an IBM of forest growth. Input information are a tree inventory, and allometries predicting tree height and crown dimensions from trunk diameter accounting for individual variation around the allometric trend.

provide realistic representations of forest structure, virtual ALS data can be produced and tested before using empirically observed canopies (Fassnacht *et al.*, 2018; Knapp *et al.*, 2018).

IV. Bayesian merging of data in IBMs

One efficient way to merge data and models is offered by Bayesian approaches such as Approximate Bayesian Computation (ABC), a widespread method in biological and ecological applications (Beaumont, 2010; Hartig *et al.*, 2011). Fig. 4 illustrates the inference of crown allometry parameters based on ABC. In qualitative terms, the approach is as follows: large numbers of simulations are performed with variations in crown allometry parameters (the prior in Bayesian statistics), the resulting virtual canopies are then compared with an empirically observed canopy (through statistics such as canopy height; Fig. 3) and, finally, the parameter values of the best-performing simulations are selected (the posterior). Inference on tree allometries is therefore turned into a parameter optimisation problem, and uncertainty around the parameter estimate reflects how informative is the data regarding a

particular allometry. In the example given in Fig. 4, the inference is considerably improved by using ALS data in addition to ground data, providing more precise estimates for allometry parameters across diameter-size classes.

When harmonising high-dimensional data, as obtained from ALS and IBMs such as TROLL, some issues emerge. Inferences can be markedly different, depending on how virtual and empirical canopies are compared, and dimension reduction and cross-validation techniques are needed to find an appropriate set of statistics (Csilléry *et al.*, 2012; Nunes & Prangle, 2015). But even when summary statistics are well chosen, a pattern (for example a virtual canopy) can be the result of several parameter combinations or ways to represent processes (for example allometries). In this case, inference methods such as ABC are not well posed. This type of uncertainty, usually referred to as 'equifinality' (Luo *et al.*, 2009), cannot always be avoided, but it can be mitigated. Additional data sources can help to narrow down the parameter space (Fig. 4). Furthermore, it is desirable to implement mechanistic models over statistical ones, because mechanistic simulations are restricted to a generally smaller universe of possibilities. They therefore




Fig. 4 Crown radius allometries inferred by Approximate Bayesian Computation (ABC) in TROLL. (a) Shape of the crown-radius allometry included in the TROLL model. Two posterior distributions of crown-radius allometry are shown, one constrained by ground data only (orange), the other also constrained by data derived from airborne laser scanning (ALS, red). The thick lines represent the posterior mode, the coloured areas the 70% highest-density intervals, and the dotted lines the extent of the prior distribution. (b, c) Cuts through the allometric distribution at 0.5 m in trunk diameter (DBH), for both simulations, with priors indicated by dotted lines. The addition of ALS data in ABC inference considerably narrows down the crown allometry parameters. The inference is based on 20 000 simulations, with a posterior composed of the best 200 simulations. Summary statistics included tree diameter-size distributions and ALS-derived canopy-height distributions. The overlap between simulated and empirical distributions was quantified and, to determine the posterior, we used the rejection scheme implemented in the R package *ABC* (Csilléry *et al.*, 2012) with an acceptance rate of 1%.

complement machine-learning techniques that are increasingly popular across science, including forest modelling, but are especially prone to the equifinality problem.

Another approach to narrow down the parameter space for models such as TROLL is the construction of an initial canopy state whose spatial arrangement is consistent with both the mechanistic principles of TROLL and the ALS-derived canopy structure. One method to produce such an initial state consists in sequentially assigning trees to spatial positions such that they receive enough light and that their size matches ALS observations. A space-filling rule is then iterated until all available space in the scene has been filled by trees (Taubert *et al.*, 2015). This initialisation can be useful to explore the range of validity of forest structure parameters (canopy gaps, crown exposure) and, therefore, yields both priors for the IBM and an evaluation of summary statistics. The IBM can then use this information to focus on ecological dynamics and provide distributions for tree trunks, crown dimensions and heights that represent a predictive check on ecological inferences and a new prior for the parameterisation of DGVMs (Fig. 1).

V. Challenges and perspectives

In this paper we argue that the explicit merging of plant allometry, forest observations, and individual-based modelling contributes to a unified vision of forest ecology. A fully spatially explicit IBM, when used for Bayesian data-model fusion, can inform quantities such as crown size and shape that are difficult to measure in dense canopies, but to which spatially explicit models are highly sensitive. The approach also helps to gain an understanding of ecological

processes, as it captures the fine-grained structure of forest canopies. It could, therefore, better explain tree regeneration and simulate the dynamics of nontree life forms, including lianas and epiphytes or even canopy-dwelling animals. This challenge is one of the greatest in biodiversity research today (Singer *et al.*, 2016). Further ecological insights can be gained regarding submodels, such as the tree growth equations implemented in gap models (Shugart *et al.*, 2018), or the autotrophic respiration equation (Atkin *et al.*, 2015). The obtained information could then constrain the parameters of physiological models that are usually prescribed in DGVMs (but see Wang *et al.*, 2017), and offer a direct benchmark of upscaling simplifications of canopy structure, such as the perfect plasticity approximation (Purves *et al.*, 2008).

Where the focus of DGVMs has traditionally been on satellite data, forest IBMs have instead been developed for and from ground inventories, with trunk diameters and their growth the main predictors of all simulated ecological processes. The remote-sensing revolution calls, however, for a new paradigm in forest modelling, including new data sets, and new approaches to model building. This mirrors the larger change of direction in global forest research in which remotely sensed metrics are increasingly used to predict ground metrics (Jucker *et al.*, 2017). This also represents a timely challenge because spaceborne missions such as GEDI and BIOMASS will acquire global forest structure datasets, but it is likely that a correct interpretation of these datasets will require an explicit linkage with models (Fisher *et al.*, 2018). Model upscaling raises the question of spatial model transferability. It is crucial to test this by validating the model at places where it has not been calibrated.

The assimilation of global remote-sensing data will be greatly helped by recent advances in computing technology that have shifted the limits of what forest extent can be simulated at tree level (Shugart *et al.*, 2015). For calibration, which requires 1000s of simulations for data-model fusion, computational cost can be reduced by classic assimilation techniques (Hurtt *et al.*, 2004). Future increases in computational power and emulators will further speed up inference (Fer *et al.*, 2018), turning IBMs into Bayesian data integrators that create a common vision of forest functioning and structure and the allometric relationships that link both across scales.

Complementary to this effort is the need to explain allometries from evolutionary optimisation arguments, for plant form (Enquist & Niklas, 2002), plant function (Wolf *et al.*, 2016), and forest structure (Farrion *et al.*, 2016). A better fundamental knowledge on allometric relationships can only improve our confidence in the parameters and simplify model calibration. This can only be achieved if theory is consistent with the known constraints of plant physiology. In return, data-model fusion, as explained here, provides a strong validation for theory.

Acknowledgements

This work has benefitted from 'Investissement d'Avenir' grants managed by Agence Nationale de la Recherche (CEBA, ref. ANR-10-LABX-25-01; ANAEE-France: ANR-11-INBS-0001, TULIP: ANR-10-LABX-0041), from CNES, and from ESA (CCI-Biomass). This work was granted access to the HPC resources of CALMIP supercomputing centre in Toulouse under allocation p17013.

ORCID

Jérôme Chave  <https://orcid.org/0000-0002-7766-1347>
 Fabian Jörg Fischer  <https://orcid.org/0000-0003-2325-9886>
 Isabelle Maréchaux  <https://orcid.org/0000-0002-5401-0197>

References

- Atkin OK, Bloomfield KJ, Reich PB, Tjoelker MG, Asner GP, Bonal D, Bönisch G, Bradford MG, Cernusak LA, Cosio EG *et al.* 2015. Global variability in leaf respiration in relation to climate, plant functional types and leaf traits. *New Phytologist* 206: 614–636.
- Beaumont MA. 2010. Approximate Bayesian computation in evolution and ecology. *Annual Review of Ecology, Evolution, and Systematics* 41: 379–406.
- Chave J, Réjou-Méchain M, Búrquez A, Chidumayo E, Colgan MS, Delitti WBC, Duque A, Eid T, Fearnside PM, Goodman RC *et al.* 2014. Improved allometric models to estimate the aboveground biomass of tropical trees. *Global Change Biology* 20: 3177–3190.
- Chen G, Hobbie SE, Reich PB, Yang Y, Robinson D. 2019. Allometry of fine roots in forest ecosystems. *Ecology Letters* 22: 322–331.
- Coomes DA, Lines ER, Allen RB. 2011. Moving on from Metabolic Scaling Theory: hierarchical models of tree growth and asymmetric competition for light. *Journal of Ecology* 99: 748–756.
- Csilléry K, François O, Blum MGB. 2012. Abc: an R package for approximate Bayesian computation (ABC). *Methods in Ecology and Evolution* 3: 475–479.
- DeAngelis DL, Grimm V. 2014. Individual-based models in ecology after four decades. *F1000Prime Reports* 6: 39.
- Disney M. 2019. Terrestrial LiDAR: a three-dimensional revolution in how we look at trees. *New Phytologist* 222: 1736–1741.
- Duncanson L, Dubayah R. 2018. Monitoring individual tree-based change with airborne lidar. *Ecology and Evolution* 8: 5079–5089.
- Enquist BJ, Niklas KJ. 2002. Global allocation rules for patterns of biomass partitioning in seed plants. *Science* 295: 1517–1520.
- Farrion CE, Bohlman SA, Hubbell S, Pacala SW. 2016. Dominance of the suppressed: power-law size structure in tropical forests. *Science* 351: 155–157.
- Fassnacht FE, Latifi H, Hartig F. 2018. Using synthetic data to evaluate the benefits of large field plots for forest biomass estimation with LiDAR. *Remote Sensing of Environment* 213: 115–128.
- Fayolle A, Ngomanda A, Mbasi M, Barbier N, Bocko Y, Boyemba F, Couteron P, Fonton N, Kamdem N, Katembo J *et al.* 2018. A regional allometry for the Congo basin forests based on the largest ever destructive sampling. *Forest Ecology and Management* 430: 228–240.
- Fer I, Kelly R, Moorcroft PR, Richardson AD, Cowdery EM, Dietze MC. 2018. Linking big models to big data: efficient ecosystem model calibration through Bayesian model emulation. *Biogeosciences* 15: 5801–5830.
- Ferraz A, Saatchi S, Mallet C, Meyer V. 2016. Lidar detection of individual tree size in tropical forests. *Remote Sensing of Environment* 183: 318–333.
- Fisher RA, Koven CD, Anderegg WR, Christoffersen BO, Dietze MC, Farrion CE, Holm JA, Hurtt GC, Knox RG, Lawrence PJ *et al.* 2018. Vegetation demographics in Earth System Models: a review of progress and priorities. *Global Change Biology* 24: 35–54.
- Hartig F, Calabrese JM, Reineking B, Wiegand T, Huth A. 2011. Statistical inference for stochastic simulation models—theory and application. *Ecology Letters* 14: 816–827.
- Fischer R, Bohn F, Dantas de Paula M, Dislich C, Groeneveld J, Gutiérrez AG, Kazmierczak M, Knapp N, Lehmann S, Paulick S *et al.* 2016. Lessons learned from applying a forest gap model to understand ecosystem and carbon dynamics of complex tropical forests. *Ecological Modelling* 326: 124–133.
- Hurtt GC, Dubayah R, Drake J, Moorcroft PR, Pacala SW, Blair JB, Fearon MG. 2004. Beyond potential vegetation: combining lidar data and a height-structured model for carbon studies. *Ecological Applications* 14: 873–883.
- Jucker T, Bouriaud O, Coomes DA. 2015. Crown plasticity enables trees to optimize canopy packing in mixed-species forests. *Functional Ecology* 29: 1078–1086.
- Jucker T, Caspersen J, Chave J, Antin C, Barbier N, Bongers F, Dalponte M, van Ewijk KY, Forrester DI, Haeni M *et al.* 2017. Allometric equations for integrating remote sensing imagery into forest monitoring programmes. *Global Change Biology* 23: 177–190.
- Knapp N, Fischer R, Huth A. 2018. Linking lidar and forest modeling to assess biomass estimation across scales and disturbance states. *Remote Sensing of Environment* 205: 199–209.
- Levick SR, Asner GP. 2013. The rate and spatial pattern of treefall in a savanna landscape. *Biological Conservation* 157: 121–127.
- Lines ER, Zavala MA, Purves DW, Coomes DA. 2012. Predictable changes in aboveground allometry of trees along gradients of temperature, aridity and competition. *Global Ecology and Biogeography* 21: 1017–1028.
- Luo Y, Weng E, Wu X, Gao C, Zhou X, Zhang L. 2009. Parameter identifiability, constraint, and equifinality in data assimilation with ecosystem models. *Ecological Applications* 19: 571–574.
- Maréchaux I, Chave J. 2017. An individual-based forest model to jointly simulate carbon and tree diversity in Amazonia: description and applications. *Ecological Monographs* 87: 632–664.
- Medvigy D, Wofsy SC, Munger JW, Hollinger DY, Moorcroft PR. 2009. Mechanistic scaling of ecosystem function and dynamics in space and time: ecosystem Demography model version 2. *Journal of Geophysical Research: Biogeosciences* 114: G01002.
- Momo Takoudjou S, Ploton P, Sonké B, Hackenberg J, Griffon S, Coligny F, Kamdem NG, Libalah M, Mofack GII, Le Mogueudé G *et al.* 2017. Using terrestrial laser scanning data to estimate large tropical trees biomass and calibrate allometric models: a comparison with traditional destructive approach. *Methods in Ecology and Evolution* 9: 905–916.
- Mori S, Yamaji K, Ishida A, Prokushkin SG, Masyagina OV, Hagihara A, Rafiqul Hoque ATM, Suwa R, Osawa A, Nishizono T *et al.* 2010. Mixed-power scaling of whole-plant respiration from seedlings to giant trees. *Proceedings of the National Academy of Sciences, USA* 107: 1447–1451.

- Muller-Landau HC, Condit RS, Chave J, Thomas SC, Bohlman SA, Bunyavejchewin S, Davies S, Foster R, Gunatilleke S, Gunatilleke N *et al.* 2006. Testing metabolic ecology theory for allometric scaling of tree size, growth and mortality in tropical forests. *Ecology Letters* 9: 575–588.
- Nelson R, Krabill W, Tonelli J. 1988. Estimating forest biomass and volume using airborne laser data. *Remote Sensing of Environment* 24: 247–267.
- Niklas KJ. 1994. Morphological evolution through complex domains of fitness. *Proceedings of the National Academy of Sciences, USA* 91: 6772–6779.
- Nunes MA, Prangle D. 2015. abctools: an R package for tuning approximate Bayesian computation analyses. *R Journal* 7: 1–16.
- Olson ME, Soriano D, Rosell JA, Anfodillo T, Donoghue MJ, Edwards EJ, León-Gómez C, Dawson T, Camarero Martínez JJ, Castorena M *et al.* 2018. Plant height and hydraulic vulnerability to drought and cold. *Proceedings of the National Academy of Sciences, USA* 115: 7551 LP-7556.
- Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Lewis SL, Canadell JG *et al.* 2011. A large and persistent carbon sink in the world's forests. *Science* 333: 988 LP-993.
- Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P, Mommer L. 2012. Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist* 193: 30–50.
- Purves DW, Lichstein JW, Strigul N, Pacala SW. 2008. Predicting and understanding forest dynamics using a simple tractable model. *Proceedings of the National Academy of Sciences, USA* 105: 17018–17022.
- Rödig E, Cuntz M, Heinke J, Rammig A, Huth A. 2017. Spatial heterogeneity of biomass and forest structure of the Amazon rain forest: linking remote sensing, forest modelling and field inventory. *Global Ecology and Biogeography* 26: 1292–1302.
- Rogers A, Medlyn BE, Dukes JS, Bonan G, von Caemmerer S, Dietze MC, Kattge J, Leakey ADB, Mercado LM, Prentice IC. 2017. A roadmap for improving the representation of photosynthesis in Earth system models. *New Phytologist* 213: 22–42.
- Sato H, Itoh A, Kohyama T. 2007. SEIB-DGVM: a new Dynamic Global Vegetation Model using a spatially explicit individual-based approach. *Ecological Modelling* 200: 279–307.
- Schreier H, Loughheed J, Tucker C, Leckie D. 1985. Automated measurements of terrain reflection and height variations using an airborne infrared laser system. *International Journal of Remote Sensing* 6: 101–446.
- Scoffoni C, Albuquerque C, Brodersen CR, Townes SV, John GP, Cochard H, Buckley TN, McElrone AJ, Sack L. 2017. Leaf vein xylem conduit diameter influences susceptibility to embolism and hydraulic decline. *New Phytologist* 213: 1076–1092.
- Seidl R, Spies TA, Rammer W, Steel EA, Pabst RJ, Olsen K. 2012. Multi-scale drivers of spatial variation in old-growth forest carbon density disentangled with Lidar and an individual-based landscape model. *Ecosystems* 15: 1321–1335.
- Sellers PJ, Dickinson RE, Randall DA, Betts AK, Hall FG, Berry JA, Collatz GJ, Denning AS, Mooney HA, Nobre CA *et al.* 1997. Modeling the exchanges of energy, water, and carbon between continents and the atmosphere. *Science* 275: 502–509.
- Shugart HH, Asner GP, Fischer R, Huth A, Knapp N, Le Toan T, Shuman JK. 2015. Computer and remote-sensing infrastructure to enhance large-scale testing of individual-based forest models. *Frontiers in Ecology and the Environment* 13: 503–511.
- Shugart HH, Wang B, Fischer R, Ma J, Fang J, Yan X, Huth A, Armstrong AH. 2018. Gap models and their individual-based relatives in the assessment of the consequences of global change. *Environmental Research Letters* 13: 033001.
- Singer A, Johst K, Banitz T, Fowler MS, Groeneveld J, Gutiérrez AG, Hartig F, Krug RM, Liess M, Matlack G *et al.* 2016. Community dynamics under environmental change: how can next generation mechanistic models improve projections of species distributions? *Ecological Modelling* 326: 63–74.
- Smith B, Wärlind D, Arneth A, Hickler T, Leadley P, Siltberg J, Zaehle S. 2014. Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences* 11: 2027–2054.
- Sullivan MJP, Lewis SL, Hubau W, Qie L, Baker TR, Banin LF, Chave J, Cuni-Sanchez A, Feldpausch TR, Lopez-Gonzalez G *et al.* 2018. Field methods for sampling tree height for tropical forest biomass estimation. *Methods in Ecology and Evolution* 9: 1179–1189.
- Taubert F, Jahn MW, Dobner H-J, Wiegand T, Huth A. 2015. The structure of tropical forests and sphere packings. *Proceedings of the National Academy of Sciences, USA* 112: 15125–15129.
- Wang H, Prentice IC, Keenan TF, Davis TW, Wright IJ, Cornwell WK, Evans BJ, Peng C. 2017. Towards a universal model for carbon dioxide uptake by plants. *Nature Plants* 3: 734–741.
- Wenger SJ, Olden JD. 2012. Assessing transferability of ecological models: an underappreciated aspect of statistical validation. *Methods in Ecology and Evolution* 3: 260–267.
- Wolf A, Anderegg WRL, Pacala SW. 2016. Optimal stomatal behavior with competition for water and risk of hydraulic impairment. *Proceedings of the National Academy of Sciences, USA* 113: E7222–E7230.
- Wolf A, Ciais P, Bellassen V, Delbart N, Field CB, Berry JA. 2011. Forest biomass allometry in global land surface models. *Global Biogeochemical Cycles* 25: GB3015.



About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit www.newphytologist.com to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit www.newphytologist.com