

## **Volatile organic compounds of six French Dryopteris species: natural odorous and bioactive resources**

Didier Froissard, Sylvie Rapior, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois, Françoise Fons

► **To cite this version:**

Didier Froissard, Sylvie Rapior, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois, et al.. Volatile organic compounds of six French Dryopteris species: natural odorous and bioactive resources. Natural Product Communications , SAGE Publications, 2014, 9 (1), pp.137-140. 10.1177/1934578X1400900138 . hal-02194220

**HAL Id: hal-02194220**

**<https://hal.umontpellier.fr/hal-02194220>**


Submitted on 25 Jul 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all  
Aspects of Natural Products Research



Volume 9. Issue 1. Pages 1-144. 2014  
ISSN 1934-578X (printed); ISSN 1555-9475 (online)  
[www.naturalproduct.us](http://www.naturalproduct.us)

**EDITOR-IN-CHIEF****DR. PAWAN K AGRAWAL**

Natural Product Inc.  
7963, Anderson Park Lane,  
Westerville, Ohio 43081, USA  
agrawal@naturalproduct.us

**EDITORS****PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry,  
University of Granada,  
Campus de Fuente Nueva, s/n, 18071, Granada, Spain  
afbarre@ugr.es

**PROFESSOR ALESSANDRA BRACA**

Dipartimento di Chimica Bioorganica e Biofarmacia,  
Università di Pisa,  
via Bonanno 33, 56126 Pisa, Italy  
braca@farm.unipi.it

**PROFESSOR DEAN GUO**

State Key Laboratory of Natural and Biomimetic Drugs,  
School of Pharmaceutical Sciences,  
Peking University,  
Beijing 100083, China  
gda5958@163.com

**PROFESSOR YOSHIHIRO MIMAKI**

School of Pharmacy,  
Tokyo University of Pharmacy and Life Sciences,  
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan  
mimakiy@ps.toyaku.ac.jp

**PROFESSOR STEPHEN G. PYNE**

Department of Chemistry  
University of Wollongong  
Wollongong, New South Wales, 2522, Australia  
spyne@uow.edu.au

**PROFESSOR MANFRED G. REINECKE**

Department of Chemistry,  
Texas Christian University,  
Forts Worth, TX 76129, USA  
m.reinecke@tcu.edu

**PROFESSOR WILLIAM N. SETZER**

Department of Chemistry  
The University of Alabama in Huntsville  
Huntsville, AL 35809, USA  
wsetzer@chemistry.uah.edu

**PROFESSOR YASUHIRO TEZUKA**

Institute of Natural Medicine  
Institute of Natural Medicine, University of Toyama,  
2630-Sugitani, Toyama 930-0194, Japan  
tezuka@inm.u-toyama.ac.jp

**PROFESSOR DAVID E. THURSTON**

Department of Pharmaceutical and Biological Chemistry,  
The School of Pharmacy,  
University of London, 29-39 Brunswick Square,  
London WC1N 1AX, UK  
david.thurston@pharmacy.ac.uk

**HONORARY EDITOR****PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,  
University of Portsmouth,  
Portsmouth, PO1 2DT U.K.  
axuf64@dsl.pipex.com

**ADVISORY BOARD**

Prof. Viqar Uddin Ahmad  
Karachi, Pakistan

Prof. Giovanni Appendino  
Novara, Italy

Prof. Yoshinori Asakawa  
Tokushima, Japan

Prof. Roberto G. S. Berlinck  
São Carlos, Brazil

Prof. Anna R. Bilia  
Florence, Italy

Prof. Maurizio Bruno  
Palermo, Italy

Prof. César A. N. Catalán  
Tucumán, Argentina

Prof. Josep Coll  
Barcelona, Spain

Prof. Geoffrey Cordell  
Chicago, IL, USA

Prof. Fatih Demirci  
Eskişehir, Turkey

Prof. Dominique Guillaume  
Reims, France

Prof. Ana Cristina Figueiredo  
Lisbon, Portugal

Prof. Cristina Gracia-Viguera  
Murcia, Spain

Prof. Duvvuru Gunasekar  
Tirupati, India

Prof. Hisahiro Hagiwara  
Niigata, Japan

Prof. Kurt Hostettmann  
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga  
Mexico, D. F., Mexico

Prof. Leopold Jirovetz  
Vienna, Austria

Prof. Vladimir I Kalinin  
Vladivostok, Russia

Prof. Niel A. Koorbanally  
Durban, South Africa

Prof. Chiaki Kuroda  
Tokyo, Japan

Prof. Hartmut Laatsch  
Göttingen, Germany

Prof. Marie Laccaille-Dubois  
Dijon, France

Prof. Shoen-Sheng Lee  
Taipei, Taiwan

Prof. Imre Mathe  
Szeged, Hungary

Prof. Ermino Murano  
Trieste, Italy

Prof. M. Soledade C. Pedras  
Saskatoon, Canada

Prof. Luc Pieters  
Antwerp, Belgium

Prof. Peter Proksch  
Düsseldorf, Germany

Prof. Phila Raharivelomanana  
Tahiti, French Polynesia

Prof. Luca Rastrelli  
Fisciano, Italy

Prof. Stefano Serra  
Milano, Italy

Prof. Monique Simmonds  
Richmond, UK

Dr. Bikram Singh  
Palampur, India

Prof. John L. Sorensen  
Manitoba, Canada

Prof. Johannes van Staden  
Scottsville, South Africa

Prof. Valentin Stonik  
Vladivostok, Russia

Prof. Winston F. Tinto  
Barbados, West Indies

Prof. Sylvia Urban  
Melbourne, Australia

Prof. Karen Valant-Vetschera  
Vienna, Austria

**INFORMATION FOR AUTHORS**

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

**To Subscribe:** Natural Product Communications is a journal published monthly. 2014 subscription price: US\$2,395 (Print, ISSN# 1934-578X); US\$2,395 (Web edition, ISSN# 1555-9475); US\$2,795 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

## Volatile Organic Compounds of six French *Dryopteris* Species: Natural Odorous and Bioactive Resources

Didier Froissard<sup>a</sup>, Sylvie Rapior<sup>b</sup>, Jean-Marie Bessi re<sup>c</sup>, Alain Fruchier<sup>d</sup>, Bruno Buatois<sup>c</sup> and Fran oise Fons<sup>b\*</sup>

<sup>a</sup>Laboratoire de Botanique, Facult  de Pharmacie de Limoges, 2 rue du Docteur Raymond Marcland, F-87025 Limoges Cedex, France

<sup>b</sup>Laboratoire de Botanique, Phytochimie et Mycologie, Facult  de Pharmacie (Universit  Montpellier 1), UMR 5175 CEFE, B.P. 14 491, 15 avenue Charles Flahault, F-34093 Montpellier Cedex 5, France

<sup>c</sup>UMR 5175 CEFE, 1919 Route de Mende, F-34293 Montpellier Cedex 5, France

<sup>d</sup>ENSCM, UMR 5253, 8 Rue de l' cole Normale, F-34296 Montpellier Cedex 5, France

didier.froissard@unilim.fr

Received: July 12<sup>th</sup>, 2013; Accepted: November 5<sup>th</sup>, 2013

Aerial parts of six *Dryopteris* species collected in France were investigated for volatile organic compounds (VOC) for the first time. Fifty-three biosynthesized VOC from the shikimic, lipidic and terpenic pathways were identified using gas chromatography/mass spectrometry. Many bioactive polyketide compounds as filicinic derivatives (from 8.5 to 23.5%) and phloroglucinol derivatives (from 8.2 to 53.8%) with various pharmacological activities were detected in high amount from five analysed *Dryopteris* species, in particular *D. oreades* and *D. borrieri*, i.e., propionylfilicinic acid (> 10% in *D. affinis* and *D. ardechensis*) and 2,6-dihydroxy-4-methoxy-3-methylbutyrophenone (aspidinol) (19.1% and 14.6% in *D. oreades* and *D. borrieri*, respectively). Several terpenic derivatives with a low odor threshold were identified, i.e., carota-5,8-diene (from 2.5 to 18.4%: floral, woody or fresh bark note), (*E*)-nerolidol (> 10% for *D. borrieri* and *D. cambrensis*; floral or woody odor),  $\alpha$ -selinene (> 7% for *D. ardechensis*; woody-spicy odor), and aristolene (12.8% in *D. affinis*; flower, sweet odor). The main isoprenoid derivatives were 4-hydroxy-5,6-epoxyionol, 3-oxo- $\alpha$ -ionol and 4-oxo-7,8-dihydro- $\beta$ -ionone (essentially in *D. remota*), whereas the main aromatic compound was 4-hydroxy-3-methoxyacetophenone (20.6% and 12.6% in *D. cambrensis* and *D. borrieri*, respectively) and the main lipid derivative was 1-octen-3-ol with a mushroom-like odor (from 0.4 to 8.3%). *Dryopteris* species resources are of great interest as a reservoir of odorous and bioactive compounds.

**Keywords:** Fern, Volatile Organic Compounds, Carota-5,8-diene, (*E*)-Nerolidol, Phloroglucinols, Filicinic derivatives, 1-Octen-3-ol, Biological activities.

*Dryopteridaceae* is a subcosmopolitan family including approximately one thousand species and with only two native genera in Western Europe: *Dryopteris* Adans and *Polystichum* Roth [1a]. *Dryopteris affinis* group is constituted in Western Europe by *D. affinis* (R. Lowe) Fras.-Jenk., *D. borrieri* (Newm.) Newm. ex Oberh. & Tavel, *D. cambrensis* (Fras.-Jenk.) Beitel & W.R. Buck, the uncommon *D. pseudodisjuncta* (Tavel ex Fras.-Jenk.) Fras.-Jenk. and a very recently reported species, i.e., the rare *D. lacunosa* S. Jess., Zenner, Ch. Stark & Bujnoch [1b,1c].

*Dryopteris* species have been mainly investigated for phloroglucinol derivatives due to their various pharmacological activities for humans and many ethnoveterinary uses, i.e., schistosomicidal, antimicrobial, antitumor, anti-inflammatory and antioxidant properties [2a-2m].

Few *Dryopteris* species are known having an odor. *D. villarii* (Bellardi) Schinz & Thell. and *D. mindshelkensis* Pavlov have a balsamic odor when crushed [3a]. *D. aemula* (Aiton) O. Kuntze and *D. crispifolia* Rasbach, Reichstein & Vida smell like newly mown hay, whereas *D. fragrans* (L.) Schott has a spicy odor when dry [3b]. Very few data have been reported previously on the volatile organic compounds (VOC) of *Dryopteris* species and none from the *D. affinis* group. *D. filix-mas* (L.) Schott contains in particular (*E*)-nerolidol (38.7%) with floral, woody or fresh bark odor [4a], acylfilicinic acids and 1-octen-3-ol [4b]. *D. dilatata* (Hoffm.) A. Gray odor is characterized by (*E*)-2-hexenal and (*Z*)-3-hexenal, responsible for the "green odor" and 1-octen-3-ol [4c], whereas the

odor of *D. fragrans* is based on terpenic derivatives such as 10-hydroxy-15-oxo- $\alpha$ -cadinol, albicanyl acetate,  $\alpha$ -cadinene and albicanol [4d].

Fresh aerial parts of three main species from the *D. affinis* group, i.e., *D. affinis*, *D. borrieri*, and *D. cambrensis*, and three related species, *D. oreades* Fomin, *D. ardechensis* Fraser-Jenkins and *D. remota* (A. Braun ex D ll) Druce were investigated for their volatile profiles using GC-MS. The VOC were identified as previously reported [4b, 4c, 4e, 4f, 5a].

Fifty-three components biosynthesized from the shikimic, lipidic and terpenic pathways were identified from the concentrated diethyl ether extracts of the six *Dryopteris* species (Table 1).

*D. affinis* was dominated by a large number of terpenic derivatives (49.5%), i.e., carota-5,8-diene (14.7%; floral, woody or fresh bark note), aristolene (12.8%; flower, sweet odor) [5b], (*E*)-nerolidol (5.9%; with floral odor), aristola-1(10),8-diene (3.4%),  $\alpha$ -selinene (3%; amber type odor or sweet-woody, slightly peppery odor [5c]),  $\beta$ -selinene (2.5%; mild, woody, warm, herbaceous peppery odor) and eremophilene (2.4%). It also contained polyketide derivatives (36.4%) as filicinic compounds (19.7%; [5d]), i.e., propionylfilicinic acid (11%) and acetylfilicinic acid (6.7%), and phloroglucinol compounds (16.7%). The main lipid derivative was 1-octen-3-ol (6.9%; mushroom-like odor), well-known for many mushrooms [5e-5g], and previously reported for ferns, including *D. filix-mas* and *D. dilatata*, horsetails and Angiosperms [4b, 4c, 5a, 5h].

**Table 1:** Percentage of volatile organic compounds<sup>a</sup> in fresh aerial parts of *Dryopteris* species.

Compounds	RI <sup>b</sup>	<i>D. affinis</i>	<i>D. borrieri</i>	<i>D. cambrensis</i>	<i>D. oreades</i>	<i>D. ardechensis</i>	<i>D. remota</i>
<b>Lipid derivatives</b>		<b>9.1</b>	<b>5.7</b>	<b>4.4</b>	<b>3.4</b>	<b>3.2</b>	<b>20.7</b>
1-Octen-3-ol	982	6.9	4.8	4.4	3.0	0.4	8.3
3-Octanol	1000	1.6	0.9		0.3	0.9	2.5
3-Hexenoic acid	1017					0.3	3.1
2-Hexenoic acid	1025					0.5	2.1
Nonanal	1106	0.7			0.1	0.3	2.6
Decanal	1205					0.8	2.0
<b>Polyketide compounds</b>		<b>36.4</b>	<b>45.5</b>	<b>22.3</b>	<b>62.3</b>	<b>31.6</b>	<b>0.0</b>
Acetylfilicinic acid	1443	6.7	2.9	4.9			
Propionylfilicinic acid	1546	11.0	7.7	4.4	0.5	10.1	
Butyrylfilicinic acid	1615		1.9	3.1	4.3	4.4	
Valeroylfilicinic acid	1622	2.1	1.6	1.1	2.5	5.2	
NI <sub>c</sub>	1686		2.7		1.2	3.8	
= <b>Sub-total filicinic compounds</b>		<b>(19.8)</b>	<b>(16.8)</b>	<b>(13.5)</b>	<b>(8.5)</b>	<b>(23.5)</b>	<b>(0.0)</b>
2,4,6-Trimethoxy-propiofenone	1855				6.0		
2,6-Dihydroxy-4-methoxypropiofenone	1882		0.3		3.3		
2,6-Dihydroxy-3-methyl-4-methoxypropiofenone	1942		6.7		3.1		
2,6-Dihydroxy-4-methoxybutyrophenone (desaspidinol)	1984				2.7		
NI	2028				6.7		
2,6-Dihydroxy-4-methoxy-3-methylbutyrophenone (aspidinol)	2038		14.6	1.8	19.1		
2,6-Dihydroxy-4-methoxyvalerophenone	2091	2.9	7.2	7.0	12.9	8.2	
NI	2140	13.8					
= <b>Sub-total phloroglucinol compounds</b>		<b>(16.7)</b>	<b>(28.8)</b>	<b>(8.8)</b>	<b>(53.8)</b>	<b>(8.2)</b>	<b>(0.0)</b>
<b>Aromatic compounds</b>		<b>1.4</b>	<b>18.4</b>	<b>24.8</b>	<b>0.2</b>	<b>5.7</b>	<b>6.6</b>
Benzaldehyde	962		1.1			0.8	1.8
Benzyl alcohol	1043		0.7			2.5	2.9
Phenylethanal	1046	1.0	0.7		0.1		
Cinnamaldehyde	1164		0.7				
4-Hydroxy-3-methoxystyrene	1293		0.5	1.3		2.5	1.8
4-Hydroxy-3-methoxybenzaldehyde (vanilline)	1385		1.2	1.3			
2,4,6-Trihydroxybutyrophenone (phlorobutyrophenone)	1552				0.1		
4-Hydroxy-3-methoxyacetophenone	1560	0.4	12.6	20.6			
4-Hydroxy-3-methoxybenzoic acid (vanillic acid)	1576		0.6	1.7			
4-Hydroxy-3-methoxycinnamaldehyde	1694		0.4				
<b>Terpenic compounds</b>		<b>49.4</b>	<b>21.2</b>	<b>32.8</b>	<b>26.3</b>	<b>48.5</b>	<b>0.6</b>
Linalool	1104	1.3				2.7	0.6
Thymoquinone	1240		2.0	1.2			
Daucene	1380	0.4				0.8	
β-Elemene	1391		1.0	0.8		2.3	
Aristolene	1419	12.8				0.4	
γ-Elemene	1427	2.6				2.3	
Aristola-1(10),8-diene	1458	3.4			2.1	6.6	
Carota-5,8-diene	1473	14.7	2.5	12.7	18.4	9.3	
β-Selinene	1487	2.5					
Eremophilene	1489	2.4	3.1	4.4	0.9	4.4	
Valencene	1494	0.3	0.6			1.1	
α-Selinene	1496	3.0			0.5	7.8	
Germacrene A	1501		0.6	1.5			
7-Epi-α-selinene	1519					3.1	
(E)-Nerolidol	1556	5.9	11.5	12.2	4.3	7.9	
<b>Isoprenoid derivatives</b>		<b>3.7</b>	<b>9.2</b>	<b>15.7</b>	<b>7.7</b>	<b>11.0</b>	<b>72.1</b>
NI	1335						1.4
2,3-Dehydro-α-ionol	1384						3.1
3,4-Dehydro-7,8-dihydro-β-ionone	1413						1.3
2,3-Dehydro-α-ionone	1428						1.0
β-Ionone	1480		0.2			1.1	0.5
Dihydroactinidiolide	1505		1.1				
NI	1599			0.2			10.4
NI	1608						5.4
NI	1615						3.4
4-Hydroxy-7,8-dihydro-β-ionone	1632		1.1	3.1			3.6
4-Hydroxy-β-ionone	1638				1.6	1.4	3.6
3-Oxo-α-ionol	1645				2.0		5.8
3-Oxo-α-ionone	1652		1.1	0.9		1.2	0.8
4-Oxo-β-ionone	1663		0.4				0.8
4-Hydroxy-5,6-epoxyionol	1672						19.1
4-Oxo-7,8-dihydro-β-ionone	1675		1.8	2.5			8.1
NI	1692	1.1	3.5	9.0	4.1	7.3	2.1
NI	1708	0.5					
NI	1717	1.0					
NI	1746	1.2					1.8

<sup>a</sup> Relative percentage of the VOC based on the GC-MS chromatographic area.<sup>b</sup> RI = Retention Indices on SLB<sup>TM</sup>-5MS column (Supelco).<sup>c</sup> NI = Not identified.

The volatile pattern of *D. borrieri* was based on polyketide derivatives (45.5%), mainly phloroglucinols (28.8%), i.e., 2,6-dihydroxy-4-methoxy-3-methylbutyrophenone, also named aspidinol (14.6%), 2,6-dihydroxy-4-methoxyvalerophenone (7.2%)

and 2,6-dihydroxy-3-methyl-4-methoxypropiofenone (6.7%), as well as filicinic compounds (16.8%), in particular propionylfilicinic acid (7.7%). The investigation indicated a lower amount of terpenic derivatives (21.2%), i.e., (*E*)-nerolidol (11.5%) and eremophilene

(3.1%), aromatic components (18.4%), i.e., 4-hydroxy-3-methoxyacetophenone (12.6%), and 1-octen-3-ol (4.8%). It should be noted that low amounts of strong odorous components were detected as benzaldehyde (bitter almond odor; [5i]), cinnamaldehyde (cinnamon odor), vanillin, vanillic acid (sweet cream with vanilla aroma) and dihydroactinidiolide (fruity odor [5j]), also found in *Equisetum arvense* L. [5a].

The broad spectrum of volatile components identified in *D. cambrensis* contributed to its complex smell, including isoprenoid, polyketide, aromatic and terpenic derivatives and also an odorous lipid derivative, 1-octen-3-ol. The terpenic profile (32.8%) was based on carota-5,8-diene (12.7%), (*E*)-nerolidol (12.2%) and eremophilene (4.4%). The fern also contained a high level of aromatic components (24.8%), mainly 4-hydroxy-3-methoxyacetophenone (20.6%). Table 1 lists the polyketide derivatives (22.3%) divided into filicinic compounds (13.5%), i.e., acetylfilicinic acid (4.9%), propionylfilicinic acid (4.4%) and butyrylfilicinic acid (3.1%), as well as phloroglucinols (8%), mainly 2,6-dihydroxy-4-methoxyvalerophenone (7%) and aspidinol (1.8%). The main isoprenoids (15.7%) were  $\beta$ -ionone derivatives, i.e., 4-hydroxy-7,8-dihydro- $\beta$ -ionone (3.1%) and 4-oxo-7,8-dihydro- $\beta$ -ionone (2.5%), already found in *Dryopteris dilatata* and *Phegopteris connectilis* (Michx) Watt, and in *Gymnocarpium dryopteris* (L.) Newman, respectively [4c].

The volatile content of *D. oreades* was mainly dominated by polyketide derivatives (62.3%), essentially phloroglucinols (53.8%) as 2,6-dihydroxy-4-methoxy-3-methylbutyrophenone (aspidinol; 19.1%), 2,6-dihydroxy-4-methoxyvalerophenone (12.9%), 2,4,6-trimethoxypropiophenone (6%) and 2,5-dihydroxy-4-methoxybutyrophenone (desaspidinol; 2.7%), as well as filicinic components (8.8%), in particular butyrylfilicinic acid (4.3%). It should be noted that aspidinol or desaspidinol are well-known for their anthelmintic, anti-tumor and antibacterial properties at very low concentration [6a,6b,2e]. The main terpenes were carota-5,8-diene (18.4%) and (*E*)-nerolidol (4.3%), and the major lipid derivative was still 1-octen-3-ol (3%).

The major VOCs from *D. ardechensis* were terpenic derivatives (48.5%), i.e., carota-5,8-diene (9.3%), (*E*)-nerolidol (7.9%),  $\alpha$ -selinene (7.8%), aristola-1(10),8-diene (6.6%) and linalool (2.7%; orange flower odor [6c]). It should be noted that eremophilene, valencene,  $\alpha$ - and  $\beta$ -selinene and 7-epi- $\alpha$ -selinene have the same biosynthetic pathway; several of them have antifungal or insecticidal properties [6d-6g]. The polyketide pathway (31.6%) produced three filicinic compounds (23.5%), namely propionylfilicinic acid (10.1%), valeroylfilicinic acid (5.2%) and butyrylfilicinic acid (4.4%), and only one phloroglucinol compound, i.e., 2,6-dihydroxy-4-methoxyvalerophenone (8.2%). Several isoprenoid derivatives, such as 4-hydroxy- $\beta$ -ionone (1.4%), 3-oxo- $\alpha$ -ionone (1.2%) and  $\beta$ -ionone (1.1%) with a violet flower odor [6c] were also identified. Six lipid derivatives (3.2%), such as 3-octanol, decanal and 1-octen-3-ol completed the VOC content of *D. ardechensis*.

## References

- [1] (a) Prelli R, Boudry M. (2001) *Les fougères et plantes alliées de France et d'Europe occidentale*, Belin Publisher, Paris; (b) Fraser-Jenkins CR. (2007) The species and subspecies in the *Dryopteris affinis* group. *Fern Gazette*, **18**, 1-26; (c) Jessen S, Bujnoch W, Zenner G, Stark C. (2011) *Dryopteris lacunosa* – eine neue Art des *Dryopteris affinis*-Aggregats (*Dryopteridaceae*, Pteridophyta). *Kochia*, **5**, 9-31.
- [2] (a) Bharate SP. (2006) Phloroglucinol compounds of natural origin. *Natural Product Reports*, **23**, 558-591; (b) Gao Z, Ali Z, Zhao J, Qiao L, Lei H, Lu Y, Khan IA. (2008) Phytochemical investigation of the rhizomes of *Dryopteris crassirhizoma*. *Phytochemistry Letters*, **1**, 188-190; (c) Githiori JB, Athanasiadou S, Thamsborg SM. (2006) Use of plants in novel approaches for control of gastrointestinal helminths in livestock with emphasis on small ruminants. *Veterinary Parasitology*, **139**, 308-320; (d) Jahan N, Ahmad M, Mehjabeen, Zia-Ul-Haq M, Alam SM, Qureshi M. (2010) Antimicrobial screening of some medicinal plants of Pakistan. *Pakistan Journal of Botany*, **42**, 4281-4284; (e) Kapadia GJ, Tokuda H, Konoshima

*D. remota* showed the highest amount of isoprenoid derivatives (72%), including eleven identified compounds. The VOC profile was widely dominated by 4-hydroxy-5,6-epoxyionol (19.1%), 4-oxo-7,8-dihydro- $\beta$ -ionone (8.1%) and 3-oxo- $\alpha$ -ionol (5.8%; spicy odor). It also contained several odorous C<sub>8</sub> to C<sub>10</sub> derivatives (20.7%), such as 1-octen-3-ol (8.3%), 3-hexenoic acid (3.1%; honey odor somewhat waxy fruity and herbal [6h]), nonanal (2.6%; orange and green scent [5j]), 3-octanol (2.5%; mushroom-like odor and buttery) used in the food, flavor, cosmetic, pharmaceutical, tobacco and perfume industries [6i], and 2-hexenoic acid (2.1%), with a powerful fruity odor. Benzyl alcohol (2.9%; walnut flavor [6j]) and benzaldehyde (1.8%; bitter almond odor [4f]) emphasized the great interest of *D. remota* for aroma and flavor companies. Regarding the eight *Dryopteris* species investigated for VOC (fresh aerial part) in Table 1 and previously by the authors [4b,4c], *D. remota* and *D. dilatata* do not contain any filicinic derivatives, phloroglucinols or (*E*)-nerolidol.

This paper emphasizes that ferns are novel resources for natural compounds. Table 1 demonstrates that *Dryopteris* species can generate a broad spectrum of VOC for both odorous and bioactive ingredients. Within the former, terpenic compounds with floral, fruity or spicy notes, i.e., linalool, (*E*)-nerolidol, and ionone derivatives are the main fragrant components required for aroma applications. Within the last, phloroglucinol and filicinic derivatives are of various biological interests for the pharmaceutical as well as the cosmetic and hygiene products industries. Because only aerial parts have been used, intensive culture of ferns may be developed to extract biomolecules from the natural resources without any plant destruction. Natural components can also be used for hemisynthesis of di- to polymeric bioactive derivatives. *Dryopteris* species resources are indeed of great interest as candidates for bioactive aroma ingredients and for the discovery of new drugs with various therapeutic applications due to their potential antioxidant, antibacterial and antitumor promoting properties [7a-7d].

**Plant material:** Fresh aerial parts of ferns were collected as follows: *Dryopteris affinis*: 14/07/2009, Gimel les Cascades (Corrèze), *D. borrieri* and *D. cambrensis* 30/05/2010, Saint Etienne Vallée Française, (Ardèche). *D. oreades* 13/07/2011, Murat-le-Quaire, (Puy de Dôme), *D. ardechensis* and *D. remota*: 31/08/2010, Botanical Garden of Strasbourg. Voucher specimens are deposited at the Laboratory of Botany (Faculty of Pharmacy, Limoges, France).

**Plant part and GC-MS analyses:** Fresh aerial parts of 6 *Dryopteris* species were treated and investigated for volatile organic compounds as previously reported [4b,4c,5a].

**Acknowledgments** - The authors greatly thank the Botanical Garden of Strasbourg for providing *D. remota* and *D. ardechensis*. The GC-MS analyses were carried out at the Plate-forme d'analyses chimiques en écologie (PACE) / Labex CeMEB (Montpellier, France).

- T, Takasaki M, Takayasu J, Nishino H. (1996) Anti-tumor promoting activity of *Dryopteris* phlorophenone derivatives. *Cancer Letters*, **105**, 161-165; (f) Lee JS, Miyashiro H, Nakamura N, Hattori M. (2008) Two new triterpenes from the rhizome of *Dryopteris crassirhizoma*, and inhibitory activities of its constituents on human immunodeficiency virus-1 protease. *Chemical & Pharmaceutical Bulletin*, **56**, 711-714; (g) Lee HB, Kim JC, Lee SM. (2009) Antibacterial activity of two phloroglucinols, flavaspidic acids AB and PB, from *Dryopteris crassirhizoma*. *Archives of Pharmacal Research*, **32**, 655-659; (h) Lu C, Zhang HY, Ji J, Wang GX. (2012) *In vivo* anthelmintic activity of *Dryopteris crassirhizoma*, *Kochia scoparia*, and *Polygala tenuifolia* against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitology Research*, **110**, 1085-1090; (i) Magalhães LG, Kapadia GJ, Da Silva Tonuci LR, Caixeta SC, Parreira NA, Rodrigues V, Da Silva Filho AA. (2010) *In vitro* schistosomicidal effects of some phloroglucinol derivatives from *Dryopteris* species against *Schistosoma mansoni* adult worms. *Parasitology Research*, **106**, 395-340; (j) Penttilä A, Sundman J. (1964) On natural and synthetic homologues of *Dryopteris* phloroglucinol derivatives. *Acta Chemica Scandinavica*, **18**, 344-352; (k) Singh IP, Soare LC, Ferdeş M, Stefanov S, Denkova Z, Nicolova R, Denev P, Bejan C, Păunescu A. (2012) Antioxidant activity, polyphenols content and antimicrobial activity of several native pteridophytes of Romania. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, **40**, 53-57; (l) Socolsky C, Domínguez L, Asakawa Y, Bardón A. (2012) Unusual terpenylated acylphloroglucinols from *Dryopteris wallichiana*. *Phytochemistry*, **80**, 115-122; (m) Sun Y, Gao C, Luo M, Wang W, Gu C, Zu Y, Li J, Efferth T, Fu Y. (2013) Aspidin PB, a phloroglucinol derivative, induces apoptosis in human hepatocarcinoma HepG2 cells by modulating PI3K/Akt/GSK3 $\beta$  pathway. *Chemico-Biological Interactions*, **201**, 1-8.
- [3] (a) Moret JL. (1994) Excursion pteridologique et paludologique en suisse orientale du 25 au 28 août 1993 (ag, zg, sg, ai, ar, sz). *Bulletin du Cercle Vaudois de Botanique*, **23**, 43-51; (b) Jones DL. (1987) *Encyclopedia of ferns*. Lothian, Port Melbourne, Australia.
- [4] (a) Joichi A, Nakamura Y, Haze S, Ishikawa T, Atoji H, Nishida T, Sakurai K. (2013) Volatile constituents of blue-coloured hybrid tea rose flowers. *Flavour and Fragrance Journal*, **28**, 180-187; (b) Fons F, Froissard D, Bessière JM, Buatois B, Rapior S. (2010) Biodiversity of volatile organic compounds from five French ferns. *Natural Product Communications*, **5**, 1655-1658; (c) Froissard D, Fons F, Bessière JM, Buatois B, Rapior S. (2011) Volatiles of French ferns and “fougère” scent in perfumery. *Natural Product Communications*, **6**, 1723-1726; (d) Shen ZB, Luo WY, Yan YS, Zhu JF. (2006) Study on terpene of *Dryopteris fragrans* L. *Journal of Chinese Medicinal Materials*, **29**, 334-335; (e) Rapior S, Fons F, Bessière JM. (2000) The fenugreek odor of *Lactarius helvus*. *Mycologia*, **92**, 305-308; (f) Rapior S, Kanska G, Guillot J, Andary C, Bessière JM. (2000) Volatile composition of *Laetiporus sulphureus*. *Cryptogamie, Mycologie*, **21**, 67-72.
- [5] (a) Fons F, Froissard D, Bessière JM, Fruchier A, Buatois B, Rapior S. (2013) Volatile composition of six horsetails: prospects and perspectives. *Natural Product Communications*, **8**, 509-512; (b) Liu CJ, Zhang J, Zhou ZK, Hua ZT, Wan HY, Xie YH, Wang ZW, Deng L. (2013) Analysis of volatile compounds and identification of characteristic aroma components of *Toona sinensis* (A. Juss.) Roem. using GC-MS and GC-O. *Food and Nutrition Sciences*, **4**, 305-314; (c) Pino JA, Marbot R, Carlos Vázquez C. (2002) Characterization of volatiles in Costa Rican guava [*Psidium friedrichsthalianum* (Berg) Niedenzu] fruit. *Journal of Agricultural and Food Chemistry*, **50**, 6023-6026; (d) Lounasmaa M, Karjalainen A, Widen CJ, Huhtikangas A. (1971) Mass spectral studies on some naturally occurring phloroglucinol derivatives. Part I. The mass spectra of filicinic acid and its acetyl, propionyl and butyryl derivatives. *Acta Chemica Scandinavica*, **25**, 3428-3440; (e) Rapior S, Breheret S, Talou T, Pélissier Y, Bessière JM. (2002) The anise-like odor of *Clitocybe odora*, *Lentinellus cochleatus* and *Agaricus essettei*. *Mycologia*, **94**, 373-376; (f) Rapior S, Breheret S, Talou T, Pélissier Y, Milhau M, Bessière JM. (1998) Volatile components of fresh *Agrocybe aegerita* and *Tricholoma sulfureum*. *Cryptogamie, Mycologie*, **19**, 15-23; (g) Rapior S, Fons F, Bessière JM. (2003) Volatile flavor constituents of *Lepista nebularis* (Clouded Clitocybe). *Cryptogamie, Mycologie*, **24**, 159-166; (h) Fons F, Rapior S, Gargadennec A, Andary C, Bessière JM. (1998) Volatile components of *Plantago lanceolata* (Plantaginaceae). *Acta Botanica Gallica*, **145**, 265-269; (i) Fons F, Rapior S, Fruchier A, Saviuc P, Bessière JM. (2006) Volatile composition of *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum*. *Cryptogamie, Mycologie*, **27**, 45-55; (j) Fons F, Rapior S, Eyssartier G, Bessière JM. (2003) Volatile compounds in the *Cantharellus*, *Craterellus* and *Hydnum* genera. *Cryptogamie, Mycologie*, **24**, 367-376.
- [6] (a) Bosman AA, Combrinck S, Roux-van der Merwe R, Botha BM, McCrindle R. (2004) Isolation of an anthelmintic compound from *Leucosidea sericea*. *South African Journal of Botany*, **70**, 509-511; (b) Wang W, Zeng YH, Osman K, Shinde K, Rahman M, Gibbons S, Mu Q. (2010) Norlignans, acylphloroglucinols, and a dimeric xanthone from *Hypericum chinense*. *Journal of Natural Products*, **73**, 1815-1820; (c) Noguero-Pato R, González-Barreiro C, Cancho-Grande B, Santiago JL, Martínez MC, Simal-Gándara J. (2012) Aroma potential of Brancellao grapes from different cluster positions. *Food Chemistry*, **132**, 112-124; (d) Bağcı E, Baser KHC, Kurkuoğlu M, Babac T, Celik S. (1999) Study of the essential oil composition of two subspecies of *Abies cilicica* (Ant. et Kotschy) Carr. from Turkey. *Flavour and Fragrance Journal*, **14**, 47-49; (e) Miyazawa M, Teranishi A, Ishikawa Y. (2003) Components of the essential oil from *Petasites japonicas*. *Flavour and Fragrance Journal*, **18**, 231-233; (f) Troncoso C, Perez C, Hernandez V, Sanchez-Olate M, Rios D, San Martin A, Becerra J. (2013) Induction of defensive response in *Eucalyptus globulus* plants and its persistence in vegetative propagation. *Natural Product Communications*, **8**, 397-400; (g) Wang SY, Wu CL, Chug FH, Chien SC, Kuo YH, Shyur LF, Chang ST. (2005) Chemical composition and antifungal activity of essential oil isolated from *Chamaecyparis formosensis* Matsum. wood. *Holzforchung*, **59**, 295-299; (h) The Good Scents Company (2011) <http://www.thegoodscentscompany.com/>; (i) Cho IH, Kim SY, Choi HK, Kim YS. (2006) Characterization of aroma-active compounds in raw and cooked pine-mushrooms (*Tricholoma matsutake* Sing.). *Journal of Agricultural and Food Chemistry*, **54**, 6332-6335; (j) Larsen M, Poll L. (1990) Odour thresholds of some important aroma compounds in raspberries. *Zeitschrift Lebensmittel-Untersuchung Forschung*, **191**, 129-131.
- [7] (a) Singh IP, Sidana J, Bansal P, Foley WJ. (2009) Phloroglucinol compounds of therapeutic interest: global patent and technology status. *Expert Opinion Therapeutic Patents*, **19**, 847-856; (b) Socolsky C, Hernández MA, Bardón A. (2012) Fern acylphloroglucinols: structure, location, and biological effects. In *Studies in Natural Products Chemistry*, Vol. **38**, Elsevier Science and Technology, Oxford, UK. 105-157; (c) Widén CJ, Fraser-Jenkins CR, Reichstein T, Sarvela J. (2001) A survey of phenolic compounds in *Dryopteris* and related fern genera. Part III. Phloroglucinol derivatives in subgenera *Erythrovaria* and *Nephrocystis* and related genera (Pteridophyta, *Dryopteridaceae*). *Annales Botanici Fennici*, **38**, 99-138; (d) Franchi GG, Ferri S. (1988) Localization of the active principles of the male fern (*Dryopteris filix-mas* (L.) Schott.) by fluorescence microscopy. *Pharmacological Research Communications*, **20**, 135-138.

<b>Antiplatelet Aggregation Effects of Phenanthrenes from <i>Calanthe arisanensis</i></b> Chia-Lin Lee, Ming-Hon Yen, Fang-Rong Chang, Chin-Chung Wu and Yang-Chang Wu	83
<b><i>In Vivo</i> Anti-inflammatory Activity of Some Naturally Occurring <i>O</i>- and <i>N</i>-Prenyl Secondary Metabolites</b> Francesco Epifano, Salvatore Genovese, Serena Fiorito, Roberto della Loggia, Aurelia Tubaro and Silvio Sosa	85
<b>Phomopsolides and Related Compounds from the Alga-associated Fungus, <i>Penicillium clavigerum</i></b> Andrea A. Stierle, Donald B. Stierle, Grant G. Mitman, Shea Snyder, Christophe Antezak and Hakim Djaballah	87
<b>Qualitative Identification of Dibenzoylmethane in Licorice Root (<i>Glycyrrhiza glabra</i>) using Gas Chromatography-Triple Quadrupole Mass Spectrometry</b> Marisela D. Mancia, Michelle E. Reid, Evan S. DuBose, James A. Campbell and Kimberly M. Jackson	91
<b>Anti-<i>L. donovani</i> Activity in Macrophage/Amastigote Model of Palmarumycin CP<sub>18</sub> and its Large Scale Production</b> Humberto E. Ortega, Eliane de Morais Teixeira, Ana Rabello, Sarah Higginbotham and Luis Cubilla-Rios	95
<b>Medelamine C, A New <math>\omega</math>-Hydroxy Alkylamine Derivative from Endophytic <i>Streptomyces</i> sp. YIM 66142</b> Ju-Cheng Zhang, Ya-Bin Yang, Hao Zhou, Tian-Feng Peng, Fang-Fang Yang, Li-Hua Xu and Zhong-Tao Ding	99
<b>Enzyme-treated <i>Asparagus officinalis</i> Extract Shows Neuroprotective Effects and Attenuates Cognitive Impairment in Senescence-accelerated Mice</b> Takuya Sakurai, Tomohiro Ito, Koji Wakame, Kentaro Kitadate, Takashi Arai, Junetsu Ogasawara, Takako Kizaki, Shogo Sato, Yoshinaga Ishibashi, Tomonori Fujiwara, Kimio Akagawa, Hitoshi Ishida and Hideki Ohno	101
<b>Anticancer Activity of Binary Toxins from <i>Lysinibacillus sphaericus</i> IAB872 against Human Lung Cancer Cell Line A549</b> Wenjuan Luo, Cuicui Liu, Ruijuan Zhang, Jianwei He and Bei Han	107
<b>The Use of Cycleave PCR for the Differentiation of the Rejuvenating Herb Species <i>Pueraria candollei</i> (White Kwao Khruea), <i>Butea superba</i> (Red Kwao Khruea), and <i>Mucuna macrocarpa</i> (Black Kwao Khruea), and the Simultaneous Detection of Multiple DNA Targets in a DNA Admixture</b> Suchaya Wiriyakarun, Shu Zhu, Katsuko Komatsu and Suchada Sukrong	111
<b>Chemical Compositions and Antimicrobial Activity of the Essential Oils of <i>Hornstedtia havilandii</i> (Zingiberaceae)</b> Siti Erneyanti Hashim, Hasnah Mohd Sirat and Khong Heng Yen	119
<b>Chemical Composition, Antioxidant and Antimicrobial Activity of Essential Oil and Extracts of <i>Tragopogon graminifolius</i>, a Medicinal Herb from Iran</b> Mohammad Hosein Farzaei, Roja Rahimi, Farideh Attar, Farideh Siavoshi, Parastoo Saniee, Mannan Hajimahmoodi, Tahmineh Mirnezami and Mahnaz Khanavi	121
<b>Antinociceptive and Anti-edematous Activities of the Essential Oils of Two Balkan Endemic <i>Laserpitium</i> Species</b> Višnja Popović, Silvana Petrović, Maja Tomić, Radica Stepanović-Petrović, Ana Micov, Milica Pavlović-Drobac, Maria Couladis and Marjan Niketić	125
<b>Chemical Composition of the Essential Oil from <i>Croton kimosorum</i>, an Endemic Species to Madagascar</b> Delphin J. R. Rabehaja, Harilala Ihandriharison, Panja A. R. Ramanolaina, Rakotonirina Benja, Suzanne Ratsimamanga-Urverg, Ange Bighelli, Joseph Casanova and Félix Tomi	129
<b>Intraspecific Variability of the Essential Oil of <i>Cladanthus mixtus</i> from Morocco</b> Anass Elouaddari, Abdelaziz El Amrani, Jamal JamalEddine, José G. Barroso, Luis G. Pedro and Ana Cristina Figueiredo	133
<b>Volatile Organic Compounds of six French <i>Dryopteris</i> Species: Natural Odorous and Bioactive Resources</b> Didier Froissard, Sylvie Rapior, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois and Françoise Fons	137
<b>Essential Oil Compositions of Two Populations of <i>Salvia samuelssonii</i> Growing in Different Biogeographical Regions of Jordan</b> Ammar Bader, Pier Luigi Cioni, Nunziatina De Tommasi and Guido Flamini	141



# Natural Product Communications

## 2014

Volume 9, Number 1

### Contents

<u>Original Paper</u>	<u>Page</u>
<b>New Guaian-type Sesquiterpene from <i>Wikstroemia indica</i></b> Mamoru Kato, Yu-Min He, Dya Fita Dibwe, Feng Li, Suresh Awale, Shigetoshi Kadota and Yasuhiro Tezuka	1
<b>Differences in the Chemical Composition of <i>Arnica montana</i> Flowers from Wild Populations of North Italy</b> Maria Clauser, Nicola Aiello, Fabrizio Scartezzini, Gabriella Innocenti and Stefano Dall'Acqua	3
<b>A New Dolabellane Diterpenoid and a Sesquignan from <i>Aglaia odorata</i> var. <i>microphyllina</i></b> Shuai Liu, Wei Yang, Shou-Bai Liu, Hui Wang, Zhi-Kai Guo, Yan-Bo Zeng, Wen-Hua Dong, Wen-Li Mei and Hao-Fu Dai	7
<b>New Diterpenes from <i>Azorella spinosa</i></b> Luis Astudillo, Margarita Gutiérrez, Luisa Quesada, Aurelio San-Martín, Luis Espinoza and Patricio Peñailillo	9
<b>A New Diterpenoid from the Aerial Parts of <i>Andrographis paniculata</i></b> Chun-Hua Wang, Wen Li, Rui-Xia Qiu, Miao-Miao Jiang and Guo-Qiang Li	13
<b>Isolation of a New Anti-inflammatory 20, 21, 22, 23, 24, 25, 26, 27-Octanorecurbitacin-type Triterpene from <i>Ibervillea sonorae</i></b> Angel Jardón-Delgado, Gil Alfonso Magos-Guerrero and Mariano Martínez-Vázquez	15
<b>Determination of Triterpenic Acids and Screening for Valuable Secondary Metabolites in <i>Salvia</i> sp. Suspension Cultures</b> Sibylle Kümmitz, Christiane Haas, Atanas I. Pavlov, Doris Geib, Roland Ulber, Thomas Bley and Juliane Steingroewer	17
<b>Inhibitory Effect of the Plant <i>Clusia fluminensis</i> against Biological Activities of <i>Bothrops jararaca</i> Snake Venom</b> Eduardo Coriolano de Oliveira, Maria Carolina Anholeti, Thaisa Francielle Domingos, Camila Nunes Faioli, Eladio Flores Sanchez, Selma Ribeiro de Paiva and André Lopes Fuly	21
<b>Chiral Resolution and Absolute Configuration of 3<math>\alpha</math>,6<math>\beta</math>-Dicinnamoyloxytropene and 3<math>\alpha</math>,6<math>\beta</math>-Di(1-ethyl-1<i>H</i>-pyrrol-2-ylcarbonyloxy)tropene, Constituents of <i>Erythroxylum</i> Species</b> Marcelo A. Muñoz, Solange Arriagada and Pedro Joseph-Nathan	27
<b>Aporphine Alkaloids of <i>Cinnamomum mollissimum</i> and their Bioactivities</b> Fatin Fasahah Masnon, Najmah PS Hassan and Farediah Ahmad	31
<b>Antifungal Activity of Metabolites from the Marine Sponges <i>Amphimedon</i> sp. and <i>Monanchora arbuscula</i> against <i>Aspergillus flavus</i> Strains Isolated from Peanuts (<i>Arachis hypogaea</i>)</b> Cynthia Arevabini, Yasmin D. Crivellini, Mariana H. de Abreu, Tamires A. Bitencourt, Mário F. C. Santos, Roberto G. S. Berlinck, Eduardo Hajdu, René O. Belebóni, Ana L. Fachin and Mozart Marins	33
<b>Synthesis of Sepiapterin-C via Hydrolysis of 6-Ethynylpteridine</b> Winston Nxumalo and Andrew Dinsmore	37
<b>Flavonoids Produced by Tissue Culture of <i>Dracaena cambodiana</i></b> Hui Wang, Guanyong Luo, Jiayuan Wang, Haiyan Shen, Ying Luo, Haofu Dai and Wenli Mei	39
<b>Determination of Catechins from <i>Elephantorrhiza elephantina</i> and <i>Pentanisia prunelloides</i> using Voltammetry and UV spectroscopy</b> Smart J. Mpofo, Omotayo A. Arotiba, Lerato Hlekelele, Derek T. Ndinteh and Rui W.M. Krause	41
<b>In vitro Antioxidant Activity, Phenolic Compounds and Protective Effect against DNA Damage Provided by Leaves, Stems and Flowers of <i>Portulaca oleracea</i> (Purslane)</b> Rúben Silva and Isabel S. Carvalho	45
<b>In Vitro Antiviral Activity of a Series of Wild Berry Fruit Extracts against Representatives of <i>Picornia</i>-, <i>Orthomyxo</i>- and <i>Paramyxoviridae</i></b> Lubomira Nikolaeva-Glomb, Luchia Mukova, Nadya Nikolova, Ilian Badjakov, Ivayla Dincheva, Violeta Kondakova, Lyuba Doumanova and Angel S. Galabov	51
<b>Induction of Apoptosis and Cell Cycle Arrest in Human Colon Carcinoma Cells by <i>Corema album</i> Leaves</b> Antonio J. León-González, Margaret M. Manson, Miguel López-Lázaro, Inmaculada Navarro and Carmen Martín-Cordero	55
<b>How to Deal with Nomenclatorial Ambiguities of Trivial Names for Natural Products? – A Clarifying Case Study Exemplified for "Corymbosin"</b> Vatsavaya Ramabharathi and Wolfgang Schuehly	57
<b>Chromatographic Analysis and Antioxidant Capacity of <i>Tabernaemontana catharinensis</i></b> Aline A. Boligon, Mariana Piana, Thiago G. Schawnz, Romaiiana P. Pereira, João B. T. Rocha and Margareth L. Athayde	61
<b>Simultaneous Determination of 13 Chemical Marker Compounds in Gwakhyangjeonggi-san, a Herbal Formula, with Validated Analytical Methods</b> Jung-Hoon Kim, Hyeun-Kyoo Shin and Chang-Seob Seo	65
<b>Single Crystal X-ray Diffraction, Spectroscopic and Mass Spectrometric Studies of Furanocoumarin Peucedanin</b> Magdalena Bartnik, Marta Arczewska, Anna A. Hoser, Tomasz Mroczek, Daniel M. Kamiński, Kazimierz Głowniak, Mariusz Gagoś and Krzysztof Woźniak	71
<b>8-Hydroxycudraxanthone G Suppresses IL-8 Production in SP-C1 Tongue Cancer Cells</b> Arlette S. Setiawan, Roosje R. Owen, Supriatno, Willyanti Soewondo, Sidik and Unang Supratman	75
<b>Antiausterity Activity of Arctigenin Enantiomers: Importance of (2<i>R</i>,3<i>R</i>)-Absolute Configuration</b> Suresh Awale, Mamoru Kato, Dya Fita Dibwe, Feng Li, Chika Miyoshi, Hiroyasu Esumi, Shigetoshi Kadota, and Yasuhiro Tezuka	79

Continued inside backcover