



HAL
open science

The Methylobacterium nodulans / Crotonalaria podocarpa symbiosis: a classic process for an original model

Adeline Renier, Sylvie Rapior

► To cite this version:

Adeline Renier, Sylvie Rapior. The Methylobacterium nodulans / Crotonalaria podocarpa symbiosis: a classic process for an original model. Colloque d'Ecologie Microbienne, Oct 2007, La Grande Motte, France. 2007. hal-02267064

HAL Id: hal-02267064

<https://hal.umontpellier.fr/hal-02267064>

Submitted on 19 Aug 2019


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The *Methylobacterium nodulans* / *Crotalaria podocarpa* symbiosis: a classic process for an original model

A. Renier^{1*}, S. Rapior²

¹Laboratoire des Symbioses Tropicales et Méditerranéennes (LSTM), UMR 113, Campus International de Baillarguet, 34398 Montpellier cedex 5, France.
²Laboratoire de Botanique, Phytochimie et Mycologie, UMR 5175 CEFE, Faculté de Pharmacie, 34093 Montpellier cedex 5, France.
 * adeline.renier@mpl.lrd.fr



Plants belonging to the *Crotalaria* genus are tropical leguminous mainly making symbiosis with *Bradyrhizobium*. Nevertheless, some *Methylobacterium* spp. strains were isolated from root nodules of *Crotalaria* as well as *Lotononis* (*Fabaceae*, *Crotalariae*) (1).

Fig 1. *Crotalaria* plants inoculated with either *Methylobacterium nodulans* ORS 2060 or *Bradyrhizobium* sp. ORS 1810. A: *Crotalaria glaucoides*; B: *C. perrottetii*; C: *C. podocarpa*.

Methylobacterium strains were only isolated from three *Crotalaria* species, i.e., *C. glaucoides* (Fig 1A), *C. perrottetii* (Fig 1B) and *C. podocarpa* (Fig 1C), and were described as a single novel species: *Methylobacterium nodulans*. The main feature of this original bacterial symbiotic partner is its ability to oxidize methanol, a methylophilic property based on the presence of methanol dehydrogenase. During the symbiosis between *M. nodulans* and *C. podocarpa*, it has been shown that the bacterial methylophilic property plays a major role in the symbiotic process (2).






Fig 2. Comparison of growth (5 weeks after inoculation) of *Crotalaria podocarpa*.

Nothing is known on the molecular dialogue occurring between *M. nodulans* and *C. podocarpa*. Commonly, the symbiosis between rhizobial soil bacteria and legumes was described as a multi-step process mediated by signal molecules produced from both two partners: exudation of phenolic compounds (1) able to induce the transcription of bacterial *nod* genes (2) leading to the biosynthesis of a bacterial signal, the nodulation factors (3).





Recently Giraud et al. (3) described photosynthetic *Bradyrhizobium* ORS 278 able to nodulate *Aeschynomene indica* L. (*Fabaceae*) in a Nod factor-independent pathway, opening the possibility that rhizobia could use alternative signals to nodulate legumes.

Which signal molecules are involved in the symbiosis between *Methylobacterium nodulans* and *Crotalaria podocarpa* ?




INDUCTION OF NOD GENES


To monitor the nodulation genes expression, six flavonoids were investigated for their ability to induce the *nodA* promoter of the recombinant strain *M. nodulans* ORS2060 *nodA-LacZ*.




◆ **Apigenin** and **kaempferol** act as strong *nod* gene inducers on strain ORS 2060



◆ For **luteolin**, a 3 times decrease of activity is observed when compared to **apigenin**. More drastically, **quercetin** has no induction effect when compared to **kaempferol**.



◆ **Genistein** and **daidzein** are not able to induce *nod* gene in *M. nodulans*



NODULATION GENES OPERON

A 7,13-kb region in the inserted DNA fragment, showing a positive hybridization signal to the *nodA* probe from *M. nodulans* ORS 2060 was fully sequenced (Fig 4). The analysis of the nucleotide sequence revealed the existence of six entire open reading frames (ORFs).




Fig 4. Functions of the different Nod proteins encoded by the genes belonging to the XmaI fragment.

NODULATION FACTORS

In order to prepare Nod factors, *M. nodulans* ORS 2060 was grown in the presence of apigenin. After induction, Nod factors were extracted from the culture supernatant and purified. HPLC analyses revealed the presence of only two fractions (Fig 6), then analysed by LC/MS.




Fig 6. HPLC profile of a Nod factor extract from an apigenin-inducing *M. nodulans* ORS 2060 culture supernatant.

The positive ion spectra of compounds identified in fraction 1 show two DP5 glucosamine sulfated forms as C_{18:1} (vaccenic acid) and C_{16:0} (palmitic acid) in proportion 10/1 (Fig 7). No Nod factor was found in fraction 2.




Fig 7. Structure of the major lipochito-oligosaccharide identified produced by *M. nodulans* ORS 2060.

◆ *Methylobacterium nodulans* is able to distinguish between flavonoids (flavone and flavonol) and isoflavonoids (isoflavone and isoflavonol). Thus, the attachment of the B ring to C-2 is of crucial importance for induction.

Commonly apigenin and surprisingly kaempferol, usually found as suppressor, are highlighted as to be the most powerful flavonoid inducers of the *nodA* promoter of *M. nodulans*.

◆ The presence of *nod* genes is checked and a DNA fragment containing 8 nodulation genes *nod DABCIJHQ* have been identified.

◆ *M. nodulans* produces one major Nod factor structure identified as to be Nod Mn-V(C_{18:1}S), suggesting a classic symbiosis model for *M. nodulans* / *C. podocarpa* association.

(1) Sy A, Giraud E, Jourand P, et al. (2001) Methylophilic *Methylobacterium* bacteria nodulate and fix nitrogen in symbiosis with legumes. *Journal of Bacteriology* **183**, 214-220.
 (2) Jourand P, Renier A, Rapior S, et al. (2005) Role of methylophilicity during symbiosis between *Methylobacterium nodulans* and *Crotalaria podocarpa*. *Molecular Plant-Microbe Interactions* **18**, 1061-1068.
 (3) Giraud E, Moulin L, Vallenet D, et al. (2007) Legumes symbioses: absence of nod genes in photosynthetic bradyrhizobia. *Science* **316**, 1307-1312.