



HAL
open science

Photoreduction of triplet thioxanthone derivative by azolium tetraphenylborate: a way to photogenerate N-heterocyclic carbenes

Thi Kim Hoang Trinh, Fabrice Morlet-Savary, Julien Pinaud, Patrick Lacroix-Desmazes, Corine Reibel, Cécile Joyeux, Didier Le Nouen, Rémi Métivier, Arnaud Brosseau, Valérie Héroguez, et al.

► To cite this version:

Thi Kim Hoang Trinh, Fabrice Morlet-Savary, Julien Pinaud, Patrick Lacroix-Desmazes, Corine Reibel, et al.. Photoreduction of triplet thioxanthone derivative by azolium tetraphenylborate: a way to photogenerate N-heterocyclic carbenes. *Physical Chemistry Chemical Physics*, 2019, 21 (31), pp.17036-17046. <10.1039/c9cp03098k>. <hal-02285796>

HAL Id: hal-02285796

<https://hal.umontpellier.fr/hal-02285796v1>

Submitted on 12 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Photoreduction of Triplet Thioxanthone Derivative by Azolium Tetraphenylborate: a Way to Photogenerate N-Heterocyclic Carbenes

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Thi Kim Hoang Trinh,^{a,b} Fabrice Morlet-Savary,^{a,b} Julien Pinaud,^c Patrick Lacroix-Desmazes,^c Corine Reibel,^c Cécile Joyeux,^d Didier Le Nouen,^d Rémi Métivier,^e Arnaud Brosseau,^e Valérie Héroguez,^f Abraham Chemtob^{†a,b}

Although N-heterocyclic carbenes (NHCs) have brought profound changes in catalytic organic synthesis, their generation generally requires inert atmosphere and harsh conditions. To overcome these limitations, an air-stable NHC photogenerator has been developed involving two mild components: 1,3-bis(mesityl)imidazolium tetraphenylborate (IMesH⁺BPh₄⁻) and electronically excited isopropylthioxanthone (ITX). In this study, the photochemical mechanism is investigated through the accurate identification of the transient species and photoproducts. Electron transfer reaction between the excited triplet state of ITX and BPh₄⁻ is demonstrated as being the primary photochemical step. Nanosecond laser spectroscopy shows an efficient quenching and the formation of the expected ITX radical anion. The oxidized borane species is not observed, suggesting that this short-lived species could dissociate very rapidly to give phenyl radical – successfully identified using electron paramagnetic resonance – and triphenylborane. As regards the final photoproducts, ¹H and ¹³C NMR spectroscopy support the formation of the targeted NHC, 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene (IMes), suggesting the occurrence of a subsequent proton transfer reaction between ITX radical anion and imidazolium cation (IMesH⁺). Gas chromatography-mass spectrometry reveals three other products: biphenyl, isopropylthioxanthone and ITX. Their formation can be reconciled with a 2-step mechanism of photoinduced electron/proton transfer reactions. ¹¹B NMR spectroscopy demonstrates that the main organoboron photoproduct is diphenylborinic acid formed by oxidation of BPh₃. Due to its Lewis acidity, Ph₂BOH can react with IMes to yield an NHC-boron adduct.

Introduction

Of high importance in photochemistry are arylborates, which are four-coordinate boron compounds bearing a tetrahedral geometry, a formal negative charge, and at least one aryl substituent. Although their photochemical reactivity has been studied since the late 1970s,^{1,2} progress in this field has been made at the cost of intense debates,³ and many unknowns still remain.⁴ Photoreactivity of arylborates is organized around two major types of reactions (**Figure 1**): 1/ *direct photolysis* and 2/ *photoinduced electron transfer (PET)*, in this latter case, the borate anion acts as one electron reducing agent. Recently, other reactions such as photochromic isomerization or photoelimination were described for arylborate molecules with an N,C- or a C,C-chelate backbone.⁵

^a Institut de Science des Matériaux de Mulhouse, IS2M UMR 7361 CNRS, Université de Haute-Alsace, Mulhouse, France.

^b Université de Strasbourg, Strasbourg, France.

^c ICGM, Université de Montpellier, CNRS, ENSCM, Montpellier, France.

^d Laboratoire d'Innovation Moléculaire et Applications, LIMA, Université de Haute-Alsace, France.

^e Laboratoire PPSM, ENS Cachan, CNRS, Université Paris-Saclay, 94235, Cachan, France.

^f Université de Bordeaux, CNRS, Bordeaux INP, LCPO, UMR 5629, F-33600, Pessac, France.

[†] Electronic Supplementary Information (ESI) available. See

DOI: 10.1039/x0xx00000x




Figure 1. Photochemical reactivity of arylborate species. Route 1 is based on direct photolysis while Route 2 is a photoinduced electron transfer (PET) involving an electron acceptor (EA) sensitizer. The asterisk designates an excited state.

Observing that intermediate anion species could abstract acidic protons from water and alcohol,^{7,8} Sun *et al.* proposed in 2008 a tetraphenylborate salt containing as cation the conjugated acid of a nitrogen base (BH^+) such as bicyclic guanidine TBD (TBDH^+).⁶ The base was photogenerated through photoinduced proton abstraction of BH^+ during the photolysis at 254 nm. For example, $\text{TBDH}^+\text{BPh}_4^-$ was used as photobase generator (PBG) for the organocatalyzed ring-opening polymerization of cyclic esters⁶ or thiol-epoxy polymerization.⁹ Several authors added an aryl ketone derivative to extend absorption to near UV-Vis range via a supposed triplet-triplet energy transfer.^{10–13}

Due to their relatively low oxidation potential,¹⁴ arylborates such as Ph_4B^- or triarylmethylborate salts (Ph_3BR^-) were also used as electron donor in *inter-* and *intra-PET reactions* (route 2) with cationic or neutral electron acceptor (EA) sensitizers such as carbocyanine,¹⁵ fullerene,¹⁶ coumarine,¹⁷ fluorone,¹⁸ benzophenone derivatives.¹⁹ A boranyl radical is formed, which in turn undergoes rapid cleavage yielding a triphenylborane and a phenyl (or alkyl) radical. Only alkyl radicals have been applied for initiation of radical polymerizations,¹⁷ or alkylation of the EA.²⁰

Recently, we proposed another application for photoinitiated oxidation of arylborates, the photogeneration of N-heterocyclic carbene (NHC) from an azolium arylborate salt ($\text{NHCH}^+\text{BPh}_4^-$) (**Figure 2**).²¹ When paired with an EA sensitizer such as isopropylthioxanthone (ITX), an electron transfer from the borate (Ph_4B^-) to the excited triplet state of ITX ($^3\text{ITX}^*$) may occur as described above. Unlike the previous case, the azolium cation (NHCH^+) enables a subsequent step of proton abstraction with ITX radical anion $\text{ITX}^{\bullet-}$ (**1**) to take place, yielding the expected free NHC (**2**).²¹ Our recent investigations proved that it was possible to form IMes and its saturated analogue SIMes, the two most employed NHCs.²¹ The utility of this two-component NHC photogenerator $\text{NHCH}^+\text{BPh}_4^-/\text{ITX}$ was also demonstrated in photopolymerization reactions to form polyurethane, polyester and polynorbornene.²² Indeed, although NHCs have

brought profound changes in catalytic organic synthesis,^{23–25} they generally require inert atmosphere and harsh conditions for their generation. As a result, an air-stable system based on mild components, able to generate NHC on demand and on simple UV exposure has the potential to significantly simplify implementation of the broad spectrum of NHC-catalyzed reactions. While it is true that thermally latent NHC precursors also exist,^{26,27} photochemically produced NHCs have net advantages including ambient temperature reaction, thermal stability and the possibility to finely tune NHC concentration upon adjusting the energetic dosage.

Though the mechanistic hypothesis for photoreduction of thioxanthone derivative by arylborates is basically acceptable, conclusive evidences in support of a coupled electron transfer/proton abstraction are required. With exception of the photogenerated NHC molecule (**2**), little is known about the transient species – $^3\text{ITX}^*$, $\text{ITX}^{\bullet-}$ (**1**), ITXH^\bullet (**3**), $\text{Ph}_4\text{B}^\bullet$ (**4**), Ph^\bullet (**5**) – which are supposed to form and the other putative products such as Ph_3B (**6**) and the various thioxanthone derivatives. To provide conclusive evidences in support of our mechanism, it is important to clearly establish the identity of transient species and photoproducts. The structure of the organoboron compound(s) formed is particularly important given BPh_3 is likely to form ate-complexes with NHCs, with strong implications for carbene reactivity.²⁸ Indeed, trivalent boron species typically have Lewis acid characteristics, while NHCs behave as Lewis base. Motivated by this situation, this paper investigates the identity of transient species and photoproducts generated by the irradiation of 1,3-bis(mesityl)imidazolium tetraphenylborate ($\text{IMesH}^+\text{BPh}_4^-$, **Figure 2**) with ITX to form the NHC 1,3-bis(mesityl)imidazol-2-ylidene (IMes). To this purpose, a range of techniques was used: nanosecond laser spectroscopy and electron paramagnetic resonance (EPR) for the detection of the transient species, as well as gas chromatography-mass spectrometry (GC-MS) and ^{11}B NMR spectroscopy for the identification of the photoproducts.




Figure 2. Photochemical pathway for the generation of NHC from a mixture of $\text{NHCH}^+\text{BPh}_4^-$ and ITX.

Experimental

Materials

1,3-Bis(2,4,6-trimethylphenyl)imidazolium chloride (IMesH⁺Cl⁻, 98.0 %, TCI), sodium tetraphenylborate (NaBPh₄, 99.5 %, TCI), N-tert-butyl- α -phenylnitron (PBN, 98.0 %, TCI), diphenyl(2,4,6-trimethylbenzoyl)phosphine oxide (TPO, 98 %, TCI), 2-isopropylthioxanthone (ITX, analytical standard, Aldrich), 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene (IMes, 97%, Aldrich) and carbon disulfide (CS₂, anhydrous, Aldrich) were used as received unless otherwise mentioned. Acetonitrile-*d*₃ (ACN-*d*₃, 99.8 % D, Aldrich), tetrahydrofuran-*d*₈ (THF-*d*₈, 99.5 % D, Eurisotop), acetonitrile (ACN, HPLC grade, VWR) and ethanol (EtOH, HPLC grade, VWR) were dried over 4 Å molecular sieves before use.

Synthesis of IMesH⁺BPh₄⁻

The IMesH⁺BPh₄⁻ was synthesized as described previously.²¹ ¹H NMR (300 MHz, THF-*d*₈), IMesH⁺BPh₄⁻: δ_{ppm} : 2.03 (s, 12H, 4 \times *o*-ArCH₃), 2.37 (s, 6H, 2 \times *p*-ArCH₃), 6.62 – 6.66 (t, 4H, *J* = 6 Hz, 4 \times ArH), 6.75 – 6.80 (t, 8H, *J* = 7.5 Hz, 8 \times ArH), 7.13 (s, 4H, 4 \times ArH), 7.15 (s, 2H, 2 \times NCH) 7.22 – 7.26 (br, 8H, 2 \times 8 \times ArH), and 8.70 (s, 1H, NCHN).

Characterization methods

Nanosecond laser spectroscopy. All nanosecond time-resolved transient absorption spectra were performed in a 1 cm optical path length and monitored by Edinburgh Instruments LP920 laser flash photolysis spectrometer. The solutions were excited at 90° from the probe beam by a Q-switched nanosecond Nd/YAG laser (λ_{exc} = 355 nm, 8 ns pulse duration; energy reduced down to 5 mJ per pulse from Continuum (Surelite II– 10). A filter was used to remove the residual excitation light at 355 nm (Schott GG385). As preliminary step, all acetonitrile solutions were purged with nitrogen for 5 min prior to measurement. The decay of the triplet-triplet state absorption of ITX (1 \times 10⁻⁴ M) at 600 nm was measured and follows a first-order kinetics. The temporal change of triplet concentration [³ITX*] can be represented in equation (1):

$$[{}^3\text{ITX}^*](t) = [{}^3\text{ITX}^*](0) \times e^{-t/\tau_0} \quad (1)$$

where τ_0 is the lifetime of the ITX triplet without quencher.

$$\text{Abs.}(t, 600 \text{ nm}) = \text{Abs.}(0, 600 \text{ nm}) \times e^{-t/\tau_0} \quad (2)$$

Exponential fitting of the experimental decay *Abs.*(*t*, 600 nm) provides τ_0 (Equation (2)). In presence of a triplet quencher (IMesH⁺BPh₄⁻, NaBPh₄ or IMesH⁺Cl⁻), a pseudo-first-order kinetic equation can be applied to extract a triplet

lifetime τ that depends on quencher concentration [Q]. Note that all the triplet lifetimes were determined by using a double exponential equation in the time range 0 – 40 μ s the longer time-constant being due to the formation of the ITX radical anion at the monitored wavelength. By measuring τ at different [Q], the quenching rate constants, k_q , can be obtained by the linear plot based on the Stern-Volmer equation (3).²⁹

$$\tau^{-1} = \tau_0^{-1} + k_q[\text{Q}] \quad (3)$$

Electron paramagnetic resonance (EPR). EPR measurements were performed with a Bruker Elexsys E500 spectrometer with X band frequency in continuous wave (around 9.8 GHz) at room temperature. Spectra were recorded with a modulation amplitude of 1 G, a modulation frequency of 100 kHz and a microwave power of ~ 2 mW. Both Bruker WIN-EPR and SimFonia software were used to note the spectra and carry out the simulation afterwards. In a typical experiment, an acetonitrile solution (1 mL) containing IMesH⁺BPh₄⁻ (9.18 mg, 5 equiv.), ITX (1.27 mg, 3 equiv.) and PBN (0.54 mg, 1 equiv.) was degassed prior to transfer into an aqueous EPR cell. The cell was irradiated for different times (0 s, 30 s and 60 s) with a 365 nm LED light-guide (LC-L1V3, Hamamatsu, 65 mW·cm⁻²) then an EPR spectrum was acquired.

¹H NMR and ¹³C NMR spectroscopy. All ¹H NMR and ¹³C NMR spectra were recorded in appropriate deuterated solvents with tetramethylsilane (TMS) as the internal reference on a Varian Mercury 300 MHz. In a typical measurement, IMesH⁺BPh₄⁻ (9.18 mg, 0.015 mmol, 3 equiv.) and ITX (1.27 mg, 0.005 mmol, 1 equiv.) were charged into a borosilicate NMR tube. The mixture was dissolved in 0.3 mL of acetonitrile-*d*₃ (0.03 M, relative to IMesH⁺BPh₄⁻). Before exposure to a 365 nm LED light spot during 5 min (see details in previous section), the NMR tube was capped with a rubber septum and degassed with N₂. The as-irradiated tube was then analyzed by ¹H NMR. Excess amount of CS₂ (0.02 mL) was added subsequently into the tube. It caused a sudden color change from yellow to red followed by a gradual precipitation. The suspension media was kept for 24 h, and the solid was collected by vacuum filtration and air dried for 24 h. ¹H NMR and ¹³C NMR measurements were performed in DMSO-*d*₆ to confirm the formation of the IMes-CS₂ zwitterion adduct.

¹¹B NMR spectroscopy. All ¹¹B NMR were recorded in THF-*d*₈ with BF₃·Et₂O as the internal reference on a Varian Bruker Avance NEO 500 MHz. All solutions were prepared under argon using a glove box technique. Prior to utilization, quartz NMR tubes were dried under vacuum overnight at 80°C. The THF-*d*₈ solvent was dried over molecular sieves.

Gas chromatography-mass spectrometry (GC-MS). The GC instrument is a Shimadzu GC-2010 model completed with the QP-2010 mass spectrometer system. Helium is the gas vector, used at the constant linear velocity of $25 \text{ cm}\cdot\text{s}^{-1}$. The sample was injected into a Supelco® BP-X5 column (0.15 mm of diameter, 25 m of length and $0.25 \mu\text{m}$ of film thickness) by using split mode with a ratio of 75:25. The temperature of the injector is maintained at 330°C . The column temperature was heated from 70°C to 340°C in two subsequent steps: the first heating state to 250°C with heating rate of $15^\circ\text{C}\cdot\text{min}^{-1}$ and the second to 340°C at the rate of $4^\circ\text{C}\cdot\text{min}^{-1}$. Then, the sample was ionized using an electronic ionization source heated at 200°C . The mass spectrum was recorded from 4 min to end of the program with scan mode from 50 m/z to 600 m/z . Mass Spectra of each peak were compared with spectra of NIST05 and NIST05s databases to identify compounds. In a typical experiment, 2 mL of acetonitrile solution including $\text{IMesH}^+\text{BPh}_4^-$ (1.71 mg, 0.003 mmol, 3 equiv.) and ITX (0.24 mg, 0.001 mmol, 1 equiv.) were placed into a UV quartz cuvette (1 cm optical path) and closed by a rubber septum. The cuvette was degassed with N_2 during 5 min then immediately exposed to a 365 nm LED spot light ($65 \text{ mW}\cdot\text{cm}^{-2}$) for 5 min while keeping the stirring. Finally, the irradiated solution was transfer into Agilent 2 mL glass vial with a screw top (PTFE septum) under inert condition for GC-MS analysis.

Results and discussion

Assignment of transient species

Nanosecond laser spectroscopy

To identify the intermediate species formed in the 2-component system $\text{ITX}/\text{IMesH}^+\text{BPh}_4^-$ (Figure 2), excitation transfer reactions of ITX with different quenchers were recorded by means of a nanosecond laser spectroscopy. Laser flash exposure was performed at 355 nm in deaerated and anhydrous acetonitrile, where only the sensitizer (ITX) absorbs the light. Given the short lifetime of ITX singlet state as well as its high intersystem crossing quantum yield,^{30,31} ITX is assumed to interact mainly via its triplet state. As can be seen in Figure 3a, the transient absorption spectra of ITX alone shows an absorption maximum at 630 nm, fully consistent with the ITX triplet ($^3\text{ITX}^*$) in acetonitrile reported in the literature.^{29,30,32–34} Slow decrease of the transient absorption was attributed to the deactivation process of the $^3\text{ITX}^*$ with a lifetime of $5.7 \mu\text{s}$ at an initial ITX concentration of 10^{-4} M . To study the effect of the azolium cation, the reactivity of the triplet state was evaluated in presence of $\text{IMesH}^+\text{Cl}^-$ since no quenching is expected from Cl^- anion.³⁵ The absorption at 630 nm was weakly quenched when this imidazolium salt was added to a deoxygenated solution of ITX. A low rate constant ($4.7 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$) was determined by constructing a Stern-Volmer plot of the triplet lifetime as a function of $[\text{IMesH}^+\text{Cl}^-]$, with the triplet being monitored at 600 nm (see experimental section for details). A typical decay experimental trace for the triplet decay of ITX in presence of $\text{IMesH}^+\text{Cl}^-$ is given in Figure 4 (trace a), showing clean first-order kinetics. The existence of




Figure 3. Transient absorption spectra of: (a) $[\text{ITX}] = 10^{-4} \text{ M}$ and (b) $\text{ITX}/\text{NaBPh}_4 = 6 \times 10^{-3} \text{ M}$ for 100 ns – 90 μs delay time in N_2 -saturated acetonitrile with excitation wavelength at 355 nm. Delta OD is a change in optical density.

very weak interactions is consistent with the fact that photooxidation of $^3\text{ITX}^*$ by IMesH^+ is thermodynamically unfavorable ($\Delta G_{\text{et}} = +0.74 \text{ eV}$)²¹ and that the triplet energy transfer between ITX ($E_{\text{T}} = 2.77 \text{ eV}$) and IMesH^+ ($E_{\text{T}} = 3.34 \text{ eV}$) exhibits a fairly endothermic character (Fig. S1 in Supporting Information). It can thus be concluded that the azolium cation plays a minor role in the primary photochemical reaction.

Similarly, triplet quenching experiment was carried out with NaBPh_4 to investigate the specific role of the tetraarylborate anion given that Na^+ cannot quench or donate proton. As can be seen in Figure 4 (trace b), the decay rate was much faster, illustrating the more efficient quenching of $^3\text{ITX}^*$ by BPh_4^- . Additionally, it can be noticed that the decay trace does not return to zero in this case; deviation to first order kinetics was attributed to the fact that the triplet absorption probably overlaps with another species. To clarify this point, Figure 3b shows the temporal evolution of the transient absorption spectrum of $\text{ITX}/\text{NaBPh}_4$. After 100 ns, it seems that the narrow 630 nm peak due to triplet absorption has disappeared and has been replaced by a new broad absorption in the region of 560 – 680 nm. Subsequent spectra show the slow decay of this broad peak. We hypothesized that this transient species could be the radical anion $\text{ITX}^{\cdot-}$ generated after photoreduction of $^3\text{ITX}^*$ by BPh_4^- . To support our assignment, we rely first on the fact that the electron transfer for this reaction is slightly exothermic ($\Delta G_{\text{et}} = -0.13$




Figure 4. Decays of ${}^3\text{ITX}^*$ at 600 nm in the presence and absence of: (a) $\text{IMesH}^+\text{Cl}^-$, (b) NaBPh_4 , (c) NaBPh_4 (under O_2) and (d) $\text{IMesH}^+\text{BPh}_4^-$ ($[\text{ITX}] = 10^{-4}$ M, $[\text{IMesH}^+\text{Cl}^-] = [\text{NaBPh}_4] = [\text{IMesH}^+\text{BPh}_4^-] = 6 \times 10^{-3}$ M, respectively, recorded in N_2 -saturated acetonitrile).

eV),²¹ because of the high oxidation potential of the donor anion BPh_4^- . Second, triplet energy transfer, which is the second possible interactions process between ITX ($E_T = 2.77$ eV) and BPh_4^- ($E_T = 3.64$ eV) (Fig. S1), is not energetically favorable. Third, our transition absorption spectrum strongly resembles that of thioxanthone radical anion observed by Schuster *et al.*, who reported also a similar broad absorption at 650 nm.^{36,37} To the best of our knowledge, the spectrum of $\text{ITX}^{\bullet-}$ has not been previously reported. Because of the absorption overlap between ${}^3\text{ITX}^*$ and $\text{ITX}^{\bullet-}$ species, the quenching rate ($4.1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$) was obtained by analyzing only the first short component of the transient absorption decay. Additionally, the triplet transient of ITX in the presence of BPh_4^- was quickly quenched if an oxygen-saturated solution is used, with a typical rate constant in order of $10^9 \text{ M}^{-1} \text{ s}^{-1}$ (Figure 4, trace c).^{38,39} This lends additional confidence to the involvement of ${}^3\text{ITX}^*$ and the operation of a charge transfer mechanism. However, there is no additional band in the transient spectrum that might be attributed to the boranyl radical **4** or other boron species derived from the oxidation of tetraphenyl borate anion. In borate salts, it is well established that back electron is not significant but that boranyl radical can dissociate very rapidly and irreversibly (see next section).⁷ Therefore, the failure to observe $\text{Ph}_4\text{B}^{\bullet}$ (**4**) indicates that a very short-lived character¹⁶ as this radical probably dissociates or rearranges during the laser pulse.

Having investigated the individual role of IMesH^+ and BPh_4^- , quenching experiment was then carried out using $\text{IMesH}^+\text{BPh}_4^-$ (Figure 4, trace d). The similarity of quenching rate obtained by this quencher ($4.9 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$) and NaBPh_4 ($4.1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$) is well indicative that the photoreduction of ITX by BPh_4^- is the primary photochemical process. Additionally, a broad feature characteristic of $\text{ITX}^{\bullet-}$ was also observed in the region of 560 – 680 nm (Fig. S2). However, there is no additional absorption band in the 380 – 430 nm region that could be assigned to a thioxanthyl ketyl radical **3** (ITXH^{\bullet}) derived from subsequent proton transfer step of $\text{ITX}^{\bullet-}$ with IMesH^+ .²⁹ This result can be understood on the basis of a much slower proton transfer step compared to electron




Figure 5. EPR spectra of radical adduct from a solution: (a) $\text{ITX} - \text{NaBPh}_4 - \text{PBN}$ and (b) $\text{ITX} - \text{IMesH}^+\text{BPh}_4^- - \text{PBN}$ after 60 s of irradiation. $[\text{ITX}] = 5 \times 10^{-3}$ M, $[\text{IMesH}^+\text{BPh}_4^-] = 1.5 \times 10^{-2}$ M, $[\text{NaBPh}_4] = 1.5 \times 10^{-2}$ M and $[\text{PBN}] = 3 \times 10^{-3}$ M in acetonitrile.

transfer. The only indication for protonation of the radical anion **1** is the shorter lifetime of $\text{ITX}^{\bullet-}$ in presence of $\text{IMesH}^+\text{BPh}_4^-$ ($29.5 \mu\text{s}$) compared to NaBPh_4 ($40.1 \mu\text{s}$). A final quenching experiment was also carried out using IMes , the targeted NHC. A significant acceleration of the decay at 600 nm is observed, leading to an increased quenching rate ($7.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$). Such a result is consistent with the stronger donor properties of IMes ($E_{\text{ox}} = -0.84$ eV) compared to BPh_4^- ($E_{\text{ox}} = 0.94$ eV).²² It also highlights a more complex mechanism where photogenerated NHC can also interact with triplet ITX and compete with BPh_4^- for charge transfer.

Electron paramagnetic resonance (EPR)

As indicated in the previous section, failure to observe the boranyl radical (**4**) in the nanosecond experiment may indicate that this short-lived species dissociates very rapidly to give the phenyl radical (**5**) and triphenylborane (**6**). Spin trapping and EPR detection (with the trapping reagent α -phenyl-*N*-tert-butyl nitron, PBN) were employed to detect the free radical Ph^{\bullet} formed during in situ irradiation of ITX in presence of tetraphenyl borate anion. In Figure 5, the experimental EPR spectra from $\text{ITX}/\text{NaBPh}_4/\text{PBN}$ (a) and $\text{ITX}/\text{IMesH}^+\text{BPh}_4^-/\text{PBN}$ (b) are shown. Similar in both instances, the EPR spectra show a strong triple-doublet signal suggestive of a single paramagnetic species and gradually increasing with irradiation time (see Fig. S3). Hyperfine coupling constants of the spin adduct that were retrieved from simulation ($a_{\text{N}} = 14.7$ G and $a_{\text{H}} = 2.7$ G, Fig. S4) are in agreement with a PBN spin adduct of the phenyl radical. In addition, these values correspond reasonably well to previously reported values for these spin adduct in literature.^{16,19} This result supports that the transition state of oxidized BPh_4^- could dissociate into Ph^{\bullet} and Ph_3B .^{16,40} Moreover, it is noted that the detection of this phenyl radical **5** is generally very challenging,¹⁶ and failures are generally reported in the literature for both electrochemically⁷ and photochemically⁴¹ induced oxidation of BPh_4^- . By contrast, *n*-butyl radical was more easily detected from a similar oxidation of Ph_3BBu^- species.⁴¹




Figure 6. ^1H NMR spectra change of ITX (0.07 M) and $\text{IMesH}^+\text{BPh}_4^-$ (0.21 M) in ACN-d_3 prior to UV exposure (a) and after 5 min irradiation (b).

Assignment of photoproducts

^1H and ^{13}C NMR spectroscopy

Using ^1H NMR spectroscopy, we found that irradiation of an oxygen-free acetonitrile- d_3 solution containing $\text{IMesH}^+\text{BPh}_4^-$ (1 equiv.) and ITX (3 equiv.) at 365 nm resulted in a complete deprotonation at C2 position (H_b , $\delta = 8.70$ ppm) of the imidazolium cation (**Figure 6**).²¹ This result clearly suggests the generation of IMes (2). Further evidence was given upon adding CS_2 to the as-irradiated medium. A red precipitate forms immediately that was straightforwardly assigned to IMes-CS_2 using ^1H and ^{13}C NMR analysis (Fig. S5). With these results and the identification of $\text{ITX}^{\bullet-}$ described above, it is thus possible to formulate a tentative second step where NHC are created through proton abstraction of IMesH^+ cation by $\text{ITX}^{\bullet-}$ (**Figure 2**). Two methods to determine NHC concentration that were described in previous publications (acid/base titration²¹ and ^1H NMR analysis of NHC-carbodiimide adduct²²) revealed a similar IMesH^+ conversion of 50 % after 5 min irradiation, which is in conflict with NMR data. A partial conversion means that IMes coexists in the reaction medium with its conjugated acid IMesH^+ and may form H-bonded bis(carbene)-proton adduct.^{42–44} Formation of this $\text{C}\cdots\text{H}-\text{C}$ type adduct has been already observed spectroscopically, e.g. UV, NMR.⁴⁴ In our case, evidence for the formation of bis(carbene)-proton adduct is reflected by the slight shift to lower resonance of methylene protons $\text{H}_{e'}$ ($\delta = 2.07$ ppm) and $\text{H}_{d'}$ ($\delta = 2.37$ ppm) observed after irradiation (trace b) compared to the initial state (trace a). Similar upfield shift was evidenced when adding 1 equiv. of free IMes to 9 equiv. of $\text{IMesH}^+\text{BPh}_4^-$ (Fig. S6). In this spectrum, the bridging proton H_b ($\delta = 8.70$ ppm) of the $\text{C}\cdots\text{H}-\text{C}$ complex disappeared entirely due to H-bonding. This confirms that protonation degree cannot be used to quantitatively assess the NHC yield.

GC-MS

To move forward in the identification of other photoproducts, the freshly irradiated photolysis medium was analyzed by GC-MS. Three products were detected: biphenyl,




Figure 7. Side-reactions involving ITX ketyl radical.

isopropylthioxanthene and ITX (Fig. S7 and Table S1). Biphenyl was obtained in good yield and its formation was also observed in other (photo)oxidation experiments of BPh_4^- anion by iron (II) complex,⁴⁰ 1,4-dicyanonaphthalene⁷ or coumarin derivative.⁴¹ However, the formation of biphenyl upon coupling of phenyl radical (detected by EPR) is unlikely. Labelling studies^{7,41} revealed indeed that two phenyl groups of the biphenyl come from the same borate, suggesting that the biphenyl was formed from carbons that were bound to boron. It is therefore reasonable to assume that biphenyl comes from triphenylborane. The most plausible explanation is that Ph^\bullet reacts rapidly with Ph_3B to produce Ph-Ph and $\text{Ph}_2\text{B}^\bullet$.⁴⁰ The fate of the boron radical will be discussed in the ^{11}B NMR section. Since it does not require the addition of oxygen in the reaction medium, biphenyl is thus a primary photochemical product.

In contrast, isopropylthioxanthene and ITX result from the ensuing ground-state chemistry and might derive from ITXH^\bullet (3), the thioxanthyl ketyl radical that is formed after the protonation process of $\text{ITX}^{\bullet-}$, and that we were not able to detect by nanosecond spectroscopy. As described in **Figure 7**, the ketyl radical 3 may undergo two types of reaction: first, a *disproportionation*, giving ITX and isopropylthioxanthol;^{45,46} second, a *dimerization* leading to dithioxanthyl pinacol derivative.⁴⁷ However, except ITX, none of these thioxanthone reduction products were detected by GC-MS. This is apparently due to their thermal instability in the GC injection port as established by Schuster *et al.*³⁶ Isopropylthioxanthol is known to disproportionate upon heating to form ITX and isopropylthioxanthene. Additionally, pinacol can be easily oxidized under air to regenerate ITX.^{36,46} This set of reactions is consistent with the formation of ketyl radical 3 and the thioxanthone products detected in GC-MS. Note also that 1-phenyl-1,4-cyclohexadiene is not detected while it is a major product of the direct photolysis of BPh_4^- in presence of H donors.¹ Its absence confirms that energy transfer from ITX to BPh_4^- is not the predominant mechanism.

dissociation and the subsequent formation of the more stable IMes-CS₂ zwitterionic adduct. By this means, a free boron species was released whose value of ¹¹B chemical shift at 45.2 ppm can be useful for its assignment. Observed⁵³ and calculated ¹¹B NMR spectra⁵⁴ are consistent with the exact chemical shift of diphenylborinic acid Ph₂BOH. Consequently, the singlet signal at 1.1 ppm could be ascribed to IMes-Ph₂BOH adduct present in the photolysis solution. However, the second weak signal at 5.9 ppm could not be assigned. Notably, this second boron-containing complex is not able to form ate-complex with IMes due to the absence of change in the chemical resonance when CS₂ is introduced. To provide further evidence for this assignment, Ph₂BOH was generated in situ by moderate hydrolysis of the Ph₃B in non-dried THF-*d*₈ upon exposing the solution in the NMR tube to atmosphere.⁵⁵ This solution revealed two broad signals centered at 44.9 ppm (Ph₂BOH) and 50.6 ppm (residual Ph₃B) in the ¹¹B NMR spectrum (Figure 9). Addition of IMes into the medium caused the two signals to shift at -9 ppm (Ph₃B-IMes adduct) and 1.1 ppm (IMes-Ph₂BOH adduct). This latter is visible in the spectrum **b** of Figure 8, lending further confidence in our assignment. The last issue to be settled concerns the mechanism leading to the formation of Ph₂BOH. As described in Figure 10, we postulate that the formation of Ph₂BOH may relate to the autooxidation of Ph₃B species. Reaction of Ph₃B with Ph• yields Ph₂B• and Ph₂. Ph₂B• can react subsequently with traces of oxygen (even if argon purged NMR tube were used) to form the peroxy radical Ph₂BOO•. New reaction with Ph₃B might form the peroxide species Ph₂BOOBPh₂, easily decomposed in Ph₂BO•. Ph₂BOH could be formed by hydrogen abstraction with proton donor or after hydrolysis with water.

Conclusions

The photochemical mechanism underlying the release of NHC upon irradiation of IMesH⁺BPh₄⁻ with ITX has been clarified. As summarized in Figure 11, the first step is an electron transfer ($k_q = 4.9 \times 10^7 \text{ M}\cdot\text{s}^{-1}$) between electronically excited ITX and borate anion BPh₄⁻ acting as electron donor to yield the ion pair IMesH⁺/ITX⁻ and the boranyl radical Ph₄B•. This latter is subjected to a rapid decomposition into Ph• and Ph₃B. A subsequent reaction of the ion pair is proton abstraction of IMesH⁺ by ITX⁻ to form the IMes (NHC) and the ketyl radical (ITXH•). Of high importance for NHC reactivity was the fate of organoboron products. Ph₃B and Ph• were proved to undergo a multi-step oxidation reaction, yielding biphenyl and diphenylborinic acid. Ph₂BOH was found to react with the photogenerated IMes to give an IMes-Ph₂BOH ate-complex. While it is difficult to avoid the formation of NHC-boron adduct, our recent investigation showed that such complex did not prevent the activity of its NHC photogenerator in ring-opening methathesis polymerization or for synthesis of polyurethane,²² but could have a detrimental effect on other reactions. Knowledge about the precise photochemical mechanism offers now the possibility for improving the photoinitiating system in a rational way, and create a new generation of NHC photogenerator.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support by French National Research Agency (ANR program: DS0304 2016, contract number: ANR-16-CE07-0016) is gratefully acknowledged.




Figure 11. General mechanistic pathway involved in the formation of IMes during the photolysis of a mixture ITX/ IMesH⁺BPh₄⁻.

Notes and references

- J. L. R. Williams, J. C. Doty, P. J. Grisdale, R. Searle, T. H. Regan, G. P. Happ and D. P. Maier, Boron photochemistry. I. Irradiation of sodium tetraarylborates in aqueous solution, *J. Am. Chem. Soc.*, 1967, **89**, 5153–5157.
- A. Pelter, R. T. Pardasani and P. Pardasani, The Photochemistry of Boron Compounds, *Tetrahedron*, 2000, **56**, 7339–7369.
- J. Radtke, S. K. Mellerup, M. Bolte, H.-W. Lerner, S. Wang and M. Wagner, Aryl Insertion vs Aryl–Aryl Coupling in C,C-Chelated Organoborates: The “Missing Link” of Tetraarylborate Photochemistry, *Org. Lett.*, 2018, **20**, 3966–3970.
- W. G. Santos, J. Pina, D. H. Burrows, M. D. E. Forbes and D. R. Cardoso, New insight into the photophysics and reactivity of trigonal and tetrahedral arylboron compounds, *Photochem. Photobiol. Sci.*, 2016, **15**, 1124–1137.
- T. Baumgartner and F. Jaekle, *Main Group Strategies Towards Functional Hybrid Materials*, John Wiley & Sons, New Jersey, 2018, pp. 87–92.
- X. Sun, J. P. Gao and Z. Y. Wang, Bicyclic Guanidinium Tetraphenylborate: A Photobase Generator and A Photocatalyst for Living Anionic Ring-Opening Polymerization and Cross-Linking of Polymeric Materials Containing Ester and Hydroxy Groups, *J. Am. Chem. Soc.*, 2008, **130**, 8130–8131.
- J. D. Wilkey and G. B. Schuster, Irradiation of tetraphenylborate does not generate a borene anion, *J. Org. Chem.*, 1987, **52**, 2117–2122.
- J. L. R. Williams, J. C. Doty, P. J. Grisdale, T. H. Regan, G. P. Happ and D. P. Maier, Boron photochemistry. II. Irradiation of sodium tetraarylborates in alcohol solutions, *J. Am. Chem. Soc.*, 1968, **90**, 53–55.
- Y. H. Zhao, D. Vuluga, L. Lecamp and F. Burel, Photoinitiated thiol–epoxy addition for the preparation of photoinduced self-healing fatty coatings, *RSC Adv.*, 2016, **6**, 32098–32105.
- Y. Jian, Y. He, Y. Sun, H. Yang, W. Yang and J. Nie, Thiol–epoxy/thiol–acrylate hybrid materials synthesized by photopolymerization, *J. Mater. Chem. C*, 2013, **1**, 4481.
- S. Chatani, T. Gong, B. A. Earle, M. Podgórski and C. N. Bowman, Visible-Light Initiated Thiol–Michael Addition Photopolymerization Reactions, *ACS Macro Lett.*, 2014, **3**, 315–318.
- J. Shin, H. Matsushima, C. M. Comer, C. N. Bowman and C. E. Hoyle, Thiol–Isocyanate–Ene Ternary Networks by Sequential and Simultaneous Thiol Click Reactions, *Chem. Mater.*, 2010, **22**, 2616–2625.
- D. Perrot, C. Croutxé-Barghorn and X. Allonas, UV-curable thio-ether-urethane network with tunable properties, *J. Polym. Sci. Part A: Polym. Chem.*, 2016, **54**, 3119–3126.
- S. Murphy and G. B. Schuster, A Kinetic Method for Determination of Redox Potentials: Oxidation of Tetraarylborates, *J. Phys. Chem.*, 1995, **99**, 511–515.
- S. Chatterjee, P. D. Davis, P. Gottschalk, M. E. Kurz, B. Sauerwein, X. Yang and G. B. Schuster, Photochemistry of carbocyanine alkyltriphenylborate salts: intra-ion-pair electron transfer and the chemistry of boranyl radicals, *J. Am. Chem. Soc.*, 1990, **112**, 6329–6338.
- T. Konishi, Y. Sasaki, M. Fujitsuka, Y. Toba, H. Moriyama and O. Ito, Persistent C60 anion-radical formation via photoinduced electron transfer from tetraphenylborate and triphenylbutylborate, *J. Chem. Soc. Perkin Trans. 2*, 1999, **3**, 551–556.
- R. Popielarz, A. M. Sarker and D. C. Neckers, Applicability of Tetraphenylborate Salts as Free Radical Initiators, *Macromolecules*, 1998, **31**, 951–954.
- A. Y. Polykarpov, S. Hassoon and D. C. Neckers, Tetramethylammonium Tetraorganylborates as Coinitiators with 5,7-Diiodo-3-butoxy-6-fluorone in Visible Light Polymerization of Acrylates, *Macromolecules*, 1996, **29**, 8274–8276.
- S. Hassoon, A. Sarker, A. Y. Polykarpov, M. A. J. Rodgers and D. C. Neckers, Photoinduced Inter- and Intra-Ion-Pair Electron Transfer Reactions in N-(p-Benzoylbenzyl)-N,N,N-tri-n-butylammonium Triphenyl-n-butylborate and -Gallate Salts, *J. Phys. Chem.*, 1996, **100**, 12386–12393.
- J. Y. Lan and G. B. Schuster, Photoalkylation of dicyanoarenes with alkyltriphenylborate salts, *J. Am. Chem. Soc.*, 1985, **107**, 6710–6711.
- J. Pinaud, T. K. H. Trinh, D. Sauvanier, E. Placet, S. Songsee, P. Lacroix-Desmazes, J.-M. Becht, B. Tarabls, J. Lalevée, L. Pichavant, V. Héroguez and A. Chemtob, In Situ Generated Ruthenium-Arene Catalyst for Photoactivated Ring-Opening Metathesis Polymerization through Photolabile N-Heterocyclic Carbene Ligand, *Chem. Eur. J.*, 2018, **24**, 337–341.
- T. K. H. Trinh, J. P. Malval, F. Morlet-Savary, J. Pinaud, P. Lacroix-Desmazes, C. Reibel, V. Héroguez and A. Chemtob, Mixture of azolium tetraphenylborate with isopropylthioxanthone: a new class of N-heterocyclic carbene (NHC) photogenerator for polyurethane, polyester and ROMP polymers synthesis, *Chem. Eur. J.*, 2019, DOI: doi.org/10.1002/chem.201901000.
- M. Scholl, S. Ding, C. W. Lee and R. H. Grubbs, Synthesis and Activity of a New Generation of Ruthenium-Based Olefin Metathesis Catalysts Coordinated with 1,3-Dimesityl-4,5-dihydroimidazol-2-ylidene Ligands, *Org. Lett.*, 1999, **1**, 953–956.
- N. E. Kamber, W. Jeong, S. Gonzalez, J. L. Hedrick and R. M. Waymouth, N-Heterocyclic Carbenes for the Organocatalytic Ring-Opening Polymerization of ϵ -Caprolactone, *Macromolecules*, 2009, **42**, 1634–1639.
- J. Raynaud, W. N. Ottou, Y. Gnanou and D. Taton, Metal-free and solvent-free access to α,ω -heterodifunctionalized poly(propylene oxide)s by N-heterocyclic carbene-induced ring opening polymerization, *Chem. Commun.*, 2010, **46**, 3203.
- B. Bantu, G. M. Pawar, U. Decker, K. Wurst, A. M. Schmidt and M. R. Buchmeiser, CO₂ and Sn^{II} Adducts of N-Heterocyclic Carbenes as Delayed-Action Catalysts for Polyurethane Synthesis, *Chem. Eur. J.*, 2009, **15**, 3103–3109.
- S. Naumann, F. G. Schmidt, W. Frey and M. R. Buchmeiser, Protected N-heterocyclic carbenes as latent pre-catalysts for the polymerization of ϵ -caprolactone, *Polym. Chem.*, 2013, **4**, 4172.
- J. Zhang, N. Pidlynyi, M. Nieger, J. C. Namyslo and A. Schmidt, Zwitterionic borane adducts of N-heterocyclic carbenes from mesomeric betaines of uracil, *Org. Biomol. Chem.*, 2014, **12**, 2737–2744.
- V. Charlot, A. Ibrahim, X. Allonas, C. Croutxé-Barghorn and C. Delaite, Photopolymerization of methyl methacrylate: effects of photochemical and photonic parameters on the chain length, *Polym. Chem.*, 2014, **5**, 6236–6243.
- J. Christmann, X. Allonas, C. Ley, A. Ibrahim and C. Croutxé-Barghorn, Triazine-Based Type-II Photoinitiating System for Free Radical Photopolymerization: Mechanism, Efficiency, and Modeling, *Macromol. Chem. Phys.*, 2017, **218**, 1600597.
- J. Christmann, S. Shi, A. Ibrahim, C. Ley, C. Croutxé-Barghorn, M. Bessières and X. Allonas, Mechanistic Investigation of a Dual Bicyclic Photoinitiating System for Synthesis of Organic–Inorganic Hybrid Materials, *J. Phys. Chem. B*, 2017, **121**, 1972–1981.
- A. Allushi, C. Kutahya, C. Aydogan, J. Kreutzer, G. Yilmaz and Y. Yagci, Conventional Type II photoinitiators as activators for photoinduced metal-free atom transfer radical polymerization, *Polym. Chem.*, 2017, **8**, 1972–1977.
- X. Allonas, C. Ley, C. Bibaut, P. Jacques and J. P. Fouassier, Investigation of the triplet quantum yield of thioxanthone by time-resolved thermal lens spectroscopy: solvent and population lens effects, *Chem. Phys. Lett.*, 2000, **322**, 483–490.

- 34 E. Andrzejewska, D. Zych-Tomkowiak, M. Andrzejewski, G. L. Hug and B. Marciniak, Heteroaromatic Thiols as Co-initiators for Type II Photoinitiating Systems Based on Camphorquinone and Isopropylthioxanthone, *Macromolecules*, 2006, **39**, 3777–3785.
- 35 H. Shizuka and H. Obuchi, Anion-induced triplet quenching of aromatic ketones by nanosecond laser photolysis, *J. Phys. Chem.*, 1982, **86**, 1297–1302.
- 36 S. F. Yates and G. B. Schuster, Photoreduction of triplet thioxanthone by amines: charge transfer generates radicals that initiate polymerization of olefins, *J. Org. Chem.*, 1984, **49**, 3349–3356.
- 37 J. C. Scaiano, Intermolecular photoreductions of ketones, *J. Photochem.*, 1973, **2**, 81–118.
- 38 Amirzadeh, G. and Schnabel, W., On the photoinitiation of free radical polymerization-laser flash photolysis investigations on thioxanthone derivatives, *Makromol Chem*, 1981, **182**, 2821–2835.
- 39 M. Aydin, N. Arsu, Y. Yagci, S. Jockusch and N. J. Turro, Mechanistic Study of Photoinitiated Free Radical Polymerization Using Thioxanthone Thioacetic Acid as One-Component Type II Photoinitiator, *Macromolecules*, 2005, **38**, 4133–4138.
- 40 Y. Nishida, Y.-M. Lee, W. Nam and S. Fukuzumi, Autocatalytic Formation of an Iron(IV)–Oxo Complex via Scandium Ion-Promoted Radical Chain Autoxidation of an Iron(II) Complex with Dioxygen and Tetraphenylborate, *J. Am. Chem. Soc.*, 2014, **136**, 8042–8049.
- 41 A. M. Sarker, Y. Kaneko and D. C. Neckers, Photochemistry and photophysics of novel photoinitiators: N,N,N-tributyl-N-(4-methylene-7-methoxycoumarin) ammonium borates, *J. Photochem. Photobiol. Chem.*, 1998, **117**, 67–74.
- 42 O. Hollóczki, Unveiling the peculiar hydrogen bonding behavior of solvated N-heterocyclic carbenes, *Phys. Chem. Chem. Phys.*, 2016, **18**, 126–140.
- 43 M. Thomas, M. Brehm, O. Hollóczki and B. Kirchner, How Can a Carbene be Active in an Ionic Liquid?, *Chem. Eur. J.*, 2014, **20**, 1622–1629.
- 44 A. J. I. Arduengo, S. F. Gamper, M. Tamm, J. C. Calabrese, F. Davidson and H. A. Craig, A Bis(carbene)-Proton Complex: Structure of a C-H-C Hydrogen Bond, *J. Am. Chem. Soc.*, 1995, **117**, 572–573.
- 45 S. Dadashi-Silab, C. Aydogan and Y. Yagci, Shining a light on an adaptable photoinitiator: advances in photopolymerizations initiated by thioxanthenes, *Polym. Chem.*, 2015, **6**, 6595–6615.
- 46 F. G. Kny-Jones and A. M. Ward, LXXVI.—Investigations on the bivalency of carbon. Part III. Some experiments on xanthhydrol, dixanthhydrol ether, and xanthhydrol chloride, *J. Chem. Soc.*, 1930, **0**, 535–542.
- 47 W. A. Green, *Industrial Photoinitiators: A Technical Guide*, CRC Press, Boca Raton, 1st edn, 2010, pp. 69–70.
- 48 S. T. Murphy, C. Zou, J. B. Miers, R. M. Ballew, D. D. Dlott and G. B. Schuster, Tetraarylborates {[Ar] 4B-}: estimation of oxidation potentials and reorganization energies from electron-transfer rates, *J. Phys. Chem.*, 1993, **97**, 13152–13157.
- 49 B. P. Sullivan, W. J. Dressick and T. J. Meyer, Photoelectrochemical cell based on the ion pair 1,1'-dimethyl-4,4'-bipyridinium bis(tetraphenylborate). Photocurrents from the direct irradiation of a donor-acceptor complex, *J. Phys. Chem.*, 1982, **86**, 1473–1478.
- 50 P. A. Chase, A. L. Gille, T. M. Gilbert and D. W. Stephan, Frustrated Lewis pairs derived from N-heterocyclic carbenes and Lewis acids, *Dalton Trans.*, 2009, **35**, 7179–7188.
- 51 P. A. Chase and D. W. Stephan, Hydrogen and Amine Activation by a Frustrated Lewis Pair of a Bulky N-Heterocyclic Carbene and B(C₆F₅)₃, *Angew. Chem. Int. Ed.*, 2008, **47**, 7433–7437.
- 52 J. J. Eisch, J. H. Shah and M. P. Boleslawski, Skeletal rearrangements of arylborane complexes mediated by redox reactions: thermal and photochemical oxidation by metal ions, *J. Organomet. Chem.*, 1994, **464**, 11–21.
- 53 S. Arkhipenko, Approaches to Novel B-N Chemistry at the Boundary of Frustrated Lewis Pairs and Bifunctional Catalysis, PhD Thesis Durh. Univ., 2017, E-Thesis Online. DOI: 10.15128/et0012213.
- 54 H. S. Rzepa, S. Arkhipenko, E. Wan, M. T. Sabatini, V. Karaluka, A. Whiting and T. D. Sheppard, An Accessible Method for DFT Calculation of ¹¹B NMR Shifts of Organoboron Compounds, *J. Org. Chem.*, 2018, **83**, 8020–8025.
- 55 N. Miyaura, in *Science of Synthesis: Houben-Weyl Methods of Molecular Transformations*, ed. D. E. Kaufmann and D. S. Matteson, Georg Thieme Verlag, Stuttgart, 1st edn, 2014, vol. 6, pp. 677–678.