



HAL
open science

Transmissible cancer and the evolution of sex

Frédéric Thomas, Thomas Madsen, Mathieu Giraudeau, Dorothée Missé,
Rodrigo Hamede, Orsolya Vincze, François Renaud, Benjamin Roche, Beata
Ujvari

► **To cite this version:**

Frédéric Thomas, Thomas Madsen, Mathieu Giraudeau, Dorothée Missé, Rodrigo Hamede, et al..
Transmissible cancer and the evolution of sex. PLoS Biology, 2019, 17 (6), pp.e3000275. 10.1371/jour-
nal.pbio.3000275 . hal-02469626

HAL Id: hal-02469626

<https://hal.umontpellier.fr/hal-02469626>

Submitted on 6 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

ESSAY


Transmissible cancer and the evolution of sex

Frédéric Thomas¹*, Thomas Madsen², Mathieu Giraudeau¹, Dorothée Misse¹, Rodrigo Hamede³, Orsolya Vincze^{4,5}, François Renaud¹, Benjamin Roche¹, Beata Ujvari^{2,3}

1 CREEC/UMR CNRS 5290-IRD 224-Université de Montpellier, Montpellier, France, **2** Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Waurin Ponds, Victoria, Australia, **3** School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia, **4** Hungarian Department of Biology and Ecology, Evolutionary Ecology Group, Babeş-Bolyai University, Cluj-Napoca, Romania, **5** Department of Tisza Research, MTA Centre for Ecological Research-DRI, Debrecen, Hungary

✉ These authors contributed equally to this work.

* rederic.thomas2@ird.fr



Abstract

The origin and subsequent maintenance of sex and recombination are among the most elusive and controversial problems in evolutionary biology. Here, we propose a novel hypothesis, suggesting that sexual reproduction not only evolved to reduce the negative effects of the accumulation of deleterious mutations and processes associated with pathogen and/or parasite resistance but also to prevent invasion by transmissible selfish neoplastic cheater cells, henceforth referred to as transmissible cancer cells. Sexual reproduction permits systematic change of the multicellular organism's genotype and hence an enhanced detection of transmissible cancer cells by immune system. Given the omnipresence of oncogenic processes in multicellular organisms, together with the fact that transmissible cancer cells can have dramatic effects on their host fitness, our scenario suggests that the benefits of sex and concomitant recombination will be large and permanent, explaining why sexual reproduction is, despite its costs, the dominant mode of reproduction among eukaryotes.

OPEN ACCESS

Citation: Thomas F, Madsen T, Giraudeau M, Misse D, Hamede R, Vincze O, et al. (2019) Transmissible cancer and the evolution of sex. *PLoS Biol* 17(6): e3000275. <https://doi.org/10.1371/journal.pbio.3000275>

Published: June 6, 2019

Copyright: © 2019 Thomas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: FT, RH and BU were supported by the ANR (Blanc project TRANSCAN) and by the CNRS (INEE). OV was supported by the Romanian Ministry of Research and Innovation in the form of an Exploratory Research Grant (PN-III-P4-ID-PCE-2016-0404) and a mobility grant (PN-III-P1-1.1-MC-2018-1986). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: HLA, human leukocyte antigen.

Provenance: Not commissioned; externally peer reviewed.

One of the greatest enigma in evolutionary biology is the high prevalence (>99%) of sexual reproduction among eukaryotes [1,2]. Because sexual reproduction requires males that do not produce offspring, an asexual population should consequently reproduce faster than a sexual one [3]. Asexual individuals also benefit from maintaining co-adapted gene complexes and avoid costs involved in mate acquisition [4]. Despite this, the high prevalence of sexual reproduction in the natural world indirectly suggests that the selective forces behind the evolution of sex must be strong and pervasive.

Among the most prominent hypotheses that have been put forward to explain the evolution and maintenance of sexual reproduction, the Fisher–Muller hypothesis proposes that sex may rapidly generate multiple novel advantageous alleles [5–7]. Sexual reproduction will also reduce the deleterious effects of Muller's ratchet, i.e., the build-up and accumulation of deleterious mutations in asexual organisms [8]. Another and probably the most famous hypothesis concerning the benefits of sexual reproduction suggests that recombination create novel

genotypes that are able to resist pathogen and/or parasite infections (i.e., the Red Queen hypothesis) thereby maintaining host fitness despite endlessly evolving virulent pathogens/parasites [9,10]. Several empirical studies support this hypothesis [11–13], e.g., in facultative sexual crustacean *Daphnia magna*, sexually produced offspring were twice as resistant to parasites infecting the parents than asexual ones [14].

All the current hypotheses proposed to explain the evolution of sexual reproduction converge toward the idea that sexual reproduction is beneficial because the genetic diversity it creates provides significant evolutionary advantages to counteract infectious agents, enhance individual intra- and interspecific competition abilities, and alleviate the effects of ongoing fluctuations in environmental conditions [15]. However, what remains unclear is that occasional sex, rather than obligate, could presumably provide the above evolutionary benefits: according to most models, organisms that engage in sexual reproduction only sporadically seem to have the best of both worlds (e.g., [16,17]). Therefore, despite 50 years of research, the selective forces maintaining obligate sex are still not fully understood. Here, we argue that sex has been, and is still, favoured by selection because in contrast to asexual reproduction, it permits to reduce the fitness costs imposed by an ancestral enemy still present: transmissible malignant cell lines.

Multicellular organisms are societies of cooperating clonal cells that have emerged independently on several occasions approximately 1 billion years ago [18,19]. The primary benefit of multicellularity included the division of labor and specialization by differentiated cells [20]. The evolution of multicellular organisms, metazoans, required that individual cells had to forgo their own reproductive interests, i.e., shifting the Darwinian unit of selection from individual cells to the benefit of the entire multicellular community, i.e. the organism. However, one of the first challenges faced by asexual metazoans was, as for any cooperative system (e.g., [21]), the risk of exploitation by internal cheater cells, i.e., cancer cells [22]. Because uncontrolled proliferation of cancer cells is an ubiquitous phenomenon of metazoans [23], it has been proposed to have appeared during the transition from unicellularity to multicellularity [19]. Consequently, the first asexual multicellular organisms did not only have to deal with their own cheater cells but also to evolve adaptations preventing the colonization by infectious malignant cells coming from other individuals. Because anticancer defenses were presumably basic in the first multicellular organisms, both self and infectious cell lines were the major natural enemies. A mile stone in the evolution of metazoans was therefore to counteract and, if possible, to prevent the negative effects of internal cancer cells as well as those caused by non-self invaders, such as viruses, bacteria, parasites, as well as somatic and germ cell parasitism (e.g., in ascidians [24–26]) as well as transmissible cancer cells. These interactions ultimately resulted in the evolution of different evolved defense mechanisms (e.g., different branches and aspects of the immune system) across the animal kingdom. However, in order to reduce the deleterious effects of transmissible cancer cells, the metazoan immune system had to acquire an ability to differentiate between the former and healthy somatic cells.

While the ultimate fate of the vast majority of present malignant cancer cells is to perish with the death of their host, transmissible cancer cells have during the last decades been shown to occur in both invertebrates and vertebrates, often resulting in a massive increase of host mortality (reviewed in [27]). This transmission can involve direct routes such as interindividual aggression (e.g., biting), sexual interactions, and passive transport of transmissible cancer cells. The ability of some of these transmissible cancer cells to avoid immune recognition appears to emanate from a combination of reduced host genetic diversity (that could be the result of bottlenecks and small effective population size [28]) and an ability of the transmissible cancer to down-regulate their antigenic epitopes [27]. Other albeit rare transmissions of cancer cells have been observed in humans from mother to fetus [29,30], i.e., in which the maternally

derived neoplastic cells in the infant had deleted human leukocyte antigen (HLA) alleles suggesting a possible mechanism for immune evasion [30]. Moreover, neoplastic leukemia cells arising in one monozygotic twin, having a single or monochorionic placenta, have been shown to transmit to the co-twin via intraplacental anastomoses [31,32], highlighting the impact of genetic similarity in the successful transmission of cancerous cells. Other organisms, such as the basal metazoan hydra, when reproducing asexually have also shown occurrence of vertical interindividual transmission of tumors [33].

Because clonal reproduction leads to organisms that are identical, we propose that (1) malignant cells produced by the first multicellular organisms were likely to be well adapted to other (identical) organisms, including direct descendants; (2) it was difficult for the victim organisms to recognize (and hence eliminate) transmissible cancer cells that were almost identical to normal somatic cells (i.e., immune evasion). An efficient way to prevent this was to be different from other individuals, and also to produce unique offspring. Organisms adopting sexual reproduction, conversely to clonal ones, form gametes, mix those together, and create progeny with an entirely novel genome. This both limits the chance for clonal infectious malignant cell lines to be already adapted to a novel host and increase the chance that victim organisms can immediately detect the colonization by a transmissible malignant cell, i.e., malignant cells are this time perceived as foreign allograft. Therefore, sexual reproduction could have evolved as an adaptive trait to prevent horizontal and/or vertical transmission of cancer cells (Fig 1).

If the transmission of cancerous cells was a major factor in the evolution of sexual reproduction, the negative effects of such cheaters may also affect “super organisms,” such as social insect colonies, the queen being the gonads, the workers being the somatic cells, a system comparable to that of a multicellular organism [34]. Some social asexual ants and honey bees do develop asexual cheater workers that abandon reproductive self-restraint and reproduce at the expense of other colonies, hence adopting a behavior comparable to selfish transmissible cheater cells in a multicellular organism [35,36]. Moreover, analogous to transmissible cancerous cells, these cheater workers often invade other colonies with devastating consequences for the colony [35,36]. Just like host defense mechanisms (i.e., immune responses), the colonies of superorganisms police against the cheater workers, whereby the queen and/or the workers inhibit the reproduction of cheaters, by either attacking and mutilating recalcitrant workers or by consuming their eggs [37,38].

Empirical testing of theories that explain the evolutionary origin(s) of sex is often complex. However, several observations seem to support our hypothesis:

- i. Although sexual processes undoubtedly antedated multicellularity, no successful transition to multicellularity (i.e., when cancer emerged [33]) has avoided a tight connection with the sexual process [39,40]. Multicellularity may even set the stage for the overall diversity of sexual complexity throughout the Tree of Life [41].
- ii. Species that are not affected by cancer, like prokaryotes and unicellular eukaryotes, should intermittently revert to asexual reproduction. Accordingly, bacteria and archaea reproduce primarily through asexual reproduction, usually by binary fission, with some genetic exchange and recombination occurring occasionally through horizontal gene transfer [42]. The majority of protists and fungi reproduce asexually via fissioning, budding, or spore production [43].
- iii. Unlike animals, plants rarely develop cancer, potentially due to fundamental differences between plant and animal cellular structures, development, and physiology (reviewed in [44]). Plant cells possess rigid cell walls (containing hemicellulose fibers, pectin




Fig 1. Asexual versus sexual reproduction and the transmission of malignant cells. (A) Asexual reproduction maintains high levels of interindividual similarity within a population, and this phenomenon increases the risk of vertical and horizontal transmission of malignant cells. (B) By blending genetics, sexual reproduction produces greater genetic diversity in a population, and as such, limits the transmission of cancer cells across individuals in the population. Genetic diversity facilitates the detection of the invading non-self cells and also limits the chances that the transmissible cancer cells are preadapted to the new host. Thus, cancer cells regularly emerge (e.g., red tumor) in individuals, but unless a “perfect storm” is present, as in the Tasmania devil/devil facial tumor disease system [27], malignant cells fail to be transmitted.

<https://doi.org/10.1371/journal.pbio.3000275.g001>

polysaccharides, and lignin) that maintain strict cellular structure and prevent uncontrolled cell growth. Plant cells also rarely accumulate enough mutations that would lead to cancer, due to their stem cells being hypersensitive to DNA damage and being ready for apoptosis in response to genetic abnormalities. The locomotion of tumor cells is also limited because plants rely on an acellular vascular system (i.e., the xylem and phloem), not on cellular circulatory systems such as blood or lymph vessels. Although plants can occasionally develop tumors, they occur much less often than in animals; they are not metastatic and certainly not as lethal [44,45]. Although plant reproductive strategies are highly diverse [46], many plants exhibit dual reproductive modes, producing both sexual and asexual offspring, being capable of vegetative reproduction (via rhizomes, runners, tubers, bulbils, etc.) and/or of asexual seed production [47,48].

- iv. According to our theory, most asexual species should have a recent evolutionary history, whereas ancient asexual species should possess special adaptations to reduce the deleterious effects of cancer. Close to 50% of asexual lineages have been estimated to be <500,000 years old [49], whereas the remaining 50% of lineages consist of the “evolutionarily scandalous” organisms, such as oribatid mites, darwinulid ostracods, and bdelliid rotifers, that have persisted for millions of generations [49]. The latter species have indeed been shown to be resistant to mutagens such as radiation and heavy metals [50–53], which indicates high resistance to oncogenic processes and selection of tumor-suppressor mechanisms that enable the survival of these ancient asexual lineages.
As a corollary, one might predict that recently evolved asexual species should be affected by cancer at higher frequency than their sexual conspecifics, unless they also have evolved efficient anticancer defenses. Further studies would be necessary to test these hypotheses.
- v. Multicellular eukaryotes that are strongly impacted by malignant cell emergence and proliferation should mostly have obligate sexual reproduction. Obligate sex is indeed the dominant mode of reproduction in many lineages of complex eukaryotes [1].
- vi. Transmissible cancers should be rare in species practising sexual reproduction. Although we probably underestimate their prevalence [54], only 4 cases of transmissible cancers are currently known in the wild, supporting the idea that the evolution of transmissible cancer in sexually reproducing species is very rare and occurs only under very particular conditions (e.g., the “perfect storm hypothesis” [27]).

There are different possibilities to experimentally test our hypothesis. For instance, we predict that in organisms reproducing both by sexually and asexually, a shift toward more sexual reproduction should be observed following the emergence and progression of malignant cells. Hydra has the ability to switch between sexual and asexual reproduction and the propensity to develop tumors, therefore it could be a good candidate to test this hypothesis [33]. In accordance with our hypothesis, parental tumors are almost systematically transmitted to daughter polyps of hydra when reproduction is asexual (i.e., budding results in the vertical transmission of tumors), whereas offspring resulting from sexual reproduction are tumor free [33]. Demonstrating that tumor-bearing hydra, compared with healthy ones, preferentially reproduce sexually would provide support to our hypothesis. Because tumors can also be experimentally transplanted between polyps, this biological system also offers the possibility to test whether transplanted tumors establish better when recipient polyps are identical to the donor with tumors, compared with when they are different individuals.

Constant progress in animal cloning [55] could also help to evaluate the risk of cancer cell transmission associated with asexual reproduction. We predict that the likelihood of mother to fetus malignant cell transmission will be higher when the implanted embryos (e.g., in mammals) are genetically identical to their mother, compared with embryos that have originated from another female or are the mother’s natural embryos.

Comparative oncology approaches could also provide in depth analyses of the difference in anticancer defenses between recent and ancient asexual species, as well as in comparison with their sexual relatives. From a theoretical perspective, our hypothesis could be tested through developing new theoretical frameworks based on mathematical models so far used to elucidate the “Red Queen” hypothesis [56,57]. However, current mathematical models applied to host–pathogen interactions rarely consider parasite diversity. In the case of our hypothesis, future theoretical extensions will need to consider the fact that host diversity generated by sexual reproduction decreases the probability of cancer cell transmission and thus *de facto* reduces the diversity of the cancer cells that can be transmitted, which concomitantly allows the host’s

immune system to be more efficient in eliminating them. Although including "parasite diversity" is generally mathematically challenging, combining recent theoretical developments in studying multistrain pathogens [56] with the Red Queen models would be an interesting first attempt.

Concluding remarks

Although selfish neoplastic cells are omnipresent cheaters in all multicellular organisms [58], the among-individual transmission of such cells requires a "perfect storm" in sexual reproducing organisms with an optimal confluence of multiple host and tumor cell traits [27]. A major constraint of such transmissions requires an ability of transmissible cancerous cells to evade immunological histocompatibility barriers. Because asexual reproduction results in clonal, often identical organisms, asexual organisms and their progeny would be susceptible to the invasion of clonal transmissible cancer cells. Conversely, due to its enhanced among-individual genetic heterogeneity and concomitant increased ability to detect non-self cells, sexual reproduction should significantly reduce the risk of among-individual transmission of such cancerous cells. Given the ubiquity of oncogenic processes in the multicellular world together with the diversity of potential transmission routes, sexual reproduction, despite its associated costs, may consequently have been favored as a less risky, more profitable option to produce viable offspring, i.e., less subjective to transmissible cancers. To our knowledge, this selective scenario for the initial evolution of sex across the Tree of Life is novel. As illustrated, e.g., with the human twin example above, it also explains its continued maintenance despite the significant evolutionary costs. Also, the experimental approaches we proposed above should permit the evaluation of the critical role transmissible cancers play in shaping animal reproductive strategies.

In conclusion, we propose that the prevalence of sex in eukaryotes is a ghost of a past apogee of transmissible cancerous cell lines in the first asexual multicellular organisms. Although natural selection found a way to radically reduce the prevalence of transmissible cancer, sexual organisms are, however, still subjected to the deleterious effects of internal malignant cells. We hope that this paper will pave the way for a novel research direction on the evolutionary enigma of sex.

References

1. Bell G. *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. Berkeley, CA: University of California Press.; 1982.
2. Hartfield M, Keightley PD. Current hypotheses for the evolution of sex and recombination. *Integrative zoology*. 2012; 7(2):192–209. Epub 2012/06/14. <https://doi.org/10.1111/j.1749-4877.2012.00284.x> PMID: 22691203.
3. Maynard Smith J. *The evolution of sex*. Cambridge, UK: Cambridge University Press.; 1978.
4. Lehtonen J, Jennions MD, Kokko H. The many costs of sex. *Trends in Ecology & Evolution*. 2012; 27(3):172–8. <https://doi.org/10.1016/j.tree.2011.09.016> PMID: 22019414
5. Fisher RA. *The genetical theory of natural selection*. Oxford, UK: The Clarendon Press.; 1930.
6. Muller HJ. Some genetic aspects of sex. *The American Naturalist*. 1932; 66(703):118–38. <https://doi.org/10.1086/280418>
7. Williams GC. *Sex and evolution*. Princeton, N.J: Princeton University Press; 1975.
8. Muller HJ. The relation of recombination to mutational advance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 1964; 1(1):2–9. [https://doi.org/10.1016/0027-5107\(64\)90047-8](https://doi.org/10.1016/0027-5107(64)90047-8).
9. Hamilton WD, Axelrod R, Tanese R. Sexual reproduction as an adaptation to resist parasites (a review). *Proceedings of the National Academy of Sciences*. 1990; 87(9):3566.
10. Jaenike J. A hypothesis to account for the maintenance of sex in populations. *Evolutionary Theory*. 1978; 3:191–194. <https://www.mn.uio.no/cees/english/services/van-valen/evolutionary-theory/volume->

[3/vol-3-no-4-pages-191-194-j-jaenike-an-hypothesis-to-account-for-the-maintenance-of-sex-within-populations.pdf](#)

11. Decaestecker E, Gaba S, Raeymaekers JAM, Stoks R, Van Kerckhoven L, Ebert D, et al. Host–parasite ‘Red Queen’ dynamics archived in pond sediment. *Nature*. 2007; 450:870. <https://doi.org/10.1038/nature06291> PMID: 18004303
12. Jokela J, Dybdahl Mark F, Lively Curtis M. The maintenance of sex, clonal dynamics, and host-parasite coevolution in a mixed population of sexual and asexual snails. *The American Naturalist*. 2009; 174(S1):S43–S53. <https://doi.org/10.1086/599080> PMID: 19441961
13. Vergara D, Jokela J, Lively CM. Infection dynamics in coexisting sexual and asexual host populations: support for the Red Queen hypothesis. *The American Naturalist*. 2014; 184(S1):S22–S30. <https://doi.org/10.1086/676886> PMID: 25061675
14. Auld SKJR Tinkler SK, Tinsley MC. Sex as a strategy against rapidly evolving parasites. *Proceedings of the Royal Society B: Biological Sciences*. 2016; 283(1845).
15. Colegrave N. The evolutionary success of sex. *Science & Society Series on Sex and Science*. EMBO reports. 2012; 13(9):774–8. Epub 2012/08/07. <https://doi.org/10.1038/embor.2012.109> PMID: 22868665.
16. Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on neutral molecular variation. *Genetics*. 1993; 134(4):1289. PMID: 8375663
17. D’Souza TG, Michiels NK. The costs and benefits of occasional Sex: theoretical predictions and a case study. *Journal of Heredity*. 2010; 101(suppl_1):S34–S41. <https://doi.org/10.1093/jhered/esq005> PMID: 20212007
18. Aktipis CA, Boddy AM, Jansen G, Hibner U, Hochberg ME, Maley CC, et al. Cancer across the tree of life: cooperation and cheating in multicellularity. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2015; 370(1673):20140219. <https://doi.org/10.1098/rstb.2014.0219> PMID: 26056363.
19. Szathmáry E, Maynard Smith J. The major evolutionary transitions. *Nature* 1995; 374(6519): 227–232. <https://doi.org/10.1038/374227a0> PMID: 7885442
20. Smith JM, Szathmáry E. *The major transitions in evolution*: OUP Oxford; 1997.
21. West SA, Griffin AS, Gardner A. Evolutionary explanations for cooperation. *Current Biology*. 2007; 17(16):R661–R72. <https://doi.org/10.1016/j.cub.2007.06.004> PMID: 17714660
22. Grosberg RK, Strathmann RR. The evolution of multicellularity: a minor major transition? *Annual Review of Ecology, Evolution, and Systematics*. 2007; 38(1):621–54. <https://doi.org/10.1146/annurev.ecolsys.36.102403.114735>
23. Madsen T, Arnal A, Vittecoq M, Bernex F, Abadie J, Labrut S, et al. Chapter 2—Cancer prevalence and etiology in wild and captive animals. In: Ujvari B, Roche B, Thomas F, editors. *Ecology and Evolution of Cancer*: Academic Press; 2017. p. 11–46.
24. Buss LW. Somatic cell parasitism and the evolution of somatic tissue compatibility. *Proceedings of the National Academy of Sciences of the United States of America*. 1982; 79(17):5337–41. <https://doi.org/10.1073/pnas.79.17.5337> PMID: 6957867.
25. Stoner DS, Weissman IL. Somatic and germ cell parasitism in a colonial ascidian: Possible role for a highly polymorphic allorecognition system. *Proceedings of the National Academy of Sciences*. 1996; 93(26):15254. <https://doi.org/10.1073/pnas.93.26.15254> PMID: 8986797
26. Weiss RA. Open questions: knowing who’s who in multicellular animals is not always as simple as we imagine. *BMC Biology*. 2018; 16(1):115. <https://doi.org/10.1186/s12915-018-0582-8> PMID: 30322384
27. Ujvari B, Gatenby RA, Thomas F. The evolutionary ecology of transmissible cancers. *Infection, Genetics and Evolution*. 2016; 39:293–303. <https://doi.org/10.1016/j.meegid.2016.02.005> PMID: 26861618
28. Ujvari B, Klaassen M, Raven N, Russell T, Vittecoq M, Hamede R, et al. Genetic diversity, inbreeding and cancer. *Proceedings of the Royal Society B: Biological Sciences*. 2018; 285(1875):20172589. <https://doi.org/10.1098/rspb.2017.2589> PMID: 29563261
29. Greaves M, Hughes W. Cancer cell transmission via the placenta. *Evolution, Medicine, and Public Health*. 2018; 2018(1):106–15. <https://doi.org/10.1093/emphe/eoy011> PMID: 29765597
30. Isoda T, Ford AM, Tomizawa D, van Delft FW, De Castro DG, Mitsuiki N, et al. Immunologically silent cancer clone transmission from mother to offspring. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106(42):17882–5. Epub 2009/10/12. <https://doi.org/10.1073/pnas.0904658106> PMID: 19822752.
31. Clarkson B, Boyse E. Possible explanation of the high concordance for acute leukemia in monozygotic twins. *The Lancet*. 1971; 297(7701):699–701. [https://doi.org/10.1016/S0140-6736\(71\)92705-X](https://doi.org/10.1016/S0140-6736(71)92705-X).

32. Greaves MF, Maia AT, Wiemels JL, Ford AM. Leukemia in twins: lessons in natural history. *Blood*. 2003; 102(7):2321. <https://doi.org/10.1182/blood-2002-12-3817> PMID: 12791663
33. Domazet-Lošo T, Klimovich A, Anokhin B, Anton-Erxleben F, Hamm MJ, Lange C, et al. Naturally occurring tumours in the basal metazoan Hydra. *Nature Communications*. 2014; 5:4222. <https://doi.org/10.1038/ncomms5222> PMID: 24957317
34. Hölldobler B, Wilson EO. *The superorganism: the beauty, elegance, and strangeness of insect societies*. New York, USA: W. W. Norton Press; 2008.
35. Dobata S, Sasaki T, Mori H, Hasegawa E, Shimada M, Tsuji K. Persistence of the single lineage of transmissible 'social cancer' in an asexual ant. *Molecular Ecology*. 2010; 20(3):441–55. <https://doi.org/10.1111/j.1365-294X.2010.04954.x> PMID: 21155912
36. Goudie F, Oldroyd BP. Thelytoky in the honey bee. *Apidologie*. 2014; 45(3):306–26. <https://doi.org/10.1007/s13592-013-0261-2>
37. Foster KR, Ratnieks FL. Convergent evolution of worker policing by egg eating in the honeybee and common wasp. *Proceedings Biological sciences*. 2001; 268(1463):169–74. <https://doi.org/10.1098/rspb.2000.1346> PMID: 11209887.
38. Ratnieks FLW, Visscher PK. Worker policing in the honeybee. *Nature*. 1989; 342(6251):796–7. <https://doi.org/10.1038/342796a0>
39. Domazet-Lošo T, Tautz D. Phylostratigraphic tracking of cancer genes suggests a link to the emergence of multicellularity in metazoa. *BMC biology*. 2010; 8:66–. <https://doi.org/10.1186/1741-7007-8-66> PMID: 20492640.
40. Speijer D, Lukeš J, Eliáš M. Sex is a ubiquitous, ancient, and inherent attribute of eukaryotic life. *Proceedings of the National Academy of Sciences of the United States of America*. 2015; 112(29):8827–34. Epub 2015/07/20. <https://doi.org/10.1073/pnas.1501725112> PMID: 26195746.
41. Hanschen ER, Herron MD, Wiens JJ, Nozaki H, Michod RE. Multicellularity Drives the Evolution of Sexual Traits. *The American Naturalist*. 2018; 192(3):E93–E105. <https://doi.org/10.1086/698301> PMID: 30125231
42. Arber W. *Horizontal Gene Transfer among Bacteria and Its Role in Biological Evolution*. Life (Basel, Switzerland). 2014; 4(2):217–24. <https://doi.org/10.3390/life4020217> PMID: 25370194.
43. Otto Sarah P. The evolutionary enigma of sex. *The American Naturalist*. 2009; 174(S1):S1–S14. <https://doi.org/10.1086/599084> PMID: 19441962
44. Doonan JH, Sablowski R. Walls around tumours—why plants do not develop cancer. *Nature Reviews Cancer*. 2010; 10:794. <https://doi.org/10.1038/nrc2942> PMID: 20966923
45. Doonan J, Hunt T. Why don't plants get cancer? *Nature*. 1996; 380(6574):481–2. <https://doi.org/10.1038/380481a0> PMID: 8606760
46. C H Barrett S. Barrett SCH. The evolution of plant sexual diversity. *Nat Rev Gen* 3: 274–284 2002. 274–84 p.
47. Eckert CG. The loss of sex in clonal plants. *Evolutionary Ecology*. 2001; 15(4):501–20. <https://doi.org/10.1023/A:1016005519651>
48. Richards A. Breeding systems in flowering plants and the control of variability. *Folia Geobotanica*. 1996; 31(3):283–293.
49. Neiman M, Meirmans S, Meirmans PG. What can asexual lineage age tell us about the maintenance of sex? *Annals of the New York Academy of Sciences*. 2009; 1168(1):185–200. <https://doi.org/10.1111/j.1749-6632.2009.04572.x> PMID: 19566708
50. Krisko A, Leroy M, Radman M, Meselson M. Extreme anti-oxidant protection against ionizing radiation in bdelloid rotifers. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(7):2354–7. Epub 2012/01/26. <https://doi.org/10.1073/pnas.1119762109> PMID: 22308443.
51. Luo W, Verweij RA, van Gestel CAM. Toxicity of Pb contaminated soils to the oribatid mite *Platynothrus peltifer*. *Ecotoxicology*. 2015; 24(5):985–90. <https://doi.org/10.1007/s10646-015-1439-3> PMID: 25757529
52. Skubala P, Zaleski T. Heavy metal sensitivity and bioconcentration in oribatid mites (Acari, Oribatida): Gradient study in meadow ecosystems. *Science of The Total Environment*. 2012; 414:364–72. <https://doi.org/10.1016/j.scitotenv.2011.11.006> PMID: 22134027
53. Vandenbroecke L, Vanfleteren J, Martens K, Schön I. Hurdles in investigating UVB damage in the putative ancient asexual *Darwinula stevensoni* (Ostracoda, Crustacea) 2013. 106–18 p.
54. Ujvari B, Gatenby RA, Thomas F. Transmissible cancers, are they more common than thought? *Evolutionary applications*. 2016; 9(5):633–4. <https://doi.org/10.1111/eva.12372> PMID: 27398093.
55. Cibelli J, Gurdon J, Wilmut I, Jaenisch R, Lanza R, West MD, et al. *Principles of Cloning (Second Edition)*. San Diego: Academic Press; 2014 2014/01/01/.

56. Roche B, Drake JM, Rohani P. An agent-based model to study the epidemiological and evolutionary dynamics of Influenza viruses. *BMC bioinformatics*. 2011;12:87-. <https://doi.org/10.1186/1471-2105-12-12> PMID: [21450071](https://pubmed.ncbi.nlm.nih.gov/21450071/).
57. Salathé M, Kouyos RD, Bonhoeffer S. The state of affairs in the kingdom of the Red Queen. *Trends in Ecology & Evolution*. 2008; 23(8):439–45. <https://doi.org/10.1016/j.tree.2008.04.010>.
58. Thomas F, Vavre F, Tissot T, Vittecoq M, Giraudeau M, Bernex F, et al. Cancer is not (only) a senescence problem. *Trends in Cancer*. 2018; 4(3):169–72. <https://doi.org/10.1016/j.trecan.2018.01.002> PMID: [29506667](https://pubmed.ncbi.nlm.nih.gov/29506667/)