



Natural resistance to cancers: a Darwinian hypothesis to explain Peto's paradox

Benjamin Roche, Michael Hochberg, Aleah F Caulin, Carlo C Maley, Robert Gatenby, Dorothée Missé, Frederic Thomas

► To cite this version:

Benjamin Roche, Michael Hochberg, Aleah F Caulin, Carlo C Maley, Robert Gatenby, et al.. Natural resistance to cancers: a Darwinian hypothesis to explain Peto's paradox. *BMC Cancer*, 2012, 12, pp.387. 10.1186/1471-2407-12-387 . hal-02502607

HAL Id: hal-02502607

<https://hal.umontpellier.fr/hal-02502607>

Submitted on 9 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

DEBATE

Open Access

Natural resistance to cancers: a Darwinian hypothesis to explain Peto's paradox

Benjamin Roche^{1,8*}, Michael E Hochberg^{2,3,8}, Aleah F Caulin⁴, Carlo C Maley⁵, Robert A Gatenby⁶, Dorothée Misse^{7,8} and Frédéric Thomas^{7,8}

Abstract

Background: Peto's paradox stipulates that there is no association between body mass (a surrogate of number of cells and longevity) and cancer prevalence in wildlife species. Resolving this paradox is a very promising research direction to understand mechanisms of cancer resistance. As of present, research has been focused on the consequences of these evolutionary pressures rather than of their causes.

Discussion: Here, we argue that evolution through natural selection may have shaped mechanisms of cancer resistance in wildlife species and that this can result in a threshold in body mass above which oncogenic and tumor suppressive mechanisms should be increasingly purified and positively selected, respectively.

Summary: We conclude that assessing wildlife species in their natural ecosystems, especially through theoretical modeling, is the most promising way to understand how evolutionary processes can favor one or the other pathway. This will provide important insights into mechanisms of cancer resistance.

Background

One of the challenges to multicellular organisms is the deregulation of the cooperative interactions between component cells that natural selection has taken long periods to produce. Cancer is one large class of such multicellular deregulation often resulting in death [1], and poses the puzzle of how or whether it is opposed by natural selection. Although assessing natural selection in human populations is challenging, indications of how natural selection may have moulded human cancer incidence can be assessed by comparing incidences of cancers in wildlife species [2]. Specifically, we need to understand the mechanisms underlying the so-called "Peto's paradox"—the absence of a correlation across species between cancer and body size or longevity [2,3], Figure 1. If each dividing cell in a multicellular organism has the same probability of initiating a malignant neoplasm, then all else being equal, the more cells an

organism has, the greater the chance of cancer emerging. Thus, blue whales which, at adulthood, may weigh more than 100 metric tons, should be at least 1000 times more likely to develop a cancer than humans. Moreover, transitions to malignancy are expected to increase with the number of cell divisions—that is with organism lifespan. Numerous studies have shown correlations between longevity and body size, making Peto's paradox all the more difficult to resolve [2]. How do large, long-lived species overcome the burden of cancer? We [2] hypothesized that organisms evolving large body sizes and long life spans, concomitantly evolved mechanisms to offset increased cancer risk. These may include slower somatic cell turnover, redundancy of tumor suppressor genes, more efficient immune systems, better suppression of inflammation or resistance to oncogenic viruses. Currently, no clear consensus has emerged about which of these mechanisms, if any, are the products of natural selection. It is therefore crucial to focus more attention on the evolutionary causes of Peto's paradox.

Discussion

Here, we propose that oncogenic or tumor suppressive mechanisms should be increasingly purified or positively selected, respectively, above a given threshold in body

* Correspondence: benjamin.roche@ird.fr

¹IRD, UMMISCO (UMI IRD/UPMC), 32, avenue Henry Varagnat 93143 Bondy Cedex France

²CREEC, Université Montpellier 2, CC 06003, 95 rue de la Galéra, 34095 Montpellier Cedex 5, France

Full list of author information is available at the end of the article




Figure 1 Summary of the current knowledge on Peto's paradox according to the spare data available reviewed in [2]. Body mass seems to be a relevant surrogate for cancer prevalence at an intra-species level, but not an inter-species level.

mass (Figure 1). There is correlative evidence for a within-species increase in cancer frequency with body mass [4]. For instance, cohort studies in humans have shown that leg lengths 3–4 mm above the average is associated with a 80% higher risk of non-smoking cancers (Figure 1; reference 4, but see 5). Despite the prediction that more cells in a body should be associated with a higher probability of cancer, current knowledge indicates that cancers do not increase across wildlife species as a function of body mass (incidences are typically between 20 and 40%, 2).

We suggest that these contrasting patterns are produced by large wildlife species undergoing stronger selection against cell deregulation and for tumor suppression than smaller ones. Analyzing the selective pressures within a species' ecosystem (i.e., biotic and abiotic environments) will inform if natural selection targets oncogenic and/or tumor suppressive mechanisms. Identifying patterns across species in cancer resistance evolution would be an important insight into resolving Peto's paradox.

We need to understand the selective (biotic and abiotic) landscapes in which species evolved and continue to evolve. Indeed, natural selection is the product of how environments favor specific heritable phenotypes. Cancer vulnerability amongst wildlife species is likely to have been shaped by natural selection, depending on which fitness-reducing risks predominate (somatic diseases including cancer, infectious and parasitic diseases, predation and adverse environmental conditions). For instance, small rodents *in natura* may succumb to cancer, but only if they do not first die from any one of numerous other causes, such as predators, infectious

diseases, or environmental vagaries such as floods, temperature extremes, etc. Natural selection will tend to promote resistance to sources of mortality prior to reproduction, meaning that for blue whales to grow so large and live so long, they need to both develop defenses against predators and resistance to somatic diseases like cancer. There is clearly a chicken-and-egg problem here, since changes in body size, longevity and life history strategy will alter the selective influence of different mortality factors, including the probability of cancer emergence.

Studying this complexity (i.e., numerous environmental factors acting and interacting in opposite directions and/or with reciprocal effects (Figure 2)) requires the development of a theoretical approach. Adopting a quantitative framework, such as adaptive dynamics [5], widely applied to understanding the evolution of pathogens and life-history traits, can help understand how different biotic and abiotic selective pressures affect trait evolution and especially those involved in cancer protection.

Since wildlife species are subjected to a large variety of selective pressures and are found in a diverse range of habitats, it should be possible to use comparative genomics [6] to understand how proto-oncogenes and TSGs covary with certain environmental characteristics. Indeed, comparing genomic regions of interest for cancer research, e.g., proto-oncogenes or Tumor Suppressor Genes widespread in mammals, according to the biotic and abiotic environments where these species are found can give important insights into how these classes of genes have been shaped by natural selection as a function of the environment.




Figure 2 The network of interactions between cancer and various variables acting on individuals and species in ecosystems. Arrows represent direct (e.g., oncogenic pathogens cause cancer) or evolutionary (e.g., predation exert selective pressure on parasite virulence) responses. Arrows are in two colors for clarity (blue for unidirectional relationships, dark for reciprocal, or potentially reciprocal, relationships).

Addressing these considerations is undoubtedly relevant for human populations living in different environmental conditions (e.g., presence or absence of pathogens). Indeed, evidence suggests that many human populations lack alleles with enhanced protection against certain cancers, possibly because their short life-spans have precluded selection for those alleles [4,7,8].

Summary

Given the complex nature of interactions arbitrating cancer (Figure 2), mathematical and statistical approaches will be necessary to tease apart causal mechanisms and their interactions [9-19], especially for the optimal use of chemotherapies [20]. In particular, quantitative approaches will help understand the selective forces acting on oncogene dynamics in animals in nature. Although cancer is an important cause of morbidity and mortality in several wildlife species [21], data are difficult to collect because most wild animals live and die unseen [21]. In this context, quantitative approaches will be of considerable interest to understand and predict wildlife cancers, and to see to what extent insights can instruct on the prevention of cancer in humans.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BR, MEH and FT conceived the study, BR, MEH, AFC, CCM RAG, DM, and FT wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work is a contribution from the "Centre de Recherche Ecologique et Evolutive sur le Cancer" (CREEC) and the PEPII program funded by the CNRS.

BR, MEH and FT thank the RNSC (Réseau National des Systèmes Complexes) for financial support.

Author details

Author details
1IRD, UMMISCO (UMI IRD/UPMC), 32, avenue Henry Varagnat 93143 Bondy Cedex France. 2ISEM, Institut des Sciences de l'Evolution, Université Montpellier 2, CNRS Case Postal 65, Place Eugène Bataillon 34095 Montpellier cedex 5 France. 3Santa Fe Institute 1399 Hyde Park Road Santa Fe NM 87501 USA. 4Genomics and Computational Biology Graduate Program, University of Pennsylvania Philadelphia PA 19104 USA. 5Center for Evolution and Cancer, Helen Diller Family Comprehensive Cancer Center and Department of Surgery, University of California San Francisco San Francisco CA 94143 USA. 6Departments of Radiology and Integrated Mathematical Oncology, the Moffitt Cancer Center 12902 Magnolia Dr Tampa FL 33612 USA. 7IRD, MIVEGEC (UMR CNRS/IRD/UM1) 911 Ave. Agropolis BP 64501, FR-34394 Montpellier cedex 5 France. 8CREEC, Université Montpellier 2, CC 06003, 95 rue de la Galéra, 34095 Montpellier Cedex 5, France.

Received: 30 April 2012 Accepted: 27 August 2012

Published: 3 September 2012

References

- Greaves M: Darwinian medicine: a case for cancer. *Nature Rev Canc* 2007, 7:213–221.
 - Caulin AF, Maley CC: Peto's paradox: evolution's prescription for cancer prevention. *TREE* 2011, 26:175–182.
 - Peto R, Roe FJ, Lee PN, Levy L, Clack J: Cancer and ageing in mice and men. *Br J Cancer* 1975, 32:411–426.
 - Albanes D, Winic M: Are cell number and cell proliferation risk factors for cancer? *JNCI J Natl Cancer Inst* 1988, 80(10):772–775.
 - Diekmann O: A beginner's guide to adaptive dynamics. *Banach Center Publ* 2003, 63:47–86.
 - Hardison RC: Comparative genomics. *PLoS Biol* 2003, 1(2):e58.
 - Calabrese P, Shibata D: A simple algebraic cancer équation: calculating how cancers may arise with normal mutation rates. *BMC Cancer* 2010, 10:3.
 - Walker B, Figgs LW, Zahm SH: Differences in cancer incidence, mortality and survival between African Americans and whites. *Environ Health Perspect* 1995, 103(Suppl 8):275–281.
 - Le Souëf PN, Goldblatt J, Lynch NR: Evolutionary adaptation of inflammatory immune responses in human beings. *Lancet* 2000, 356:242–244.

10. O'Byrne KJ, Dalgleish AG: Evolution, immune response, and cancer. *Lancet* 2000, **356**:1033.
11. Maley CC, Forrest S: Exploring the relationship between neutral and selective mutations in cancer. *Artif Life* 2000, **6**(4):325–345.
12. Luebeck EG, Moolgavkar SH: Multistage carcinogenesis and the incidence of colorectal cancer. *Proc Natl Acad Sci U S A* 2002, **99**(23):15095–15100.
13. Nunney L: The population genetics of multistage carcinogenesis. *Proc Biol Sci* 2003, **270**(1520):1183–1191.
14. Calabrese P, Tavare S, Shibata D: Pretumor progression: clonal evolution of human stem cell populations. *Am J Pathol* 2004, **64**(4):1337–1346.
15. Beerewinkel N, Antal T, Dingli D, Traulsen A, Kinzler KW, Velculescu VE, Vogelstein B, Nowak MA: Genetic progression and the waiting time to cancer. *PLoS Comput Biol* 2007, **3**(11):e225.
16. Meza R, Jeon J, Moolgavkar SH, Luebeck EG: Age-specific incidence of cancer: phases, transitions, and biological implications. *Proc Natl Acad Sci U S A* 2008, **105**(42):16284–16289.
17. Maley CC, Lewis W, Reid BJ: Has cancer sculpted the genome? In *Modeling linkage and the role of tetraploidy in neoplastic progression*. In *Multiscale Cancer Modeling*. Edited by Deisbock T, Stamakos GS. Boca Raton, FL: Chapman & Hall/CRC Press; 2010.
18. Sprouffske K, Pepper JW, Maley CC: Accurate reconstruction of the temporal order of mutations in neoplastic progression. *Cancer Prevention Research* 2011, **4**(7):1135–1144.
19. Gatenby RA, Maini PK: Mathematical oncology: cancer summed up. *Nature* 2003, **421**:321.
20. Silva AS, Gatenby RA: A theoretical quantitative model for evolution of cancer chemotherapy resistance. *Biol Direct* 2010, **5**:25.
21. McAloose D, AL N: Wildlife cancer: a conservation perspective. *Nature Rev Canc* 2009, **9**:517–526.

doi:10.1186/1471-2407-12-387

Cite this article as: Roche et al.: Natural resistance to cancers: a Darwinian hypothesis to explain Peto's paradox. *BMC Cancer* 2012 **12**:387.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

