



HAL
open science

Rhizosphere bacteria are more strongly related to plant root traits than fungi in temperate montane forests: insights from closed and open forest patches along an elevational gradient

Luis Merino-Martín, Robert Griffiths, Hyun Gweon, Clément Furget-Bretagnon, Anna Oliver, Zhun Mao, Yves Le Bissonnais, Alexia Stokes

► To cite this version:

Luis Merino-Martín, Robert Griffiths, Hyun Gweon, Clément Furget-Bretagnon, Anna Oliver, et al.. Rhizosphere bacteria are more strongly related to plant root traits than fungi in temperate montane forests: insights from closed and open forest patches along an elevational gradient. *Plant and Soil*, 2020, 450 (1-2), pp.183-200. 10.1007/s11104-020-04479-3 . hal-02534567

HAL Id: hal-02534567

<https://hal.umontpellier.fr/hal-02534567v1>

Submitted on 18 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rhizosphere bacteria are more strongly related to plant root traits than fungi in temperate montane forests: insights from closed and open forest patches along an elevational gradient

Article

Accepted Version

Merino-Martín, L., Griffiths, R. I., Gweon, H. S., Furget-Bretagnon, C., Oliver, A., Mao, Z., Le Bissonnais, Y. and Stokes, A. (2020) Rhizosphere bacteria are more strongly related to plant root traits than fungi in temperate montane forests: insights from closed and open forest patches along an elevational gradient. *Plant and Soil*, 450 (1-2). pp. 183-200. ISSN 0032-079X doi: <https://doi.org/10.1007/s11104-020-04479-3> Available at <http://centaur.reading.ac.uk/89885/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s11104-020-04479-3>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Rhizosphere bacteria are more strongly related to plant root traits than fungi in**
2 **temperate montane forests: insights from closed and open forest patches along an**
3 **elevational gradient**

4 **Author list / address list:**

5
6 Luis Merino-Martín^{1,2,3}, Robert I. Griffiths³, Hyun S. Gweon^{3,4}, Clément Furget-Bretagnon¹,
7 Anna Oliver³, Zhun Mao¹, Yves Le Bissonnais⁵, Alexia Stokes¹.

8
9 ¹ University Montpellier, Amap, INRAE, Cirad, Cnrs, Ird, Montpellier, France.

10 ² C.N.R.S., UMR CEFE, Montpellier, France.

11 ³ Centre for Ecology & Hydrology, Crowmarsh Gifford, Wallingford, Oxfordshire, United Kingdom.

12 ⁴ Harborne Building, School of Biological Sciences, University of Reading, United Kingdom

13 ⁵ INRAE, UMR1221 LISAH, Montpellier, France.

14
15 *Corresponding author. Tel.: +33(0)789288168. E-mail address: luismerinomartin@gmail.com
16 (*L. Merino-Martín*).

17 For submission to *Plant and Soil*

18 **Article type:** Full paper.

19 **Word count:**

Total word count (excluding summary, references and legends):	6541	No. of figures:	5
Summary:	219	No. of Tables:	4
Introduction:	1190	No of Supporting Information files:	8 (Fig. S1, S2; Table S1–S5, Notes S1)
Materials and methods:	3024		
Results:	845		
Discussion:	1249		
Conclusions:	161		
Acknowledgements:	64		

20

21

22 **Summary**

23 **Aims:**

24 Heterogeneous canopies in temperate montane forests affect microclimate and soil characteristics,
25 with important effects on soil microbial communities and related processes. Here, we studied the
26 interactions between plant root traits and soil bacterial and fungal communities in closed forest
27 and gaps in a mixed forest along an elevational gradient in the French Alps (1400, 1700 and 2000
28 m).

29 **Methods:**

30 Samples were separated into three fractions (plant root endosphere, rhizosphere and bulk soil), to
31 further investigate the influence of plant zones on microbial communities. Bacterial (16S) and
32 fungal (ITS) biodiversity was determined using high throughput sequencing, along with standard
33 measures of soil, litter and root traits.

34 **Results:**

35 We found that (i) microbial community diversity was higher in gaps than in closed forest because
36 of increased root trait diversity and density; (ii) open versus closed forest patches affected
37 phylogenetic dispersion despite differences in elevations with phylogenetic clustering in closed
38 forest; (iii) the interaction between root traits and microbial communities was stronger for
39 rhizosphere and endosphere compartments than for bulk soil and (iv) bacterial community
40 composition was better explained by root traits than for fungi.

41 **Conclusions:**

42 Our findings highlight the importance of open gaps versus closed forest patches and associated
43 root traits affecting microbial community structure, particularly for bacterial assemblages that
44 exhibited a stronger interaction with root traits than for fungi.

45 **Key words:** Bacteria; closed forest; fungi; gaps; ITS; phylogenetic clustering; root traits; soil
46 properties; 16S.

47 **Introduction**

48 Naturally heterogeneous canopies in forests create a mosaic of belowground responses in the
49 activity and composition of microbial communities, that are linked primarily to the presence of
50 specific plant species, as well as heterogeneity in microclimate and soil characteristics (Lladó et
51 al. 2018; Muscolo et al. 2014). Disturbance in closed forests causes gaps that can alter local soil
52 moisture, solar radiation and air and soil temperatures (de Freitas and Enright 1995; Gray et al.
53 2002). Understorey species then dominate in the gap, until tree growth causes the canopy to close
54 again. Recent evidence shows that gap size alters soil microbial community structure (Muscolo
55 et al. 2014; Yang et al. 2017a; Yang et al. 2017b) but it is not clear if these changes were primarily
56 due to differences in the composition of plant species and/or to modifications in the local
57 microclimate. Individual trees also structure soil microbial communities (Bach et al. 2010)
58 through: (i) a modification in soil physicochemical parameters mostly via changes in litter
59 quantity and quality (Baldrian 2017; Dukunde et al. 2019; Prescott and Grayston 2013; Saetre and
60 Bååth 2000), (ii) root traits and root exudates in the rhizosphere (Colin et al. 2017; Lladó et al.
61 2018), in addition to (iii) the modifications in microclimatic conditions mentioned above. Here,
62 we aim to identify whether gaps in naturally heterogeneous forests affect bacterial and fungal
63 community structure through changes in microclimate, soil physicochemical parameters or
64 vegetation, with a specific focus on root traits of both trees and understory species. To achieve
65 our objective, we use an elevational gradient to observe if shifts in soil microbial diversity
66 between gaps and closed forests are reflected in diverse climatic and soil conditions (but with
67 minimal differences in soil texture).

68 Studying forest microbial diversity and structure along climate and soil gradients is challenging
69 because factors can co-vary (McCain and Grytnes 2010). Nevertheless, elevational gradients
70 permit the study of several abiotic factors, since major changes in these factors are found along
71 relatively short distances (Körner 2007; Ren et al. 2018). Based on bacterial taxonomic diversity,
72 contradicting results have been found in temperate regions, with taxonomic richness varying
73 along elevational gradients; either decreasing (Bryant et al. 2008), showing a hump-backed

74 relationship (Singh et al. 2012), or not showing any relationship (Shen et al. 2014). Fungal
75 richness has been observed to either decrease at high elevation (Bahram et al. 2012; Kernaghan
76 and Harper 2001), or have a hump-backed shape along the gradient (Miyamoto et al. 2014).
77 Several studies have included both fungal and bacterial communities, but trends varied (Ren et
78 al. 2018; Shen et al. 2014; Siles and Margesin 2016). These contradicting results are possibly
79 because different environmental variables along the elevational gradient affect community
80 assemblages (Coince et al. 2014). Here, our approach is to examine microbial diversity in gaps
81 and closed forest along an elevational gradient, to determine if the patterns observed in response
82 to local differences are repeated along the elevational gradient.

83 Any modifications in soil microbial community assemblages will have a cascade of effects on
84 soil structure and nutrient cycling, including: (i) soil aggregation and aggregate stability (Baumert
85 et al. 2018; Chenu and Sotzky 2002) and (ii) decomposition processes (Kohout et al. 2018;
86 Schneider et al. 2012). Gaps in forests might be beneficial to microbial communities through
87 increases in their biomass and activity (Muscolo et al. (2014); Yang et al. (2017b)). If
88 microorganisms are more abundant and active in gaps, soil aggregate stability and litter
89 decomposition should be enhanced. However, these processes are also linked to the plant species
90 present and their chemical and physiological traits (Grigulis et al. 2013; Poirier et al. 2018).
91 Prescott and Grayston (2013) found that the main factors associated to differences in microbial
92 communities in litter, forest floors and soil were pH and base cation content of the litter and
93 whether the trees were broadleaf or coniferous. Besides, Brant et al. (2006) revealed that in forest
94 ecosystems of Oregon, Pennsylvania, and Hungary, root carbon (C) inputs exerted a larger control
95 on microbial community composition than litter inputs. Plant species diversity and root density
96 are also greater in gaps compared to closed forests (Mao et al. 2015) due to a higher presence of
97 shrubs, forbs and grasses, benefitting the activity of rhizosphere microbial communities
98 (Kuzyakov and Blagodatskaya 2015). Therefore, even though a higher input of litter is expected
99 in closed forests, the increase in root density and diversity, and associated litter, could result in
100 more diverse and active microbial communities in gaps.

101 Approaches for studying microbial diversity along gradients have changed from taxonomic to
102 phylogenetic in recent years (Parks and Beiko 2013). Phylogenetic clustering processes are
103 observed when evolutionarily related organisms coexist due to restricting abiotic or biotic
104 conditions. Several studies have found that phylogenetic clustering increases with elevation for
105 bacterial communities (Bryant et al. 2008; Wang et al. 2012; Zhang et al. 2018). However, this
106 filtering (i.e. environmental selection against certain species) is not only mediated by abiotic
107 factors along the gradient but also by biotic interactions (Goberna et al. 2014a; Mayfield and
108 Levine 2010) and so, there might be a role of gaps and closed forests in phylogenetic patterns. To
109 the best of our knowledge, there is no study of the effects of gaps and closed forests in temperate
110 forests, where changes in root diversity and density and microclimatic conditions are expected to
111 influence strongly the bacterial phylogenetic patterns.

112 Ecological habitat (i.e. rhizosphere or bulk soil) is the main factor structuring bacterial
113 communities (Uroz et al. 2010), due primarily to the supply of rhizodeposits released from live
114 roots (Philippot et al. 2013; Shi et al. 2012). Rhizodeposits include root cells and tissues, exudates,
115 mucilage, volatiles and soluble lysates that are sloughed-off as a root grows through soil (Uren
116 2000). However, recent studies have shown that the structure of fungal communities are not
117 related to these rhizodeposits, but are strongly related to tree species (Urbanová et al. 2015; Uroz
118 et al. 2016) mainly through litter quality that affects community composition of saprotrophic and
119 ectomycorrhizal (ECM) fungi (Aponte et al. 2013; Prescott and Grayston 2013). If root traits
120 differ between gaps and closed forest, modifications in bacterial communities in particular, should
121 be observed within the rhizosphere compared to bulk soil.

122 Using an elevational gradient to observe if patterns of microbial diversity and structure between
123 gaps and closed forest are repeated along the gradient, we aim at addressing four hypotheses.
124 First, we hypothesize that microbial community diversity will be higher in gaps than in closed
125 forest because of increased root trait diversity and density. Second, we expect to observe
126 phylogenetic clustering in closed forest due to reduced root trait diversity and density compared
127 to gaps. Third, we hypothesize that the interaction between root traits and microbial communities

128 will be stronger for rhizosphere and endosphere compartments than for microbial communities
129 inhabiting bulk soil. Finally, we expect that bacterial communities will be more strongly affected
130 by ecological habitat (bulk soil, rhizosphere or endosphere) than fungal communities. These
131 modifications to patterns in community assemblage should be repeated along the elevational
132 gradient, because plant communities should have a greater effect on structuring microbial
133 communities than abiotic factors, because of the habitat quality that they provide.

134 **Materials and Methods:**

135 Study site

136 Field sites are located near Chamrousse, Isère, French Alps (45°6'N, 5°54'E). Three mixed,
137 mature, naturally regenerated forests of Norway spruce (*Picea abies* (L.) Karst.), Silver fir (*Abies*
138 *alba* Mill.), European beech (*Fagus sylvatica* L.) and Mountain pine (*Pinus uncinata* Ramond ex
139 DC.) growing at elevations of 1400 (Prémol), 1700 (Bachat-Bouloud) and 2000 m a.s.l. (near
140 Achard lake, at the treeline) were sampled to assess the effects of changes along elevation for soil
141 physicochemical properties, microbial communities, litter and root traits. Plant species
142 composition shifts from 1400 m to 2000 m with broad-leaved species almost absent above 1700
143 m. *F. sylvatica* is one of the dominant species at 1400 m and is not present at 1700 and above. *A.*
144 *alba*, *P. abies*, and *P. uncinata* are the most dominant species at 1400 m, 1700 m, and 2000 m,
145 respectively. *P. abies* is the only species present at all elevations (Mao et al. 2015; Wang et al.
146 2018b). At the three sites, the forest shows spatial heterogeneity with closed canopy forests and
147 open canopy areas (>10 m diameter) formed through tree mortality after disturbances such as
148 storm or tree felling. Vegetation composition in open canopy areas also changes with elevation
149 with *Gallium rotundifolium*, *Lysimachia nemorum* and *Luzula nivea* being the most dominant
150 species at 1400 m and *Rhododendron ferrugineum* and *Vaccinium myrtillus* at 1700 and 2000 m
151 (Mao et al. 2015; Prieto et al. 2015).

152 As species composition changed along the elevation gradient, a sampling design comparing
153 closed forest and gaps at each site was selected. This approach is a mean of standardization of the
154 effect of vegetation through the comparison between closed forest and gaps along the elevational

155 gradient. This patch-gap mosaic approach has been used to discern between abiotic and biotic
156 mechanisms underlying the coexistence of phylogenetically related bacteria in a dryland
157 environment (Goberna et al. 2014) but to our knowledge, has never been used to study microbial
158 communities along environmental gradients. A detailed description of selected sites can be found
159 in Wang et al. (2018a) and Mao et al. (2015) and a complete description of species and abundance
160 of herbs and trees in closed forest and gaps at the three elevations can be found in Table S1.

161 The study sites possess similar soil type, and differences between the sites are mainly caused by
162 climatic conditions and the type of vegetation. Soils are acidic at all sites, ranging from (a)
163 “Cambisols (Hyperdystric)” according to the World Reference Base for Soil Resources (IWG
164 2007), above green schist and with an abundant water supply at 1400 m, to (b) “Cambisols
165 (Humic, Hyperdystric)”, above the crystalline formation at 1700 m, and to (c) “Epileptic
166 Umbrisols (Hyperdystric)”, above the crystalline formation at 2000 m (Joud 2006; Mao et al.
167 2012). More details on the study site are available in Mao et al. (2013, 2015) and Wang et al.
168 (2018b).

169 The air and soil temperatures (10 cm depth) in two closed forests and two gaps were assessed in
170 the three elevations from September 28th, 2010 to March 3rd, 2014 (Table S2, Fig. S2) with a
171 portable thermistor thermometer (HI-93510N Hanna Instruments) in soil trenches adequately
172 covered by insulation. Additionally, data from Wang et al. (2018a) were used to include soil water
173 potential (ψ) in the microclimate and climate assessment (Table S2). These authors used electrical
174 resistance blocks (WaterMark, IRROMETER Company, Inc., USA) installed in one gap and one
175 closed forest at 1400 and 1700 m (but not at 2000 m as equipment was stolen) from September
176 10th, 2012 to November 18th, 2013.

177 Sampling and storage

178 In each location, three different paired plots (gap *versus* closed forest) with representative patches
179 of closed forests and gaps were chosen (Fig. 1). The conditions for the selection of these pairs
180 were: (i) in closed forest, at least three adult trees of the dominant canopy species were present in

181 a cluster of trees that had a diameter >8 m, and (ii) gaps comprised an open area with no trees
182 and a diameter greater than 8 m. In closed forests, samples were taken from between trees whilst
183 in gaps, samples were taken in the middle of the gap in order to avoid as much as possible the
184 influence surrounding forest. Four different surface soil samples (0-10 cm) were collected in each
185 plot using sterilised material: (i) a soil cylinder for fine root analysis, (ii) a plastic bag with 10 to
186 20 g of soil for microbial analysis, (iii) a sample of leaf debris from the surface (hereafter termed
187 'litter') and (iv) one 0.75 dm³ container of soil for measurements of aggregate stability. A total of
188 18 samples of each type were collected: 3 elevations x 6 samples at each site three in gaps and
189 three in closed forests). Of each of these 18 samples collected for root analysis, 3 sample fractions
190 were separated: root, rhizosphere and bulk soil making a total of 54 samples. Soil samples were
191 collected during July 7th - 10th, 2014, at the peak of the summer season when microbial activity at
192 the three elevations is expected to be at its maximum.

193 Soil samples for root, litter and microbial analysis were kept in a freezer at -20°C until analyses
194 were performed. Soil samples for aggregate stability tests were air-dried in the laboratory until
195 they were processed approximately a month after they were collected.

196 Soil physicochemical properties

197 Soil was sieved at 2 mm after air drying and the soil fraction <2 mm was used to assess
198 physicochemical properties. Soil samples were sent to Natural Resource Management (Berkshire,
199 UK). Soil pH was measured in water as 1:2.5 extract. Soil organic matter content was determined
200 via loss-on-ignition at 500 °C (Dean 1974). Total nitrogen (N) and carbon (C) were determined
201 via the DUMAS method (Shea and Watts 1939). Available potassium (K) and magnesium (Mg)
202 were determined through ammonium nitrate extraction and available phosphorus (P) was
203 measured via Olsen (extraction on 0.5 M sodium bicarbonate, (Olsen et al. 1954)). Soil texture
204 was determined by laser-diffraction analysis (McCave et al. 1986). The soil sample was
205 previously digested in hydrogen peroxide solution to destroy the organic matter and sodium
206 hexametaphosphate to release the bound clay particles.

207 Aggregate stability was determined by the fast wetting standard method, ISO/CD 10930,
208 developed by Le Bissonnais (1996). This methodology is appropriate to compare the behaviour
209 of a large range of soils during rapid wetting mimicking heavy rainstorms in summer. Initially, 5
210 g of aggregates (3-5 mm) were gently immersed in 50 ml of deionized water for 10 min; water
211 was then removed with a pipette and the soil material was transferred to a 50- μm sieve previously
212 immersed in ethanol. The 50 μm sieve immersed in ethanol was gently moved five times to
213 separate fragments smaller and bigger than 50 μm . The $>50 \mu\text{m}$ fraction was collected, oven-dried
214 and gently dry-sieved by hand on a column of six sieves: 2000, 1000, 500, 200, 100 and 50 μm .
215 The mass percentage of each size fraction was calculated, and the aggregate stability was
216 expressed by computation of the mean weight diameter (MWD).

217 Fine roots and litter

218 Soil cylinders and litter samples were defrosted. Litter samples were gently washed and rinsed
219 with deionised water and dried at 40°C until constant weight. Root samples were also washed
220 gently with deionised water and divided in two subsamples: 1) a representative subsample that
221 was selected for scanning and later drying in the oven at 40°C (n = 18) and the 2) remaining root
222 material of the sample that was dried at 40°C until constant weight (n = 18). Both subsamples
223 were weighed after drying.

224 Roots selected for morphological measurement were stained with methylene blue (1 g L⁻¹) to
225 increase the contrast and allow the detection of fine roots. Then, roots were placed in a tray with
226 deionised water and scanned (Epson© V700 perfection) at a resolution of 1200 dpi. Analysed
227 roots were then recovered, and oven dried at 40 °C and weighed to obtain dry mass. Root images
228 were analysed with the WinRhizo® software (Pro version 2007, Regent Instrument, Quebec,
229 Canada) using the automatic thresholding option and Lagarde's mode with a filter identifying
230 roots when length was five times the width. Total root length and the length of roots in seven
231 diameter classes (width 0 mm to 1 mm in 0.2 mm classes and 1 mm to 2 mm in 0.5 mm classes)
232 were measured with the software. Specific root length (SRL) was calculated as the ratio between

233 total root length and root dry mass. The percentages of very fine (VFR, diameter < 0.2 mm) and
234 fine (FR, 0.2 < diameter < 1 mm) roots were defined as the ratio of length in the concerned root
235 classes to total root length (Miller and Jastrow 1990). Total root mass density (RMD) was
236 calculated by the ratio of total root dry mass and the soil volume extracted. Total root length
237 density (RLD) was calculated by the ratio of total root length and the soil volume extracted. Total
238 root dry mass was calculated as the sum of the dry mass of roots selected for morphological
239 analysis and those of the remaining roots. Root dry matter content (RDMC) was calculated as the
240 ratio of the root dry weight and root fresh weight.

241 The concentrations of water soluble compounds (cellulose and lignin; mg g⁻¹) in root and litter
242 samples (n = 18 for litter and n = 6 for roots as replicates were combined due to limited sample
243 amount), were obtained by the Van Soest Method (1963) with a Fibersac fibre analyser (Ankom,
244 Macedon, USA). Root and litter C and N concentrations (n = 18 for each) were measured using
245 an elemental analyser (Thermo-Finnigan EA1112, Milan, Italy).

246 Soil microbial communities

247 The protocol for cleaning roots and obtaining root, rhizosphere and bulk soil samples was
248 performed following Bulgarelli et al. (2012, 2015). Briefly, loose soil was manually removed
249 from the root system and stored as the bulk soil sample. Roots were collected in 50 ml falcon
250 tubes containing 10 ml PBS-S buffer (130 mM NaCl, 7 mM Na₂HPO₄, 3mM NaH₂PO₄, pH 7.0,
251 0.02 % Silwet L-77) and washed for 20 minutes at 180 rev min⁻¹ on a shaking platform. These
252 roots were transferred to a new falcon tube and subjected to a second washing treatment (20
253 minutes at 180 rpm in 3 ml PBS-S buffer). The soil suspensions collected in the falcon tubes after
254 the first and second washing treatments were combined, centrifuged at 4000g for 20 min and the
255 pellet, considered as the rhizosphere sample, was frozen and stored at -20°C until further
256 processing. Double-washed roots were then transferred to a new falcon tube with 3 ml PBS and
257 sonicated for 2 minutes at 160 W to enrich for microbes living in close association with root
258 tissues. Roots were removed from PBS-S, rinsed in a fresh volume of 10 ml PBS-S buffer and

259 ground with a mortar and pestle in liquid nitrogen. Pulverised roots (considered as the ‘root’
260 sample) were collected in 15 ml falcon tubes and stored at -20°C until further processing.

261 Total DNA was extracted from soil (0.25 g) and the rhizosphere and root fractions (0.25 g when
262 possible and the entire material available when quantity was less than 0.25 g). DNA extraction
263 was performed using PowerSoil®-htp96 Well Soil DNA Isolation Kit according to
264 manufacturer’s instructions (MOBIO Laboratories, UK).

265 Bacterial and fungal community biodiversity was assessed using Illumina amplicon sequencing
266 of 16S rRNA genes (bacteria) and the Internal transcribed spacer (ITS) region (fungi) to
267 phylogenetically identify responsive taxa. A phylogenetic analysis was also performed for
268 bacterial communities. Amplicon libraries were constructed according to the dual indexing
269 strategy of Kozich et al. (2013), with each primer consisting of the appropriate Illumina adapter,
270 an 8-nt index sequence, a 10-nt pad sequence, a 2-nt linker and the gene specific primer. For 16S,
271 the V3-V4 hypervariable regions of the 16S rRNA gene was targeted using primers based upon
272 the universal primer sequence 341F and 806R. For ITS, region 2 (ITS2) was amplified utilising
273 the fITS7 (forward) and ITS4 (reverse) primer sequences described in Ihrmark et al. (2012).
274 Additional methodological details of Illumina sequencing are described in Notes S1.

275 Sequenced 16S rRNA paired-end reads were joined using PEAR (Zhang et al. 2014), quality
276 filtered using FASTX tools (hannonlab.cshl.edu), length filtered with the minimum length of
277 300bps, presence of PhiX and adapters were checked and removed with BBTools
278 (jgi.doe.gov/data-and-tools/bbtools/), and chimeras were identified and removed with
279 VSEARCH_UCHIME_REF (Rognes et al. 2016) using Greengenes Release 13_5 (at 97%)
280 (DeSantis et al. 2006b). Singletons were removed and the resulting sequences were clustered into
281 operational taxonomic units (OTUs) with VSEARCH_CLUSTER (Rognes et al. 2016) at 97%
282 sequence identity (Tindall et al. 2010). Representative sequences for each OTU were
283 taxonomically assigned by RDP Classifier with the bootstrap threshold of 0.8 or greater (Wang
284 et al. 2007) using the Greengenes Release 13_5 (full) (DeSantis et al. 2006b) as the reference.

285 Unless stated otherwise, default parameters were used for the steps listed. ITS2 sequences were
286 processed using the PIPITS pipeline (Gweon et al. 2015), where OTUs were taxonomically
287 assigned against the UNITE database (Release 31.01.2016, Koljalg et al. (2013).

288 Statistical analysis

289 A four-step statistical procedure was performed to determine the interrelationships between the
290 different variables under study: (i) examine the variations on environmental variables (soil
291 properties and root and litter traits) in closed forest versus gaps along the elevational gradient; (ii)
292 non-metric multi-dimensional scaling (NMDS) data ordinations of microbial data and test of the
293 effect of closed forest versus gaps and sample fraction (root, rhizosphere and bulk soil) on their
294 structure (Permutational Multivariate Analysis of Variance, PERMANOVA) along the
295 elevational gradient; (iii) study of correlation between microbial community structure and
296 environmental variables (Spearman correlations and distance-based redundancy analysis,
297 dbRDA); and (iv) study of taxonomic and phylogenetic diversity and divergence among closed
298 forest and gaps along the elevation for the three sample fractions (bulk soil, rhizosphere and root).

299 Air and soil temperatures, soil physicochemical properties and root traits were analysed by
300 Analysis of Covariance (ANCOVA) and Student's t-tests for closed forests *versus* gaps (hereafter
301 termed “tree-gap”). The tree-gap explanatory variable was treated as a factor and elevation was
302 included as a covariate. All variables tested fulfilled ANCOVA assumptions except for SRL that
303 was transformed and RLD, soil P, litter hemicellulose N, C, and C:N that were analysed by
304 Kruskal Wallis tests and Wilcoxon Rank Sum Tests. Soil water potential data were analysed with
305 Friedman rank sum tests and post hoc comparisons were performed using Nemenyi multiple tests.

306 Similarities/dissimilarities between microbial communities were displayed using NMDS of Bray–
307 Curtis dissimilarity for bacterial and fungal OTUs matrices. To indicate similarities between
308 treatments on the NMDS configuration, the points on the NMDS ordination were overlapped with
309 polygons and spider diagrams indicating elevation, tree-gap and soil fraction. PERMANOVA
310 tests (adonis R function) were performed for the bacteria and fungi OTUs matrices as the response

311 and the three different factors (tree-gap and soil fraction) as the explanatory variables and
312 elevation as a covariate.

313 The relationships between the microbial community composition and the soil physicochemical
314 variables, root and litter traits were tested using Spearman correlations between these variables
315 and alpha (Shannon diversity index) and beta diversities (NMDS first and second axis) and a db-
316 RDA. In order to select the environmental parameters to be included in the constrained ordination,
317 an initial db-RDA including all parameters was performed followed by a stepwise model selection
318 using Generalized Akaike Information Criterion (AIC, ordistep function with a backward
319 direction). Finally, the db-RDA analysis was performed only for the variables obtained. ANOVA
320 tests were performed on the final constrained ordination to confirm that the first two axes and the
321 environmental variables and the final constrained ordination were significant.

322 Additionally, the dbRDA analyses were computed for each of the sample fractions and each of
323 the three environmental matrices (soil, root and litter traits) to study the variance explained by
324 these environmental variables and the bacterial and fungal composition.

325 Apart from the traditional taxonomic approach, a phylogenetic approach was also used as it is a
326 useful method to measure biodiversity incorporating phylogenetic difference between species
327 (Webb 2000). The phylogenetic approach relies on homologous genes (genes that are derived
328 from a common ancestor). Here, we sequenced the ITS2 region for fungi which is not a
329 homologous gene, and 16S rRNA gene for bacterial communities (which is homologous).
330 Therefore, the phylogenetic approach was only used for bacterial communities. A phylogenetic
331 tree was constructed based on 16S rRNA representative sequences. The sequences were aligned
332 with PyNAST (Caporaso et al. 2010) to the Greengenes reference database (DeSantis et al.
333 2006a). FastTree (Price et al. 2010) was then used on the resulting alignment to produce a
334 maximum-likelihood phylogenetic tree by iterative rearrangement of branches with generalized
335 time-reversible (GTR) models of nucleotide evolution. This phylogenetic tree was used to
336 generate the distance matrices and obtain two main indexes: one to measure phylogenetic

337 diversity (Faith's Phylogenetic Diversity (Faith (1992), hereafter PD) and a second one to
338 measure phylogenetic divergence, the standardized mean nearest taxon distance (hereafter
339 ses.MNTD). Faith's Phylogenetic Diversity is a measure of the total phylogenetic branch length
340 that joins the basal node to the tips of all species in the sample. Mean nearest taxon distance
341 (MNTD) is an estimate of the mean phylogenetic relatedness between each OTU in a bacterial
342 community and its nearest relative. The standardized mean nearest taxon distance (ses.MNTD)
343 can be used to test for phylogenetic clustering or over dispersion as it is an index that compensates
344 for random processes in the observed phylogenetic community structure. The standardized effects
345 of MNTD were obtained comparing the observed phylogenetic relatedness to the pattern obtained
346 by community randomizations holding community species richness constant (runs = 999,
347 iterations = 1000). Finally, Analysis of Covariance (ANCOVA) tests were performed for these
348 indexes as response variables, tree-gap and sample fraction as explanatory variables and elevation
349 as a covariate. The variables H, PD and ses.MNTD were transformed with a Johnson, inverse and
350 Tukey's Ladder of Powers transformations to meet ANCOVA assumptions.

351 All analyses were performed in RStudio Version 1.0.136 (RStudio Team 2016) using the *vegan*
352 (Oksanen et al. 2016), *picante* (Kembel et al. 2010), and *ade4* (Dray and Dufour 2007) packages.

353 **Results**

354 Relationships between environmental variables and bacterial and fungal communities.

355 The results of the relationships between soil physicochemical properties, root and litter traits
356 (Tables S3 and S4, Figure 2) and microbial community composition are shown in the db-RDA
357 (Figure 3) and Spearman correlations with alpha and beta diversities (Table 2). The final db-RDA
358 analysis (Figure 3) shows only the environmental parameters that were selected through stepwise
359 model selection. ANCOVA tests performed on the final constrained ordination confirmed that the
360 first two axes, the environmental variables and the final constrained ordination were significant.
361 Bacterial alpha diversity was positively correlated with SRL and negatively correlated with
362 RDMC and root N (Table 2). Additionally, bacterial beta diversity was positively correlated with

363 soil C:N, MRD and root C and negatively correlated to SRL, VFR, RLD and litter hemicellulose.
364 There was a clear separation between closed forests and gaps for bacteria (Figure 3), with
365 aggregates from closed forests having greater stability (MWD), higher root dry matter content
366 (RDMC), soil C and C:N. Gaps were associated to higher VFR and SRL.

367 No correlations were found between any of the parameters measured and the alpha diversity of
368 fungi. Nevertheless, numerous soil properties (sand, SOC, N, C, C:N) and root traits (MRD, N, C
369 and cellulose) were positively correlated with fungal beta diversity whilst other traits were
370 negatively correlated with fungal beta diversity (clay, SRL, VFR, RLD and litter hemicellulose).
371 As for bacterial communities, the composition of fungal communities was different between
372 closed forests and gaps particularly for elevations 1700 and 2000 but not for 1400 m (Figure 3).
373 For fungi, the environmental variables showed similar trends than those obtained for bacteria
374 although more factors were significant (higher MRD, root C, soil N and litter lignin for closed
375 forests and higher RLD, RMD, litter C, N and hemicellulose content for gaps).

376 Data on the climate along the elevation (Table S2, Figure S2) showed that elevation decreased air
377 and soil temperatures and water potential in gaps and closed forests with a gradual decrease of
378 soil and air temperatures for closed forests and a greater decrease between 1700 and 2000 m for
379 gaps. Soil and air temperatures in gaps were higher than in closed forest at the mid elevation (1700
380 m) but not at the lowest and highest elevations (1400 and 2000 m).

381 The dbRDA analyses computed for the sample fractions and environmental matrices (soil, root
382 and litter traits) showed that the variance of bulk soil communities (bacterial and fungal) was
383 mainly explained by soil properties but not by root or litter traits (adjusted $R^2=0.34$ and 0.13
384 respectively, Table 1). Notably, the variance of rhizosphere and root bacterial communities
385 explained by root traits was significant (adjusted $R^2=0.28$ and 0.23 respectively) compared to
386 fungal communities with no significant relationships ($p > 0.05$). Only soil properties significantly
387 explained the variance of soil and rhizosphere fungal communities, with lower explained variance
388 than in bacterial communities (Table 1).

389 Effects of closed forests-gaps and elevation gradient on the structure of bacterial and fungal
390 communities.

391 NMDS ordinations showed that bacterial communities were markedly distinct between closed
392 forests and gaps along the elevation gradient (Figure 4). Conversely, fungal communities did not
393 show this robust distinction between closed forests and gaps, but the elevational effect was
394 evident. These findings were supported by the PERMANOVA performed for the bacterial and
395 fungal species matrices, which showed that the main factors structuring bacterial communities
396 were tree-gap ($R^2 = 0.122$; $p = 0.001$; Table 3) and sample fraction ($R^2 = 0.137$; $p = 0.001$), and
397 the main factors structuring fungal communities were elevation ($R^2 = 0.085$; $p = 0.001$) and tree-
398 gap ($R^2 = 0.062$; $p = 0.001$). Remarkably, sample fraction did not significantly structured fungal
399 communities ($p = 0.163$).

400 Bacterial and fungal composition, diversity and divergence among closed forests and gaps.

401 Most sequences found in the 16S dataset were members of the phyla Proteobacteria,
402 Acidobacteria, Actinobacteria and Verrucumicrobia (Fig. S1). Within the Proteobacteria,
403 Alphaproteobacteria was highly abundant in the three fractions. For fungal communities (ITS
404 region), the sequences most predominant were the Basidiomycota and Ascomycota (Fig. S1).
405 Agaricomycetes was particularly dominant across the three soil fractions and showed a slightly
406 higher presence in bulk soil when compared to rhizosphere and root fractions.

407 Tree-gap ($p = 0.003$) and elevation ($p < 0.001$) significantly modified taxonomic diversity
408 (Shannon's diversity index, H) for bacterial communities but not for fungal communities (Table
409 4, Figure 5). Namely, bacterial H was higher in gaps than in closed forests, and H in soil and
410 rhizosphere communities noticeably increased from 1700 to 2000 m while H in root communities
411 increased from 1400 to 1700 m. Tree-gap ($p < 0.001$) and elevation ($p = 0.033$) significantly
412 modified bacterial Phylogenetic diversity (PD) for bacterial communities. More specifically, a
413 lower PD was found in closed forest ($p < 0.001$; Figure 5). Regarding phylogenetic divergence,
414 bacterial communities showed negative values indicating phylogenetic clustering (Figure 5) in all

415 situations. Tree-gap modified significantly ses.MNTD ($p = 0.030$; Table 4) with a particularly
416 evident effect of closed forests in reducing the bacterial ses.MNTD in rhizosphere and root
417 communities and at higher elevations.

418 **Discussion**

419 Our results highlight the importance of plant root traits for the structure and diversity of bacterial
420 communities since bacterial community composition was better explained by root traits than for
421 fungi. The significance of canopy heterogeneity (i.e. open versus closed forest patches) on root
422 traits was demonstrated, along with the structuring impact on associated microbial communities.

423 We hypothesized that microbial community diversity would be higher in gaps than in closed forest
424 due to increased root trait diversity and root density in gaps. These differences in root properties
425 between closed forest and gaps were supported by our results that showed higher SRL and RLD
426 in gaps. The closed forest and gaps influenced significantly both bacterial and fungal community
427 structure and diversity (Table 1, Figures 3, 4). In agreement with our hypothesis, a general
428 increase in bacterial taxonomic diversity was observed in gaps (Table 4) with a marked effect in
429 the endosphere fraction (Figure 5). Besides the effect of root traits on microbial diversity, an
430 increase in soil temperature in gaps is generally positively related to an increase in soil microbial
431 activities (Muscolo et al. 2007). Here, we found that soil temperature in gaps at a depth of 10 cm
432 at 1700 m was higher (mean temperature of 5.27°C in closed forest and 6.68°C in gaps), which
433 could also partially explain the observed increase in taxonomic diversity in gaps. Since microbial
434 communities and roots are major biotic contributors to soil aggregation (Lehmann et al. 2017),
435 we expected a higher soil aggregate stability in gaps. Contrary to our expectations, soil in closed
436 forests had a higher aggregate stability. Thus, our results highlight the importance of other factors
437 for aggregate stability, such as RDMC, soil C and C:N ratio that were significantly higher in
438 closed forest and are important contributors to soil aggregate stability (Gale et al. 2000; Tisdall
439 and Oades 1982).

440 Although we found significant effects of closed forest and gaps on the structure of fungal
441 communities along the elevational gradient (Figure 2, Table 1), the effect on fungal taxonomic
442 diversity was not significant (Table 4). These results are in line with results from Collins et al.
443 (2018), who found a high spatial variation of fungal diversity and abundance compromising the
444 predictive power of vegetation and soil properties. This non-significant effect of closed forest and
445 gaps along the elevational gradient on fungal taxonomic diversity could be related to (i) plant-
446 microbe interactions and their specificity (Brundrett 2002; Uroz et al. 2016),(ii) the capacity of
447 dispersion of fungi that could make them very variable across all situations and (iii) the
448 complexities of the Fungi kingdom, as it is an extensive category and successional changes in
449 communities could be masked when using diversity indices. Additionally, we acknowledge that
450 we did not distinguish the presence of saprotrophic, ectomycorrhizal and arbuscular mycorrhizal
451 fungi, which could explain the lack of trends observed for fungal communities.

452 We observed a phylogenetic clustering in all situations for bacterial communities (negative
453 ses.MNTD values, Figure 5). This observed clustering for bacteria agrees with current consensus
454 as previous studies showed that bacterial communities tend to contain lower taxonomic diversity
455 and are more likely to be phylogenetically clustered than expected by chance (Bryant et al. 2008;
456 Horner-Devine and Bohannan 2006). Consistent with our second hypothesis, we found a lower
457 ses.MNTD in closed forest compared to gaps. This finding indicates the existence of phylogenetic
458 clustering in these environments and suggests the presence of environmental conditions that may
459 cause clustering in closed forest compared to gaps. However, we did not observe this lower
460 ses.MNTD in closed forest compared to gaps at 1400 m, which could be related to the dominance
461 of broadleaved species at this elevation (Prescott and Grayston 2013). Similar results were found
462 by Goberna et al. (2014b) in drylands when comparing vegetation patches to bare soil. These
463 authors found that bacterial communities in gaps were phylogenetically clustered compared to
464 vegetation patches. Goberna et al. (2014) suggested that traits related to environmental stress
465 tolerance are conserved under resource limited conditions (gaps in drylands), while under
466 environments with a high availability of resources (vegetation patches) competitive exclusion of

467 poorly competitive clades becomes significant (Mayfield and Levine 2010). Consequently, these
468 findings agree with our results if we consider that in this study, resource limited conditions could
469 be found in closed forest compared to gaps since vegetation in gaps had thinner roots with lower
470 RDMC and greater SRL, and higher litter C, N and hemicellulose. Elevation altered the effect of
471 patches on the divergence of bacterial communities, that is likely due to the increase in dominant
472 coniferous species in closed forest and the difference in microclimatic conditions at lower
473 elevations. We found an increase of soil C in closed forest at higher elevations related to the lower
474 decomposability of coniferous litter (Jonard et al. 2017). However, we did not observe the
475 expected decrease in N and acidification of the soil (Hornung 1985).

476 With the three-stratum sampling approach (soil, rhizosphere and root), we have shown the key
477 role of root properties on microbial communities from bulk soil to root endosphere. We have also
478 highlighted the strength of this sampling approach to adequately identify processes or
479 environmental variables affecting community assemblage. We hypothesized that microbial
480 communities from root and rhizosphere would be greatly affected by root traits compared to
481 microbial communities in the soil. The dbRDA analyses (Table 2) demonstrated a strong
482 relationship between root traits and microbial communities for bacteria, showing that the variance
483 of rhizosphere and root bacterial communities was significantly explained by root traits. A study
484 on root traits along a boreal-temperate forest gradient also demonstrated strong relationships
485 between absorptive root morphology and fungal and bacterial communities with better
486 correlations in rhizosphere samples than in bulk soils (Ostonen et al. 2017). Thus, according to
487 our last hypothesis, the influence of root traits in the rhizosphere and endosphere on bacterial
488 communities was higher than for fungal communities. These results were confirmed with
489 Spearman correlations showing that bacterial Shannon diversity was affected by root traits (SRL,
490 RDMC and N content), but this was not the case for fungi. Similar results in elevational gradients
491 were obtained by Ren et al. (2018) who found that plant diversity modified bacterial but not fungal
492 diversities, as was previously suggested by several authors (Shen et al. 2014; Siles and Margesin
493 2016) indicating that fungi respond to multiple variables (Jarvis et al. 2015; Ren et al. 2018). This

494 strong association between root traits and bacterial communities, but not for fungal communities,
495 could be related to the limited capacity of dispersion of bacterial communities as compared to
496 fungal communities with their hyphal growth and branching. According to this, we found that
497 several soil properties were correlated with the beta diversity of fungal communities but not for
498 bacterial communities. Besides, we found that root traits were correlated with both bacterial and
499 fungal beta diversities, mainly SRL, VFR, MRD, RLD and C content, highlighting the importance
500 of fine absorptive roots and root C for microbial communities (Bardgett et al. 2014; de Graaff et
501 al. 2010; Liu et al. 2018).

502 We found a remarkably high variance of both bacterial and fungal communities explained by soil
503 properties in the dbRDA analysis. This result is in agreement with the proposed hierarchy
504 concerning the contribution of soil and plant species on microbial communities' structure and
505 composition (Bulgarelli et al. 2012; Lareen et al. 2016; Philippot et al. 2013). In other words, soil
506 physicochemical properties determine the composition of the soil microbiome, whereas root traits
507 and exudates can gradually alter the soil microbiome (Bever et al. (2012); van der Putten et al.
508 (2013).

509 **Conclusions**

510 We found that bacterial community composition was better explained by root traits than for fungi.
511 As expected, this interaction between microbial communities and root traits was more intense in
512 communities isolated from rhizosphere and roots than for the bulk soil. In addition, we found that
513 canopy heterogeneity (i.e. closed forest versus gaps) along the elevation gradient structured
514 bacterial and fungal communities and modified bacterial phylogenetic diversity that decreased in
515 closed forest. Finally, we found that bacterial phylogenetic dispersion was higher in gaps along
516 the elevation gradient. Our results highlight the importance of incorporating (i) root traits, (ii)
517 canopy forest heterogeneity (closed forest versus gaps in this case), and (iii) soil-rhizosphere-
518 endosphere sample fractions in studies along environmental gradients for the correct
519 understanding of factors affecting microbial community assemblage. Overall, our findings

520 highlight the importance of soil properties for bulk soil and rhizosphere microbial communities
521 and the importance of root traits for rhizosphere and root endosphere bacterial communities but
522 not for fungal communities.

523 **Acknowledgements**

524 LMM was funded with a Marie Curie IEF fellowship (FP7 European program, ref. 626666/2013).
525 Funding for CFB was provided by the French and Mexican governments (ECOPICS project,
526 ANR-16-CE03-0009 and CONACYT-273659). Many thanks to Hervé Rey (CIRAD, France),
527 Francois Pailler (INRA France) and Patricia Tabernerero for their help with field and laboratory
528 work. Thanks are due to the Mairie de Chamrousse for access to fieldsites.

529 **Author Contribution**

530 LMM, RIG, AS, YLB designed, carried out the experimental research and analysed the data. HSG,
531 CFB, AO carried out the experimental research and analysed the data. LMM, RIG, AS wrote the
532 manuscript. All authors edited the manuscript.

533 **Supporting Information**

534 **Fig. S1.** Phylum and class of bacterial and fungal communities along the elevation and in gaps
535 and closed forest.

536 **Fig. S2.** Soil temperature data in gaps and closed forest, over time and for the three elevations.

537 **Table S1** Dominant species and abundance of herbs and trees in gaps and closed forest along the
538 elevation gradient.

539 **Table S2** Climate and microclimate data along the elevation gradient in gaps and closed forest.

540 **Table S3** Soil physicochemical properties along the elevation gradient in gaps and closed forest.

541 **Table S4** Root and litter traits along the elevation gradient in gaps and closed forest.

542 **Table S5.** Abbreviations used in this paper

543 **Notes S1** Additional methodological details on Illumina amplicon sequencing of 16S rRNA genes
544 and the ITS region.

Tables:

Table 1. Partition of variance in constrained ordination distance-based Redundancy Analysis (dbRDA) for the three sample fractions and the three set of environmental variables (soil properties, root and litter traits). The degrees of freedom (Df), proportion of the variance explained by each model (R^2), adjusted R^2 (Adj. R^2) and its significance (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, ns = not significant) are shown.

Fraction	Bulk soil				Rhizosphere				Root			
	Df	R^2	Adj. R^2		Df	R^2	Adj. R^2		Df	R^2	Adj. R^2	
Soil properties	11	0.79	0.34	**	11	0.81	0.39	**	11	0.76	0.24	*
Root traits Bacteria	10	0.66	0.10	ns	10	0.73	0.28	*	10	0.71	0.23	*
Litter traits	6	0.40	0.04	ns	6	0.45	0.12	ns	6	0.39	0.02	ns
Soil properties	11	0.73	0.13	*	11	0.70	0.15	***	11	0.80	0.07	ns
Root traits Fungi	10	0.65	0.06	ns	10	0.61	0.06	ns	10	0.71	0.00	ns
Litter traits	6	0.39	0.02	ns	6	0.36	0.02	ns	6	0.47	0.07	*

Table 2. Spearman's correlation coefficients between bacterial and fungal alpha diversity (H: Shannon index) and beta diversity (NMDS1 and NMDS2), and soil properties, root and litter traits (****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05). Refer to Table S5 for abbreviations.

		Bacteria			Fungi		
		H	NMDS1	NMDS2	H	NMDS1	NMDS2
Soil	Sand	0.13	0.14	0.30*	-0.14	-0.12	0.72****
	Silt	-0.2	0.1	-0.23	-0.05	0.30*	-0.49***
	Clay	0.04	-0.37**	-0.30*	0.2	-0.16	-0.68****
	pH	0.18	-0.24	-0.06	-0.25	-0.49***	0.17
	P	0.22	-0.30*	-0.23	0.17	-0.33*	-0.12
	Potassium	0.22	0.11	0.40**	-0.15	-0.15	0.50***
	Magnesium	0.06	0.25	0.49***	-0.07	0.19	0.41**
	SOC	-0.03	0.43**	0.41**	-0.34*	0.18	0.64****
	N	0.1	0.27	0.38**	-0.30*	-0.06	0.58****
	C	0.01	0.40**	0.42**	-0.31*	0.07	0.69****
	C:N	-0.29*	0.63****	0.25	-0.16	0.38**	0.55****
Roots	SRL	0.50***	-0.76****	-0.06	-0.02	-0.66****	-0.21
	VFR	0.2	-0.53****	-0.23	-0.19	-0.54****	-0.28
	FR	0	0.08	0.2	0.16	0.19	0.09
	MRD	-0.45***	0.77****	0.09	0.07	0.70****	0.21
	RDMC	-0.55****	0.37**	-0.17	-0.2	0.44**	-0.13
	RMD	0.1	-0.25	-0.17	-0.23	-0.37**	-0.05
	RLD	0.47***	-0.71****	-0.1	-0.06	-0.70****	-0.21
	N	-0.55****	0.42**	-0.36**	-0.14	0.64****	-0.36*
	C	-0.37**	0.57****	-0.01	-0.2	0.53****	-0.02
	C:N	0.39**	-0.21	0.43**	0.09	-0.42**	0.42**
	lignin	-0.15	0.17	0.19	-0.42	-0.05	0.21
	cellulose	-0.35	0.02	-0.22	0.31	0.56*	-0.45
	hemicellulose	0.18	-0.42	-0.25	0.35	-0.040	-0.07
Litter	N	0.40**	-0.25	0.28*	0.01	-0.34*	0.21
	C	0.25	0.05	0.29*	0.09	0.06	0.22
	C:N	-0.25	0.24	-0.15	0.02	0.28	-0.04
	lignin	0.15	0.07	0.27	-0.24	-0.01	0.22
	cellulose	-0.37**	0.40**	-0.12	0.11	0.45***	-0.14
	hemicellulose	0.23	-0.62****	-0.32*	0.07	-0.51****	-0.35*

Table 3. Effects of tree-gap, elevation and sample fraction and their interactions on the structure of bacterial and fungal communities assessed with PERMANOVA. The degrees of freedom (Df), sum of squares (sum of sqs), mean of squares (mean of sqs), the F. statistic, the proportion of the variance explained by each model (R^2) and probability (P) are shown.

Factors	Bacteria						Fungi					
	Df	Sum of sqs	Mean of sqs	F	R^2	P(>F)	Df	Sum of sqs	Mean of sqs	F	R^2	P(>F)
Tree-gap	1	1.0542	1.05417	8.7707	0.122	0.001	1	1.2443	1.24429	3.3684	0.062	0.001
Elevation	1	0.5703	0.57034	4.7452	0.066	0.001	1	1.7022	1.70222	4.608	0.085	0.001
Samplefraction	2	1.1862	0.59308	4.9344	0.137	0.001	2	0.8555	0.42776	1.158	0.043	0.163
Elevation:Tree-gap	1	0.3944	0.3944	3.2814	0.045	0.003	1	0.9148	0.91483	2.4765	0.046	0.001
Elevation:Samplefraction	2	0.2742	0.13708	1.1405	0.032	0.249	2	0.4946	0.24729	0.6694	0.025	0.994
Tree-gap:Samplefraction	2	0.2578	0.12892	1.0726	0.030	0.34	2	0.4086	0.2043	0.553	0.020	1
Elevation:Tree-gap:Samplefraction	2	0.2495	0.12474	1.0378	0.029	0.366	2	0.4484	0.22419	0.6069	0.022	0.998
Residuals	39	4.6875	0.12019		0.540		38	14.0373	0.3694		0.698	
Total	50	8.6741			1		49	20.1057			1	

Table 4. Analysis of covariance (ANCOVA) table showing the effects of tree-gap, elevation and sample fraction on bacterial and fungal taxonomic (H) and phylogenetic (PD) diversities and phylogenetic divergence (ses.MNTD). The degrees of freedom (Df), the F statistic and P values are shown. Refer to Table S5 for abbreviations.

		Bacteria		Fungi		
		F	P value	F	P value	
H	Tree-gap	1	10.25	0.003	0.048	0.829
	Elevation	1	14.89	<0.001	0.02	0.889
	Samplefraction	2	0.52	0.599	3.231	0.051
	Elevation:Tree-gap	1	0.432	0.515	2.102	0.155
	Elevation:Samplefraction	2	0.142	0.868	0.48	0.622
	Tree:Samplefraction	2	1.099	0.343	1.524	0.231
	Elevation:Tree-gap:Samplefraction	2	0.559	0.576	0.049	0.952
	PD	Tree-gap	1	20.7	<0.001	
Elevation		1	4.879	0.033		
Samplefraction		2	0.479	0.623		
Elevation:Tree-gap		1	0.141	0.710		
Elevation:Samplefraction		2	0.257	0.775		
Tree:Samplefraction		2	1.075	0.351		
Elevation:Tree-gap:Samplefraction		2	0.55	0.582		
ses.MNTD		Tree-gap	1	5.106	0.030	
	Elevation	1	1.864	0.180		
	Samplefraction	2	3.117	0.056		
	Elevation:Tree-gap	1	1.681	0.202		
	Elevation:Samplefraction	2	0.506	0.607		
	Tree:Samplefraction	2	1.237	0.301		
	Elevation:Tree-gap:Samplefraction	2	0.033	0.968		




Figure 1. a) Location of the plots in the three elevations (1400, 1700 and 2000m) where samples were taken (Map data: Google, Image ©2017 Digital Globe); b) pictures showing the three sites where root, soil and litter were collected for analysis and location of pairs of gaps (open circles) and closed forest (closed circles). Distances (in m), between each plot are indicated.




Figure 2. A selection of root and litter physical traits: (a) specific root length (SRL), (b) root length density (RLD); and chemical traits: (c) root nitrogen (N), (d) root carbon (C), (e) litter hemicellulose and (f) litter cellulose in gaps (white bars) and closed forest (green bars) along the elevation gradient. Boxplots represent the minimum, maximum, median, first quartile and third quartile in the data set.




Figure 3. Graphs of dbRDA constrained ordinations of bacterial and fungal species matrices with convex hull polygons containing plots of gaps (white) and closed forest (green) and significant variables obtained by automatic backward stepwise model building. Data are shown for roots (○), rhizosphere (□) and bulk soil (△) fractions at 1400 m (white symbols), 1700 m (grey symbols) and 2000 m (black symbols).




Figure 4. NMDS ordinations of a) bacterial and b) fungal species matrices with convex hull polygons containing plots of the two locations (gap and closed forest white and green shaded respectively) and spider diagrams linking plots with the same elevation (2000 and 1700 m tags are overlapped for bacteria). Sample fractions are shown with different symbols (\circ : root, \square : rhizosphere, \triangle : bulk soil).




Figure 5. Changes in a) bacterial taxonomic diversity (Shannon's diversity), b) phylogenetic diversity and c) phylogenetic clustering (standardized effect size of MNTD versus null communities, ses.MNTD) in gaps (white bars) and closed forest (green bars), along the elevational gradient for the three sample fractions (root, rhizosphere and soil). Boxplots represent the minimum, maximum, median, first quartile and third quartile in the data set.

Supporting Information

Table S1. Dominant species and abundance of herbs and trees in gaps and closed forest along the elevation gradient. Abundance of herbs and trees is given as a percentage foliar cover. Age to last disturbance (Dist) is the time (in years) to the last disturbance (tree thinning). Taken from Prieto et al. (2015).

Elevation	Land use type	Dominant species	Herbs (% cover)	Trees (% cover)	Dist (years)
1400	Gap	<i>Galium rotundifolium</i> L.,	89	0	5
		<i>Lysimachia nemorum</i> L.,			
		<i>Luzula nivea</i> (Nath.) DC			
1400	Closed forest	<i>Abies alba</i> Mill.,	10	90	40
		<i>Picea abies</i> (L.) H. Karst.,			
		<i>Fagus sylvatica</i> L.			
1700	Gap	<i>Luzula nivea</i> (Nath.) DC	55	0	5
		<i>Rhododendron ferrugineum</i> L.,			
		<i>Vaccinium myrtillus</i> L.			
1700	Closed forest	<i>Picea abies</i> (L.) H. Karst.,	0	90	40
		<i>Abies alba</i> Mill.			
2000	Gap	<i>Rhododendron ferrugineum</i> L.,	50	0	5
		<i>Vaccinium myrtillus</i> L.			
2000	Closed forest	<i>Pinus uncinata</i> Ramond ex. DC.,	10	90	40
		<i>Picea abies</i> (L.) H. Karst.			

Table S2. Climate and microclimate data (mean \pm standard error) along the elevation gradient in gaps and closed forest. P-values of analysis of covariance (ANCOVA) tests are shown for air and soil temperatures and Friedman tests for soil water potential. Number of repeated measures is shown between brackets. (**P < 0.001, *P < 0.01, *P < 0.05). Post hoc comparisons performed with Nemenyi multiple tests.

	1400		1700		2000		Tree	Elevation	Elevation*	Tree	
	Gap	Closed forest	Gap	Closed forest	Gap	Closed forest	p	p	p		
Air temperature (°C)	8.24 \pm 1.04 (68)	7.73 \pm 1.01 ns (66)	8.59 \pm 1.04 (67)	7.44 \pm 0.93 ** (66)	7.33 \pm 0.94 (51)	6.96 \pm 0.94 ns (56)	0.08	ns	<0.001	***	0.91 ns
Soil temperature 10 cm depth (°C)	6.48 \pm 0.57 (68)	6.36 \pm 0.57 ns (66)	6.68 \pm 0.63 (67)	5.27 \pm 0.46 * (66)	5.64 \pm 0.63 (51)	5.42 \pm 0.55 ns (56)	:0.00	***	<0.001	***	0.36 ns
Soil water potential 20 cm depth (kpa)	-10.67 \pm 2.54 (11)	-59.15 \pm 17.31 ns (11)	-13.88 \pm 2.43 (11)	-81.23 \pm 24.79 ns (11)	-	-	0.13	ns	0.37	ns	-

Table S3. Soil physicochemical properties along the elevation gradient in gaps and closed forest. p-values from analysis of covariance (ANCOVA) tests are shown. Two sample t-test significant results at $p < 0.05$ are shown with the symbol "+" for gap versus closed forest. Kruskal-Wallis and Wilcoxon test were performed for p. Refer to Table S5 for abbreviations.

	1400		1700		2000		Tree	Elevation	Elevation*Tree
	Gap	Closed forest	Gap	Closed forest	Gap	Closed forest	p	p	p
MWD (mm)	3.37 ± 0.03	3.35 ± 0.02	3.33 ± 0.02	3.33 ± 0.03	3.02 ± 0.03	3.31 ± 0.03	0.102	0.008	0.028
pH	4.43 ± 0.09	4.80 ± 0.13	5.03 ± 0.11	4.90 ± 0.08	5.27 ± 0.07	5.03 ± 0.07	0.900	0.022	0.147
P (mg/l)	4.60 ± 0.50	5.47 ± 0.35	3.13 ± 0.09	2.73 ± 0.12	9.40 ± 0.62	3.73 ± 0.09	0.148	0.005	-
Potassium (mg l⁻¹)	30.73 ± 5.66	31.80 ± 1.33	44.97 ± 5.27	50.77 ± 2.68	49.43 ± 3.64	63.53 ± 6.02	0.465	0.022	0.238
Magnesium (mg l⁻¹)	44.53 ± 5.75	33.57 ± 3.48	69.43 ± 6.71	94.20 ± 6.83	32.37 ± 3.23	67.60 ± 5.80	0.167	0.416	0.250
Sand (% w w⁻¹)	35.67 ± 2.13	44.00 ± 0.29	40.67 ± 0.83	58.33 ± 2.19	53.67 ± 1.59	46.67 ± 0.33	0.133	0.066	0.117
Silt (% w w⁻¹)	35.67 ± 0.73	31.67 ± 0.17	31.00 ± 0.50	24.00 ± 1.26	24.33 ± 0.73	31.00 ± 0.58	0.571	0.037	0.039
Clay (% w w⁻¹)	28.67 ± 1.42	24.33 ± 0.44	28.33 ± 0.33	17.67 ± 0.93	22.00 ± 0.87	22.33 ± 0.88	0.031	0.171	0.404
SOC (% w w⁻¹)	12.00 ± 0.32	11.53 ± 1.53	15.33 ± 0.40	22.70 ± 0.77	12.63 ± 1.86	22.17 ± 1.29	0.028	0.081	0.087
N (% w w⁻¹)	0.34 ± 0.01	0.33 ± 0.05	0.47 ± 0.01	0.50 ± 0.01	0.42 ± 0.07	0.64 ± 0.03	0.185	0.024	0.097
TC (% w w⁻¹)	5.37 ± 0.12	5.53 ± 0.75	7.47 ± 0.16	11.37 ± 0.42	6.37 ± 0.92	12.43 ± 0.16	0.004	0.010	0.029
C : N	15.71 ± 0.44	16.70 ± 0.10	15.97 ± 0.16	22.70 ± 0.64	15.33 ± 0.17	19.50 ± 0.64	0.002	0.380	0.346

Table S4. Root and litter traits along the elevation gradient in gaps and closed forest. P-values of analysis of variance (ANOVA) tests or Kruskal Wallis are shown. Different letters show post-hoc Tukey honestly significant difference (HSD) results for elevation. Two sample t-test significant results at $p < 0.05$ are shown with the symbol "+" for gap versus closed forest. Refer to Table S5 for abbreviations.

	1400		1700		2000		Tree p	Elevation p	Elevation*Tree p
	Gap	Closed forest	Gap	Closed forest	Gap	Closed forest			
Root traits									
SRL (m gr ⁻¹)	22.61 ± 4.21	8.16 ± 1.28	59.20 ± 4.49	10.36 ± 3.49	83.76 ± 28.40	11.66 ± 0.83	<0.001	0.005	0.037
VFR (%)	40.65 ± 1.80	49.69 ± 5.30	57.86 ± 3.48	39.29 ± 6.30	54.76 ± 4.05	45.41 ± 4.39	0.146	0.343	0.088
FR (%)	52.01 ± 3.86	34.66 ± 5.79	40.52 ± 3.30	46.83 ± 4.52	43.65 ± 3.22	44.81 ± 4.67	0.393	0.847	0.063
Mean Root Diameter (mm)	0.40 ± 0.03	0.53 ± 0.02	0.26 ± 0.02	0.52 ± 0.09	0.27 ± 0.03	0.46 ± 0.02	<0.001	0.030	0.499
RDMC (gr gr ⁻¹)	0.37 ± 0.01	0.40 ± 0.01	0.37 ± 0.04	0.39 ± 0.02	0.33 ± 0.01	0.37 ± 0.02	0.076	0.138	0.623
RMD (g m ⁻³)	228.65 ± 56.37	272.16 ± 37.30	332.37 ± 44.57	211.41 ± 31.25	406.38 ± 128.18	341.92 ± 65.62	0.086	0.402	0.434
RLD (km m ⁻³)	4.70 ± 0.13	2.30 ± 0.67	19.30 ± 1.22	2.05 ± 0.47	40.01 ± 18.25	4.05 ± 1.01	0.001	0.264	-
Lignin (mg g ⁻¹)	46.04	54.56	63.37	54.67	50.36	59.74	-	-	-
Cellulose (mg g ⁻¹)	27.90	22.61	18.75	21.82	19.61	16.16	-	-	-
Hemicellulose (mg g ⁻¹)	21.97	17.67	17.47	19.00	31.05	16.87	-	-	-
N (mg g ⁻¹)	0.85 ± 0.03	1.01 ± 0.04	0.73 ± 0.09	0.78 ± 0.02	0.72 ± 0.01	0.71 ± 0.10	0.200	0.003	0.170
C (mg/g)	45.67 ± 1.15	46.88 ± 1.43	44.63 ± 0.19	45.82 ± 0.14	42.78 ± 0.92	44.85 ± 1.44	0.022	0.077	0.662
C : N	54.01 ± 3.03	46.39 ± 1.52	62.89 ± 8.12	58.92 ± 1.41	59.13 ± 1.58	65.16 ± 7.85	0.643	0.026	0.176
Litter traits									
Lignin (mg g ⁻¹)	47.10 ± 2.45	45.57 ± 2.66	49.11 ± 1.74	51.53 ± 3.71	48.79 ± 2.81	51.21 ± 1.38	0.595	0.164	0.442
Cellulose (mg g ⁻¹)	21.57 ± 2.15	27.67 ± 0.66	20.87 ± 2.11	22.73 ± 1.95	18.54 ± 0.89	20.61 ± 1.79	0.023	0.007	0.231
Hemicellulose (mg g ⁻¹)	20.50 ± 1.47	17.93 ± 2.74	22.49 ± 1.41	15.84 ± 1.77	26.34 ± 2.52	17.16 ± 1.13	0.004	0.751	-
N (mg g ⁻¹)	1.16 ± 0.02	0.84 ± 0.06	0.95 ± 0.21	0.92 ± 0.21	1.70 ± 0.53	1.12 ± 0.10	0.122	0.240	-
C (mg g ⁻¹)	40.97 ± 1.13	38.72 ± 2.18	37.93 ± 3.56	43.09 ± 1.15	52.63 ± 12.92	40.41 ± 1.18	0.825	0.864	-
C : N	35.44 ± 0.42	46.87 ± 5.68	44.12 ± 10.09	53.96 ± 15.69	33.05 ± 4.88	36.68 ± 2.56	0.058	0.386	-

549 **Table S5.** Abbreviations used in this paper

Variable	abbreviation	units
Soil properties		
Mean weight diameter	MWD	mm
P	Phosphorus	mg·l ⁻¹
Soil organic carbon	SOC	% w·w ⁻¹
Total carbon	TC	% w·w ⁻¹
Root traits		
Specific root length	SRL	m·g ⁻¹
Very fine roots	VFR	%
Fine roots	FR	%
Mean Root Diameter	MRD	mm
Root dry matter content	RDMC	mg·g ⁻¹
Root mass density	RMD	g·m ⁻³
Root length density	RLD	km·m ⁻³
General abbreviations		
Nitrogen	N	% w·w ⁻¹
Carbon	C	% w·w ⁻¹
C-to-N ratio	C:N	none
Shannon's diversity	H	none
Faith's Phylogenetic Diversity	PD	none
Standardized mean nearest taxon distance	ses.MNTD	none




Fig. S1. Phylum and class of bacterial and fungal communities along the elevation and in gaps and closed forest. Proportion of presence within sampling unit. Only the most abundant classes are shown (for which the sum of proportions for all sites were greater than one). **Zygomycota* assignation here was before their later reclassification among *Glomeromycota* and several subphyla *incertae sedis* (Hibbett et al. 2007).




Fig. S2. Soil temperature data in gaps and closed forest, over time and for the three elevations: a) soil temperature at a depth of 10 cm and b) soil temperature at a depth of 40 cm.

Notes S1 Additional methodological details on Illumina amplicon sequencing of 16S rRNA genes and the ITS region.

Amplicons were generated using a high-fidelity DNA polymerase (Q5 Taq, New England Biolabs) and pooled. PCR was conducted on 20 ng of template DNA employing an initial denaturation of 30 seconds at 95 °C, followed by cycles (25 for 16S and 30 for ITS) of 30 seconds at 95 °C, 30 seconds at 52 °C and 2 minutes at 72 °C. A final extension of 10 minutes at 72 °C was also included to complete the reaction. Amplicon sizes were determined using an Agilent 2200 TapeStation system (~550bp:16S; ~350-425: ITS; ~650:18S) and libraries normalized using SequalPrep Normalization Plate Kit (Thermo Fisher Scientific). Library concentration was calculated using a SYBR green quantitative PCR (qPCR) assay with primers specific to the Illumina adapters (Kappa, Anachem). Libraries were sequenced at a concentration of 5.4 pM with a 0.6 pM addition of an Illumina generated PhiX control library. Sequencing runs, generating 2 x 300 bp, reads were performed on an Illumina MiSeq using V3 chemistry. The read 1 (R1), read 2 (R2) and index sequencing primers used were also gene specific: R1 = sequence of the combined pad, linker and forward primer (e.g. 314F; 16S) or; R2 = sequence of the combined pad, linker and reverse primer (e.g. 806R; 16S); I = reverse complement of the R2 primer.

- 551 Aponte C, García LV, Marañón T (2013) Tree species effects on nutrient cycling and soil biota: A
552 feedback mechanism favouring species coexistence. *For Ecol Manage* 309:36-46
- 553 Bach LH, Grytnes J-A, Halvorsen R, Ohlson M (2010) Tree influence on soil microbial community
554 structure. *Soil Biol Biochem* 42:1934-1943
- 555 Bahram M, Polme S, Koljalg U, Zarre S, Tedersoo L (2012) Regional and local patterns of
556 ectomycorrhizal fungal diversity and community structure along an altitudinal gradient in the
557 Hyrcanian forests of northern Iran. *New Phytol* 193:465-473
- 558 Baldrian P (2017) Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiol Rev* 41:109-
559 130
- 560 Bardgett RD, Mommer L, De Vries FT (2014) Going underground: root traits as drivers of ecosystem
561 processes. *Trends Ecol Evol* 29:692-699
- 562 Baumert VL, Vasilyeva NA, Vladimirov AA, Meier IC, Kögel-Knabner I, Mueller CW (2018) Root
563 Exudates Induce Soil Macroaggregation Facilitated by Fungi in Subsoil. *Front Environ Sci* 6
- 564 Bever JD, Platt TG, Morton ER (2012) Microbial population and community dynamics on plant roots and
565 their feedbacks on plant communities. *Annu Rev Microbiol* 66:265-283
- 566 Brant JB, Myrold DD, Sulzman EW (2006) Root controls on soil microbial community structure in forest
567 soils. *Oecologia* 148:650-659
- 568 Brundrett MC (2002) Coevolution of roots and mycorrhizas of land plants. *New Phytol* 154:275-304
- 569 Bryant JA, Lamanna C, Morlon H, Kerkhoff AJ, Enquist BJ, Green JL (2008) Microbes on
570 mountainsides: Contrasting elevational patterns of bacterial and plant diversity. *PNAS*
571 105:11505-11511
- 572 Bulgarelli D, Garrido-Oter R, Münch Philipp C, Weiman A, Dröge J, Pan Y, McHardy Alice C, Schulze-
573 Lefert P (2015) Structure and Function of the Bacterial Root Microbiota in Wild and
574 Domesticated Barley. *Cell Host & Microbe* 17:392-403
- 575 Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P,
576 Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-
577 Lefert P (2012) Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial
578 microbiota. *Nature* 488:91-95
- 579 Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, Knight R (2010) PyNAST: a
580 flexible tool for aligning sequences to a template alignment. *Bioinformatics* 26:266-267
- 581 Chenu C, Sotzky G (2002) Interactions Between Microorganisms and Soil Particles. In: Interactions
582 between soil particles and microorganisms. IUPAC series of Applied Chemistry. pp 3-40.
- 583 Coince A, Cordier T, Lengellé J, Defossez E, Vacher C, Robin C, Buée M, Marçais B (2014) Leaf and
584 Root-Associated Fungal Assemblages Do Not Follow Similar Elevational Diversity Patterns.
585 *PLOS ONE* 9:e100668
- 586 Colin Y, Nicolitch O, Van Nostrand JD, Zhou JZ, Turpault MP, Uroz S (2017) Taxonomic and functional
587 shifts in the beech rhizosphere microbiome across a natural soil toposequence. *Sci Rep* 7:9604
- 588 Collins CG, Stajich JE, Weber SE, Pombubpa N, Diez JM (2018) Shrub range expansion alters diversity
589 and distribution of soil fungal communities across an alpine elevation gradient. *Mol Ecol*
- 590 de Freitas CR, Enright NJ (1995) Microclimatic differences between and within canopy gaps in a
591 temperate rainforest. *International Journal of Biometeorology* 38:188-193
- 592 de Graaff MA, Classen AT, Castro HF, Schadt CW (2010) Labile soil carbon inputs mediate the soil
593 microbial community composition and plant residue decomposition rates. *New Phytol* 188:1055-
594 1064
- 595 Dean WE (1974) Determination of carbonate and organic matter in calcareous sediments and sedimentary
596 rocks by loss on ignition; comparison with other methods. *J Sediment Res* 44:242-248
- 597 DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,
598 Andersen GL (2006a) Greengenes, a chimera-checked 16S rRNA gene database and workbench
599 compatible with ARB. *Appl Environ Microbiol* 72:5069-5072
- 600 DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,
601 Andersen GL (2006b) Greengenes, a Chimera-Checked 16S rRNA Gene Database and
602 Workbench Compatible with ARB. *Applied and Environmental Microbiology* 72:5069-5072
- 603 Dray S, Dufour A-B (2007) The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal*
604 *of Statistical Software* 22:1-20
- 605 Dukunde A, Schneider D, Schmidt M, Veldkamp E, Daniel R (2019) Tree Species Shape Soil Bacterial
606 Community Structure and Function in Temperate Deciduous Forests. *Front Microbiol* 10
- 607 Faith DP (1992) Conservation evaluation and phylogenetic diversity. *Biological Conservation* 61:1-10

- 608 Gale WJ, Cambardella CA, Bailey TB (2000) Root-derived carbon and the formation and stabilization of
609 aggregates. *Soil Sci Soc Am J* 64:201-207
- 610 Goberna M, García C, Verdú M (2014a) A role for biotic filtering in driving phylogenetic clustering in
611 soil bacterial communities. *Global Ecol Biogeogr* 23:1346-1355
- 612 Goberna M, Navarro-Cano JA, Valiente-Banuet A, García C, Verdú M (2014b) Abiotic stress tolerance
613 and competition-related traits underlie phylogenetic clustering in soil bacterial communities.
614 *Ecol Lett* 17:1191-1201
- 615 Gray AN, Spies TA, Easter MJ (2002) Microclimatic and soil moisture responses to gap formation in
616 coastal Douglas-fir forests. *Can J For Res* 32:332-343
- 617 Grigulis K, Lavorel S, Krainer U, Legay N, Baxendale C, Dumont M, Kastl E, Arnoldi C, Bardgett RD,
618 Poly F, Pommier T, Schloter M, Tappeiner U, Bahn M, Clément J-C (2013) Relative
619 contributions of plant traits and soil microbial properties to mountain grassland ecosystem
620 services. *J Ecol* 101:47-57
- 621 Gweon HS, Oliver A, Taylor J, Booth T, Gibbs M, Read DS, Griffiths RI, Schonrogge K (2015) PIPITS:
622 an automated pipeline for analyses of fungal internal transcribed spacer sequences from the
623 Illumina sequencing platform. *Methods Ecol Evol* 6:973-980
- 624 Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, Huhndorf S, James T, Kirk
625 PM, Lücking R, Thorsten Lumbsch H, Lutzoni F, Matheny PB, McLaughlin DJ, Powell MJ,
626 Redhead S, Schoch CL, Spatafora JW, Stalpers JA, Vilgalys R, Aime MC, Aptroot A, Bauer R,
627 Begerow D, Benny GL, Castlebury LA, Crous PW, Dai Y-C, Gams W, Geiser DM, Griffith
628 GW, Gueidan C, Hawksworth DL, Hestmark G, Hosaka K, Humber RA, Hyde KD, Ironside JE,
629 Kõljalg U, Kurtzman CP, Larsson K-H, Lichtwardt R, Longcore J, Miądlikowska J, Miller A,
630 Moncalvo J-M, Mozley-Standridge S, Oberwinkler F, Parmasto E, Reeb V, Rogers JD, Roux C,
631 Ryvarden L, Sampaio JP, Schüßler A, Sugiyama J, Thorn RG, Tibell L, Untereiner WA, Walker
632 C, Wang Z, Weir A, Weiss M, White MM, Winka K, Yao Y-J, Zhang N (2007) A higher-level
633 phylogenetic classification of the Fungi. *Mycol Res* 111:509-547
- 634 Horner-Devine MC, Bohannan BJM (2006) Phylogenetic clustering and overdispersion in bacterial
635 communities. *Ecology* 87:S100-S108
- 636 Hornung M (1985) Acidification of soils by trees and forests. *Soil Use Manag* 1:24-27
- 637 Ihrmark K, Bodeker IT, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y, Stenlid J,
638 Brandstrom-Durling M, Clemmensen KE, Lindahl BD (2012) New primers to amplify the fungal
639 ITS2 region—evaluation by 454-sequencing of artificial and natural communities. *FEMS*
640 *Microbiol Ecol* 82:666-677
- 641 IWG W (2007) World reference base for soil resources 2006, first update 2007. FAO, Rome
- 642 Jarvis SG, Woodward S, Taylor AF (2015) Strong altitudinal partitioning in the distributions of
643 ectomycorrhizal fungi along a short (300 m) elevation gradient. *New Phytol* 206:1145-1155
- 644 Jonard M, Nicolas M, Coomes DA, Caignet I, Saenger A, Ponette Q (2017) Forest soils in France are
645 sequestering substantial amounts of carbon. *Sci Total Environ* 574:616-628
- 646 Joud D (2006) Guide pour identifier les stations forestières de Rhône-Alpes—Synthèse pour les Alpes du
647 Nord et les montagnes de l'Ain. In: Rhône-Alpes C (ed). p 132.
- 648 Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, Blomberg SP, Webb CO
649 (2010) Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26:1463-1464
- 650 Kernaghan G, Harper KA (2001) Community structure of ectomycorrhizal fungi across an
651 alpine/subalpine ecotone. *Ecography* 24:181-188
- 652 Kohout P, Charvátová M, Štursová M, Mašínová T, Tomšovský M, Baldrian P (2018) Clearcutting alters
653 decomposition processes and initiates complex restructuring of fungal communities in soil and
654 tree roots. *ISME J*
- 655 Koljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST, Bruns TD,
656 Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Duenas M, Grebenc
657 T, Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking R, Martin
658 MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Poldmaa
659 K, Saag L, Saar I, Schussler A, Scott JA, Senes C, Smith ME, Suija A, Taylor DL, Telleria MT,
660 Weiss M, Larsson KH (2013) Towards a unified paradigm for sequence-based identification of
661 fungi. *Mol Ecol* 22:5271-5277
- 662 Körner C (2007) The use of 'altitude' in ecological research. *Trends Ecol Evol* 22:569-574
- 663 Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD (2013) Development of a dual-index
664 sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq
665 Illumina sequencing platform. *Appl Environ Microbiol* 79:5112-5120
- 666 Kuzyakov Y, Blagodatskaya E (2015) Microbial hotspots and hot moments in soil: Concept & review.
667 *Soil Biol Biochem* 83:184-199

668 Lareen A, Burton F, Schafer P (2016) Plant root-microbe communication in shaping root microbiomes.
669 Plant Mol Biol 90:575-587

670 Le Bissonnais Y (1996) Aggregate stability and assessment of soil crustability and erodibility: I. Theory
671 and methodology. Eur J Soil Sci 47:425-437

672 Lehmann A, Zheng W, Rillig MC (2017) Soil biota contributions to soil aggregation. Nat Ecol Evol

673 Liu J, Ngoc Ha V, Shen Z, Dang P, Zhu H, Zhao F, Zhao Z (2018) Response of the rhizosphere microbial
674 community to fine root and soil parameters following Robinia pseudoacacia L. afforestation.
675 Appl Soil Ecol 132:11-19

676 Lladó S, López-Mondéjar R, Baldrian P (2018) Drivers of microbial community structure in forest soils.
677 Appl Microbiol Biotechnol 102:4331-4338

678 Mao Z, Jourdan C, Bonis M-L, Pailler F, Rey H, Saint-André L, Stokes A (2013) Modelling root
679 demography in heterogeneous mountain forests and applications for slope stability analysis.
680 Plant Soil 363:357-382

681 Mao Z, Saint-André L, Genet M, Mine FX, Jourdan C, Rey H, Courbaud B, Stokes A (2012) Engineering
682 ecological protection against landslides in diverse mountain forests: Choosing cohesion models.
683 Ecol Eng 45:55-69

684 Mao Z, Wang Y, Jourdan C, Cécillon L, Nespoulous J, Rey H, Saint-André L, Stokes A (2015)
685 Characterizing Above- and Belowground Carbon Partitioning in Forest Trees along an
686 Altitudinal Gradient using Area-Based Indicators. Arct Antarct Alp Res 47:59-69

687 Mayfield MM, Levine JM (2010) Opposing effects of competitive exclusion on the phylogenetic structure
688 of communities. Ecol Lett 13:1085-1093

689 McCain CM, Grytnes J-A (2010) Elevational Gradients in Species Richness. In: Encyclopedia of Life
690 Sciences (ELS). John Wiley & Sons, Ltd.

691 McCave IN, Bryant RJ, Cook HF, Coughanowr CA (1986) Evaluation of a laser-diffraction-size analyzer
692 for use with natural sediments. J Sediment Res 56:561-564

693 Miller RM, Jastrow JD (1990) Hierarchy of root and mycorrhizal fungal interactions with soil
694 aggregation. Soil Biology and Biochemistry 22:579-584

695 Miyamoto Y, Nakano T, Hattori M, Nara K (2014) The mid-domain effect in ectomycorrhizal fungi:
696 range overlap along an elevation gradient on Mount Fuji, Japan. Isme j 8:1739-1746

697 Muscolo A, Bagnato S, Sidari M, Mercurio R (2014) A review of the roles of forest canopy gaps. J For
698 Res 25:725-736

699 Muscolo A, Sidari M, Mercurio R (2007) Influence of gap size on organic matter decomposition,
700 microbial biomass and nutrient cycle in Calabrian pine (*Pinus laricio*, Poiret) stands. For Ecol
701 Manage 242:412-418

702 Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB,
703 Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H (2016) vegan: Community
704 Ecology Package. R package version 2.4-0. <https://CRAN.R-project.org/package=vegan>.

705 Olsen SR, Cole CV, Watanabe FS, Dean LA, United S, Department of A (1954) Estimation of available
706 phosphorus in soils by extraction with sodium bicarbonate. U.S. Dept. of Agriculture,
707 Washington, D.C.

708 Ostonen I, Truu M, Helmisaari HS, Lukac M, Borken W, Vanguelova E, Godbold DL, Lohmus K, Zang
709 U, Tedersoo L, Preem JK, Rosenvald K, Aosaar J, Armolaitis K, Frey J, Kabral N, Kukumagi M,
710 Leppalammi-Kujansuu J, Lindroos AJ, Merila P, Napa U, Nojd P, Parts K, Uri V, Varik M, Truu
711 J (2017) Adaptive root foraging strategies along a boreal-temperate forest gradient. New Phytol
712 215:977-991

713 Parks DH, Beiko RG (2013) Measures of phylogenetic differentiation provide robust and complementary
714 insights into microbial communities. ISME J 7:173-183

715 Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH (2013) Going back to the roots: the
716 microbial ecology of the rhizosphere. Nat Rev Microbiol 11:789-799

717 Poirier V, Roumet C, Angers DA, Munson AD (2018) Species and root traits impact macroaggregation in
718 the rhizospheric soil of a Mediterranean common garden experiment. Plant Soil 424:289-302

719 Prescott CE, Grayston SJ (2013) Tree species influence on microbial communities in litter and soil:
720 Current knowledge and research needs. For Ecol Manage 309:19-27

721 Price MN, Dehal PS, Arkin AP (2010) FastTree 2 – Approximately Maximum-Likelihood Trees for
722 Large Alignments. PLOS ONE 5:e9490

723 Prieto I, Roumet C, Cardinael R, Dupraz C, Jourdan C, Kim JH, Maeght JL, Mao Z, Pierret A, Portillo N,
724 Rounsard O, Thammahacksa C, Stokes A, Cahill J (2015) Root functional parameters along a
725 land - use gradient: evidence of a community - level economics spectrum. J Ecol 103:361-373

726 Ren C, Zhang W, Zhong Z, Han X, Yang G, Feng Y, Ren G (2018) Differential responses of soil
727 microbial biomass, diversity, and compositions to altitudinal gradients depend on plant and soil
728 characteristics. *Sci Total Environ* 610-611:750-758

729 Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open source tool for
730 metagenomics. *PeerJ* 4:e2584

731 RStudio Team (2016) RStudio: Integrated Development for R. RStudio Inc., Boston, MA URL

732 Saetre P, Bååth E (2000) Spatial variation and patterns of soil microbial community structure in a mixed
733 spruce–birch stand. *Soil Biol Biochem* 32:909-917

734 Schneider T, Keiblinger KM, Schmid E, Sterflinger-Gleixner K, Ellersdorfer G, Roschitzki B, Richter A,
735 Eberl L, Zechmeister-Boltenstern S, Riedel K (2012) Who is who in litter decomposition?
736 Metaproteomics reveals major microbial players and their biogeochemical functions. *ISME J*
737 6:1749-1762

738 Shea F, Watts CE (1939) Dumas method for organic nitrogen. *Industrial & Engineering Chemistry*
739 *Analytical Edition* 11:333-334

740 Shen C, Liang W, Shi Y, Lin X, Zhang H, Wu X, Xie G, Chain P, Grogan P, Chu H (2014) Contrasting
741 elevational diversity patterns between eukaryotic soil microbes and plants. *Ecology* 95:3190-
742 3202

743 Shi S, O’Callaghan M, Jones EE, Richardson AE, Walter C, Stewart A, Condrón L (2012) Investigation
744 of organic anions in tree root exudates and rhizosphere microbial communities using in situ and
745 destructive sampling techniques. *Plant Soil* 359:149-163

746 Siles JA, Margesin R (2016) Abundance and Diversity of Bacterial, Archaeal, and Fungal Communities
747 Along an Altitudinal Gradient in Alpine Forest Soils: What Are the Driving Factors? *Microb*
748 *Ecol* 72:207-220

749 Singh D, Takahashi K, Kim M, Chun J, Adams JM (2012) A hump-backed trend in bacterial diversity
750 with elevation on Mount Fuji, Japan. *Microb Ecol* 63:429-437

751 Tindall BJ, Rossello-Mora R, Busse HJ, Ludwig W, Kämpfer P (2010) Notes on the characterization of
752 prokaryote strains for taxonomic purposes. *International journal of systematic and evolutionary*
753 *microbiology* 60:249-266

754 Tisdall JM, Oades JM (1982) Organic matter and water-stable aggregates in soils. *J Soil Sci* 33:141-163

755 Urbanová M, Šnajdr J, Baldrian P (2015) Composition of fungal and bacterial communities in forest litter
756 and soil is largely determined by dominant trees. *Soil Biol Biochem* 84:53-64

757 Uren NC (2000) Types, amounts, and possible functions of compounds released into the rhizosphere by
758 soil-grown plants. In: *The rhizosphere*. CRC Press, pp 35-56.

759 Uroz S, Buée M, Murat C, Frey-Klett P, Martin F (2010) Pyrosequencing reveals a contrasted bacterial
760 diversity between oak rhizosphere and surrounding soil. *Environ Microbiol Rep* 2:281-288

761 Uroz S, Oger P, Tisserand E, Cébron A, Turpault MP, Buée M, De Boer W, Leveau JHJ, Frey-Klett P
762 (2016) Specific impacts of beech and Norway spruce on the structure and diversity of the
763 rhizosphere and soil microbial communities. *Sci Rep* 6:27756

764 van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB, Fukami T, Kardol P, Klironomos
765 JN, Kulmatiski A, Schweitzer JA, Suding KN, Van de Voorde TFJ, Wardle DA (2013) Plant–
766 soil feedbacks: the past, the present and future challenges. *J Ecol* 101:265-276

767 Van Soest PJ (1963) Use of detergents in the analysis of fibrous feeds. 2. A rapid method for the
768 determination of fiber and lignin. *Journal of the Association of Official Agricultural Chemists*
769 46:829-835

770 Wang J, Soininen J, He J, Shen J (2012) Phylogenetic clustering increases with elevation for microbes.
771 *Environ Microbiol Rep* 4:217-226

772 Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for Rapid Assignment of
773 rRNA Sequences into the New Bacterial Taxonomy. *Applied and Environmental Microbiology*
774 73:5261-5267

775 Wang Y, Kim JH, Mao Z, Ramel M, Paillet F, Perez J, Rey H, Tron S, Jourdan C, Stokes A (2018a) Tree
776 root dynamics in montane and sub-alpine mixed forest patches. *Ann Bot* 122:861-872

777 Wang Y, Mao Z, Bakker MR, Kim JH, Brancheriau L, Buatois B, Leclerc R, Selli L, Rey H, Jourdan C,
778 Stokes A (2018b) Linking conifer root growth and production to soil temperature and carbon
779 supply in temperate forests. *Plant Soil* 426:33-50

780 Webb CO (2000) Exploring the phylogenetic structure of ecological communities: an example for rain
781 forest trees. *The American Naturalist* 156:145-155

782 Yang B, Pang X, Hu B, Bao W, Tian G (2017a) Does thinning-induced gap size result in altered soil
783 microbial community in pine plantation in eastern Tibetan Plateau? *Ecology and evolution*
784 7:2986-2993

785 Yang Y, Geng Y, Zhou H, Zhao G, Wang L (2017b) Effects of gaps in the forest canopy on soil microbial
786 communities and enzyme activity in a Chinese pine forest. *Pedobiologia* 61:51-60
787 Zhang J, Kobert K, Flouri T, Stamatakis A (2014) PEAR: a fast and accurate Illumina Paired-End reAd
788 mergeR. *Bioinformatics* 30:614-620
789 Zhang Q, Goberna M, Liu Y, Cui M, Yang H, Sun Q, Insam H, Zhou J (2018) Competition and habitat
790 filtering jointly explain phylogenetic structure of soil bacterial communities across elevational
791 gradients. *Environ Microbiol* 20:2386-2396
792