



HAL
open science

Ventilator-induced diaphragmatic dysfunction

Basil Petrof, Samir Jaber, Stefan Matecki

► **To cite this version:**

Basil Petrof, Samir Jaber, Stefan Matecki. Ventilator-induced diaphragmatic dysfunction. *Current Opinion in Critical Care*, 2010, 16 (1), pp.19-25. 10.1097/MCC.0b013e328334b166 . hal-02546264

HAL Id: hal-02546264

<https://hal.umontpellier.fr/hal-02546264>

Submitted on 30 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ventilator-induced diaphragmatic dysfunction

Basil J. Petrof^a, Samir Jaber^b and Stefan Matecki^c

^aMeakins-Christie Laboratories and Respiratory Division, McGill University Health Center and Research Institute, Montreal, Quebec, Canada, ^bIntensive Care Unit, Anesthesia and Critical Care Department B, Saint Eloi Teaching Hospital, Equipe soutenue par la Région et l'Institut National de la Santé et de la Recherche Médicale 25, Université Montpellier 1, Centre Hospitalier Universitaire Montpellier and ^cClinical Physiology Center, Arnaud de Villeneuve Teaching Hospital, Equipe soutenue par la Région et l'Institut National de la Santé et de la Recherche Médicale 25, Université Montpellier 1, Centre Hospitalier Universitaire Montpellier, Montpellier, France

Correspondence to Basil J. Petrof, MD, Royal Victoria Hospital, Room L4-11, 687 Pine Avenue West, Montreal, QC H3A 1A1, Canada
Tel: +1 514 934 1934x35946;
fax: +1 514 843 1695; e-mail: basil.petrof@mcgill.ca

Purpose of review

Diaphragmatic function is a major determinant of the ability to successfully wean patients from mechanical ventilation. There is increasing recognition of a condition termed ventilator-induced diaphragmatic dysfunction. The purpose of the present review is to present evidence that mechanical ventilation can itself be a cause of diaphragmatic dysfunction, to outline our current understanding of the cellular mechanisms responsible for this phenomenon, and to discuss the implications of recent research for future therapeutic strategies.

Recent findings

Many critically ill patients demonstrate diaphragmatic weakness. A large body of evidence from animal models, and more limited data from humans, indicates that mechanical ventilation can cause muscle fiber injury and atrophy within the diaphragm. Current data support a complex underlying pathophysiology involving oxidative stress and the activation of several intracellular proteolytic pathways involved in degradation of the contractile apparatus. This includes the calpain, caspase, and ubiquitin–proteasome systems. In addition, there is a simultaneous downregulation of protein synthesis pathways. Studies in animal models suggest that future therapies may be able to specifically target these processes, whereas for the time being current preventive measures in humans are primarily based upon allowing persistent diaphragmatic activation during mechanical ventilation.

Summary

Diaphragmatic dysfunction is common in mechanically ventilated patients and is a likely cause of weaning failure. Recently, there has been a great expansion in our knowledge of how mechanical ventilation can adversely affect diaphragmatic structure and function. Future studies need to better define the evolution and mechanistic basis for ventilator-induced diaphragmatic dysfunction in humans, in order to allow the development of mechanical ventilation strategies and pharmacologic agents that will decrease the incidence of ventilator-induced diaphragmatic dysfunction.

Keywords

diaphragmatic fatigue, diaphragmatic function, mechanical ventilation, respiratory muscles, weaning failure

Introduction

Mechanical ventilation is a two-edged sword. Although it is life-saving in patients with respiratory failure, mechanical ventilation is also associated with numerous potential complications. For example, although many patients in the ICU are placed on mechanical ventilation as supportive therapy for acute lung injury, there is substantial evidence that the institution of mechanical ventilation itself can paradoxically create and sustain damage to the lungs [1]. The recognition of this phenomenon, known as ventilator-induced lung injury (VILI), eventually led clinicians to adopt ventilator strategies designed to mitigate its occurrence, and this practice has now been linked to improved clinical outcomes in

patients with acute respiratory distress syndrome [2,3]. In the past several years, a similar paradigm has been emerging with regard to potential adverse effects of mechanical ventilation on the ventilatory muscles. This entity was originally termed ventilator-induced diaphragmatic dysfunction (VIDD) by Vassilakopoulos and Petrof [4], although it may well involve other respiratory muscles as well [5,6].

Diaphragmatic weakness appears to be very common in patients undergoing mechanical ventilation [7–9]. This is of major importance, as diaphragmatic function plays a crucial role in determining the ability of patients to be successfully weaned from the ventilator [10,11]. Furthermore, difficulties in weaning patients from mechanical

ventilation account for a large proportion of time spent in the ICU [12]. Taken together, these facts suggest that VIDD has the potential to have a major impact on clinical practice and the utilization of healthcare resources. In the present review, we will briefly summarize the existing evidence for VIDD in animal and human studies. We will then review the most recent information about the cellular mechanisms responsible for VIDD, and discuss the possible implications of these findings for future therapeutic strategies.

Brief review of evidence for ventilator-induced diaphragmatic dysfunction

In experimental animals, the ability of the intact diaphragm to generate pressure is reduced by 40–50% within a few days of instituting ‘controlled’ mechanical ventilation, which permits little or no spontaneous diaphragmatic activity [13–15]. Endurance of the diaphragm also appears to be adversely affected [6], with a reduced ability to sustain diaphragmatic force in the face of an inspiratory resistive load [13]. It should be pointed out that for the purposes of the present review, VIDD refers to changes in diaphragmatic function that arise from alterations outside of the central or peripheral nervous systems. Hence, in animal models of VIDD, nervous impulse transmission at the levels of the phrenic nerve and the neuromuscular junction remain normal [15], and the contractility of isolated (i.e., removed from their neural input) diaphragmatic strips is severely reduced along a similar time course and magnitude to that observed in the intact animal [14,16–19]. Taken together, the above findings indicate that the deleterious effects of mechanical ventilation upon diaphragmatic function are primarily the result of changes that occur within the muscle fibers *per se*. This is consistent with the fact that in the majority of patients with weaning difficulties, the level of neural input to the diaphragm is actually increased, but force generation remains decreased nonetheless [20**]. It is also clear that the loss of diaphragmatic force-generating capacity cannot be ascribed to atrophy alone, as many studies have shown that the force loss is persistent even after correcting for any reductions in muscle cross-sectional area [4,14,16–19].

Beyond decreased diaphragmatic strength, a number of histological and biochemical changes have been described in the diaphragms of animals with VIDD. These include the muscle fiber atrophy [5,13,16,17,19,21], which appears to be the result of decreased protein synthesis [22,23*] as well as increased protein breakdown [21,24–27]; muscle fiber remodeling, as indicated by changes in the expression of multiple structural [28] and muscle-specific proteins such as myosin heavy chain, MyoD, and myogenin [17,29]; and signs of muscle fiber injury, including disrupted myofibrils, increased numbers of vacuolar structures, and abnormal mitochondria [5,14,24,30].

In human individuals, there is more limited but nonetheless compelling evidence for the development of VIDD. In a postmortem analysis of neonates, diffuse diaphragmatic muscle fiber atrophy was found in patients who received ventilatory assistance for 12 days or more immediately before death, whereas such changes were not present in extradiaphragmatic muscles from the same patients or diaphragms of infants ventilated for 7 days or less [31]. More recently, in a landmark study, Levine *et al.* [32**] evaluated diaphragm biopsy specimens from adult brain-dead organ donors who had undergone mechanical ventilation for variable periods of time (18–69 h) prior to organ harvest, and compared them with specimens obtained from control patients who were undergoing thoracic surgery for benign lesions or localized lung cancer (mechanical ventilation for 2–3 h). In comparison with the controls, biopsy specimens from the organ donor group showed a decreased cross-sectional area of slow-twitch and fast-twitch fibers (atrophy), decreased glutathione levels (suggesting increased oxidative stress), and greater expression of active caspase-3 and the E3 ubiquitin ligases atrogenin-1, and muscle RING-finger protein-1 (MuRF-1; implicated in muscle proteolysis). Several of these results have recently been confirmed by another similarly designed study, which additionally reported rapidly progressive contractile dysfunction and diaphragmatic injury in mechanically ventilated humans [33]. Taken together, the above findings are remarkably similar to those that have been documented in animal models of VIDD.

AQ2

It should be noted that mechanical ventilation constitutes a rather unique form of muscle ‘disuse’, in the sense that the diaphragm is at the same time mechanically unloaded, electrically quiescent, and subjected to changes in myofiber length by cyclical lung inflation or positive end-expiratory pressure (PEEP). The diaphragm itself is also unique. It is normally exposed to a negative pressure environment along its pleural surface that can potentially serve as a stretch-like hypertrophic stimulus [34], which is removed by the application of positive-pressure ventilation. In addition, the diaphragm is more active than most other skeletal muscles (30–40% of the time, 24 h/day). All of these factors may help to explain the very rapid diaphragmatic atrophy and force loss observed during mechanical ventilation. It has been estimated that 12 h of mechanical ventilation in rats is approximately equivalent to 96 h of locomotor muscle unloading in terms of the muscle wasting responses that are induced [35].

Major cellular mechanisms of ventilator-induced diaphragmatic dysfunction

Our current understanding of the cellular mechanisms causing VIDD is derived primarily from animal models. Numerous studies have shown that mechanical

AQ1

ventilation is associated with an increase in markers of oxidative stress in the diaphragm [21,32^{••},36–38]. The onset of oxidative modifications is rapid, occurring within 6 h of instituting mechanical ventilation in rats [36]. Expression levels of antioxidant enzymes or free radical scavengers such as glutathione, superoxide dismutase, catalase, heme oxygenase-1, and others are variably decreased [28,37,38] or increased [28,39[•]], with the latter presumably being an attempt to limit oxidant-mediated injury. Interestingly, the cellular targets of diaphragmatic protein oxidation may involve elements of the contractile machinery such as myosin and actin [36]. Moreover, treatment with an antioxidant (the vitamin E analogue Trolox; Hoffman-La Roche, Nutley, New Jersey, USA) during mechanical ventilation mitigated diaphragmatic proteolysis and also prevented the loss of diaphragmatic force [40]. This same antioxidant also attenuated diaphragmatic fiber atrophy in a manner that was independent of alterations in insulin-like growth factor-1 (IGF1)-phosphoinositide-3 kinase (PI3K)-Akt signaling or ubiquitin ligase expression [41], despite the association of these pathways with diaphragmatic atrophy [19,25]. Collectively, the findings strongly suggest that mechanical ventilation-induced oxidative stress plays a major role in atrophy as well as the intrinsic contractile impairment of the diaphragm found in VIDD.

Recent studies have begun to search for the main sources of the reactive oxygen species (ROS) generated in the diaphragm during mechanical ventilation. There is no evidence for increased diaphragmatic inflammatory cell infiltration [32^{••},42] or a major upregulation of proinflammatory cytokine gene expression in the diaphragm [28]. Nitric oxide synthases also do not appear to be involved [43]. NADPH oxidase, which can generate superoxide within muscle fibers [44], was only found to be minimally increased in the mechanically ventilated diaphragm [45]. Xanthine oxidase, another potential source of superoxide, is also upregulated in the diaphragm during mechanical ventilation, and its inhibition modestly improved diaphragmatic contractility but failed to have an impact upon atrophy [46]. Given the above, mitochondria could be the main source of excessive ROS production in the diaphragm during mechanical ventilation. In this regard, mitochondria isolated from mechanically ventilated rats release significantly more ROS and also exhibit biochemical evidence of oxidative damage [39[•]]. Furthermore, abnormalities of diaphragmatic mitochondrial respiration have been found in several animal species during mechanical ventilation [5,39[•],47]. Interestingly, these findings are also consistent with morphological evidence of damaged mitochondria in the diaphragm after mechanical ventilation [5].

There is good evidence that the calpain, caspase, and ubiquitin–proteasome proteolysis pathways all play sig-

nificant roles in the development of mechanical ventilation-induced atrophy. Calpains are calcium-dependent proteases that have long been known to be capable of degrading cytoskeletal proteins in muscle [48], whereas recognition of the involvement of caspases in muscle atrophy is more recent [49]. Myofilament proteins, which constitute approximately two-thirds of bulk muscle protein, must first be partially cleaved and disassembled in order to be processed and degraded by the ubiquitin–proteasome system [50]. This initial step of proteolytic release of myofilaments from their native state can be accomplished by either calpains or caspases, both of which are upregulated in the diaphragm during mechanical ventilation [21,26,27]. Interestingly, the calpain and caspase systems demonstrate considerable crosstalk and can be mutually reinforcing [51]. The role of lysosomal proteases is less clear, but cathepsin activity is also increased [27]. Intriguingly, vacuolar structures in the mechanically ventilated diaphragm [5,30] also suggest the possibility of autophagy, another lysosomally mediated process recently shown to be involved in skeletal muscle atrophy associated with increased oxidative stress [52[•]].

Several studies have shown that myofiber atrophy and impaired contractility in the mechanically ventilated diaphragm can be uncoupled from one another [14,46,53]. For example, administration of oxypurinol (xanthine oxidase inhibitor) improved diaphragmatic contractility but not atrophy [46]. Conversely, allowing intermittent trials of spontaneous breathing between periods of mechanical ventilation was able to prevent mechanical ventilation-induced atrophy, but reductions in diaphragmatic contractility were persistent [53]. These observations suggest that the mechanisms responsible for atrophy and intrinsic contractile dysfunction are not identical, although they are likely to be linked. For instance, one possibility is that the disassembly of actomyosin complexes by calpains or caspases, which is a prerequisite for myofiber atrophy via the proteasome, also plays a critical role in causing intrinsic contractile impairment of the diaphragm. Moreover, calpains and caspases are activated by ROS and also have the ability to mediate activation of the proapoptotic protein Bid [54–57]. This could potentially lead to mitochondrial damage and drive further ROS generation, thereby sustaining a vicious cycle. The histological evidence of mitochondrial damage and particularly the presence of myofibrillar disarray, which has been reported in several studies and significantly correlated with abnormal contractile function of the diaphragm [5,14,24,30], are consistent with the above hypothesis.

Potential therapeutic strategies and future directions


Animal studies indicate that VIDD is alleviated (but not completely prevented) when using partial support modes

of mechanical ventilation, in which a significant degree of diaphragmatic effort is permitted [53,58,59]. From a purely practical standpoint, it appears logical to permit as much diaphragmatic activity as possible, as long as this allows for adequate patient comfort and gas exchange. However, further study is needed to ascertain the optimal level of diaphragmatic effort and to determine whether the specific method of promoting diaphragmatic effort during mechanical ventilation (e.g., spontaneous breathing trials, assist-control, pressure-support, newer modes such as NAVA, etc.) has any impact upon the risk of developing VID. In addition, it is important to recognize that even when using partial support modes of mechanical ventilation or allowing for intermittent periods of spontaneous breathing, studies have found evidence of persistent oxidative stress [59] as well as a substantial residual deficit of diaphragmatic force production, even in the absence of atrophy [53,58]. Taken together, the above findings suggest that other measures designed to target the specific cellular pathways involved in muscle injury, may be required in order to completely prevent or reverse VID.

With regard to pharmacological interventions, treatment with the vitamin E analogue Trolox during mechanical ventilation prevents the loss of diaphragmatic contractility and attenuates atrophy in rats over the short-term [40]. It is also interesting to note that in a study of critically ill surgical patients, an antioxidant supplement containing vitamins E and C was reported to reduce the duration of mechanical ventilation in comparison with nonsupplemented patients [60]. Although these findings are encouraging and support the use of antioxidants to prevent VID, these agents can also have deleterious as well as nonspecific effects. For example, it appears that the beneficial effects of apocynin on VID in rats may be more a function of its ability to upregulate a calpain inhibitor (calpastatin) rather than any specific antioxidant action [45].

The proteolytic systems activated during VID are also logical targets for therapeutic intervention. In this regard, a single administration of leupeptin (inhibitor of calpain/cathepsin) at the onset of mechanical ventilation, not only blocked atrophy, but also prevented intrinsic contractile

Figure 1 Cellular mechanisms implicated in ventilator-induced diaphragmatic dysfunction



Several of the mechanistic pathways implicated in the development of VID are illustrated. Protein synthesis pathways are downregulated (indicated by dashed lines). In addition, ROS generated from several possible sources can activate downstream proteolytic pathways involved in myofiber injury and atrophy, including the calpain and caspase systems; these have the potential to be mutually reinforcing and to drive further ROS production. The role of lysosomally-mediated proteolysis, and particularly autophagy, remains to be explored. IGF-1, insulin-like growth factor-1; MV, mechanical ventilation; PI3K, phosphoinositide-3 kinase; ROS, reactive oxygen species.

impairment in the rat diaphragm [27]. Somewhat surprisingly, acute high-dose corticosteroid administration was also found to prevent calpain upregulation and mitigate VIDD in rats [61[•]], although corticosteroids are unlikely to be a viable treatment for VIDD given their association with acute thick filament loss and other forms of myopathy [62]. Less surprising is the fact that neuromuscular blocking agents can synergize with mechanical ventilation to exacerbate VIDD, with attendant increases in activation of the calpain and ubiquitin–proteasome systems [63,64]. Interestingly, recent lessons from the sepsis literature suggest that inhibition of the proteasome pathway may not be an effective way to prevent the loss of contractile force associated with enhanced proteolysis in the diaphragm [65[•]]. These findings are consistent with the notion that events upstream of the proteasome (e.g., caspase-mediated or calpain-mediated myofilament dissociation [23[•]] and other forms of injury) are more likely to be responsible for the early reductions in diaphragmatic force-generating capacity in VIDD, but this hypothesis needs to be confirmed.

Finally, there are many questions of both clinical and scientific importance that need to be addressed by future studies. For example, what are the effects of mechanical ventilation on calcium homeostasis? Many of the processes implicated in VIDD, including injury, can be triggered by increases in intracellular calcium, but there is no direct information available on this point. Furthermore, although investigations to date have elegantly dissected the relationships of several cellular pathways to muscle fiber atrophy, the mechanisms leading to injury and contractile dysfunction remain obscure. There is also very little information about how VIDD is influenced by underlying conditions commonly found in mechanically ventilated patients, such as hyperglycemia [66] and sepsis [67], or by the preexisting state of the muscle (e.g., if fatigued or injured) prior to initiating mechanical ventilation. Most importantly, data are sorely lacking in human patients regarding most aspects of the VIDD phenomenon, including the time course for development of diaphragmatic weakness during mechanical ventilation and its relationship to injury, atrophy, and the candidate cellular pathways implicated in the animal models discussed in this review (see Fig. 1).

Conclusion

There is now evidence that mechanical ventilation itself may be an important cause of diaphragmatic weakness, associated with a combination of mechanical ventilation-induced diaphragmatic atrophy and injury, which is collectively referred to as VIDD. It is likely that several factors (e.g., underlying disease state, infection, drug therapy, etc.) can converge with VIDD to exacerbate diaphragmatic weakness in critically ill patients. Indeed,

due to the presence of multiple confounding factors, it is not currently possible to definitively diagnose any given patient with VIDD, just as no individual patient can be assigned a diagnosis of VILI with certainty. Nevertheless, as in the case of VILI, research into the basic mechanisms underlying the phenomenon of VIDD is allowing us to develop a conceptual framework for understanding the problem and applying this knowledge to clinical practice. At present, the best approach for preventing VIDD is to avoid controlled mechanical ventilation and the use of neuromuscular blocking agents to the greatest extent possible. In the future, the challenge will be to develop strategies or modes of mechanical ventilation which decrease the likelihood of VIDD, as well as to determine the ability of pharmacological interventions such as antioxidant therapy or inhibitors of muscle proteolysis pathways, to preserve diaphragmatic function in mechanically ventilated patients.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (pp. 000–000).

- 1 Dreyfuss D, Saumon G. Ventilator-induced lung injury: lessons from experimental studies. *Am J Respir Crit Care Med* 1998; 157:294–323.
- 2 The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and acute respiratory distress syndrome. *N Engl J Med* 2000; 342:1301–1308.
- 3 Amato MB, Barbas CS, Medeiros DM, *et al.* Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338:347–354.
- 4 Vassilakopoulos T, Petrof BJ. Ventilator-induced diaphragmatic dysfunction. *Am J Respir Crit Care Med* 2004; 169:336–341.
- 5 Bernard N, Matecki S, Py G, *et al.* Effects of prolonged mechanical ventilation on respiratory muscle ultrastructure and mitochondrial respiration in rabbits. *Intensive Care Med* 2003; 29:111–118.
- 6 Capdevila X, Lopez S, Bernard N, *et al.* Effects of controlled mechanical ventilation on respiratory muscle contractile properties in rabbits. *Intensive Care Med* 2003; 29:103–110.
- 7 Laghi F, Cattapan SE, Jubran A, *et al.* Is weaning failure caused by low-frequency fatigue of the diaphragm? *Am J Respir Crit Care Med* 2003; 167:120–127.
- 8 Watson AC, Hughes PD, Louise HM, *et al.* Measurement of twitch transdiaphragmatic, esophageal, and endotracheal tube pressure with bilateral anterolateral magnetic phrenic nerve stimulation in patients in the intensive care unit. *Crit Care Med* 2001; 29:1325–1331.
- 9 Chang AT, Boots RJ, Brown MG, *et al.* Reduced inspiratory muscle endurance following successful weaning from prolonged mechanical ventilation. *Chest* 2005; 128:553–559.
- 10 Vassilakopoulos T, Zakyntinos S, Roussos C. The tension-time index and the frequency/tidal volume ratio are the major pathophysiologic determinants of weaning failure and success. *Am J Respir Crit Care Med* 1998; 158:378–385.
- 11 Harikumar G, Egberongbe Y, Nadel S, *et al.* Tension time index as a predictor of extubation outcome in ventilated children. *Am J Respir Crit Care Med* 2009 [Epub ahead of print].
- 12 Esteban A, Alia I, Ibanez J, *et al.* Modes of mechanical ventilation and weaning. A national survey of Spanish hospitals. The Spanish Lung Failure Collaborative Group. *Chest* 1994; 106:1188–1193.
- 13 Anzueto A, Peters JI, Tobin MJ, *et al.* Effects of prolonged controlled mechanical ventilation on diaphragmatic function in healthy adult baboons. *Crit Care Med* 1997; 25:1187–1190.

- 14 Sassoon CS, Caiozzo VJ, Manka A, *et al.* Altered diaphragm contractile properties with controlled mechanical ventilation. *J Appl Physiol* 2002; 92:2585–2595.
- 15 Radell PJ, Remahl S, Nichols DG, *et al.* Effects of prolonged mechanical ventilation and inactivity on piglet diaphragm function. *Intensive Care Med* 2002; 28:358–364.
- 16 Le Bourdelles G, Viies N, Boczkowski J, *et al.* Effects of mechanical ventilation on diaphragmatic contractile properties in rats. *Am J Respir Crit Care Med* 1994; 149:1539–1544.
- 17 Yang L, Luo J, Bourdon J, *et al.* Controlled mechanical ventilation leads to remodeling of the rat diaphragm. *Am J Respir Crit Care Med* 2002; 166: 1135–1140.
- 18 Powers SK, Shanely RA, Coobes JS, *et al.* Mechanical ventilation results in progressive contractile dysfunction in the diaphragm. *J Appl Physiol* 2002; 92:1851–1858.
- 19 Gayan-Ramirez G, De Paeppe K, Cadot P, *et al.* Deterimental effects of short-term mechanical ventilation on diaphragm function and IGF-I mRNA in rats. *Intensive Care Med* 2003; 29:825–833.
- 20 Tobin MJ, Laghi F, Brochard L. Role of the respiratory muscles in acute respiratory failure of COPD: lessons from weaning failure. *J Appl Physiol* 2009; 107:962–970.
- Excellent review of the pathophysiology of weaning failure in mechanically ventilated patients, and particularly of the role played by the respiratory muscles.
- 21 Shanely RA, Zergeroglu MA, Lennon SL, *et al.* Mechanical ventilation-induced diaphragmatic atrophy is associated with oxidative injury and increased proteolytic activity. *Am J Respir Crit Care Med* 2002; 166:1369–1374.
- 22 Shanely RA, Van Gammeren D, DeRuisseau KC, *et al.* Mechanical ventilation depresses protein synthesis in the rat diaphragm. *Am J Respir Crit Care Med* 2004; 170:994–999.
- 23 McClung JM, Whidden MA, Kavazis AN, *et al.* Redox regulation of diaphragm proteolysis during mechanical ventilation. *Am J Physiol Regul Integr Comp Physiol* 2008; 294:R1608–R1617.
- Animal study demonstrating that mechanical ventilation causes the release of myofilaments from their protein lattice, and that the antioxidant Trolox provides protection against this process as well as activation of the proteasome in the diaphragm.
- 24 Zhu E, Sassoon CS, Nelson R, *et al.* Early effects of mechanical ventilation on isotonic contractile properties and MAF-box gene expression in the diaphragm. *J Appl Physiol* 2005; 99:747–756.
- 25 DeRuisseau KC, Kavazis AN, Deering MA, *et al.* Mechanical ventilation induces alterations of the ubiquitin–proteasome pathway in the diaphragm. *J Appl Physiol* 2005; 98:1314–1321.
- 26 McClung JM, Kavazis AN, DeRuisseau KC, *et al.* Caspase-3 regulation of diaphragm myonuclear domain during mechanical ventilation-induced atrophy. *Am J Respir Crit Care Med* 2007; 175:150–159.
- 27 Maes K, Testelmans D, Powers S, *et al.* Leupeptin inhibits ventilator-induced diaphragm dysfunction in rats. *Am J Respir Crit Care Med* 2007; 175:1134–1138.
- 28 DeRuisseau KC, Shanely RA, Akunuri N, *et al.* Diaphragm unloading via controlled mechanical ventilation alters the gene expression profile. *Am J Respir Crit Care Med* 2005; 172:1267–1275.
- 29 Racz GZ, Gayan-Ramirez G, Testelmans D, *et al.* Early changes in rat diaphragm biology with mechanical ventilation. *Am J Respir Crit Care Med* 2003; 168:297–304.
- 30 Radell P, Edstrom L, Stibler H, *et al.* Changes in diaphragm structure following prolonged mechanical ventilation in piglets. *Acta Anaesthesiol Scand* 2004; 48:430–437.
- 31 Knisely AS, Leal SM, Singer DB. Abnormalities of diaphragmatic muscle in neonates with ventilated lungs. *J Pediatr* 1988; 113:1074–1077.
- 32 Levine S, Nguyen T, Taylor N, *et al.* Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. *N Engl J Med* 2008; 358:1327–1335.
- Landmark study in humans demonstrating preferential atrophy in the diaphragms of brain-dead patients after prolonged mechanical ventilation, together with evidence of increased oxidative stress and activation of proteolytic pathways in the diaphragm.
- 33 Jaber S, Chanques G, Jung B, *et al.* Mechanical ventilation decreases diaphragm force, induces muscular injury and stimulates proteolytic pathway: in vivo and in vitro human study (abstract). *Anesthesiology* 2009.
- 34 Yang L, Bourdon J, Gottfried SB, *et al.* Regulation of myosin heavy chain gene expression after short-term diaphragm inactivation. *Am J Physiol* 1998; 274:L980–L989.
- 35 Powers SK, Kavazis AN, McClung JM. Oxidative stress and disuse muscle atrophy. *J Appl Physiol* 2007; 102:2389–2397.
- 36 Zergeroglu MA, McKenzie MJ, Shanely RA, *et al.* Mechanical ventilation-induced oxidative stress in the diaphragm. *J Appl Physiol* 2003; 95:1116–1124.
- 37 Jaber S, Sebbane M, Koechlin C, *et al.* Effects of short vs. prolonged mechanical ventilation on antioxidant systems in piglet diaphragm. *Intensive Care Med* 2005; 31:1427–1433.
- 38 Falk DJ, DeRuisseau KC, Van Gammeren DL, *et al.* Mechanical ventilation promotes redox status alterations in the diaphragm. *J Appl Physiol* 2006; 101:1017–1024.
- 39 Kavazis AN, Talbert EE, Smuder AJ, *et al.* Mechanical ventilation induces diaphragmatic mitochondrial dysfunction and increased oxidant production. *Free Radic Biol Med* 2009; 46:842–850.
- Study showing that the mitochondria of mechanically ventilated rat diaphragms release significantly more ROS and exhibit evidence of increased oxidative injury.
- 40 Betters JL, Criswell DS, Shanely RA, *et al.* Trolox attenuates mechanical ventilation-induced diaphragmatic dysfunction and proteolysis. *Am J Respir Crit Care Med* 2004; 170:1179–1184.
- 41 McClung JM, Kavazis AN, Whidden MA, *et al.* Antioxidant administration attenuates mechanical ventilation-induced rat diaphragm muscle atrophy independent of protein kinase B (PKB Akt) signalling. *J Physiol* 2007; 585:203–215.
- 42 Van Gammeren D, Falk DJ, DeRuisseau KC, *et al.* Reloading the diaphragm following mechanical ventilation does not promote injury. *Chest* 2005; 127: 2204–2210.
- 43 Van Gammeren D, Falk DJ, Deering MA, *et al.* Diaphragmatic nitric oxide synthase is not induced during mechanical ventilation. *J Appl Physiol* 2007; 102:157–162.
- 44 Javesghani D, Magder SA, Barreiro E, *et al.* Molecular characterization of a superoxide-generating NAD(P)H oxidase in the ventilatory muscles. *Am J Respir Crit Care Med* 2002; 165:412–418.
- 45 McClung JM, Van Gammeren D, Whidden MA, *et al.* Apocynin attenuates diaphragm oxidative stress and protease activation during prolonged mechanical ventilation. *Crit Care Med* 2009; 37:1373–1379.
- 46 Whidden MA, McClung JM, Falk DJ, *et al.* Xanthine oxidase contributes to mechanical ventilation-induced diaphragmatic oxidative stress and contractile dysfunction. *J Appl Physiol* 2009; 106:385–394.
- 47 Fredriksson K, Radell P, Eriksson LI, *et al.* Effect of prolonged mechanical ventilation on diaphragm muscle mitochondria in piglets. *Acta Anaesthesiol Scand* 2005; 49:1101–1107.
- 48 Belcastro AN, Shewchuk LD, Raj DA. Exercise-induced muscle injury: a calpain hypothesis. *Mol Cell Biochem* 2003; 179:135–145.
- 49 Du J, Wang X, Miereles C, *et al.* Activation of caspase-3 is an initial step triggering accelerated muscle proteolysis in catabolic conditions. *J Clin Invest* 2004; 113:115–123.
- 50 Jackman RW, Kandarian SC. The molecular basis of skeletal muscle atrophy. *Am J Physiol Cell Physiol* 2004; 287:C834–C843.
- 51 Wang KK. Calpain and caspase: can you tell the difference? *Trends Neurosci* 2000; 23:20–26.
- 52 Dobrowolny G, Aucello M, Rizzuto E, *et al.* Skeletal muscle is a primary target of SOD1G93A-mediated toxicity. *Cell Metab* 2008; 8:425–436.
- Study performed in a murine model of defective superoxide dismutase function in skeletal muscle, which suggests an important role for lysosomally-mediated autophagy in muscle atrophy induced by oxidative stress.
- 53 Gayan-Ramirez G, Testelmans D, Maes K, *et al.* Intermittent spontaneous breathing protects the rat diaphragm from mechanical ventilation effects. *Crit Care Med* 2005; 33:2804–2809.
- 54 Bajaj G, Sharma RK. TNF-alpha-mediated cardiomyocyte apoptosis involves caspase-12 and calpain. *Biochem Biophys Res Commun* 2006; 345:1558–1564.
- 55 Chen M, Won DJ, Krajewski S, *et al.* Calpain and mitochondria in ischemia/reperfusion injury. *J Biol Chem* 2002; 277:29181–29186.
- 56 Haudek SB, Taffet GE, Schneider MD, *et al.* TNF provokes cardiomyocyte apoptosis and cardiac remodeling through activation of multiple cell death pathways. *J Clin Invest* 2007; 117:2692–2701.
- 57 Scarabelli TM, Stephanou A, Pasini E, *et al.* Different signaling pathways induce apoptosis in endothelial cells and cardiac myocytes during ischemia/reperfusion injury. *Circ Res* 2002; 90:745–748.
- 58 Sassoon CS, Zhu E, Caiozzo VJ. Assist-control mechanical ventilation attenuates ventilator-induced diaphragmatic dysfunction. *Am J Respir Crit Care Med* 2004; 170:626–632.

AQ5

AQ6

- 59** Futier E, Constantin JM, Combaret L, *et al.* Pressure support ventilation
• attenuates ventilator-induced protein modifications in the diaphragm. *Crit Care* 2008; 12:R116.

Animal study showing that pressure-support ventilation can prevent the reduced protein synthesis and accelerated proteolysis associated with controlled mechanical ventilation, but does not appear to completely abrogate increased oxidative stress in the diaphragm under these conditions.

- 60** Nathens A, Neff M, Jurkovich G, *et al.* Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. *Ann Surg* 2002; 236:814–822.

- 61** Maes K, Testelmans D, Cadot P, *et al.* Effects of acute administration of
• corticosteroids during mechanical ventilation on rat diaphragm. *Am J Respir Crit Care Med* 2008; 178:1219–1226.

Study reporting that acute high-dose corticosteroid therapy can prevent manifestations of VIDD in an animal model, possibly through the inhibition of calpain activity.

- 62** Dekhuijzen PN, Decramer M. Steroid-induced myopathy and its significance to respiratory disease: a known disease rediscovered. *Eur Respir J* 1992; 5:997–1003.

- 63** Testelmans D, Maes K, Wouters P, *et al.* Rocuronium exacerbates mechanical ventilation-induced diaphragm dysfunction in rats. *Crit Care Med* 2006; 34:3018–3023.

- 64** Testelmans D, Maes K, Wouters P, *et al.* Infusions of rocuronium and cisatracurium exert different effects on rat diaphragm function. *Intensive Care Med* 2007; 33:872–879.

- 65** Supinski GS, Vanags J, Callahan LA. Effect of proteasome inhibitors on
• endotoxin-induced diaphragm dysfunction. *Am J Physiol Lung Cell Mol Physiol* 2009; 296:L994–L1001.

Conceptually related study demonstrating that inhibition of the proteasome does not improve diaphragmatic function in a model of sepsis, in which ROS and the calpain/caspase proteolytic systems are implicated in the loss of force-generating capacity.

- 66** Callahan LA, Supinski GS. Hyperglycemia and acquired weakness in critically ill patients: potential mechanisms. *Crit Care* 2009; 13:125.

- 67** Ebihara S, Hussain SN, Danelou G, *et al.* Mechanical ventilation protects against diaphragm injury in sepsis: interaction of oxidative and mechanical stresses. *Am J Respir Crit Care Med* 2002; 165:221–228.