



**HAL**  
open science

## Adrenal gland volume measurement in septic shock and control patients: a pilot study

Stéphanie Nougaret, Boris Jung, S. Aufort, Gerald Chanques, Samir Jaber, B. Gallix

► **To cite this version:**

Stéphanie Nougaret, Boris Jung, S. Aufort, Gerald Chanques, Samir Jaber, et al.. Adrenal gland volume measurement in septic shock and control patients: a pilot study. *European Radiology*, 2010, 20, pp.2348 - 2357. 10.1007/s00330-010-1804-9 . hal-02550717

**HAL Id: hal-02550717**

**<https://hal.umontpellier.fr/hal-02550717>**

Submitted on 22 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stephanie Nougaret  
B. Jung  
S. Aupart  
G. Chanques  
S. Jaber  
B. Gallix

# Adrenal gland volume measurement in septic shock and control patients: a pilot study

S. Nougaret (✉) · S. Aupart · B. Gallix  
Department of Abdominal Imaging,  
CHU Montpellier,  
Hôpital Saint Eloi,  
80 avenue Augustin Fliche, 34295,  
Montpellier, Cedex 5, France  
e-mail: stephanienougaret@free.fr  
Tel.: +33-4-67337120  
Fax: +33-4-67337116

B. Jung · G. Chanques · S. Jaber  
Intensive Care Unit, Department of Critical  
Care and Anesthesiology: DAR B,  
CHU Montpellier,  
Hôpital Saint Eloi,  
80 avenue Augustin Fliche, 34295,  
Montpellier, Cedex 5, France

## Abstract

**Objectives:** To compare adrenal gland volume in septic shock patients and control patients by using semi-automated volumetry. **Methods:** Adrenal gland volume and its inter-observer variability were measured with tomodensitometry using semi-automated software in 104 septic shock patients and in 40 control patients. The volumes of control and septic shock patients were compared and the relationship between volume and outcome in intensive care was studied. **Results:** The mean total volume of both adrenal glands was  $7.2 \pm 2.0 \text{ cm}^3$  in control subjects and  $13.3 \pm 4.7 \text{ cm}^3$  for total adrenal gland volume in septic shock patients ( $p < 0.0001$ ). Measurement reproducibility was excellent with a concordance correla-

tion coefficient value of 0.87. The increasing adrenal gland volume was associated with a higher rate of survival in intensive care. **Conclusion:** The present study reports that with semi-automated software, adrenal gland volume can be measured easily and reproducibly. Adrenal gland volume was found to be nearly double in sepsis compared with control patients. The absence of increased volume during sepsis would appear to be associated with a higher rate of mortality and may represent a prognosis factor which may help the clinician to guide their strategy.

**Keywords** CT · Adrenal gland · Volumetry · Septic shock · Intensive care

## Introduction

Adrenal glands are often described on computed tomography (CT) as small, triangular glands located at the top of both kidneys. Their weights are considered to be about 4 g in non-stressed subjects (surgical specimen) and their volumes have never been properly evaluated by CT [1]. To date, only two studies have reported adrenal gland volumetry and this was performed manually by CT [2, 3], based on single thick slice acquisition of 3 mm. Automated and semi-automated volumetry are used successfully with other organs such as liver [4–9] or lungs [10–12] but no studies have been performed on adrenal gland. The volume of the adrenal glands can be modified by several pathological conditions (e.g. depression [2], Cushing's disease [13] and septic shock [14]).

Septic shock is defined as the occurrence of a proven infection with cardiovascular failure needing vasopressor therapy [15, 16]. It is the most common life-threatening disease occurring in intensive care units (ICUs) leading to a mortality rate of 30–50% [17]. Relative adrenal insufficiency has been described to be very common during septic shock, and appears to be an independent prognostic factor, but its diagnosis and adequate treatment remain controversial in 2010 [16, 18]. The place of adrenal gland volume measurement in the early phase of management of septic shock has never been evaluated. Therefore, the purposes of our preliminary study were:

- To measure adrenal gland volumes by CT using a manufacturer's software (Myrian, Intrasure, Montpellier, France) in non-stressed control patients

- To evaluate the inter-observer reproducibility of this measurement
- To compare adrenal gland volume in control and septic shock patients and
- To evaluate its impact on mortality

---

## Materials and methods

### Study patients

The study design was approved by the local ethics committee of anaesthesia and intensive care. Two groups of patients were included between November 2007 and December 2008. Included in the control group were 40 consecutive patients with no current medical illness who had come in for virtual colonoscopy after medical investigation for abdominal pain or constipation. Included in the septic shock group were 104 consecutive patients in septic shock who had undergone an abdominal CT within the first 48 h of septic shock as defined by Bone et al. [15]. Briefly, patients in the septic shock group had a proven infection with hypotension, despite adequate fluid resuscitation and required vasopressors [15]. All CTs were collected in the picture and archiving communication system (PACS, GE Healthcare, UK) which is a high-security database with protection measures for patient confidentiality approved by the institutional review board for maintaining and renewing patient information. Women taking oral contraceptives, patients who had used any steroid preparation in the recent past and all patients with a history of endocrinopathy or chronic steroid use were not included to decrease the selection bias. Furthermore, we did not include patients with one unique adrenal gland.

### CT of the adrenal glands

Examination of the adrenal glands was performed with 64-section spiral CT (Aquilon 64, Toshiba Medical Systems, Nasu, Japan). There was no standard protocol for the CT. In fact, the protocol was adapted according to renal function. CT examinations in the control group ( $n=40$ ) were performed without injection of contrast medium, as for virtual colonoscopy. In the septic shock group, 23 CTs were performed without intravenous contrast medium due to renal failure; 81 CTs were performed with intravenous contrast medium using a standardized protocol for critically ill patients (quadriphasic acquisition). Images were viewed on the PACS workstation for standard diagnostic interpretation. Subsequently, images were transferred to a workstation (Myrian, Intrasure, Montpellier, France) devoted entirely to adrenal volume measurements. A section thickness of 3 mm (without overlaps) was used for the volumetry. A magnification targeting factor of  $\times 2$  was used. The adrenal contour was semi-automatically traced by using the specially designed workstation

(Myrian) by one radiologist (with 5 years' experience). Care was taken to try to exclude adjacent fat. Because the shape of the adrenals can be somewhat irregular, threshold values were established to exclude adjacent fat from the volume calculated. For example, only tissue with attenuation values of 25 to 200 Hounsfield units within the region of interest was included. The CT software (Myrian, Intrasure, Montpellier, France) automatically calculated the adrenal volume by totalling the area on each slice. Most subjects had clearly defined adrenal margins. The radiologist was unaware of the clinical status of the subjects. The measurement lasted 5 min per case. Patient and control adrenals were assessed in random order. To assess the facility and the reproducibility of the technique, each adrenal gland volume measurement for normal, non-stressed patients was performed by the radiologist and an intensivist. The measurement of the adrenal gland volume in 10 control subjects was considered a learning curve for the inter-observer agreement evaluation. Inter-observer variability was determined with the semi-automated measurement by comparing right and left adrenal gland volume values measured by each reader. To avoid bias, each observer was blinded to the results of the other observer. The same two observers each retrospectively performed two semi-automatic segmentations for each patient after a time interval of at least 2 weeks. The radiologist was blinded to the outcome of the septic patients.

### Measurements

For both groups, sex, age, weight, height, body surface area and underlying disease were recorded. In the septic shock group, the admission category (medicine or surgery), main diagnosis at admission and outcome at ICU discharge were recorded. The main endpoints were: first, the evaluation of adrenal gland volume in both groups; second, the evaluation of the inter-observer correlation; and, third, the comparison of adrenal gland volume in the control group and the septic shock group, and its association with mortality.

### Statistics

Data are represented as means  $\pm$  standard deviation (SD) or median and quartiles (25th–75th) as required. Data were analysed by using Student's  $t$  test, the Mann–Whitney test and the chi-squared or Fisher's exact test after a population normality assessment using the Kolmogorov–Smirnov test. Inter-observer measurements of adrenal gland volume were correlated with the inter-observer correlation coefficient for continuous variables. Relationships between adrenal volume, and subjects' weight, height, body mass index, age and mortality were examined by using the Pearson product-moment correlation coefficient for continuous variables. The within-subject variance was calculated by using a one-way

variance analysis. The sensitivity and specificity of adrenal volume curves to predict mortality were assessed by a receiver operating characteristic (ROC) curve.  $P < 0.05$  was considered statistically significant. The statistical analysis was performed by using Medcalc Software 9.4.2.0 (Mariakerke, Belgium).

## Results

### Adrenal gland measurements in control subjects

The adrenal gland volume was measured in 40 control patients who were examined with virtual colonoscopy (control subjects). Demographic characteristics are reported in Table 1. The mean adrenal volume was  $3.4 \pm 1.0 \text{ cm}^3$  for the right adrenal gland and  $3.8 \pm 1.2 \text{ cm}^3$  for the left adrenal gland, with no significant difference. The mean total volume of both adrenal glands was  $7.2 \pm 2.0 \text{ cm}^3$ .

The total adrenal gland volume was larger in men ( $8.0 \pm 1.8 \text{ cm}^3$ ) than in women ( $6.2 \pm 1.7 \text{ cm}^3$ ) ( $p = 0.003$ ). There was no consistent correlation between adrenal volume and age ( $r^2 = 0.08$ ) and height ( $r^2 = 0.53$ ) but there were consistent positive correlations between adrenal volume and weight ( $r^2 = 0.73$ ) (Fig. 1a, b) and body surface area ( $r^2 = 0.73$ ) (Fig. 1c). After the first 10 measurements which were considered a trial period for the intensivist, the inter-observer reliability was evaluated (Table 2). The concordance correlation coefficient between the radiologist and the intensivist was 0.87 (0.76–0.93) for the total gland volume.

### Adrenal gland volume in the control group versus the septic shock group

During the study period, we included 104 consecutive septic shock patients (septic shock group) who had undergone CT within the first 48 h of the septic shock.

The characteristics of patients in septic shock are summarised in Table 1. There was no significant difference in weight ( $p = 0.11$ ), height ( $p = 0.89$ ) and body surface area ( $p = 0.16$ ) between the two groups, but subjects in the control group were significantly younger ( $p = 0.0003$ ) than patients in the septic shock group.

Whereas adrenal gland volume in the control group was correlated with weight and body surface area (Fig. 1a,c), adrenal gland volume in the septic shock group was not correlated with weight ( $r^2 = 0.04$ ) or body surface area ( $r^2 = 0.09$ ) (Fig. 2a, b). The mean total volume of both adrenal glands was  $7.2 \pm 2.0 \text{ cm}^3$  in normal subjects and  $13.3 \pm 4.7 \text{ cm}^3$  for total adrenal gland volume in septic shock patients (Fig. 3) ( $p < 0.0001$ ).

Among the 104 patients included in the septic shock group, 46 had undergone CT before the septic shock with a delay of 102 days (range 26–646). In this group of patients, 32 suffered from chronic illness (12 chronic pancreatitis, 8 cirrhosis, 10 cancers, and 2 liver transplantations). In this subgroup of 46 patients, the mean total volume of both adrenal glands was  $9.5 \pm 3.6 \text{ cm}^3$  before the onset of septic shock and  $13.4 \pm 4.8 \text{ cm}^3$  during septic shock ( $p < 0.0001$ ) (Fig. 4). In this subgroup of patients, the adrenal gland volume before septic shock ( $9.5 \pm 3.6 \text{ cm}^3$ ) was significantly larger than in the control group ( $7.2 \pm 2.0 \text{ cm}^3$ ) ( $p = 0.005$ ). Adrenal gland volume had increased between 10% and 50% in 9 patients (20%), between 50% and 100% in 7 patients (15%) and more than 100% in 10 patients (22%). There were no variations before or during septic shock in 20 patients (43%) and the adrenal gland volume had not decreased in any of the patients.

### Adrenal gland enlargement and outcome

A small adrenal volume during septic shock was associated with higher mortality. Figure 5 shows the receiver operating characteristic (ROC) curve correlating adrenal volume and ICU mortality. An optimal cutoff value of


**Table 1** Patient characteristics

	Control group ( <i>n</i> =40)	Septic shock group ( <i>n</i> =104)	<i>p</i>
Age (years, mean $\pm$ SD)	56 $\pm$ 11	65 $\pm$ 14	0.0003
Sex, <i>n</i> (%)			
Female	18 (45%)	30 (29%)	0.08
Male	22 (55%)	74 (71%)	
Body weight (kg, mean $\pm$ SD)	70 $\pm$ 12	77 $\pm$ 25	0.11
Body height (cm, mean $\pm$ SD)	169 $\pm$ 7	169 $\pm$ 8	0.89
Surface area (m <sup>2</sup> , mean $\pm$ SD)	1.8 $\pm$ 0.4	1.9 $\pm$ 0.3	0.16
Number of deaths, <i>n</i> (%)	0	33 (32%)	
Type of patient, <i>n</i> (%)			
Medicine	40 (100%)	47 (45%)	NA
Surgery	0	57 (55%)	
Type of disease, <i>n</i> (%)			
Abdominal	0	38 (36%)	NA
Pulmonary	0	29 (28%)	NA
Both	0	37 (36%)	NA

Data are represented as means  $\pm$  SD or *n* (%)

NA not available


**Fig. 1** **a** Scatterplot showing the results of regression analysis of adrenal gland volume and control patients' weight. A linear correlation between total adrenal gland volume and weight ( $r^2=0.73$ ) was found in the control group. **b** Scatterplot showing the results of regression analysis of adrenal gland volume and control patients' height. There was no linear correlation between total adrenal gland volume and height ( $r^2=0.53$ ) in the control group. **c** Scatterplot showing the results of regression analysis of adrenal gland volume and control patients' body surface area. A linear correlation between total adrenal gland volume and body surface area ( $r^2=0.73$ ) was found in the control group




**Table 2** Inter-observer correlation coefficient (ICC) for right, left and both adrenal glands

	Radiologist	Intensivist	ICC
Right adrenal gland volume, cm <sup>3</sup> (mean ± SD)	3.3±1.0	3.4±1.0	0.87 [0.76–0.93]
Left adrenal gland volume, cm <sup>3</sup> (mean ± SD)	3.7±1.1	3.9±1.3	0.83 [0.72–0.93]
Total adrenal gland volume, cm <sup>3</sup> (mean ± SD)	7.1±1.9	7.3±2.0	0.87 [0.76–0.93]

**Fig. 2** **a** Scatterplot showing the results of regression analysis of adrenal gland volume and septic shock patients' weight. There was no linear correlation between total adrenal gland volume and weight ( $r^2=0.11$ ) in the septic shock group. **b** Scatterplot showing the result of regression analysis of adrenal gland volume and septic shock patients' body surface area. There was no linear correlation between total adrenal gland volume and body surface area ( $r^2=0.16$ ) in the septic shock group



**Fig. 3** Box and whisker plot showing the results of total adrenal gland volume in the control group and in the septic shock group. Each box ranges from the 25th percentile at the lower edge to the 75th percentile at the upper edge. The median is shown as a line across the box. There are two adjacent values below and above the box: the largest value is below the upper inner limit and the smallest value is above the lower inner limit. There was an increase in total adrenal gland volume in the 104 patients who underwent CT during septic shock in comparison to the adrenal volume of the 40 control subjects (\*\* $p < 0.0001$ )




10 cm<sup>3</sup> provided a sensitivity of 69% and a specificity of 94% for the risk of mortality in the ICU. The likelihood ratio for dying with an adrenal gland volume of less than 10 cm<sup>3</sup> was evaluated to be 3.1.

In Figs. 6 and 7, we present two case reports of septic shock patients. A 62-year-old woman was examined by CT for back pain that revealed no disease except for osteoporosis. Her total adrenal gland volume was 8.5 cm<sup>3</sup> (Fig. 6a). Two months later she developed community-


acquired pneumonia with septic shock and CT performed 6 h after the onset of shock showed that her total adrenal gland volume was 9.0 cm<sup>3</sup> (Fig. 6b). She died 6 days after admission to the ICU. A 68-year-old woman was referred for CT for a colon cancer evaluation. Her total adrenal gland volume was 10.2 cm<sup>3</sup> (Fig. 7a). She was admitted to the ICU 15 days later for nosocomial peritonitis with septic shock after colectomy. The total adrenal gland volume had increased and measured 19.6 cm<sup>3</sup> (Fig. 7b).

**Fig. 4** Box and whisker plot showing the results of total adrenal gland volume for the 46 septic shock patients who underwent CT before and during septic shock and for the control group (n=40). The description of each box is the same as for Fig. 3. Total adrenal gland volume was greater during sepsis compared with CT performed before the septic shock (\*\* $p < 0.0001$ ) for the same group of patients. The adrenal gland volume was larger in those patients who were explored by CT before the septic shock than in control patients (\*\* $p = 0.005$ ; \*\*before septic shock vs control group,  $p = 0.005$ )





**Fig. 5** Receiver operating characteristic curve, between total adrenal gland volumetry and mortality, showing a sensitivity of 69% and a specificity of 94% with an AUC=0.86 for a cutoff at 10 cm<sup>3</sup>



The patient recovered from sepsis and was discharged from the ICU.


## Discussion

The main results of our study are, first, that the adrenal gland volume in control patients is  $7.2 \pm 2.0$  cm<sup>3</sup> and is positively correlated with weight and body surface area; second, the measurement is reproducible between special-

ists and intensivists with a concordance correlation coefficient of 0.87; and finally, for the first time, we report that the adrenal gland volume in septic shock patients is nearly twice that of control patients and the total adrenal gland volume enlargement seems to be positively associated with a better outcome and quicker discharge from the ICU.


There are numerous clinical applications for solid abdominal organ volumetry [19–26] but organ volume assessments are not routinely performed, partly because of the extensive user input and time required with the current

**Fig. 6 a** Three-dimensional volume-rendered abdominal CT of a 62-year-old woman who underwent a CT 2 months before the onset of septic shock, showing a normal adrenal gland volume (8.5 cm<sup>3</sup>). During the septic shock (**b**) total adrenal gland volume did not increase (9.0 cm<sup>3</sup>). Death occurred in the ICU





**Fig. 7** a Three-dimensional volume-rendered abdominal CT of a 68-year-old woman who underwent CT 2 weeks before she presented septic shock and showing a normal adrenal gland volume ( $10.2 \text{ cm}^3$ ). During septic shock (b) the total adrenal gland volume increased ( $19.6 \text{ cm}^3$ ). She was discharged from the ICU



manual techniques. Use of semi-automated and automated organ segmentation would resolve the issues related to the user input and time required for volumetric analysis and enable imaging-based organ segmentation to become a routine process. This potential has prompted diverse and rapidly evolving innovations in the field of imaging-based segmentation. Indeed, CT evaluations of liver volume do have an impact on mortality before major liver resection [6, 26, 27]. Furthermore, several studies have reported that lung volume assessments with CT may reduce the deleterious consequences of mechanical ventilation and could improve oxygenation in those critically ill patients [11, 28]. Several volumetry classification techniques have been used. These schemes include classifications of automation (manual, semi-automatic and automatic), technique (MR imaging, CT), the number of datasets (single, multiple and multispectral), and also the type of volume algorithm calculation. These methods of measuring liver volume, described in the literature, are reproducible and accurate [4–8, 28–31].

Since 1978, several groups described the radiographic appearance of normal [32–34] and enlarged [35] adrenal glands by CT. Two studies reported a manual estimation of adrenal volume by CT [2, 3]. These two studies were both carried out with sequential acquisition and volumetry was calculated manually.

In our study, semi-automated software (Intrasense, Montpellier, France) appeared to be adequate for measuring adrenal gland volume. Furthermore, after a short training period, any physician who was not a radiologist could measure the adrenal gland volume with excellent inter-observer reliability.

The mean total volume measured by CT in normal subjects was  $7.2 \pm 2.0 \text{ cm}^3$  which was larger ( $5.3 \pm 0.5 \text{ cm}^3$ ) than the volume reported by Rubin and Phillips [3] or that reported by Amsterdam et al. ( $3.5 \pm 0.5 \text{ cm}^3$ ) [2]. The difference in adrenal size among previous studies is mainly explained by the technological advances in CT imaging. The thin slice imaging we use nowadays provides greater accuracy. Also, to a lesser degree, differences in CT calibration or in tracing the borders of

the adrenals may have caused these discrepancies. Our results in normal subjects found a correlation between total adrenal gland volume and weight and body surface area (Fig. 1) similar to that discovered by Rubin and Phillips [3].

In patients with proven severe sepsis, early CT of the abdomen showed an enlargement of the adrenal glands compared with control patients. This result has only been reported in case reports [14]. Moreover, for the first time, we found that in septic shock and contrary to control patients, there was no correlation between patient weight or body surface area and adrenal gland volume (Fig. 2).

Few studies have reported morphological changes in adrenal volume apart from in endocrinopathy (e.g. Cushing's syndrome); only two studies have shown increased adrenal gland volume in depressed patients [2, 36].

The abnormalities of adrenal gland volume occurring in sepsis may be explained by the pathophysiology of the adrenal glands during sepsis. Indeed, several lines of evidence support the fact that adrenal gland response is altered during sepsis, mainly due to hypothalamic-pituitary adrenocortical dysfunction, decreased production of steroids by the adrenal gland and also due to peripheral resistance to steroids [37].

In the 46 patients who underwent CT before and during septic shock, the total adrenal gland volume had increased significantly between the two CTs. This morphological change may be related to sepsis in the same patient (Fig. 4). Indeed, like others, we found no correlation between adrenal gland volume and age [3]. We therefore speculate that sepsis may influence the adrenal gland size all by itself. Moreover, regarding the difference between control group and the septic group before the onset of sepsis, adrenal glands were significantly larger in 44% of the case group before onset of sepsis in comparison to control group. We speculated that this enlargement was due to chronic disease. In fact, 32 suffered from chronic illness (12 chronic pancreatitis, 8 cirrhosis, 10 cancers, 2 liver transplantation).

Finally, we reported that adrenal gland volume seems to be associated with the mortality rate of patients in the ICU. Indeed, the absence of adrenal gland enlargement (less than 10 cm<sup>3</sup>) was associated with a higher mortality in the ICU. Because of its positive impact on outcome, it seems reasonable to hypothesise that the increased adrenal gland volume found during sepsis may be interpreted as an adaptation rather than oedema or necrosis of the gland.

Nevertheless we should bear in mind the limitations of this study because of the retrospective design; this meant that we did not perform a sample size calculation, which represents a limitation of our study. However, the homogeneity of the control group volume distribution and the statistical significance of the volume between control and septic patients are probably not related to a type I error. Moreover, we cannot exclude the possibility of a discrepancy between CT-estimated adrenal volume and the actual gland volume. Indeed, certain errors may be attributable to a partial volume effect at the adrenal edge, or adhesion to the adjacent tissues with attenuation similar to that of the adrenal parenchyma. Although a discrepancy cannot be excluded, the inter-observer correlation was 0.87 and adrenal gland volume in control patients was similar to that in the literature [33, 36]. Furthermore, with this technology, the adrenal cortex and medulla cannot be visually separated, and it is possible that some of the variations in adrenal volume estimates may be accounted for by the adrenal medulla. Moreover, the study of adrenal CT density was not practicable because of the lack of a standardised protocol for contrast medium injection. Finally, we did not perform a millimetric analysis of the CT slices; but because this was performed in both the control and the septic shock groups, we speculated that the potential measurement error was observed in both groups. Ideally an analysis of the adrenal density without injection and/or after standard weight-related and timed enhancement should be performed.

Initial evaluation of the critically ill patients admitted to the ICU with a septic shock often includes CT for diagnosis and complication evaluation (such as acute bowel ischaemia). The technique of adrenal gland volume measurement is easy to use, reproducible, may be associated with outcome and has never been evaluated before in this indication. Furthermore, in opposition to ICU specialised scores which often need clinical and complex biological parameters (e.g. bilirubin, arterial blood gases, platelets count, plasma creatinine etc.) [38] for several days to describe a kinetic evolution, we reported that a single adrenal gland volume measurement seems to be strongly associated with outcome. Because of our results and the necessity for the intensivist to have a prognostic factor, we believe that radiologists should be aware of these findings.

---

## Conclusion

For the first time, this study demonstrates that adrenal gland volume can be measured easily with semi-automated software and that its measurement is reproducible. Also, for the first time, we observed that the total adrenal gland volume was nearly twice as high in septic shock patients as in control patients. Absence of an increased adrenal gland volume was associated with a higher mortality rate in the ICU and deserves further investigation by clinical studies.

**Acknowledgements** Dr Bonnel, M.D, Intrasense Society, Montpellier, France for logistical assistance.

Presented in part at the 2008 Annual Meeting of the French Society of Radiologists (Paris, France, October 24–28, 2008)

---

## References

1. Rouviere H (1932) Anatomie humaine descriptive et topographique. Masson, Paris
2. Amsterdam JD, Marinelli DL, Arger P, Winokur A (1987) Assessment of adrenal gland volume by computed tomography in depressed patients and healthy volunteers: a pilot study. *Psychiatry Res* 21:189–197
3. Rubin RT, Phillips JJ (1991) Adrenal gland volume determination by computed tomography and magnetic resonance imaging in normal subjects. *Invest Radiol* 26:465–469
4. Farragher SW, Jara H, Chang KJ, Hou A, Soto JA (2005) Liver and spleen volumetry with quantitative MR imaging and dual-space clustering segmentation. *Radiology* 237:322–328
5. Hiroshige S, Shimada M, Harada N et al (2003) Accurate preoperative estimation of liver-graft volumetry using three-dimensional computed tomography. *Transplantation* 75:1561–1564
6. Hoshida Y, Shiratori Y, Koike Y et al (2002) Hepatic volumetry to predict adverse events in percutaneous ablation of hepatocellular carcinoma. *Hepatology* 49:451–455
7. Nakayama Y, Li Q, Katsuragawa S et al (2006) Automated hepatic volumetry for living related liver transplantation at multisection CT. *Radiology* 240
8. Schiano TD, Bodian C, Schwartz ME, Glajchen N, Min AD (2000) Accuracy and significance of computed tomographic scan assessment of hepatic volume in patients undergoing liver transplantation. *Transplantation* 69:545–550
9. Vauthey JN, Chaoui A, Do KA et al (2000) Standardized measurement of the future liver remnant prior to extended liver resection: methodology and clinical associations. *Surgery* 127:512–519

10. Constantin JM GS, Rouby JJ, Futier E, Gallix B, Jung B, Bazin JE, Jaber S (2009) CT-scan lung morphology predicts the response to a recruitment maneuver in acute respiratory distress syndrome patients. *Crit Care* 13 (Suppl 1)
11. Helm E, Talakoub O, Grasso F et al (2009) Use of dynamic CT in acute respiratory distress syndrome (ARDS) with comparison of positive and negative pressure ventilation. *Eur Radiol* 19:50–57
12. de Hoop B, Gietema H, van Ginneken B, Zanen P, Groenewegen G, Prokop M (2009) A comparison of six software packages for evaluation of solid lung nodules using semi-automated volumetry: what is the minimum increase in size to detect growth in repeated CT examinations. *Eur Radiol* 19:800–808
13. Pojunas KW, Daniels DL, Williams AL, Thorsen MK, Haughton VM (1986) Pituitary and adrenal CT of Cushing syndrome. *AJR Am J Roentgenol* 146:1235–1238
14. Chanques G, Annane D, Jaber S, Gallix B (2007) Enlarged adrenals during septic shock. *Intensive Care Med* 33:1671–1672
15. Bone RC, Sibbald WJ, Sprung CL (1992) The ACCP-SCCM consensus conference on sepsis and organ failure. *Chest* 101:1481–1483
16. Dellinger RP, Levy MM, Carlet JM et al (2008) Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Intensive Care Med* 34:17–60
17. Jones AE, Saak K, Kline JA (2008) Performance of the Mortality in Emergency Department Sepsis score for predicting hospital mortality among patients with severe sepsis and septic shock. *Am J Emerg Med* 26:689–692
18. Sprung CL, Annane D, Keh D et al (2008) Hydrocortisone therapy for patients with septic shock. *N Engl J Med* 358:111–124
19. Kazan-Tannus JF, Dialani V, Kataoka ML et al (2007) MR volumetry of brain and CSF in fetuses referred for ventriculomegaly. *AJR Am J Roentgenol* 189:145–151
20. Kilian AK, Schaible T, Hofmann V, Brade J, Neff KW, Busing KA (2009) Congenital diaphragmatic hernia: predictive value of MRI relative lung-to-head ratio compared with MRI fetal lung volume and sonographic lung-to-head ratio. *AJR Am J Roentgenol* 192:153–158
21. Kim HJ, Kim AY, Oh ST et al (2005) Gastric cancer staging at multi-detector row CT gastrography: comparison of transverse and volumetric CT scanning. *Radiology* 236:879–885
22. Lee SM, Kim SH, Lee JM et al (2008) Usefulness of CT volumetry for primary gastric lesions in predicting pathologic response to neoadjuvant chemotherapy in advanced gastric cancer. *Abdom Imaging* 34:430–440
23. Parrish FJ (2007) Volume CT: state-of-the-art reporting. *AJR Am J Roentgenol* 189:528–534
24. Plathow C, Klopp M, Thieke C et al (2008) Therapy response in malignant pleural mesothelioma-role of MRI using RECIST, modified RECIST and volumetric approaches in comparison with CT. *Eur Radiol* 18:1635–1643
25. Prasad SR, Jhaveri KS, Saini S, Hahn PF, Halpern EF, Sumner JE (2002) CT tumor measurement for therapeutic response assessment: comparison of unidimensional, bidimensional, and volumetric techniques initial observations. *Radiology* 225:416–419
26. Zhou XP, Lu T, Wei YG, Chen XZ (2007) Liver volume variation in patients with virus-induced cirrhosis: findings on MDCT. *AJR Am J Roentgenol* 189:W153–W159
27. Dinant S, de Graaf W, Verwer BJ et al (2007) Risk assessment of posthepatectomy liver failure using hepatobiliary scintigraphy and CT volumetry. *J Nucl Med* 48:685–692
28. Das M, Ley-Zaporozhan J, Gietema HA et al (2007) Accuracy of automated volumetry of pulmonary nodules across different multislice CT scanners. *Eur Radiol* 17:1979–1984
29. Goo JM, Tongdee T, Tongdee R, Yeo K, Hildebolt CF, Bae KT (2005) Volumetric measurement of synthetic lung nodules with multi-detector row CT: effect of various image reconstruction parameters and segmentation thresholds on measurement accuracy. *Radiology* 235:850–856
30. Petrou M, Quint LE, Nan B, Baker LH (2007) Pulmonary nodule volumetric measurement variability as a function of CT slice thickness and nodule morphology. *AJR Am J Roentgenol* 188:306–312
31. Zhang J, Kang SK, Wang L, Touijer A, Hricak H (2009) Distribution of renal tumor growth rates determined by using serial volumetric CT measurements. *Radiology* 250:137–144
32. Brownlie K, Kreef L (1978) Computer assisted tomography of normal suprarenal glands. *J Comput Assist Tomogr* 2:1–10
33. El-Sherief MA, Hemmingsson A (1982) Computed tomography of the normal adrenal gland. Pathologic implications. *Acta Radiol Diagn (Stockh)* 23:433–442
34. Montagne JP, Kressel HY, Korobkin M, Moss AA (1978) Computed tomography of the normal adrenal glands. *AJR Am J Roentgenol* 130:963–966
35. Goldman SM, Gatewood OM, Walsh PC, Dornhorst A, Siegelman SS (1982) CT configuration of the enlarged adrenal gland. *J Comput Assist Tomogr* 6:276–280
36. Rubin RT, Phillips JJ, McCracken JT, Sadow TF (1996) Adrenal gland volume in major depression: relationship to basal and stimulated pituitary-adrenal cortical axis function. *Biol Psychiatry* 40:89–97
37. Marik PE, Zaloga GP (2003) Adrenal insufficiency during septic shock. *Crit Care Med* 31:141–145
38. Vincent JL, Moreno R, Takala J et al (1996) The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 22:707–710