



HAL
open science

Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa

Basile Kamgang, Sébastien Marcombe, Fabrice Chandre, Elysée Nchoutpouen, Philippe Nwane, Josiane Etang, Vincent Corbel, Christophe Paupy

► **To cite this version:**

Basile Kamgang, Sébastien Marcombe, Fabrice Chandre, Elysée Nchoutpouen, Philippe Nwane, et al.. Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. *Parasites & Vectors*, 2011, 4 (1), pp.79. 10.1186/1756-3305-4-79 . hal-03002477

HAL Id: hal-03002477

<https://hal.umontpellier.fr/hal-03002477>

Submitted on 12 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License

RESEARCH

Open Access

Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa

Basile Kamgang^{1,2}, Sébastien Marcombe¹, Fabrice Chandre¹, Elysée Nchoutpouen^{1,2}, Philippe Nwane², Josiane Etang², Vincent Corbel^{1,3} and Christophe Paupy^{1,4*}

Abstract

Background: *Aedes aegypti* (Linnaeus, 1762) and *Aedes albopictus* (Skuse, 1894) are the main vectors of dengue (DENV) and chikungunya (CHIKV) viruses worldwide. As there is still no vaccine or specific treatment for DENV and CHIKV, vector control remains the cornerstone of prevention and outbreak control. Unfortunately, vector control programs are facing operational challenges with mosquitoes becoming resistant to commonly used insecticides in several areas through the world. Throughout Central Africa no recent data are available susceptible/resistant status of either vector species since the introduction/arrival of *Ae. albopictus* in this area. We therefore studied the level of resistance of these two major vectors to insecticides commonly used in Africa for mosquito control.

Results: *Aedes aegypti* and *Ae. albopictus* were sampled in six urban localities of Cameroon (Garoua, Bertoua, Yaoundé, Bafia, Buea) and Gabon (Libreville). Larval bioassays, carried out to determine the lethal concentrations (LC₅₀ and LC₉₅) and resistance ratios (RR₅₀ and RR₉₅) suggested that both vector species were susceptible to *Bti* (*Bacillus thuringiensis var israeliensis*) and temephos. Bioassays were also performed on adults using WHO diagnostic test kits to assess phenotypic resistance to deltamethrin, DDT, fenitrothion and propoxur. These experiments showed that one population of *Ae. aegypti* (Libreville) and two populations of *Ae. albopictus* (Buea and Yaoundé) were resistant to DDT (mortality 36% to 71%). Resistance to deltamethrin was also suspected in *Ae. albopictus* from Yaoundé (83% mortality). All other field mosquito populations were susceptible to deltamethrin, DDT, fenitrothion and propoxur. No increase in the knockdown times (Kdt₅₀ and Kdt₉₅) was noted in the Yaoundé resistant population compared to other *Ae. albopictus* populations, suggesting the possible involvement of metabolic resistance to deltamethrin and DDT.

Conclusion: In view of the recent increase in dengue and chikungunya outbreaks in Central Africa, these unique comparative data on the insecticide susceptibility of *Ae. aegypti* and *Ae. albopictus* could help public health services to design more effective vector control measures.

Background

Dengue virus (DENV, *Flaviviridae*, *Flavivirus*) and chikungunya virus (CHIKV, *Togaviridae*, *Alphavirus*) are mosquito-borne viruses of medical concern in most tropical regions. With about 50-100 million reported cases annually, including 500 000 severe cases of dengue haemorrhagic fever (DHF) or dengue shock syndrome (DSS), DENV is the most prevalent mosquito-borne human virus worldwide [1]. In West and Central Africa, where DENV epidemics remained limited until 5 years ago, the

number of outbreaks or case reports has increased significantly in several areas, including Cameroon in 2006 [2,3], Gabon in 2007 [4], and West African countries such as Mali in 2008, and Cape Verde and Senegal in 2009 [5]. Similarly, CHIKV, which previously caused only sporadic outbreaks in sub-Saharan Africa [6], has recently emerged in several urban epidemic foci in Central Africa [4].

Aedes aegypti (Linnaeus, 1762) and *Aedes albopictus* (Skuse, 1894) are the main epidemic vectors of DENV and CHIKV worldwide [2,7]. Both species occur in sub-Saharan Africa: *Ae. aegypti* is native to Africa and *Ae. albopictus* has recently invaded several Central African countries [8]. Recent observations in Cameroon [9] and

* Correspondence: christophe.paupy@ird.fr

¹Institut de Recherche pour le Développement (IRD), UMR 224 MIVEGEC, BP 64501, 34394 Montpellier, France

Full list of author information is available at the end of the article

Gabon [10] indicate that these two species infest urban environments.

As there is still no vaccine or specific treatment for DENV or CHIKV, vector control remains the cornerstone of prevention and outbreak control. Conventional control strategies rely on the reduction of larval sources by eradicating water-holding containers that serve as larval habitats, and by using larvicides (e.g. temephos and *Bacillus thuringiensis var israeliensis* [*Bti*]) in natural and/or domestic breeding sites [11]. Space spraying (using pyrethroids or organophosphates) is generally used when larval source reduction fails to reduce the density of adult mosquitoes, and also during outbreak situations [12]. Unfortunately, many vector control programs are threatened by the development of insecticide resistance in *Ae. aegypti* and *Ae. albopictus* [13-16]. Resistance to multiple insecticides (e.g. pyrethroids and organophosphates) has been reported in *Ae. aegypti* in South-East Asia [17,18], South America and the Caribbean [19-21]. There have been fewer reports of insecticide resistance in *Ae. albopictus* [15,22], although a significant decrease in susceptibility to permethrin, malathion and temephos has been observed in Thailand [14]. In West and Central Africa, most available data on pesticide resistance concern *Ae. aegypti* but date back more than 30-40 years [23], while virtually no such data are available for *Ae. albopictus*, owing to its recent introduction in Africa. In order to implement effective and sustainable arbovirus vector control measures, there is an urgent need to determine the susceptibility of the two major vectors of DENV and CHIK to insecticides commonly used for mosquito control. We therefore examined the distribution and insecticide susceptibility of both larval and adult *Ae. aegypti* and *Ae. albopictus* populations in sentinel sites from Central Africa (Cameroon and Gabon) using WHOPES-approved procedures [24,25].

Materials and methods

Mosquito sampling

Aedes aegypti and *Ae. albopictus* mosquitoes were sampled as larvae or pupae in six urban localities (Figure 1): Garoua (09°16'N; 13°25'E), Bertoua (04°33'N; 13°46'E), Yaoundé (03°54'N; 12°29'E), Bafia (04°44'N; 11°11'E) and Buea (04°09'N; 09°13'E) in Cameroon in April 2007, and Libreville, Gabon (00°23'N; 09°27'E) in June 2007. An additional *Ae. aegypti* population (Bénoué) was sampled in the Bénoué National Park (08°15'N; 13°49'E). The latter population, which is sylvan and thus considered as naïve to pesticide, was used as a susceptible reference strain. Immature stages (field generation, F0) were collected from water in domestic (e.g. jars, tanks), peri-domestic (e.g. tires) and natural environments (e.g. tree holes). For each sampling site, larvae or pupae from 2-6 larval breeding places were

collected, stored in plastic boxes and transferred to insectaries for rearing to the adult stage. Once identified as *Ae. aegypti* or *Ae. albopictus*, mosquitoes of the same species and from the same locality were pooled to prevent inbreeding. Mosquitoes were reared to generation F1 for larval and adult bioassays. Mosquito populations were maintained at insectaries conditions (27°C +/- 2°C; relative humidity 80% +/- 10%) and females were fed on rabbits to complete their gonotrophic cycle.

Larval bioassay

The susceptibility of larvae to temephos (organophosphate, Sigma Aldrich-Pestanal, Seelze, Germany) and *Bti* (VectoBac, Bayer CropScience, Monheim, Germany) was assessed using standard WHO bioassays [24]. Stock solutions and serial dilutions were prepared in 95% ethanol (temephos) or mineral water (*Bti*) and were stored at +4°C. Each bioassay used 25 late-third or early-fourth instars in plastic cups with 99 ml of mineral water and 1 ml insecticide solution at the required concentration. We used 5 concentrations in the range of activity of each insecticide and 50 to 100 larvae per concentration (with two to four replicates, depending on the sample and the number of larvae available). Each bioassay included a control group which received 1 mL of ethanol (for temephos) or 1 mL of mineral water (for *Bti*) only. Tests were run at 27°C ± 2°C, and mortality was assessed after 24 h of insecticide exposure. For each mosquito strain, one replicate corresponded to a single rearing batch.

Adult bioassay

The procedure used for adult bioassays followed the standard WHO protocol [25]. Papers were impregnated using acetone solutions of insecticide and silicone oil as the carrier at the "Laboratoire de Lutte contre les Insectes Nuisibles" (LIN), Montpellier, France (WHO Collaborating Centre) using diagnostic concentrations defined for *Aedes* laboratory strain (Bora) susceptible reference strain [17]. Four technical-grade compounds representing the 4 major classes of insecticide (carbamate, organochlorine, pyrethroid and organophosphate) were used as follows: 0.3% propoxur (Bayer CropScience, Monheim, Germany), 4% DDT (Hindustan Insecticides Limited, New Delhi, India), 0.06% deltamethrin (Agrevo Environmental Health, Berkhamsted UK) and 0.5% fenitrothion (SUPELCO, Bellefonte, USA). Two to four batches of 25 non-blood-fed females (2-4 days of age) were introduced into exposure tubes containing impregnated filter papers for 60 minutes. The number of knocked-down (Kd) mosquitoes was recorded every 5 minutes. The mosquitoes were then transferred to a recovery tube containing 10% glucose solution and maintained at 27°C ± 2°C with 80% ± 10% relative humidity. Mortality was recorded 24 hours post-exposure.




Figure 1 *Aedes aegypti* and *Ae. albopictus* sampling sites in Cameroon and Gabon.

For each strain a batch of 25 mosquitoes was used as control.

Data analysis

Larval and adult mortality rates were corrected using Abbott's formula (Abbott 1925) in case of control

mortality >5% but less than 20%. Data from larval bioassays were analyzed with Win DL 2.0 software [26], which uses the iterative method of maximum likelihood to fit a linear regression between the log of the insecticide concentration and the probit of mortality. Goodness-of-fit was estimated with the Pearson χ^2 test. Win DL 2.0

provided also the slope of the regression lines and the estimates of lethal concentrations (LC_{50} and LC_{95}) with their 95% confidence intervals (CI). Resistance ratios (RR_{50} and RR_{95}) of each population compared to the Bénoué strain of *Ae. aegypti* (used as reference) were calculated as follows: $RR_{50} = LC_{50}$ assay/ LC_{50} reference; $RR_{95} = LC_{95}$ assay/ LC_{95} reference. A mosquito population was considered susceptible when RR_{50} was less than 2, potentially resistant when RR_{50} was between 2 and 5, and resistant when RR_{50} was over 5. For adult bioassays, resistant/susceptible status was defined according to WHO criteria [27]. Mosquitoes were considered susceptible if the mortality rates were greater than 97% and resistant if mortality rates were less than 80%. Mortality rates between 80-97% suggested possible resistance.

For deltamethrin and DDT, the knockdown times (KdT_{50} and KdT_{95}) and their 95% confidence intervals were estimated with Win DL 2.0 software.

Results

Larval bioassay

Larval bioassays using *Bti* showed that LC_{50} and LC_{95} for *Ae. albopictus* ranged from 0.18 mg/l for the Bénoué strain (susceptible reference strain) to 0.69 mg/L for the Yaoundé and Buea populations. Against *Ae. aegypti*, *Bti* acted in a similar range of concentrations, with LC_{50} and LC_{95} ranging from 0.18 mg/L for Garoua to 0.91 mg/L for Bafia (Table 1). RR_{50} ranged from 1.00 (Garoua) to 1.50 (Bertoua) and RR_{95} from 1.13 (Libreville) to 2.84 (Bafia). These values and their 95% CIs indicated that all *Ae. aegypti* and *Ae. albopictus* populations were similarly susceptible to *Bti*.

Against *Ae. aegypti*, bioassays showed that temephos LC_{50} values ranged from 0.004 mg/L (Garoua) to 0.009 mg/l (Bertoua) and the LC_{95} from 0.006 mg/L (Garoua) to 0.026 mg/L (Bertoua) (Table 2). RR_{50} ranged from 0.75 (Garoua) to 1.75 (Bertoua) and RR_{95} from 0.66 (Garoua) to 2.68 (Bertoua). These findings suggested that all the mosquito samples tested were susceptible to temephos. Concerning the Bertoua sample, although the RR_{95} value was over 2 indicating a suspected resistance, the IC values did not confirm this trend. LC_{50} values for *Ae. albopictus* ranged from 0.005 (Bertoua) to 0.008 mg/L (Libreville) and LC_{95} values from 0.008 (Yaoundé) to 0.017 mg/L (Libreville). RR_{50} ranged from 0.92 (Bertoua) to 1.43 mg/L (Libreville) and RR_{95} from 0.82 (Yaoundé) to 1.75 mg/L (Libreville), indicating susceptibility of all the mosquito populations to temephos.

Adult bioassay

After 24 h post-exposure, one population of *Ae. aegypti* (Libreville) and two populations of *Ae. albopictus* (Buea and Yaoundé) were resistant to DDT (mortality 36% to 71%). DDT mortality rates suggested probable resistance

in the Garoua *Ae. aegypti* population (97%) and the Bertoua *Ae. albopictus* population (80%). Probable resistance to deltamethrin was also detected in the Yaoundé *Ae. albopictus* population (80%). All *Ae. aegypti* except Yaoundé, and all *Ae. albopictus* populations were susceptible to deltamethrin, propoxur and fenitrothion (99-100%) (Table 3).

The estimated knockdown times (KdT) for *Ae. aegypti* mosquitoes exposed to deltamethrin indicated that the KdT_{50} and KdT_{95} values of the field populations were similar (overlapping 95% CIs) to those of the reference strain (Table 4). KdT_{50} values recorded in Yaoundé and Buea *Ae. albopictus* strains were higher than for the reference strain. Despite the ICs calculated for these two strains did not overlapped with the IC of the reference, the increase in KdT seemed no significant ($P > \text{or} = 0.05$). The DDT KdT_{50} for all *Ae. albopictus* samples and the KdT_{95} for *Ae. albopictus* Bertoua were significantly higher than the corresponding values for all *Ae. aegypti* samples. These differences in KdT between *Ae. aegypti* and *Ae. albopictus* may be physiological and/or species-dependent.

Discussion

All *Ae. aegypti* and *Ae. albopictus* samples collected in Cameroon and Libreville (Gabon) were susceptible to *Bti* and temephos. Resistance to *Bti* has been described in *Culex pipiens* in the USA [28] and suspected in *Aedes rusticus* in France [29], but no resistance to this pesticide has been reported to date in *Ae. aegypti* or *Ae. albopictus* worldwide. Temephos is the most widely used larvicide for dengue vector control. Resistance to this organophosphate has been recorded in *Ae. aegypti* in Asia [30-34] and South America [35-37], and in *Ae. albopictus* in Malaysia [38] and Thailand [14]. Mouchet et al. [23] observed full susceptibility of *Ae. aegypti* larvae to temephos in Cameroon and Gabon (Libreville and Yaoundé) during the 1970s. Up to now, larvicides targeting these two species have rarely been used in African countries, thus explaining the persistent susceptibility to temephos and *Bti* in this part of the world.

WHO bioassays carried out on adult mosquitoes showed DDT resistance in one *Ae. aegypti* population from Gabon (Libreville) and two *Ae. albopictus* populations from Cameroun (Buea and Yaoundé). In addition, DDT resistance was suspected in the *Ae. albopictus* sample from Bertoua. DDT resistance has been widely found in *Ae. aegypti* worldwide, including in Cameroon [23]. *Kdr* mutation of the voltage-gated sodium channel has been found to confer resistance to DDT in *Ae. aegypti* [39] but detoxifying enzymes such as glutathione S-transferases (GSTs) can also play a key role in DDT metabolism and resistance [40]. Although DDT resistance has also been recorded in *Ae. albopictus* in Thailand and

Table 1 Susceptibility of *Ae. aegypti* and *Ae. albopictus* larvae to Bti

Sample	N	Regression line	Pearson χ^2 goodness of fit	LC ₅₀ (95% CI)	LC ₉₅ (95% CI)	RR ₅₀	RR ₉₅
<i>Ae. aegypti</i>							
Garoua	493	Y = 5.67X + 4.15	0.69	0.18 (0.03-0.45)	0.36 (0.30-0.45)	1.00	1.13
Bertoua	498	Y = 3.84X + 2.13	0.62	0.27 (0.19-0.39)	0.75 (0.20-2.77)	1.50	2.34
Yaoundé	252	Y = 8.77X + 5.76	0.27	0.22 (0.20-0.44)	0.34 (0.26-0.44)	1.22	1.06
Bafia	500	Y = 2.99X + 1.75	0.54	0.25 (0.21-0.31)	0.91 (0.17-4.89)	1.39	2.84
Libreville	500	Y = 6.06X + 4.28	0.21	0.19 (0.09-0.24)	0.36 (0.32-0.46)	1.06	1.13
<i>Ae. albopictus</i>							
Yaoundé	500	Y = 1.91X - 0.04	0.64	0.19 (0.17-0.21)	0.69 (0.26-1.85)	1.05	1.06
Bertoua	477	Y = 16.31X + 9.32	0.94	0.27 (0.25-0.28)	0.34 (0.32-0.40)	1.50	1.13
Buea	375	Y = 3.08X + 2.13	0.51	0.20 (0.10-0.39)	0.69 (0.26-1.85)	1.11	1.06
Reference strain	496	Y = 7.04X + 5.11	0.40	0.18 (0.11-0.22)	0.32 (0.29-0.36)		

N: Total number of mosquitoes assayed; 50% and 95% lethal concentrations, LC₅₀ and LC₉₅, are expressed in mg/liter; 95%CI: 95% confidence interval; p > 0.05 suggests a well-fitting model, P < 0.05 suggests an invalid model population.

Table 2 Susceptibility of *Ae. aegypti* and *Ae. albopictus* larvae to Temephos

Sample	N	Regression line	Pearson χ^2 goodness of fit	LC ₅₀ (95%CI)	LC ₉₅ (95%CI)	RR ₅₀	RR ₉₅
<i>Ae. aegypti</i>							
Garoua	500	Y = 10.68X + 25.82	0.36	0.0040 (0.0037-0.0042)	0.0064 (0.0060-0.0070)	0.75	0.66
Bertoua	500	Y = 3.57X + 7.28	0.48	0.0090 (0.0048-0.0170)	0.0260 (0.0022-0.3100)	1.75	2.68
Yaoundé	253	Y = 11.36X + 25.68	0.30	0.0055 (0.0052-0.006)	0.0076 (0.0067-0.0100)	1.04	0.78
Bafia	500	Y = 7.07X + 16.04	0.006	0.0053 (0.0047-0.0084)	0.0092 (0.0068-0.0410)	1.02	0.95
Libreville	500	Y = 8.17X + 19.56	0.061	0.0043 (0.0020-0.0081)	0.0068 (0.0051-0.0080)	0.81	0.70
<i>Ae. albopictus</i>							
Bertoua	493	Y = 6.75X + 15.58	0.14	0.0049 (0.0013-0.0180)	0.0086 (0.0061-0.0120)	0.92	0.89
Yaoundé	497	Y = 8.00X + 18.38	0.30	0.0050 (0.0021-0.0110)	0.0080 (0.0064-0.0100)	0.94	0.82
Buea	375	Y = 9.57X + 20.55	0.27	0.0071 (0.0065-0.0200)	0.012 (0.0087-0.0500)	1.34	1.03
Libreville	275	Y = 4.56X + 25.68	0.11	0.0076 (0.0023-0.0240)	0.017 (0.0017-0.1700)	1.43	1.75
Reference strain	500	Y = 6.22X + 14.17	0.57	0.0053 (0.0035-0.0079)	0.0097 (0.0073-0.012)		

N: Total number of mosquitoes assayed; 50% and 95% lethal concentrations, LC₅₀ and LC₉₅, are expressed in mg/liter; 95%CI: 95% confidence interval; p > 0.05 suggests a well-fitting model, P < 0.05 suggests an invalid model population.

Table 3 *Aedes aegypti* and *Ae. albopictus* mortality rates 24 h after exposure to insecticides at diagnostic doses

	Mortality rate (%)							
	0.06% Deltamethrin		4% DDT		0.3% Propoxur		0.5% Fenitrothion	
	Assay	Control	Assay	Control	Assay	Control	Assay	Control
<i>Ae. aegypti</i>								
Garoua	100 (89)	2 (50)	96.8 (95)	8 (50)	100 (100)	8 (50)	100 (99)	2 (50)
Bertoua	100 (100)	2 (50)	-	-	-	-	100 (98)	2 (50)
Yaoundé	100 (95)	4 (50)	-	-	-	-	100 (96)	4 (50)
Bafia	100 (97)	4 (50)	-	-	-	-	100 (97)	4 (50)
Libreville	100 (97)	0 (50)	70.7 (99)	0 (50)	99 (100)	0 (50)	100 (99)	0 (50)
<i>Ae. albopictus</i>								
Bertoua	100 (80)	0 (50)	80.5 (87)	8 (50)	100 (94)	8 (50)	100 (61)	0 (50)
Yaoundé	83.3 (96)	2 (50)	36.3 (91)	12 (50)	100 (96)	12 (50)	100 (98)	2 (50)
Buea	100 (50)	4 (50)	47.0 (85)	2 (50)	100 (92)	2 (50)	100 (50)	4 (50)
Reference strain	100 (96)	2 (50)	-	-	-	-	100 (100)	2 (50)

Numbers in brackets indicate the number of mosquitoes assayed; -: sample not assayed.

Japan [41,42], the underlying mechanisms remain unclear.

This study also showed full susceptibility of *Ae. aegypti* and *Ae. albopictus* to deltamethrin, except in Yaoundé where the *Ae. albopictus* population showed

mortality rates of around 80%, strongly suggesting resistance. Pyrethroid resistance in *Ae. aegypti* is widespread [43,15] and is usually associated with an altered amino acid sequence of the target protein, the sodium channel that confers knockdown resistance when altered (the so-

Table 4 Knockdown times of *Ae. aegypti* and *Ae. albopictus* exposed to 0.06% deltamethrin and 4% DDT

Locality	<i>Ae. aegypti</i>		<i>Ae. albopictus</i>	
	KdT ₅₀ (95%CI); P*	KdT ₉₅ (95%CI); P*	KdT ₅₀ (95%CI); P*	KdT ₉₅ (95%CI); P*
	Deltamethrin		Deltamethrin	
Garoua	5.3 (4.7-5.9); P = 0.47	11.4 (9.7-14.2); P = 0.24	-	-
Bertoua	5.8 (3.3-8.2); P = 0.46	24.6 (16.6-51.4); P = 0.025	7.8 (7.3-8.3); P = 0.25	15.9 (14.4-18.0); P = 0.38
Yaoundé	6.7 (3.7-9.7); P = 0.35	22.8 (15.0-58.4); P = 0.05	11.2 (10.6-11.8); P = 0.06	17.3 (16.3-19.0); P = 0.28
Bafia	5.4 (3.3-7.4), P = 0.48	14.6 (10.1-36.9); P = 0.49	-	-
Buea	-	-	11.7 (10.2-13.3); P = 0.06	21.2 (18.1-27.4); P = 0.09
Libreville	5.9 (5.1-6.8); P = 0.45	13.4 (11.2-17.9); P = 0.40	-	-
Reference strain	5.5 (4.6-6.3)	14.5 (11.9-19.4)	-	-
	DDT		DDT	
Garoua	53.6 (49.9-59.2)	101.9 (84.0-146.0)	-	-
Bertoua	-	-	84.8 (72.8-106.9)	290.8 (200.7-523.0)
Yaoundé	-	-	77.4 (68.5-113.2)	109.2 (86.1-235.3)
Buea	-	-	68.0 (63.6-78.2)	93.9 (80.7-131.9)
Libreville	58.7 (55.8-62.7)	107.7 (94.6-129.2)	-	-

KdT: Knockdown times expressed in minutes; 95%CI: 95% confidence interval. *: P is the probability of homogeneity between the sample and the reference (Mantel-Haenszel chi square).

called *Kdr* mutation) [39,44]. Pyrethroid resistance has also been linked to an increase in metabolic process through over-transcription of detoxification genes [21]. Although the number of studies reporting insecticide resistance in *Ae. albopictus* has increased this decade [17,41,42,45,46], few have shown evidence of pyrethroid resistance [14]. The *Kdr* mutation has been found in *Ae. aegypti* [39,16] but has never been reported in *Ae. albopictus*. Further molecular and biochemical studies are needed to identify the genetic basis of pyrethroid resistance in *Ae. albopictus* in Yaoundé. The absence of cross-resistance between DDT and deltamethrin in all mosquito populations (except *Ae. albopictus* collected in Yaoundé) suggests the involvement of metabolic resistance rather than a target site modification (*Kdr* mutation) in the sodium channel. It is unclear whether cross-resistance to deltamethrin and DDT in the *Ae. albopictus* population in Yaoundé is due to common or different resistance mechanisms.

The source of DDT and deltamethrin resistance observed here in DENV and CHIKV vectors is unclear. Indeed, in Central Africa, insecticide treatment specifically targeting *Ae. aegypti* or *Ae. albopictus* is extremely limited. It is possible that insecticides used to control other insects of medical or agricultural interest exert indirect selection pressure on these two mosquito species. For example, indoor residual spraying of DDT for malaria control [47] was suspected of favouring the selection of DDT resistance in *Anopheles* [48] as well as in *Aedes* [42,45]. Contamination of larval breeding places by insecticides used in agriculture (cotton, vegetables) has also been shown to select for DDT and pyrethroid resistance in malaria vectors [48-52]. Concerning *Ae. albopictus*, which recently spread to Cameroon [8], one cannot exclude the possibility that the invading population possessed a resistance background.

Conclusion

This is the first study of the susceptibility status of *Ae. aegypti* and *Ae. albopictus* to various insecticides in Central Africa. The increase in dengue and chikungunya outbreaks in this region calls for more robust surveillance and vector control. Our observations could help to guide insecticide-based strategies, although broader monitoring of insecticide resistance in *Aedes* mosquitoes is needed. Further molecular and biochemical studies are also needed to determine the mechanisms involved in insecticide resistance among arboviruses vectors in Central Africa.

Acknowledgements

We thank Philippe Bousses for significant help with mosquito sampling. This research was supported by the French government through Agence Nationale pour la Recherche that funded the Epidengue project (ANR 05 SEST 010-01). Basile Kamgang is supported by an IRD/DSF PhD fellowship.

Author details

¹Institut de Recherche pour le Développement (IRD), UMR 224 MIVEGEC, BP 64501, 34394 Montpellier, France. ²Laboratoire de Recherche sur le Paludisme, Organisation de Coordination pour la Lutte contre les Endémies en Afrique Centrale (OCEAC), BP 288, Yaoundé, Cameroun. ³Centre de Recherche Entomologique de Cotonou (CREC), BP 2604 Cotonou, République du Bénin. ⁴Centre International de Recherches Médicales de Franceville (CIRMF), BP 769, Franceville, Gabon.

Authors' contributions

BK and CP designed the study and monitored its implementation. BK, EN and CP participated to field sampling. BK, EN, PN carried out bioassay tests, under guidance of JE. BK, SM and CP analyzed the data. BK, SM, VC and CP wrote the manuscript which was critically revised by FC. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 18 February 2011 Accepted: 15 May 2011

Published: 15 May 2011

References

1. DJ Gubler, Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol.* **10**, 100–103 (2002). doi:10.1016/S0966-842X(01)02288-0
2. C Peyrefitte, D Rousset, B Pastorino, R Pouillot, M Bessaud, F Tock, H Mansaray, O Merle, A Pascual, C Paupy, A Vessiere, P Imbert, P Tchendjou, JP Durand, H Tolou, M Grandadam, Chikungunya virus, Cameroon 2006. *Emerg Infect Dis.* **13**, 768–771 (2007)
3. R Krippner, G von Laer, First confirmed dengue-1 fever cases reported from Cameroon. *J Travel Med.* **9**, 273–274 (2002)
4. EM Leroy, D Nkoghe, B Ollomo, C Nze-Nkoghe, P Becquart, G Grard, X Pourrut, R Charrel, G Moureau, A Ndjoi-Mbiguino, X De Lamballerie, Concurrent Chikungunya and Dengue Virus Infections during Simultaneous Outbreaks, Gabon, 2007. *Emerg Inf Dis.* **15**, 591–593 (2009). doi:10.3201/eid1504.080664
5. L Franco, A Di Caro, F Carletti, O Vapalahti, C Renaudat, H Zeller, A Tenorio, Recent expansion of dengue virus serotype 3 in West Africa. *Euro Surveill.* **15**, 19490 (2010)
6. M Diallo, J Thonnon, M Traore-Lamizana, D Fontenille, Vectors of Chikungunya virus in Senegal: current data and transmission cycles. *Am J Trop Med Hyg.* **60**, 281–286 (1999)
7. CY Kow, LL Koon, FY Pang, Detection of dengue viruses in field caught male *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in Singapore by type-specific PCR. *J Med Entomol.* **38**, 475–479 (2001). doi:10.1603/0022-2585-38.4.475
8. C Paupy, H Delatte, L Bagny, V Corbel, D Fontenille, *Aedes albopictus*, an arbovirus vector: from the darkness to the light. *Microbes Infect.* **11**, 1177–1185 (2009). doi:10.1016/j.micinf.2009.05.005
9. B Kamgang, J Youta Happei, P Boisier, F Njiokou, JP Hervé, F Simard, C Paupy, Geographic and ecological distribution of dengue and chikungunya virus vectors, *Aedes aegypti* and *Aedes albopictus*, in urban environments of Cameroon (Central Africa). *Med Vet Ent.* **24**, 132–141 (2010)
10. C Paupy, B Ollomo, B Kamgang, S Moutailler, D Rousset, M Demanou, JP Hervé, EM Leroy, F Simard, Role of *Ae. albopictus* versus *Ae. aegypti* in dengue and chikungunya emergences in Central Africa: Laboratory and field evidence. *Vector Borne Zoonotic Dis.* **10**, 259–266 (2010). doi:10.1089/vbz.2009.0005
11. WHO, Report of the Scientific Working Group on dengue. (Document WHO/TDR/SWG/08, Geneva, Switzerland, World Health Organization, 2006)
12. WHO, Dengue Haemorrhagic Fever: Diagnosis, Treatment, Prevention and Control. (WHO, Geneva, 1997), 2, pp. 1–84
13. R Yaicharoen, R Kiatfuengfoo, T Chaeronviriyaphap, P Rongnoparut, Characterization of deltamethrin, resistance in field populations of *Aedes aegypti* in Thailand. *J Vect Ecol.* **30**, 144–150 (2005)
14. A Ponlawat, JG Scott, LC Harrington, Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* across Thailand. *J Med Entomol.* **42**, 821–825 (2005). doi:10.1603/0022-2585(2005)042[0821:ISOAAA]2.0.CO;2
15. H Ranson, J Burhani, N Lumjuan, WC Black, Insecticide resistance in dengue vectors. *Tropika net Journal.* **1**(1) (2010)

16. S Marcombe, A Carron, F Darriet, M Etienne, P Agnew, M Tolosa, MM Yp-Tcha, C Lagneau, A Yébakima, V Corbel, Reduced Efficacy of Pyrethroid Space Sprays for Dengue Control in an Area of Martinique with Pyrethroid Resistance. *Am J Trop Med Hyg.* **80**, 745–751 (2009)
17. N Jirakanjanakit, P Rongnoparut, S Saengtharapit, T Chareonviriyaphap, S Duchon, C Bellec, S Yoksan, Insecticide susceptible/resistance status in *Aedes* (*Stegomyia*) *aegypti* and *Aedes* (*Stegomyia*) *albopictus* (Diptera: Culicidae) in Thailand during 2003–2005. *J Econ Entomol.* **100**, 545–550 (2007). doi:10.1603/0022-0493(2007)100[545:IRSIA]2.0.CO;2
18. N Jirakanjanakit, S Saengtharapit, P Rongnoparut, S Duchon, C Bellec, S Yoksan, Trend of Temephos resistance in *Aedes* (*Stegomyia*) mosquitoes in Thailand during 2003–2005. *Environ Entomol.* **36**, 506–511 (2007). doi:10.1603/0046-225X(2007)36[506:TOTRIA]2.0.CO;2
19. SC Rawlins, R Martinez, S Wiltshire, G Legall, A comparison of surveillance systems for the dengue vector *Aedes aegypti* in Port of Spain, Trinidad. *J Am Mosq Control Assoc.* **14**, 131–136 (1998)
20. MM Rodriguez, JA Bisset, Y De Armas, F Ramos, Pyrethroid insecticide-resistant strain of *Aedes aegypti* from Cuba induced by deltamethrin selection. *J Am Mosq Control Assoc.* **21**, 437–445 (2005). doi:10.2987/8756-971X(2006)21[437:PISOAA]2.0.CO;2
21. S Marcombe, R Poupardin, F Darriet, S Reynaud, J Bonnet, C Strode, C Brengues, A Yébakima, H Ranson, V Corbel, JP David, Exploring the molecular basis of insecticide resistance in the dengue vector *Aedes aegypti*: a case study in Martinique Island (French West Indies). *BMC Genomics.* **10**, 494 (2009). doi:10.1186/1471-2164-10-494
22. J Hemingway, NJ Hawkes, L McCarroll, H Ranson, The molecular basis of insecticide resistance in mosquitoes. *Insect Biochem Mol Biol.* **34**, 653–665 (2004). doi:10.1016/j.ibmb.2004.03.018
23. J Mouchet, V Cordellier, M Germain, P Carnevale, J Barathe, C Sannier, Résistance aux insecticides d'*Aedes aegypti* et *Culex pipiens fatigans* en Afrique centrale. (WHO/VBC/72/381, 12P, Geneva, Switzerland, World Health Organization, 1972)
24. WHO, Guidelines for laboratory and field testing of mosquito larvicides. (Document WHO/CDS/WHOPES/GCDPP/13, Geneva, Switzerland, World Health Organization, 2005)
25. WHO, Guidelines for testing mosquito adulticides for indoor residual spraying and treatment of mosquito nets. (Document WHO/CDS/NTD/WHOPES/GCDPP/3, Geneva, Switzerland, World Health Organization, 2006)
26. M Giner, C Vassal, Z Kouaik, F Chiroleu, JM Vassal, Win DL version 2.0. (Paris: CIRAD-CA, U.R.B./M.A.B.I.S, 1999)
27. WHO, Test Procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticide on treated surfaces. (Document WHO/CDS/CPC/MAL/98.12, Geneva, Switzerland, World Health Organization, 1998)
28. A Paul, LC Harrington, L Zhang, JG Scott, Insecticide resistance in *Culex pipiens* from New York. *J Am Mosq Control Assoc.* **21**, 305–309 (2005). doi:10.2987/8756-971X(2005)21[305:IRICPF]2.0.CO;2
29. M Paris, G Tetreau, F Laurent, M Lelu, L Despres, JP David, Persistence of *Bacillus thuringiensis israelensis* (*Bti*) in the environment induces resistance to multiple *Bti* toxins in mosquitoes. *Pest Manag Sci.* **67**, 122–128 (2011). doi:10.1002/ps.2046
30. HL Lee, W Lime, A reevaluation of the susceptibility of field collected *Aedes* (*Stegomyia*) *aegypti* (Linnaeus) larvae to temephos in Malaysia. *Mosquito-Borne Dis Bull.* **6**, 91–95 (1989)
31. T Chareonviriyaphap, B Aum-aung, S Ratanatham, Current insecticide resistance pattern in mosquito vectors in Thailand. *Southeast Asian J Trop Med Public Health.* **30**, 184–194 (1999)
32. P Paeporn, N Komalamisra, V Deesin, Y Rongsriyam, Y Eshita, S Thongrungrat, Temephos resistance in two forms of *Aedes aegypti* and its significance for the resistance mechanism. *Southeast Asian J Trop Med Public Health.* **34**, 786–792 (2003)
33. KA Polson, C Curtis, MS Chang, JG Olson, N Chantha, SC Rawlins, Susceptibility of two Cambodian population of *Aedes aegypti* mosquito larvae to temephos during 2001. *Dengue Bull.* **25**, 79–83 (2001)
34. CD Chen, WA Nazni, HL Lee, M Sofian-Azirun, Susceptibility of *Aedes aegypti* and *Aedes albopictus* to temephos in four study sites in Kuala Lumpur City Center and Selangor State, Malaysia. *Trop Biomed.* **22**, 207–216 (2005)
35. MLG Macoris, MTM Andrighetti, V Takaku, CM Glasser, VC Garbeloto, JE Bracco, Resistance of *Aedes aegypti* from the state of Sao Paulo, Brazil to organophosphates insecticides. *Mem Inst Oswaldo Cruz.* **98**, 703–708 (2003). doi:10.1590/S0074-02762003000500020
36. GA Llinás, E Seccacini, CN Gardenal, S Licastro, Current resistance status to temephos in *Aedes aegypti* from different regions of Argentina. *Mem Inst Oswaldo Cruz.* **105**, 113–116 (2010). doi:10.1590/S0074-02762010000100019
37. MA Melo-Santos, JJ Varjal-Melo, AP Araújo, TC Gomes, MH Paiva, LN Regis, AF Furtado, T Magalhaes, ML Macoris, MT Andrighetti, CF Ayres, Resistance to the organophosphate temephos: mechanisms, evolution and reversion in an *Aedes aegypti* laboratory strain from Brazil. *Acta Trop.* **113**, 180–189 (2010). doi:10.1016/j.actatropica.2009.10.015
38. WA Nazni, MY Kamaludin, HL Lee, TAR Rogayah, I Sa'diyah, Oxidase activity in relation to insecticides resistance in vectors of public health importance. *Trop Biomed.* **17**, 69–79 (2000)
39. C Brengues, NJ Hawkes, F Chandre, L McCarroll, S Duchon, P Guillet, S Manguin, JC Morgan, J Hemingway, Pyrethroids and DDT cross-resistance in *Aedes aegypti* is correlated with novel mutations in the voltage-gated sodium channel gene. *Med Vet Entomol.* **17**, 87–94 (2003). doi:10.1046/j.1365-2915.2003.00412.x
40. N Lumjuan, L McCarroll, LA Prapanthadara, J Hemingway, H Ranson, Elevated activity of an Epsilon class glutathione transferase confers DDT resistance in the dengue vector, *Aedes aegypti*. *Insect Biochem Mol Biol.* **35**, 861–871 (2005). doi:10.1016/j.ibmb.2005.03.008
41. P Somboon, LA Prapanthadara, W Suwonkerd, Insecticide susceptibility tests of *Anopheles minimus* s.l., *Aedes aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* in northern Thailand. *Southeast Asian J Trop Med Public Health.* **34**, 87–93 (2003)
42. H Kawada, Y Maekawa, M Abe, K Ohashi, SY Ohba, M Takagi, Spatial distribution and pyrethroid susceptibility of mosquito larvae collected from catch basins in parks in Nagasaki city, Nagasaki, Japan. *Jpn J Infect Dis.* **63**, 19–24 (2010)
43. J Hemingway, H Ranson, Insecticide in insect vectors of human diseases. *Annu Rev Entomol.* **45**, 371–391 (2000). doi:10.1146/annurev.ento.45.1.371
44. K Saavedra-Rodriguez, L Urdaneta-Marquez, S Rajatileka, M Moulton, AE Flores, I Fernandez-Salas, J Bisset, M Rodriguez, PJ McCall, MJ Donnelly, H Ranson, J Hemingway, WC Black IV, A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. *Insect Mol Biol.* **16**, 785–798 (2007). doi:10.1111/j.1365-2583.2007.00774.x
45. LT Ping, R Yatiman, LP Gek, Susceptibility of adult field strains of *Aedes aegypti* and *Aedes albopictus* in Singapore to pirimiphos-methyl and permethrin. *J Am Mosq Control Assoc.* **17**, 144–146 (2001)
46. R Romi, L Toma, F Severini, M Di Luca, Susceptibility of Italian populations of *Aedes albopictus* to Temephos and to other insecticides. *J Am Mosq Control Assoc.* **19**, 419–423 (2003)
47. CF Curtis, AE Mnzava, Comparison of house spraying and insecticide-treated nets for malaria control. *Bull World Health Organ.* **78**, 1389–1400 (2000)
48. V Corbel, R N'Guessan, C Brengues, F Chandre, L Djogbenou, T Martin, M Akogbeto, JM Hougard, M Rowland, Multiple insecticide resistance mechanisms in *Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta Trop.* **101**, 207–216 (2007). doi:10.1016/j.actatropica.2007.01.005
49. RF Djouaka, AA Bakare, ON Coulibaly, MC Akogbeto, H Ranson, J Hemingway, C Strode, Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics.* **13**, 538 (2008)
50. F Chandre, F Darriet, L Manga, M Akogbeto, O Faye, J Mouchet, P Guillet, Status of pyrethroid resistance in *Anopheles gambiae* sensu lato. *Bull World Health Organ.* **77**, 230–234 (1999)
51. A Diabate, T Baldet, F Chandre, M Akogbeto, TR Guiguemede, F Darriet, C Brengues, P Guillet, J Hemingway, GJ Small, JM Hougard, The role of agricultural use of insecticides in resistance to pyrethroids in *Anopheles gambiae* s.l. in Burkina Faso. *Am J Trop Med Hyg.* **67**, 617–622 (2002)
52. A Yadouleton, T Martin, G Padonou, F Chandre, A Asidi, L Djogbenou, R Dabiré, R Aikpon, M Boko, I Glitho, M Akogbeto, Cotton pest management practices and the selection of pyrethroid resistance in *Anopheles gambiae* population in Northern Benin. *Parasit Vectors.* **4**, 60 (2011). doi:10.1186/1756-3305-4-60

doi:10.1186/1756-3305-4-79

Cite this article as: Kamgang et al.: Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. *Parasites & Vectors* 2011 4:79.