



HAL
open science

New insights into the affinities, autoecology, and habit of the Mesozoic fern *Weichselia reticulata* based on the revision of stems from Bernissart (Mons Basin, Belgium)

Candela Blanco-moreno, Anne-Laure Decombeix, Cyrille Prestianni

► To cite this version:

Candela Blanco-moreno, Anne-Laure Decombeix, Cyrille Prestianni. New insights into the affinities, autoecology, and habit of the Mesozoic fern *Weichselia reticulata* based on the revision of stems from Bernissart (Mons Basin, Belgium). *Special papers in palaeontology*, 2021, 7 (3), pp.1351-1372. 10.1002/spp2.1344 . hal-03038289

HAL Id: hal-03038289

<https://hal.umontpellier.fr/hal-03038289v1>

Submitted on 12 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3 New insights on the affinities, autoecology, and habit of the Mesozoic fern *Weichselia*
4
5 *reticulata* based on the revision of stems from Bernissart (Mons Basin, Belgium)
6
7

8 CANDELA BLANCO-MORENO¹, ANNE-LAURE DECOMBEIX² and CYRILLE PRESTIANNI³
9

10 ¹Departamento de Biología, Universidad Autónoma de Madrid, 28049 Cantoblanco (Madrid),
11
12 Spain; e-mail: candela.blanco@uam.es
13

14 ²AMAP, Université de Montpellier, CIRAD, CNRS, INRA, IRD, Montpellier, France e-mail: anne-
15
16 laure.decombeix@cirad.fr
17

18 ³OD Earth and Life, Royal Belgian Institute of Natural Sciences, Rue Vautier 29, 1000, Brussels,
19
20 Belgium; e-mail: cprestianni@naturalsciences.be
21
22
23
24

25
26 **ABSTRACT**
27

28 The Mesozoic is a key period in fern evolution, with the rise of most modern families. *Weichselia*
29
30 *reticulata* is a widely distributed Jurassic-Cretaceous fern that has been suggested to belong to
31
32 the Matoniaceae or possibly the Marattiaceae. The most accepted classic whole-plant
33
34 reconstruction for this species is based on stem and foliage material from the Barremian/Aptian
35
36 locality of Négresse (Bernissart, Mons Basin, Belgium). In this work, two of these stems are
37
38 revised and analysed by CT-scan imaging, providing new information on their internal anatomy
39
40 and external morphology. The results show that *Weichselia reticulata* has a unique anatomy,
41
42 distinct from all other extant or fossil ferns. Stem external morphology suggests adaptations to
43
44 stressful environments, as it presents scale insertions, and a thick cortex. Especially noteworthy
45
46 is the presence of structures interpreted as aerophores or nectaries at the base of putative roots
47
48 and petioles that could suggest a need for extra ventilation of the frond or interactions with
49
50 animals. While these new results do not settle the question of the systematic affinities of
51
52 *Weichselia*, they provide new information about its autoecology and allow us to test the validity
53
54 of previous whole-plant reconstructions.
55
56
57
58
59
60

1
2
3 **Key words:** *Weichselia reticulata*, Bernissart, fern anatomy, Early Cretaceous, plant
4
5 reconstruction, autecology.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Today ferns represent the second largest group of vascular plants, with more than 300 genera and several thousand species (Bierhorst, 1971; PPG I, 2016). Particularly diversified, they have explored a wide range of morphologies and anatomies through time (Tidwell and Ash, 1994). After a first evolutionary radiation in the Paleozoic, the Mesozoic is a key period for the group that sees the rise of most modern families (e.g. Niklas et al., 1983; Lehtonen et al., 2017).

Weichselia reticulata (Stokes and Webb) Fontaine is a widespread Mesozoic fern, ranging in age from the Bathonian (Middle Jurassic) to the Cenomanian (Upper Cretaceous) and geographically from around 60°N to 60°S (Blanco-Moreno et al., 2018). It was first described from the English Weald as *Pecopteris reticulata* (Stokes and Webb, 1924), based on small charred fragments of vegetative fronds that are the most frequent type of remain of this fern (Fig. 1C). Unambiguous stem material is very scarce, and it is only known by a few specimens coming from the Cran du Nord locality (Bernissart colliery, Négresse mine shaft, Belgium). This material was first studied by Bommer (1910) but it is Alvin (1971) that provided its most detailed description.

Even though *W. reticulata* was widely distributed and is often very abundant in the localities where it is present, its systematic affinities remain unclear. It is often assigned to the Matoniaceae because of the organisation of the leaves (Bommer, 1910; Alvin, 1968; Alvin, 1971; Van Konijnenburg -Van Cittert, 1993; Sender, 2015). However, based on the work of Bommer (1910) and focussing on anatomical characters only, Edwards (1933) suggested a close relation with *Paradoxopteris* Hirmer from the Early and early-Late Cretaceous of North Africa (e.g. Hirmer, 1927; Edwards, 1933; Koeniguer, 1966; El Afty et al., 2019), which is close to Marattiaceae. A few authors also suggest a relation to this family based on anatomy and fertile structures (Zeiller, 1914; Silantieva and Krassilov, 2006).

Weichselia has been recovered from fresh water deposits such as lakes or rivers (Seward, 1900; Carpentier, 1927; Daber, 1968; Barthel and Boettcher, 1978; Barale, 1979 ;

1
2
3 Harris, 1981; Aguirrezabala et al., 1985; Ross and Cook, 1995; Batten, 1998; Diéguez and
4
5 Meléndez, 2000; Gomez et al., 2008), as well as from brackish habitats such as marshes, deltas,
6
7 estuaries or lagoons (Batten, 1974; El-Khayal, 1985; Román Gómez, 1986; Shinaq and Bandel,
8
9 1998; Barale and Ouaja, 2001; Lyon et al., 2001; Gomez et al., 2002; Barale and Azar, 2004;
10
11 Almendros et al., 2005; Diez et al., 2005; Silantieva and Krassilov, 2006). Due to the xeromorphic
12
13 traits observed in this genus (Alvin, 1974), some authors suggest that *Weichselia* inhabited dune
14
15 regions in marine areas (El-Khayal, 1985), while others believe this fern was adapted to extreme
16
17 drought conditions and attribute its presence in wet areas to transport from its natural habitat
18
19 to fluvial ecosystems via flooding (Alvin, 1974). The high abundance of burnt remains of
20
21 *Weichselia* in the fossil record has also led some authors to suggest it was an early successional
22
23 plant (Barral et al., 2016).
24
25
26

27
28 To date there have been five main reconstructions of the general habit of *Weichselia*
29
30 *reticulata* (Fig. 1A). Following the first reconstruction by Bommer (1910), Daber (1968) proposed
31
32 a new one based on pedate petiole heads found in Quedlinburg (Barremian, Germany) that were
33
34 interpreted as stems with a crown of radiating pinnate fronds. Alvin (1971) refuted this
35
36 reconstruction and proposed the most accepted one to date. He described an upright stem with
37
38 petioles that are spirally arranged and putative rooting organs produced in the opposite
39
40 direction of the petioles. Sterile and fertile pinnae are separate and consist of radially disposed
41
42 primary pinnae inserted in a pedate head. More recently, Sender et al. (2015) published a new
43
44 reconstruction of this species based on a fertile pedate head from the Albian of Teruel (Spain).
45
46 They represented a vegetative frond at the apical end of the stipe and a fertile frond at the end
47
48 of a lateral ramification of the stem. Finally, Poyato-Ariza and Buscalioni (2016) suggested
49
50 another reconstruction, based on material from Las Hoyas (upper Barremian, Cuenca, Spain). In
51
52 this case, the general appearance of the plant is that of an extant tree fern, with a wide, tall,
53
54 upright stem and a crown of fronds with petiole heads bearing radially disposed pinnae at the
55
56 top.
57
58
59
60

1
2
3 To provide new information on *Weichselia*, we undertake here a revision of the external
4 morphology and anatomy of the specimens studied by Alvin (1971) using μ CT-scan imaging and
5 we calculate parameters for the reconstruction of the plant's habit. A comparison with extant
6 and fossil ferns allows us to propose (1) new insights about the possible systematic affinities of
7 *Weichselia*, (2) additional clues about its autecology, and (3) guidelines for an accurate
8 reconstruction of its habit.
9

10 11 12 13 14 15 16 17 18 19 **GEOLOGICAL SETTING**

20
21 Bernissart is located 25 km west of Mons, in southwestern Belgium, in the northern part
22 of the Mons basin (Baele et al., 2012) (Fig. 2 A). This basin corresponds to an east–west subsiding
23 zone where Meso-Cenozoic sediments accumulated. Rather small, the subsiding area (40 by 15
24 km) is mainly controlled by intrastratal dissolution of deep evaporite beds in the Mississippian
25 (Carboniferous) basement. This dissolution process also led to the creation of several sinkholes,
26 or natural pits (Spagna et al., 2012; Quinif & Licour, 2012). The natural pits acted as traps for the
27 sediment, fauna, and flora present there at that time.
28
29

30
31 Three sinkholes have been recognized around Bernissart (Fig. 2B). They are the North,
32 the South, and the Iguanodon sinkholes. The latter yielded the famous Bernissart Iguanodons.
33 They were found in a Barremian/Aptian lacustrine clay, attributed to the Sainte-Barbe Clays
34 Formation (Cornet & Schmitz, 1898; Cornet, 1927). The age of this formation has been
35 determined based on palynologic data, based on sediments from the Iguanodon Sinkhole, as
36 ranging from the middle Barremian to the earliest Aptian (Dejax et al., 2007).
37
38

39
40 The North Sinkhole was encountered in 1906 during the digging of a connecting gallery
41 at a depth of 160 m. It started from the pit n°1 (called Négresse) in direction of the north. The
42 gallery entered the sinkhole on its southern part and continued north for 164 m. The Wealden
43 sediments are represented by dark grey to blue clays containing abundant charcoalfied plant
44 remains between 109 and 126 m from the south margin of the sinkhole (Cornet, 1927; Delmer
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and Van Wichelen, 1980). The exact position of the specimens is unknown but appear to
4
5 originate from lithological unit δ (see arrow on Figure 2C) that corresponds to a dark clay with
6
7 abundant charcoalfied and lignified plant remains.
8

9
10 No detailed sedimentological analysis has been performed but the occurrence of sandy
11
12 layers interbedding the clays as well as the occurrence of larger plant fragments could indicate
13
14 a slightly more riverine environment than for the Iguanodon pit of Bernissart that is interpreted
15
16 as lacustrine (Van den Broeck, 1898) or lacustrine to swampy (Yans, 2007; Schnyder et al., 2009;
17
18 Spagna et al., 2012).
19
20
21
22

23 MATERIAL AND METHODS

24 *Material*

25
26
27 The material studied consists of stems, petioles, and rachises preserved as lignite that
28
29 were assigned to *Weichselia reticulata*. They were collected in the Cran du Nord locality
30
31 (Bernissart colliery, Négresse mine shaft, Belgium) and are housed in the Royal Belgian Institute
32
33 of Natural Sciences (RBINS) in (Brussels, Belgium). In most cases, the specimens are only slightly
34
35 compressed. We follow Alvin (1971) in his distinction between the different organs (stems,
36
37 petioles, roots) and surface features (e.g. crater-like marks, tubercles) for these specimens.
38
39
40
41
42

43 Material examined: IRSNB b 0229, 8424, 8425, 8426, 8427; RBINS-PBOT-935, 936, 955, 976,
44
45 1073, 06846-0024.
46
47
48
49

50 Additional material used in the comparisons: *Paradoxopteris* material figured by Edwards
51
52 (1933); *Matonia* R.Br. Ex Wall thin-sections from the Cornell University Plant Anatomy
53
54 Collection; *Danaea* Sm. material figured by Rolleri (2004); *Angiopteris evecta* (Forst.) Hoffm.
55
56 material figured by Blomquist (1922); *Osmunda cinnamomea* L. thin-sections from the Cornell
57
58 University Plant Anatomy Collection; *Cyathea* Sm. material figured by White and Turner (2017);
59
60

1
2
3 *Platyserium aethyopicum* Hook. material figured by Allison (1913); *Psaronius* Cotta material
4
5 figured by Morgan (1959), and Rothwell and Blickle (1982).
6

7 Institutional abbreviations: MCCM-LH, Las Hoyas collection housed at the Museo de
8
9 Paleontología de Castilla-La Mancha (Cuenca, Spain); IRSNB b and RBINS-PBOT, Palaeobotany
10
11 collection of the Royal Belgian Institute of Natural Sciences (Brussels, Belgium); NHMUK PB V,
12
13 Palaeobotany collection of the Natural History Museum (London, UK).
14
15
16
17

18 *Methods*

19
20
21 *CT-Scan data acquisition and image processing.* For the anatomical study of *Weichselia*
22
23 *reticulata* two specimens of stems of this species collected from the locality of Négresse, IRSNB
24
25 b 8424 and 8425, and housed at the RBINS, were imaged by μ CT-scan at the RBINS (Brussels,
26
27 Belgium). Scanning was completed using a RX EasyTom 150, with copper filter. For IRSNB b 8424,
28
29 images were generated at a voltage of 145 kV and a current of 231 μ A, with a set frame rate of
30
31 12.5 and 8 average frames per image. This generated 1440 images and a voxel size of 0.0609096
32
33 μ m. For IRSNB b 8425, images were generated at a voltage of 139 kV and a current of 480 μ A,
34
35 with a set frame rate of 12.5 and 7 average frames per image. This generated 1440 images and
36
37 a voxel size of 0.0816001 μ m. Reconstructions, were performed using 3X-Act software from RX
38
39 Solutions. The images obtained were reduced with the “binning” plugin in ImageJ 1.49v
40
41 (Schneider et al., 2012). Segmentation, visualization and analysis were performed using Avizo
42
43 software on the reduced images: 3D models were produced from the μ CT images, and from
44
45 these, models of the vascular system via segmentation by contrast thresholds. The central
46
47 strands of the stem in both specimens, and the petiole in IRSNB b 8425, were isolated by further
48
49 segmentation by hand in order to more clearly observe the course of the central strands, and
50
51 better identify the organ insertions. Posteriorly, the models were uploaded in Meshlab (Cignoni
52
53 et al., 2008) and 3D-Slicer (Fedorov et al., 2012) where sections of the models were performed.
54
55
56
57
58
59
60

1
2
3 Despite the obvious advantages of μ CT-imaging in terms of non-destructive methods,
4 several limitations may exist and are related to the preservation of the material and the actual
5 contrast existing between different structures. The preservation of the material studied here
6 was not always ideal for scanning, and the differences in contrast of the different tissues were
7 sometimes very small, making it difficult to observe minute anatomical details (Fig. 3D).
8 Additionally, the chosen resolution was not sufficient to study the bundle morphology and
9 structure in detail (Fig. 3D-E). These details have been observed on existing historic thin sections
10 (Fig. 3A-C; Alvin, 1971).
11
12
13
14
15
16
17
18
19
20
21
22

23 *Variables for reconstruction and comparison with other ferns.* Stem diameter, bundle size and
24 cortex thickness were measured on the specimens mentioned in the material section, except
25 for *Angiopteris evecta* which was obtained from Blomquist (1922), and *Psaronius* from Rothwell
26 and Blicke (1982). The largest measurement was recorded for each category. General
27 information on the anatomy and morphology was observed on the specimens mentioned in the
28 materials section, and from the literature. Growth habit categories, leaf area, and leaf mass per
29 area (LMA) were obtained from Peppe et al. (2014) except for Leaf area of *Cyathea* (from Arens,
30 1997). The largest measurement was recorded for each category and data were consulted at the
31 generic (not specific) level.
32
33
34
35
36
37
38
39
40
41
42

43 Three variables with biomechanical implications and interest for the reconstruction of
44 the plant habit were also calculated using the measurements obtained in Table 1 and in Blanco-
45 Moreno et al., (2019):
46
47
48
49
50
51

52 1. The leaf mass per area (LMA) was calculated following Peppe et al. (2014):
53
54
55

$$56 \text{Log LMA frond} = 4.207 + 0.252 * \log (\text{Petiole width}^4 / \text{Leaf area})$$

57
58
59
60

1
2
3 Once the maximum area of foliage the stem can bear is obtained, the number of possible leaves
4
5 can be calculated based on the mean area of a single leaf (estimated in Blanco-Moreno et al.,
6
7 2019).

8
9 The work by Brouat et al. (1998) is based on spermatophytes, so we applied the method to other
10
11 fern taxa with enough information on the number of leaves per node (*Matonia*, *Cyathea* and
12
13 *Psaronius*) in order to test it.
14
15

16 17 18 **RESULTS**

19
20 The stem surface is always covered with millimetric tubercles and is sometimes
21
22 irregularly creased (Figs 4A-B, 5G-H, and 6A). Alvin (1971) observed 5 types of scars on the
23
24 Belgian material of *Négresse* identified as: petiole bases, radicular organs, sclerenchymatous
25
26 tubercles, crater like marks, and large elliptical scars (interpreted as aerophores). We here
27
28 redescribe these structures with the help of the new information provided by the μ CT-scan
29
30 imaging, including their size and anatomy.
31
32
33
34
35

36
37 *Petiole bases*. Material. IRSNB b 8424, 8425; RBINS-PBOT-935
38

39 There is no difference between the surface of the stem and the petiole, they are all
40
41 covered by the same creases and tubercles (Fig. 4B). The maximum width of the petiole ranges
42
43 from 1.57 cm in RBINS-PBOT-935 up to 4.7 cm in IRSNB b 8425. Two petioles were observed on
44
45 specimen IRSNB b 8424 measuring 2.4 and 2.3 cm in maximum width. Petioles depart at a 40°
46
47 to 50° angle from the stem. This angle increases distally due to the curvature of the stem (Fig.
48
49 4C). Additionally, all specimens have elliptical scars associated with the base of the petiole that
50
51 will be described in detail later.
52
53

54 Anatomically, petioles are formed by many meristeles from the stem's central rings. The
55
56 meristeles in the petiole are placed in rows near the insertion (Fig. 6C) and become concentric
57
58 distally, with up to of 10 concentric rings in IRSNB b 8425 (Fig. 4A).
59
60

1
2
3
4
5 *Radicular organs.* Material. IRSNB b 8424
6
7

8 The anatomy of specimen IRSNB b 8424 shows some strands that are orientated in the
9 direction opposite to the petioles. This could be interpreted as a possible radicular organ (Fig.
10 4D). The preserved width is 3.5 cm and it is inserted at a 50° angle. Similarly to the petioles, the
11 surface is ornamented and there is an elliptical scar associated to the base of the organ (Fig. 3B).
12 The anatomy is also very similar and formed by many meristeles from the stem's central rings,
13 with a semi-circular transverse section (Fig. 4D). The only difference between petioles and the
14 putative rooting organ in IRSNB b 8424 is the direction of the vascular strands, which is opposite
15 to each other.
16
17
18
19
20
21
22
23
24
25
26
27

28 *Tubercles.* Material. IRSNB b 8424, 8425, 8426, 8427, RBINS-PBOT-935, 936, 955, 976, 1073,
29 06846-0024.
30
31

32 Tubercles are present at the surface of most stems and petioles present in the collection
33 (Fig. 4A-B). They cover the whole length of petioles up to the petiole heads, where they are
34 especially found on the adaxial side. This ornamentation is also observed on the abaxial side of
35 the broader primary rachises. When present, they completely cover the surface of the plant.
36 They are clearly visible in specimens without creases, but are also present in very irregular,
37 creased ones. They are millimetric, but variable in size and shape. When imaged by the μ CT-scan
38 they present the same density as the rest of the surface.
39
40
41
42
43
44
45
46
47
48
49

50 *Crater like marks.* Material. IRSNB b 8426.
51

52 These structures have only been observed on one specimen already figured by Alvin (Pl.
53 II, fig. 2, 1971). The specimen is very deformed, and it is not possible to determine the part of
54 the plant it belongs to. The surface is covered by tubercles, which are disrupted by these marks
55 (Fig. 4A). They are slightly depressed in relation to the rest of the layer and are elliptical to more
56
57
58
59
60

1
2
3 or less circular. The surface is smoother, lighter in colour, and shows no identifiable structures.
4
5 They measure 0.19 by 0.17 to 0.34 by 0.15 cm.
6
7
8
9

10 *Large elliptical scars.* Material. IRSNB b 8424, 8425, 8427; RBINS-PBOT-935.
11

12 These structures have been observed on three stem fragments, and two disarticulated
13 scars are present in the collection (Figs 5A and G-H). Their outline is clearly defined, and they
14 are slightly depressed in relation to the rest of the stem, they measure 0.64 to 2.8 cm by 0.73 to
15 1.54 cm. Two differentiated zones can be observed: a central elliptical to circular rugose
16 protruding area measuring 0.49 to 1.49 cm by 0.862 to 0.35; and a peripheral area that
17 surrounds the central zone, which has more or less linear yet irregular marks that radiate from
18 the central zone towards the margin of the scar (Fig. 5A, and G-H). The internal face of the scars,
19 which can be observed in the disarticulated specimens, shows no signs of the two zones
20 previously described. The surface is flat but rugose and covered by tubercles, their appearance
21 is similar to the outer part of some of the stems but the tubercles are smaller (Fig. 5A). These
22 structures are not connected to the vascular system. Indeed, where the scars are present the
23 vascular tissue is cut off and there is an area with no vascular tissue at all just underneath these
24 structures (Fig. 5B-E, and I-J).
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 These scars are present at the base of possible petioles and rooting organs, laterally
42 oriented towards the abaxial side, only on one of the sides in specimens RBINS-PBOT-935 and
43 IRSNB b 8424 (Fig. 5B), and one on each side in the case of the best preserved specimen (IRSNB
44 b 8425, Fig. 5G-I). In specimens RBINS-PBOT-935 and IRSNB b 8425 a smaller scar is also present
45 directly above (distally) the other scar mentioned.
46
47
48
49
50
51
52
53
54

55 *Stem anatomy.* Material. IRSNB b 0229, 8424 and 8425.
56

57 The preserved diameter of the stem is 7.14 cm in IRSNB b 8425, and 3.64 cm in IRSNB b
58 8424. The stems are curved in both specimens, at a 115° angle in IRSNB b 8425 and at a 145°
59
60

1
2
3 angle in IRSNB b 8424. The curvature in IRSNB b 8425 occurs at the level where a petiole is borne,
4
5 and in IRSNB b 8424 where a possible root is borne (Fig. 4C-D and 6B). No branching has been
6
7 directly observed, however, at the apical-most part of specimen IRSNB b 8424 the disposition of
8
9 the vascular bundles could suggest dichotomous branching (Fig. 6D-G).

10
11
12 The internal anatomy of the plant consists of a highly dissected polycyclic dictyostele,
13
14 organized in 9 concentric rings of irregularly shaped vascular bundles, which are generally C
15
16 shaped, and measure 0.15 cm to 0.22 cm (Fig. 6). The density of vascular tissue is very high,
17
18 bundles are present from the centre to the cortex. Alvin (1971) mentioned the presence of
19
20 secretory canals in the stem of *Weichselia reticulata*, alternating with the bundles. These
21
22 structures were observed in the thin sections but are not clearly identified in the μ CT-scans due
23
24 to the lack of resolution, although specimen IRSNB b 8424 does show some very dense spots in
25
26 a similar position to the possible canals observed in the thin sections (Fig. 3D). The possible
27
28 secretory canals observed in the thin sections consist on circular structures measuring up to 0.05
29
30 cm, with a single layer of cells in the periphery with a similar preservation to the vascular tissue
31
32 of the plant, and a dark centre (Fig. 3C).

33
34
35
36 The disposition of petioles and putative roots in the specimens studied (Fig. 7) could be
37
38 helicoidal or alternate in IRSNB b 8424. Lateral organ production observed in IRSNB b 8425 is
39
40 less frequent, and there is only one petiole, so the phyllotaxy can not be inferred.
41
42
43
44

45 *Calculations for reconstruction*

46
47
48 In order to attempt a reconstruction of the whole plant, certain characters dealing with
49
50 the morphology, architecture, and anatomy of *Weichselia reticulata* have to be analysed (Table
51
52 1). Important variables for plant reconstruction can be then calculated using approximations
53
54 obtained from living plants: the leaf mass, and maximum height of the plant, and the number of
55
56 fronds per node (constrained by Corner's Rules).
57
58
59
60

1
2
3 *Leaf mass.* The LMA for *Weichselia* is 317.2g/m² to 352.5 g/m², calculated following Peppe et al.
4
5 (2014) based on measurements on the specimens published in Blanco-Moreno et al. (2019). This
6
7 value could be an overestimate, as fronds with a large relationship between petiole width and
8
9 leaf area (PW⁴/LA) have a lower LMA than expected by the model, and in the case of *Weichselia*
10
11 PW⁴/LA is larger than all other ferns in the sample from Peppe et al. (2014). However, it must
12
13 be noted that the architecture of the *Weichselia* frond, with a pedate head, is mechanically more
14
15 similar to palmate fronds than pinnate fronds (Niklas, 1991), and the results in Peppe et al.,
16
17 (2014) show that the largest LMA correspond to palmate or simple fronds.
18
19
20
21
22

23 *Height of the plant.* The results calculated from the model are within the maximum height
24
25 measurements in the literature (Table 2), bearing in mind they are first order of magnitude
26
27 approximations. This supports the validity of the model for the estimate of the height in
28
29 *Weichselia*, which would be up to 8.57 m according to the model.
30
31
32
33
34

35 *Phyllotaxy and number of leaves per node.* Estimation of the number of fronds in the crown
36
37 following Brouat et al. (1998) results in a possible number of leaves that is similar to the values
38
39 recorded for each taxon in the literature (Table 3). The number of leaves estimated for
40
41 *Weichselia* by this method is therefore accepted as a valid estimate until more information can
42
43 be obtained and would be of 1 to 2 leaves (Table 3).
44
45
46
47
48

49 **DISCUSSION**

51 *Morpho-anatomical comparison of Weichselia with other ferns*

52
53 In order to better understand the affinities, autecology, and habit of *Weichselia*
54
55 *reticulata*, the fossil material analysed here is compared with extant and fossil fern taxa selected
56
57 because of previously suggested systematic affinities (Edwards, 1933). It includes (1)
58
59 *Paradoxopteris* which was suggested by Edwards (1933) as the stem of *Weichselia reticulata*, (2)
60

1
2
3 Marattiales due to the affinities proposed mainly by Edwards (1933) and Silantieva and Krassilov
4
5 (2006), and (3) Matoniaceae which is the most accepted sister group to *Weichselia* (e.g. Alvin
6
7 1971; Sender et al., 2015). We also include two other genera: *Cyathea* (Cyatheaceae), for a
8
9 comparison with an arborescent fern with a tall upright stem as has been suggested for
10
11 *Weichselia* in some reconstructions (e.g. Poyato Ariza et al., 2016), and *Platyserium* Desv.
12
13 (Polypodiaceae), which was found to be the only fern with a stele organisation more or less
14
15 comparable to the here studied specimens.
16
17
18
19
20

21 *External morphology and scars*

22
23 The here observed ornamentation of the stems of *Weichselia* is very similar to what is seen in
24
25 the Cyatheaceae and some Marattiaceae. The stem surface of the Cyatheaceae is covered by a
26
27 hard and impervious sclerotic cortex. The gas exchanges are only made possible by the presence
28
29 of small lenticel like tubercles filled with spongy tissue that act as pneumatodes (Bower, 1923).
30
31 Another characteristic feature of this family is the presence of scales borne upon
32
33 sclerenchymatous outgrowths that cover leaves and stems while young. Although the scales
34
35 tend to fall in later ontogenic stages, the sclerenchymatous outgrowths tend to remain and are
36
37 responsible for the characteristic ornamentation of the stem (Fig. 8B). The tubercles observed
38
39 on *Weichselia* are devoid of spongy tissue and rather appear as solid structures. They are
40
41 nevertheless much smaller than the epidermal outgrowths of Cyatheaceae to which they do not
42
43 compare favourably. Some fossil and extant Marattiaceae also present similar surface features
44
45 to the ones observed in *Weichselia*. *Angiopteris* Hoffm. and *Danaea* present scales associated
46
47 to persistent structures at the base of the petioles that resemble the tubercles here described,
48
49 although they are more dispersed (Fig. 8A).
50
51
52
53

54 The crater-like marks are only present in one specimen and don't have a set morphology
55
56 or size. They were probably produced by an external agent that damaged the plant's outermost
57
58 layer.
59
60

1
2
3 In contrast, the elliptical scars have a constant morphology and a specific position in the
4 stem. Similar structures have however been described in other fern taxa. Their organisation is
5 like that of some aquatic and wetland plant roots with radial lysigeny and wheel-shaped type
6 aerenchyma (Jung et al. 2008). The lack of vascular tissue in the area however led us to discard
7 this possibility. Alvin (1971) discussed these scars and proposed that they could represent
8 aerophores. In ferns, there are two types of aerating structures: linear aerophores, sometimes
9 termed pneumathodes, and nonlinear aerophores, sometimes termed pneumatophores or
10 lenticels in the Marattiaceae (Davies, 1991). In respect to this, the elliptical scars on *Weichselia*
11 would rather be described as lenticels or nonlinear aerophores, which are also present in the
12 Cyatheaceae (Bower, 1923) and the Marattiaceae (Smith et al., 2006). Lenticels are generally
13 very simple in ferns (Fig. 8C-D) with some exceptions such as in *Cyathea horrida* (L.) Sm. (White
14 and Turner, 2012). In this species lenticels are more complicated and are similar to the ones
15 observed in *Weichselia*. Additionally, some lenticels found on spermatophytes, with filling tissue
16 and closing layers (Evert, 2006) very much resemble the structure observed in *Weichselia*. The
17 existence of gas exchanging structures at the base of the petioles is frequently observed in ferns
18 e.g. Marattiaceae, Plagiogyraceae, Cyatheaceae or Thelypteridaceae. In all these cases, as
19 observed in *Weichselia*, the hypodermis is not interrupted (Davies, 1991). Pneumathodes in
20 Cyatheaceae and *Saccoloma* Spreng. (Dennstaedtiaceae) are related to involutions (i.e., curving
21 of the vascular system towards the centre of the axis) in the leaf traces (Bower, 1923 fig. 161,
22 162; Bower 1926 fig. 560). By contrast, in *Weichselia* they only create gaps in the vascular system
23 directly below them, and they don't seem to be related with involutions in the petiole or the
24 putative roots (Fig. 5E and G). Another organ that can be found at the base of fern pinnae that
25 somewhat resembles the elliptical scars observed in *Weichselia* are nectary glands, which are
26 for example present in the Cyatheaceae, in *Angiopteris*, and in *Platycterium* (White and Turner,
27 2012). In such structures, the vascular tissues do not enter the gland (White and Turner, 2012),
28 this corresponds to what is observed in *Weichselia*. Generally, nectary glands are found at the
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 base of second or third order pinnae and pinnules and are recurrent in the different orders of
4
5 ramification of the frond (White and Turner, 2012). No such disposition has been observed in
6
7 *Weichselia*. Figure 8B shows a possible nectary or large protruding aerophore at the base of the
8
9 petiole in *Sphaeropteris intermedia* (Mett.) R.M. Tryon, in a very similar position to the ones
10
11 observed in *Weichselia*.
12

13
14 Large radicular organs are not very frequent in ferns, although the Marattiaceae have
15
16 large roots (Bower, 1923). The anatomy of the putative root observed in *Weichselia reticulata*
17
18 has not been observed in any other fern. Other structures with negative gravitropism that could
19
20 be similar to those observed in *Weichselia* are lateral ramifications described in the Cyatheales
21
22 that grow into stolons that stabilize the tall trunks and permit the vegetative reproduction of
23
24 the plant (Hallé 1965, 1966). These buds are situated under the leaves in the Cyatheales, the
25
26 same position of the putative root in *Weichselia*. Additionally, the scarce record of fertile
27
28 remains of *Weichselia reticulata* could be related to a dominant vegetative reproduction of the
29
30 plant.
31
32
33
34
35

36
37 *Anatomy.* The anatomy of *Weichselia reticulata* is particularly divergent and appears to be
38
39 different to all ferns that have been published so far. The results of the comparison of the
40
41 Belgian material with a selection of fern taxa are summarised in Table 1, and Figures 9 and 10.
42
43 Firstly, the type of stele apart (which is a polycyclic dictyostele), it is clear that the anatomy of
44
45 *Weichselia reticulata* is very different from that of *Paradoxopteris*. The main differences reside
46
47 in the size of the vascular bundles and the presence of auxiliary strands with round morphology
48
49 that are alternate with the C-shaped meristeles in *Paradoxopteris* (Fig. 9G). Koeniguer (1966)
50
51 separates *Paradoxopteris stromeri* Hirmer into two varieties. One with large vascular bundles
52
53 which are more than 2.5 times larger than in *Weichselia*. A second with smaller bundles that
54
55 have a mean size of 0.22 to 0.28 cm, which fits within the size range of the bundles in *Weichselia*.
56
57 However, although not clearly visible in the μ CT-scan images of *Weichselia*, vascular strands are
58
59
60

1
2
3 more or less C-shaped in all rings observed in the thin sections. There is thus no evidence of the
4
5 auxiliary strands typical of *Paradoxopteris* (Fig. 3A).
6

7
8 While *Matonia* has a curved rhizome (Fig. 8E) that could be compared to the curved axes
9
10 of specimens IRSNB b 8424 and 8425, it can be distinguished easily in terms of anatomy by its
11
12 polycyclic solenostele. *Phanerosorus*, the second genus of the Matoniaceae, also presents a
13
14 polycyclic solenostele (Kramer, 1990).
15

16
17 Marattiaceae in general is one of the groups that most resembles *Weichselia*. They share
18
19 a similar stele organisation with small bundles and many strands arising from the central ones
20
21 going to the petioles and putative roots (Fig. 9). The extinct genus *Psaronius*, related to the
22
23 Marattiaceae, has a similar anatomy to *Weichselia* (Fig. 10). The polycyclic dictyostele is also
24
25 very fragmented, and is spirally arranged, presenting up to 6 cycles. However, the bundles are
26
27 much larger, and the external part of the stem of *Psaronius* is surrounded by a thick layer of
28
29 adventitious roots. The secretory canals observed in *Weichselia* have a clear epithelium and are
30
31 similar to the mucilage canals found in the petioles of *Angiopteris evecta* (West, 1915).
32
33

34
35 *Osmunda* L. and *Cyathea* are not polycyclic, and strands are larger than in *Weichselia* for
36
37 both taxa (Figs 9B and F). Finally, the general appearance of the anatomy of *Platycterium* is similar
38
39 to *Weichselia*, with a very fragmented polycyclic dictyostele and many strands forming the
40
41 petiole (Fig. 9C). However, *Platycterium* roots are not produced by the central strands and the
42
43 bundles are smaller than in *Weichselia*. It thus appears that *Weichselia* has a unique stellar
44
45 anatomy among known fossil and extant ferns.
46
47

48 49 50 *Systematic affinities*

51
52 The systematic position of *Weichselia* has been discussed since its discovery. It is quite
53
54 clear from the results herein obtained that it presents a unique combination of characters that
55
56 could suggest a relation with groups as distinct as the Marattiales or the Matoniaceae.
57
58
59
60

1
2
3 On the one hand, they present characters that are considered exclusive to the
4
5 Marattiales such as the roots originating from the central bundles, the possible secretory canals,
6
7 the sclerenchymatous tubercles on the surface and the apparent absence of endodermis (it has
8
9 not been observed in this work and was not identified by Alvin (1971)). However, Marattiales
10
11 do not generally share the same stem and frond organisation as *Weichselia*. Stems are generally
12
13 massive in the Marattiales whereas in *Weichselia* they are more elongated, and Marattiales do
14
15 not present pedate fronds.
16
17

18
19 On the other hand, the similarities with the Matoniaceae are mostly architectural. They
20
21 share similar curved stems, dichotomous branching and frond organisations (Blanco-Moreno et
22
23 al., 2019). Their anatomy however is rather different, *Weichselia* presenting a polycyclic stele
24
25 while Matoniaceae have solenostelic steles. However, Allison (1913) studying *Platycerium* has
26
27 argued that its polycyclic dictyostele could derive from the fragmentation of a polycyclic
28
29 solenostele such as the one present in *Matonia*. Moreover, their size is not comparable, as the
30
31 stems of *Matonia* have a mean thickness of 0.6 cm (Kato, 1993) and pinnae measure around 35
32
33 cm, whereas the rhizome width of *Weichselia* is around 7 cm and pinnae can measure more than
34
35 100 cm.
36
37

38
39 The very peculiar organisation of *Weichselia* renders comparison with other extant or
40
41 fossil ferns difficult. The here obtained results unfortunately do not allow for a definite
42
43 systematic assignation. Even if, at the end of this comparison, a relation with the Marattiaceae
44
45 and Psaroniaceae seem to be possible, we believe that the inclusion of *Weichselia* within the
46
47 Marattiales would be very doubtful considering the many specimens of this plant that have been
48
49 studied and published so far. A comprehensive analysis of all organs must be performed.
50
51

52 53 54 *Autecology*

55
56 The external morphology and the LMA provide some insights to the possible autecology
57
58 of *Weichselia reticulata*. Firstly, the presence of large elliptical scars at the base of the petioles,
59
60

1
2
3 if interpreted as aerophores, indicates that the stem and petiole was in need of extra ventilation.
4
5 This could be seen as an indicator of a partially submerged plant. However, most ferns have
6
7 aerophores (Davies, 1991). Large aerophores at the base of the petioles or stems in fact facilitate
8
9 gaseous exchange when the epidermis is protected by a hard sclerotic layer (e.g. Cyatheaceae)
10
11 and/or when the petioles of the juvenile fronds are covered by mucilage while they unfurl (e.g.
12
13 Palgiogyraceae). In this case, when the fronds are mature, leaf stomata take over the gas
14
15 exchanges and the aerophores cease to be functional (Davies, 1991). The morphology of the
16
17 elliptical scars of *Weichselia*, similar to lenticels with closing layers, suggests that it might have
18
19 had the function of a temporary aerophore. The presence of possible mucilage canals and the
20
21 large size of the fronds further justify for the need of extra ventilation while the frond unfurled.
22
23 The tubercles of the outer, probably sclerotic, layer of the stems, petioles and primary rachises
24
25 of *Weichselia* are similar to those covering the Cyatheaceae. They indicate that the plant might
26
27 have been covered by scales or hairs. This sclerotic outer layer might also be the cause of the
28
29 presence of aerophores.
30
31
32
33

34 If, on the other hand, the elliptical scars are interpreted as nectaries, an association with
35
36 insects in order to avoid herbivory can be suggested. Nectaries in ferns seem to indicate an
37
38 association with ants, which protect the fronds against herbivores (Koptur et al., 1982; Arens
39
40 and Smith, 1998; Koptur et al., 2013). The fact that ants most probably appeared in the Albian,
41
42 ca 110 mya (Grimaldi and Agosti, 2000), would rule out the association with these animals, at
43
44 least in the early records of *Weichselia*. However, others such as spiders, parasitoids, predatory
45
46 wasps, beetles, mirids or mites also feed on extrafloral nectar (Heil, 2015). Insects from
47
48 Bernissart have not been thoroughly studied, although they are present (Godefroit, 2012).
49
50 Terrestrial Hemiptera, beetles, wasps, and dipterans were present in the Upper Barremian
51
52 locality of Las Hoyas (Delclòs and Soriano, 2016), where *Weichselia* is the most common fern
53
54 remain.
55
56
57
58
59
60

1
2
3 Another explanation for the existence of fern nectaries has been suggested by Koptur
4
5 et al. (1982), related to the elimination of metabolic by-products without an excessive loss of
6
7 water in xerophytes. *Weichselia* has previously been interpreted as a xerophyte (e.g. Alvin, 1974;
8
9 Watson and Alvin, 1996).

10
11
12 High LMA is found in plants under high irradiance (Poorter et al., 2009), and the position
13
14 of the fronds in *Weichselia* suggests the sun would have fallen directly upon the leaves (Blanco-
15
16 Moreno et al., 2019). Fern fronds live around 2 to 3 years at most (literature cited by Karst and
17
18 Lechowicz, 2007), and a high LMA generally indicates the fronds were especially long-lived
19
20 (Tanner, 1983; Karst and Lechowicz, 2007). This suggests *Weichselia* fronds were not shed
21
22 regularly and grew at a lower relative rate (Poorter et al., 2009). This strategy is not very
23
24 concordant with the interpretation of *Weichselia* as an early successional plant (Barral et al.,
25
26 2016), but more so to a plant tolerant to fire (Watson and Alvin, 1996). *Weichselia* was most
27
28 probably a re-sprouter like *Pteridium* Gleditsch as suggested by Scott et al. (2000), and some
29
30 tree-ferns that are fire-tolerant and re-sprout from buds in their fibrous trunks after a fire (Ough
31
32 and Murphy, 2004).

33
34
35
36
37 The environmental conditions of the habitat of *Weichselia reticulata* have been much
38
39 discussed. Some authors suggest this fern was adapted to extreme drought conditions and
40
41 attribute its presence in aquatic environments to transport from its natural habitat to fluvial
42
43 ecosystems via flooding (Alvin, 1974). Others suggest these adaptations are related to high
44
45 salinity habitats, and that *Weichselia* inhabited dune regions in marine areas (El-Khayal, 1985),
46
47 or even that it was trophophyte in coastal wet ground that dried periodically (Daber, 1968). This
48
49 study does not provide keys to the environment where *Weichselia* grew, however, it must be
50
51 noted that some of the traits traditionally interpreted as xeromorphic might not be so. For
52
53 example, the thick cuticles in ferns do not correlate with low humidity, instead they seem to be
54
55 associated to cold and nutrient-poor environments (Kessler et al., 2007). Also, the presence of
56
57
58
59
60

1
2
3 scales or hairs discussed in this paper, which were probably covering the stem and rachis of
4
5 *Weichselia*, are related to protection from high insolation and not to aridity (Kessler et al., 2007).
6
7

8 9 10 *Guidelines for reconstruction*

11
12 We consider that the data herein obtained on the morphology, anatomy, and
13
14 architecture of *Weichselia reticulata* does not allow for a complete reconstruction of the habit
15
16 of the whole plant with enough confidence. However, it does shed new light on existing
17
18 reconstructions and allows us to discuss which aspects are accurate or not.
19

20
21 Previous reconstructions of *W. reticulata* (Bommer, 1911; Daber, 1968; Alvin, 1971;
22
23 Sender et al., 2015; Poyato-Ariza & Buscalioni, 2016; Fig. 1A) are mainly based on the external
24
25 morphology and on some thin sections of very fragmentary material. The CT-scan models herein
26
27 analysed provide additional information on the anatomy and allow for a more accurate
28
29 description of the stems. Although the illustrations provided in the previous reconstructions are
30
31 artistic, if both these representations and the comments provided by the authors are analysed
32
33 together, they can be discussed in the light of the new information obtained in this study.
34
35

36
37 The general habit of the plant, represented by the stem and organ insertions, differs
38
39 from one reconstruction to another. The stem is however always represented straight, upright
40
41 and aerial. The analyses here performed nevertheless do not provide direct evidence for the
42
43 stem being upright. Both specimens studied by μ CT-scan are incomplete and one side of the
44
45 stem is in each case missing. However, the results allow us to conclude that: (1) It is not possible
46
47 to know if the stem is dorsiventral or radial. (2) The presence of the elliptical scars, either if they
48
49 are aerophores or nectaries, in the basalmost part of the petioles suggests that the stem was
50
51 not buried. (3) The fact that both specimens show a curvature of the stem argues against an
52
53 upright or tall stem.
54
55

56
57 There are however arguments supporting the reconstruction of *Weichselia* with an
58
59 upright stem. These are: (1) the high density of the vascular system similar to the vascular system
60

1
2
3 in monocots such as *Dracaena marginata* Aiton (Fig. 11; Haushahn et al., 2014; Hesse et al.,
4
5 2016) and (2) the orientation and origin of the vascular system in the acroscopic part of the
6
7 petiole of IRSNB b 8425. Indeed, the analogy with *Dracaena marginata* can be further applied
8
9 to the organisation of the petiole vascularisation that does not arise directly from the central
10
11 strands and is perpendicular to the stem (Fig. 6B, 7B, and 11A). In this plant, the perpendicular
12
13 vascular strands act as “tensile cables” that transfer bending loads (Fig. 11B; Haushahn et al.,
14
15 2014). If the large fronds of *Weichselia reticulata* are inserted in a plagiotropic position, as it has
16
17 been previously suggested (Blanco-Moreno et al., 2019), on an upright stem the vasculature of
18
19 the petiole bases could similarly act as “tensile cables”. Fronds were large and most probably
20
21 heavy (according to high LMA) and would need a good mechanical support at the petiole
22
23 insertion. The asymmetric disposition of the “tensile cable-like” vascularization in the petiole
24
25 could support an upright position of the stem. Nevertheless, even if the stems were upright, the
26
27 plant would not have reached the height of the highest arborescent ferns (Recent tree ferns and
28
29 *Psaronius*) according to the calculations following Niklas’ formula (1994) (see results section).
30
31
32
33

34
35 Roots are present in all reconstructions, but their organisation does not fit what is seen
36
37 on the specimens. Root insertions are always too small. They are inserted at very large angles in
38
39 comparison to the possible rooting organ observed in specimen Pbot-00000-0127. Bommer
40
41 (1910) and Sender et al. (2015) draw them in a different position than observed in this study.
42

43
44 Petioles, although inserted at an excessively large angle, show a correct proportion in
45
46 Bommer (1910). This is also the case for Alvin (1971). They are too small in relation to the stem
47
48 width in Poyato-Ariza and Buscalioni (2016). Sender et al. (2015) suggest in their reconstruction
49
50 that the fronds occur at the end of branches. Set aside that a possible branching of the stem has
51
52 been described, petioles have definitely been observed, the pedate heads are thus not borne
53
54 directly as the termination of branches. Finally, the *Cycas* like reconstruction of Daber (1968)
55
56 does not fit any of the here discussed results and is considered as the less accurate of all
57
58 proposed reconstructions.
59
60

1
2
3 The exact disposition of the plant fronds could not be ascertained from the studied
4 specimens. The fronds are not opposite and the anatomy shows that there could be two petioles
5 close to each other, possibly three if the stem is interpreted as dorsiventral and the putative
6 root as a petiole. It is however impossible to know whether they are disposed in a half helix
7 arrangement (more plausible if the stem is dorsiventral) or in a complete helix. The calculations
8 performed following Brouat et al. (1998) suggest there could be up to two fronds per node,
9 possibly ruling out the complete helix disposition. On the other hand, the position of the
10 putative radicular organ, alternating with the petioles, and apparently associated with the
11 younger petiole in specimen IRSNB b 8424, is similar to the root disposition in upright stems
12 with helicoidally disposed leaves of the Marattiaceae (Bower, 1926), especially of the genus
13 *Danaea*.

30 CONCLUSIONS

31
32 The revision of the material from Négresse that provided the first whole plant
33 reconstruction of the emblematic, yet mysterious fern *Weichselia reticulata*, has yielded
34 interesting results. No evidence has been found to refute the assignation of the material to
35 *Weichselia reticulata*. Although anatomically distinct to all consulted fossil and extant ferns, this
36 study suggests some affinities with the Marattiales. This link is further supported by the surface
37 morphology of the stem. Similarities with the Cyatheales are interpreted as evidencing
38 similarities in their autecology. Further analysis including thin sections of this and other material
39 are necessary in order to study the secretory canals, and corroborate the absence of
40 endodermis, two characters that would more clearly link *Weichselia reticulata* to the
41 Marattiales.

42
43 A whole plant reconstruction is not possible with the material at hand, but a more
44 precise description of the external “scars” and stem anatomy has been obtained. Some of the
45 inferences by Alvin (1971) have been corroborated, such as the presence of aerophores (or
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

nectaries) and “rooting organs”. Additionally, the variables calculated for the reconstruction have provided some insights on the possible habit of *Weichselia*, which will be very valuable for future reconstructions based on more data and material. The new data obtained from the stems and fronds also allows for further interpretations of the autecology of this cosmopolitan fern.

Acknowledgements. We would like to acknowledge the team managing the μ CT-scan at the RBINS (Brussels, Belgium), especially Ulysse Lefevre for scanning and reconstructing the Y-Slices, and all his valuable help with all technical matters in this paper. We also acknowledge the Institut des Sciences de l'Evolution (Montpellier, France) and the labex CEMEB (ANR-10-LABX-0004) for the assistance in the use of the Avizo Software, and for allowing us to work on their installations. C.B.M. is supported by a FPI-UAM 2015 grant from the Universidad Autónoma de Madrid, and benefitted from a grant from the same organisation for the work in Montpellier. Collections in the Royal Belgian Institute of Natural Sciences, Brussels, were studied with the support of a grant from the SYNTHESYS Project (<http://www.synthesys.info/>), financed by the European Community Research Infrastructure Action under the FP7 Integrating Activities Program. Cyrille Prestianni is funded by the BRAIN project BR/143/A3/COLDCase in the framework of which this research has been done. Anne-Laure Decombeix is funded by AMAP (Botany and Computational Plant Architecture), a joint research unit which associates CIRAD (UMR51), CNRS (UMR5120), INRAE (UMR931), IRD (R123), and Montpellier University (UM).

DATA ARCHIVING STATEMENT

Data for this study are available in the Virtual Collections of the RBINS:

Specimen IRSNB b 8424: <http://virtualcollections.naturalsciences.be/virtual-collections/paleontology/paleobotany/weichselia-sp>

Specimen IRSNB b 8425: <http://virtualcollections.naturalsciences.be/virtual-collections/paleontology/paleobotany/weichselia>

1
2
3 The slides of the CT-scan must be requested to the curator of the RBINS Collections of
4 Paleontology, from the RBINS Scientific Survey of Heritage, currently Annelise Folie
5 (annelise.folie@naturalsciences.be).
6
7
8

9
10 [please note that the data for this paper are not yet published and these temporary links should
11 not be shared without the express permission of the author]
12
13
14
15

16 REFERENCES

17
18
19
20
21 AGUIRREZABALA, L. M., TORRES, J. A., and VIERA L. I. 1985. El Weald de Igea (Cameros-La
22 Rioja). Sedimentología, bioestratigrafía y paleoicnología de grandes reptiles (dinosaurios).
23 *Munibe Ciencias Naturales. Natur zientziak*, **37**, 111-138.
24
25
26

27
28
29
30 ALLISON, H. E. 1913. On the vascular anatomy of the rhizome of *Platynerium*. *New Phytologist*,
31 **12**, 311-321.
32
33
34

35
36 ALMENDROS, G., ZANCADA, M. C., GONZÁLEZ-VILA, F. J., LESIAK, M. A., and ÁLVAREZ-RAMIS, C.
37 2005. Molecular features of fossil organic matter in remains of the Lower Cretaceous fern
38 *Weichselia reticulata* from Przenosza basement (Poland). *Organic geochemistry*, **36**, 1108-
39 1115.
40
41
42
43
44
45

46
47 ALVIN, K. L. 1968. The spore-bearing organs of the Cretaceous fern *Weichselia* Stiehler. *Journal*
48 *of the Linnean Society of London, Botany*, **61**, 87-92.
49
50
51

52
53
54 --- 1971. *Weichselia reticulata* (Stokes et Webb) Fontaine from the Wealden of Belgium.
55 *Mémoire de l'Institut Royal des Sciences Naturelles de Belgique*, **166**, 1-33.
56
57
58
59
60

1
2
3 --- 1974. Leaf anatomy of *Weichselia* based on fusainized material. *Palaeontology*, **17**, 587-598.
4
5

6
7 ARENS, N. C. 1997. Responses of leaf anatomy to light environment in the tree fern *Cyathea*
8 *caracasana* (Cyatheaceae) and its application to some ancient seed ferns. *Palaios*, **12**, 84-94.
9
10
11

12
13 ARENS, N. C. and SMITH, A. R. 1998. *Cyathea planadae*, a remarkable new creeping tree fern
14 from Colombia, South America. *American Fern Journal*, **88**, 49-59.
15
16
17

18
19 BARALE, G. 1979. Découverte de *Weichselia reticulata* (Stokes & Webb) Fontaine emend. Alvin,
20 filicinée leptosporangiée, dans le Crétacé Inférieur de la province de Lérida (Espagne):
21 implications stratigraphiques et paléoécologiques. *Geobios*, **12**, 313-319.
22
23
24
25

26
27 BARALE, G. and AZAR, D. 2004. Cuticules végétales dans le Crétacé inférieur du Sud du Liban.
28 *Comptes Rendus Palevol*, **3**, 119–124.
29
30
31

32
33 BARALE, G. and OUAJA, M. 2001. Découverte de nouvelles flores avec des restes à affinités
34 angiospermiennes dans le Crétacé inférieur du Sud tunisien. *Cretaceous Research*, **22**, 131–
35 143.
36
37
38
39

40
41 BARRAL, A., GOMEZ, B., ZORRILLA, J.M., SERRANO, J.M., YANS, J., CAZEDEBAT, M., DAVIERO-
42 GOMEZ, V., EWIN, T.A.M. and LECUYER, C. 2016. Local-scale analysis of plant community from
43 the Early Cretaceous riparian ecosystem of Hautrage, Belgium. *Palaeogeography*,
44 *Palaeoclimatology, Palaeoecology*, **443**, 107-122.
45
46
47
48

49
50 BARTHEL, K. W. and BOETTCHER, R. 1978. Abu Ballas Formation (Tithonian/Berriasian;
51 Southwestern Desert, Egypt) a significant lithostratigraphic unit of the former "Nubian Series".
52
53
54
55
56
57
58
59
60

1
2
3 *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Histor. Geologie*, **18**, 153-
4
5 166.

6
7
8
9
10 BATTEN, D. J. 1974. Wealden Palaeoecology from the Distribution of Plant Fossils. *Proceedings*
11
12 *of the Geologists' Association*, **85**, 433-458.

13
14
15
16 --- 1998. Palaeoenvironmental implications of plant, insect and other organic-walled
17
18 microfossils in the Weald Clay Formation (Lower Cretaceous) of southeast England. *Cretaceous*
19
20 *Research*, **19**, 279-315.

21
22
23
24
25 BLANCO-MORENO, C., GOMEZ, B. and BUSCALIONI, A.D. 2018. Palaeobiogeographic and
26
27 metric analysis of the Mesozoic fern *Weichselia*. *Geobios*, **51**, 571-578.

28
29
30
31
32 BLANCO-MORENO, C., GOMEZ, B., MARUGÁN-LOBÓN, J., DAVIERO-GOMEZ, V., and
33
34 BUSCALIONI, A. D. 2019. A novel approach for the metric analysis of fern fronds: Growth and
35
36 architecture of the Mesozoic fern *Weichselia reticulata* in the light of modern ferns. *Plos One*,
37
38 **14**, p. e0219192.

39
40
41
42
43 BLOMQUIST, H.L. 1922. Vascular anatomy of *Angiopteris evecta*. *Botanical Gazette*, **73**, 181-
44
45 199.

46
47
48
49
50 BOMMER, C. 1910. Contribution à l'étude du genre *Weichselia*. *Bulletin de la Société Royale de*
51
52 *Botanique de Belgique*, **3**, 296-304.

53
54
55
56 BOWER, F.O. 1923. *The ferns (Filicales). Analytical examination of the criteria of comparison,*
57
58 *vol. I.* Cambridge University Press, Cambridge. 359 pp.

1
2
3
4
5 --- 1926. *The ferns (Filicales). The Eusporangiatae and other relatively primitive ferns, vol. II.*
6
7 Cambridge University Press, Cambridge. 344 pp.

8
9
10
11 BROUAT, C., GIBERNAU, M., AMSELLEM, L. and MCKEY, D. 1998. Corner's rules revisited:
12
13 ontogenetic and interspecific patterns in leaf–stem allometry. *The New Phytologist*, **139**, 459-
14
15 470.

16
17
18
19
20
21 CARPENTIER, A. 1929. Recherches sur les Végétaux fossils des argiles éocènes du Pays de
22
23 Bray. *Bulletin de la Société géologique de France*, **29**, 89-96.

24
25
26
27 CIGNONI, P., CALLIERI, M., CORSINI, M., DELLEPIANE, M., GANOVELLI, F. and RANZUGLIA, G.
28
29 2008. Meshlab: an open-source mesh processing tool. In SCARANO, V., DE CHIARA, R. and
30
31 ERRA, U. (eds). *Eurographics Italian chapter conference*. The Eurographics Association, **129-**
32
33 **136.**

34
35
36
37
38
39 CORNET, J. 1927. L'époque wealdienne dans le Hainaut. *Annales de la Société Géologique de*
40
41 *Belgique*, **Tome L**, 161–164.

42
43
44
45 DABER, R. 1968. A *Weichselia*—*Stiehleria*—Matoniaceae community within the Quedlinburg
46
47 Estuary of Lower Cretaceous age. *Journal of the Linnean Society London. Botany*, **61**, 75–85.

48
49
50
51
52 DAVIES, K. L. 1991. A brief comparative survey of aerophore structure within the Filicopsida.
53
54 *Botanical Journal of the Linnean Society*, **107**, 115-137.

1
2
3 DELCLÒS, X. and SORIANO, C. 2016. Insecta. In POYATO-ARIZA, F. J. and BUSCALIONI, A. (eds).
4
5 *Las Hoyas: a Cretaceous Wetland. A multidisciplinary synthesis after 25 years of research on an*
6
7 *exceptional fossil Lagerstätte from Spain*. Verlag Dr. Friedrich Pfeil, **70-88**, 262pp.

9
10
11
12 DELMER, A. and VAN WICHELEN, P. 1980. Répertoire des puits naturels connus en terrain
13
14 houiller du Hainaut. *Professional Papers*, **5**, 1–234.

15
16
17
18 DIÉGUEZ, M. C. and MELÉNDEZ N. 2000. Early Cretaceous ferns from lacustrine limestones at
19
20 Las Hoyas, Cuenca province, Spain. *Palaeontology*, **43**, 1113-1141.

21
22
23
24
25 DIEZ, J., SENDER, L., VILLANUEVA-AMADOZ, U., FERRER, J., and RUBIO, C. 2005. New data
26
27 regarding *Weichselia reticulata*: soral clusters and the spore developmental process. *Review of*
28
29 *Palaeobotany and Palynology*, **135**, 99–107.

30
31
32
33
34 DURAND, L.Z. and GOLDSTEIN, G. 2001. Growth, leaf characteristics, and spore production in
35
36 native and invasive tree ferns in Hawaii. *American Fern Journal*, **91**, 25-36.

37
38
39
40 EDWARDS, W. N. 1933. On the Cretaceous fern *Paradoxopteris* and its connexion with
41
42 *Weichselia*. *Annals of Botany*, **47**, 317-341.

43
44
45
46
47 EL ATFY, H., JASPER, A., and UHL, D. 2020. A new record of *Paradoxopteris stromeri* Hirmer
48
49 1927 (Monilophyta, incertae sedis) from the Cenomanian of Sinai, Egypt. *Review of*
50
51 *Palaeobotany and Palynology*, **273**, 104148.

52
53
54
55
56 EL-KHAYAL, A. 1985. Occurrence of a characteristic Wealden fern (*Weichselia reticulata*) in the
57
58 Wasia Formation, central Saudi Arabia. *Scripta Geologica*, **79**, 75-88.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

EVERT, R. F. 2006. *Esau's plant anatomy: meristems, cells, and tissues of the plant body: their structure, function, and development*. John Wiley & Sons, New Jersey. 600 pp.

FEDOROV, A., BEICHEL, R., KALPATHY-CRAMER, J., FINET, J., FILLION-ROBIN, J. C., PUJOL, S.,
BAUER, C., JENNINGS, D., FENNESSY, F., SONKA, M. and BUATTI, J. 2012. 3D Slicer as an image
computing platform for the Quantitative Imaging Network. *Magnetic resonance imaging*, **30**,
1323-1341.

GODEFROIT, P. 2012. *Bernissart dinosaurs and Early Cretaceous terrestrial ecosystems*. Indiana
University Press, Bloomington. 629 pp.

GOMEZ, B., THÉVENARD, F., FANTIN, M. and GIUSBERTI, L. 2002. Late Cretaceous plants from
the Bonarelli level of the Venetian Alps, northeastern Italy. *Cretaceous Research*, **23**, 671–685.

GOMEZ, B., GILLOT, T., DAVIERO-GOMEZ, V., SPAGNA, P. and YANS, J. 2008. Paleoflora from
the Wealden facies strata of Belgium: Mega- and meso-fossils of Hautrage (Mons Basin).
Memoirs of the Geological Survey, **55**, 53-60.

GRIMALDI, D. and AGOSTI, D. 2000. A formicine in New Jersey Cretaceous amber
(Hymenoptera: Formicidae) and early evolution of the ants. *Proceedings of the National
Academy of Sciences*, **97**, 13678–13683.

HALLE, F. 1965. Les stolons de la fougère arborescente *Cyathea manniana* Hooker. *Comptes
Rendus de l'Académie des Sciences*, **261**, 2935-2938.

1
2
3 --- 1966. Étude de la ramification du tronc chez quelques fougères arborescentes. *Adansonia*,
4
5 **6**, 405-424.
6
7

8
9
10 HARRIS, T. M. 1981. Burnt ferns from the English Wealden. *Proceedings of the Geologists'*
11
12 *Association*, **92**, 47-58.
13

14
15
16 HAUSHAHN, T., SPECK, T. and MASSETER, T. 2014. Branching morphology of decapitated
17
18 arborescent monocotyledons with secondary growth. *American Journal of Botany*, **101**, 754-
19
20 763.
21
22

23
24
25 HEIL, M. 2015. Extrafloral nectar at the plant-insect interface: a spotlight on chemical ecology,
26
27 phenotypic plasticity, and food webs. *Annual Review of Entomology*, **60**, 213-232.
28
29

30
31
32 HESSE, L., MASSETER, T., LEUPOLD, J., SPENGLER, N., SPECK, T. and KORVINK, J. G. 2016.
33
34 Magnetic resonance imaging reveals functional anatomy and biomechanics of a living dragon
35
36 tree. *Scientific Reports*, **6**, 32685.
37
38

39
40
41 HIRMER, M. 1927. *Handbuch der Paläobotanik*. R. Oldenburg, Berlin, 708pp.
42
43

44
45
46 JUNG, J., LEE, S. C., and CHOI, H. 2008. Anatomical patterns of aerenchyma in aquatic and
47
48 wetland plants. *Journal of Plant Biology*, **51**, 428-439.
49
50

51
52
53 KARST, A.L. and LECHOWICZ, M.J. 2007. Are correlations among foliar traits in ferns consistent
54
55 with those in the seed plants? *New Phytologist*, **173**, 306-312.
56
57
58
59
60

1
2
3 KATO, M. 1993. A taxonomic study of the genus *Matonia* (Matoniaceae). *Blumea-Biodiversity,*
4
5 *Evolution and Biogeography of Plants*, **38**, 167-172.
6
7

8
9
10 KOENIGUER, J. C. 1966. Étude paléophytogéographique du Continental Intercalaire de l'Afrique
11
12 Nord-Équatoriale. Sur de nouveaux échantillons du genre *Paradoxopteris*. *Mémoires de la*
13
14 *Société géologique de France*, **105**, 100-112.
15
16

17
18
19 KOPTUR, S., SMITH, A. R. and BAKER, I. 1982. Nectaries in some neotropical species of
20
21 *Polypodium* (Polypodiaceae): preliminary observations and analyses. *Biotropica*, **14**, 108-113.
22
23

24
25
26 KOPTUR, S., PALACIOS-RIOS M., DÍAZ-CASTELAZO, C., MACKAY W. P. and RICO-GRAY, V. 2013.
27
28 Nectar secretion on fern fronds associated with lower levels of herbivore damage: field
29
30 experiments with a widespread epiphyte of Mexican cloud forest remnants. *Annals of Botany*,
31
32 **111**, 1277-1283.
33
34

35
36
37 KRAMER, K. U. 1990. Matoniaceae. 183-185. In KRAMER, K. U. and GREEN, P. S. (eds).
38
39 *Pteridophytes and Gymnosperms*. Springer, Berlin, 404 pp.
40
41

42
43 LEHTONEN, S., SILVESTRO, D., KARGER, D. N., SCOTese, C., TUOMISTO, H., KESSLER, M., PEÑA,
44
45 C., WAHLBERG, N., and ANTONELLI, A. 2017. Environmentally driven extinction and
46
47 opportunistic origination explain fern diversification patterns. *Scientific Reports*, **7**, 1-12.
48
49

50
51
52 LYON, M. A., JOHNSON, K., WING, S., NICHOLAS, D., LACOVARA, K., and SMITH, J. 2001. Late
53
54 Cretaceous equatorial coastal vegetation: new megaflora associated with dinosaur finds in the
55
56 Bahariya oasis, Egypt. *Abstracts with programs Geological Society of America*, **3**, A-198.
57
58
59
60

- 1
2
3 MORGAN, E. J. 1959. The morphology and anatomy of American species of the genus
4
5 *Psaronius*. *Illinois biological monographs*, **27**, 1-108.
6
7
8
9
10 NIKLAS, K. J. 1978. Morphometric relationships and rates of evolution among Paleozoic
11
12 Vascular Plants. *Evolutionary Biology*, **11**, 509-543.
13
14
15
16 NIKLAS, K. J. 1991. Flexural stiffness allometries of angiosperm and fern petioles and rachises:
17
18 evidence for biomechanical convergence. *Evolution*, **45**, 734-750.
19
20
21
22
23 --- 1994. Predicting the height of fossil plant remains: an allometric approach to an old
24
25 problem. *American Journal of Botany*, **81**, 1235-1242.
26
27
28
29
30 NIKLAS, K. J., TIFFNEY, B. H. and KNOLL, A. H. 1983. Patterns in vascular land plant
31
32 diversification. *Nature* **303**, 614-616.
33
34
35
36
37 OUGH, K. and MURPHY, A. 2004. Decline in tree-fern abundance after clearfell harvesting.
38
39 *Forest Ecology and Management*, **199**, 153-163.
40
41
42
43 PEPPE, D. J., LEMONS, C. R., ROYER, D. L., WING, S. L., WRIGHT, I. J., LUSK, C. H. and RHODEN,
44
45 C. H. 2014. Biomechanical and leaf-climate relationships: A comparison of ferns and seed
46
47 plants. *American Journal of Botany*, **101**, 338-347.
48
49
50
51
52 POORTER, H., NIINEMETS, U., POORTER, L., WRIGHT, I. J. and VILLAR, R. 2009. Causes and
53
54 consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*, **182**,
55
56 565-588.
57
58
59
60

1
2
3 POYATO-ARIZA, F. J and BUSCALIONI, A. D. 2016. *Las Hoyas: a Cretaceous Wetland. A*
4
5 *multidisciplinary synthesis after 25 years of research on an exceptional fossil Lagerstätte from*
6
7 *Spain*. Verlag Dr. Friedrich Pfeil, Munich. 262 pp.
8
9

10
11
12 PPG I. 2016. A community-derived classification for extant lycophytes and ferns. *Journal of*
13
14 *Systematics and Evolution*, **54**, 563-603.
15
16

17
18 ROLLERI, C. H. 2002. Caracteres diagnósticos y taxonomía en el género *Angiopteris* Hoffm.
19
20 (Marattiaceae Bercht. & JS Presl): I, Los caracteres. *Revista del Museo de La Plata*, **15**, 23-49.
21
22

23
24 --- 2004. Revisión del género *Danaea* (Marattiaceae-Pteridophyta). *Darwiniana, nueva serie*,
25
26 **42**, 217-301.
27
28

29
30
31 ROMÁN GÓMEZ, P. 1986. Nota sobre la macroflora del Cretácico inferior de la Cuenca de
32
33 Préjano (LaRioja). *Coloquios de Paleontología*, **41**, 35-40.
34
35

36
37
38 ROSS, A. J. and COOK, E. 1995. The stratigraphy and palaeontology of the Upper Weald Clay
39
40 (Barremian) at Smokejacks Brickworks, Ockley, Surrey, England. *Cretaceous Research*, **16**, 705-
41
42 716.
43
44

45
46
47 ROTHWELL, G. W. and BLICKLE, A. H. 1982. *Psaronius magnificus* n. comb., a marattialean fern
48
49 from the Upper Pennsylvanian of North America. *Journal of Paleontology*, **56**, 459-468.
50
51

52
53
54 SCHNEIDER, C. A., RASBAND, W. S. and ELICERI, K. W. 2012. NIH Image to ImageJ: 25 years of
55
56 image analysis. *Nature methods*, **9**, 671.
57
58
59
60

1
2
3 SCOTT, A. C., CRIPPS, J. A., COLLINSON, M. E. and NICHOLS, G. J. 2000. The taphonomy of
4 charcoal following a recent heathland fire and some implications for the interpretation of fossil
5 charcoal deposits. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **164**, 1-31.
6
7
8

9
10
11
12 SENDER, L. M., VILLANUEVA-AMADOZ, U., PONS, D., DIEZ, J. B., GARCÍA-ÁVILA, M. and FERRER,
13 J. 2015. New reconstruction of *Weichselia reticulata* (Stokes et Webb) Fontaine in Ward
14 emend. Alvin, 1971 based on fertile remains from the middle Albian of Spain. *Historical*
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Biology, **27**, 460-468.

SEWARD, A. C. 1900. La flore wealdienne de Bernissart. *Mémoire du Muséum d'Histoire*
Naturelle de Belgique, **1**, 1-37.

SHINAQ, R. and BANDEL, K. 1998. The flora of an estuarine channel margin in the Early
Cretaceous of Jordan. *Freiberger Forschungshefte C 474, Paläontologie, Stratigraphie, Fazies*,
6, 39-57.

SILANTIEVA, N. and KRASSILOV, V. 2006. *Weichselia* Stiehler from Lower Cretaceous of
Makhtesh Ramon, Israel: new morphological interpretation and taxonomical affinities. *Acta*
Palaeobotanica Krakow, **46**, 119-135.

SMITH, A. R., PRYER, K. M., SCHETTPELZ, E., KORALL, P., SCHNEIDER, H. and WOLF, P. G. 2006.
A classification for extant ferns. *Taxon*, **55**, 705-731.

STOKES, C. and WEBB, P.B. 1824. Description of some fossil vegetables of the Tilgate Forest in
Sussex. *Transactions of the Geological Society of London*, **1**, 421-424.

1
2
3 TANNER, E. V. J. 1983. Leaf demography and growth of the tree-fern *Cyathea pubescens* Mett.
4
5 ex Kuhn in Jamaica. *Botanical Journal of the Linnean Society*, **87**, 213-227.
6
7

8
9
10 TIDWELL, W. D. and ASH, S. R. 1994. A review of selected Triassic to Early Cretaceous ferns.
11
12 *Journal of Plant Research*, **107**, 417-442.
13
14

15
16 VAN DEN BROECK, E. 1898. Les coupes du gisement de Bernissart. Caractères et dispositions
17
18 sedimentaires de l'argile ossifere du cran aux Iguanodons. *Bulletin de la Societé Belge de*
19
20 *Géologie de Paleontologie et de Hydrologie*, **12**, 216-243.
21
22
23

24
25 VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1993. A review of the Matoniaceae based on in situ
26
27 spores. *Review of Palaeobotany and Palynology*, **7**, 235-267.
28
29
30

31
32 WATSON, J. and ALVIN, K. L. 1996. An English Wealden floral list, with comments on possible
33
34 environmental indicators. *Cretaceous Research*, **17**, 5-26.
35
36
37

38
39 WEST, C. 1915. On the structure and development of the secretory tissues of the
40
41 Marattiaceae. *Annals of Botany*, **29**, 409-422.
42
43
44

45
46 WHITE, R.A. and TURNER, M.D. 2012. The anatomy and occurrence of foliar nectaries in
47
48 *Cyathea* (Cyatheaceae). *American Fern Journal*, **102**, 91-114.
49
50
51

52
53 WHITE, R.A., TURNER, M.D. 2017. The Comparative Anatomy of *Hymenophyllopsis* and
54
55 *Cyathea* (Cyatheaceae): a Striking Case of Heterochrony in Fern Evolution. *American Fern*
56
57 *Journal*, **107**, 30-57.
58
59
60

1
2
3 ZEILLER, R. 1914. Sur quelques plantes wealdiennes recueillies au Pérou par M. le Capitaine
4
5 Berthon. *Revue générale de Botanique*, **25**, 647-674.
6
7
8
9

10 **FIGURE AND TABLE CAPTIONS**

11
12 **FIG. 1.** A, Representation of all the reconstructions of *Weichselia reticulata* to date. B, fragment
13
14 of a pedate frond (MCCM-LH 17327) preserved as an impression from the locality of Las Hoyas
15
16 (upper Barremian of Cuenca, Spain). C, hand specimen from the locality of Tilgate Forest
17
18 (Valanginian of England) with numerous fragments of secondary pinnae (NHMUK PB V 51405).
19
20
21
22

23
24 **FIG. 2.** Localization of the locality of Bernissart. A, map of Belgium marking the Mons Basin. B,
25
26 detailed map of the locality including the different sinkholes. The red line marks the Négresse
27
28 connecting gallery (coordinates 50°28'60"N 3°38'33,2"E); S1, "puits Négresse" entrance leading
29
30 to the North sinkhole; S3, "puits Sainte Barbe" entrance leading to the Iguanodon sinkhole. C,
31
32 simplified log of the different lithological units encountered from south (0 m) to north (164 m)
33
34 in the Négresse connecting gallery. The black line in the lithological unit Y marks the point where
35
36 the orientation of the layers is changing. The arrow marks the position of the here studied
37
38 samples.
39
40
41
42

43
44 **FIG. 3.** Sections of stems. A-B, Thin section housed at the RBINS and figured by Alvin (1971)
45
46 IRSNB b 0229: A, whole section; B-C, enlargement of A x7.5; B, one bundle, C, secretory canal.
47
48 D-E, Y slices of μ CT-scans of: D, IRSNB b 8424; and E, IRSNB b 8425. Scale bar = 1 cm.
49
50
51

52
53 **FIG. 4.** Scars on the stem. A, Specimen IRSNB b 8426 showing crater-like marks and tubercles.
54
55 Scale bar = 0.5 cm. B, Specimen IRSNB b 842497. White arrow, petiole insertion with tubercles;
56
57 black arrow, possible radicular organ insertion. C, Longitudinal and transverse sections of the
58
59 petiole in specimen IRSNB b 8425. Blue, central strands of the petiole; red, central strands of the
60

1
2
3 stem. D, Longitudinal and transverse sections of the possible radicular organ of specimen IRSNB
4 b 8424. Blue, strands oriented in the opposite direction to the growth of the stem; red, central
5 strands of the stem. B-D scale bars = 1 cm.
6
7
8
9

10
11 **FIG. 5.** Elliptical scars. A, isolated elliptical scar (IRSNB b 8427) left, external view; right, internal
12 view. B-F, Vascular tissue at an elliptical scar in specimen IRSNB b 8424. B, reconstruction of
13 vascular tissue in 3D. C-D, subsequent sections of the frontal view: C, absence of vascular tissue
14 and D, irregular disposition of bundles underneath. E, transverse section. F, longitudinal section,
15 lateral view. G-J. Elliptical scars in specimen IRSNB b 8425. G-H. Photographs of the scars on the
16 specimen. I-J. Diagrams of outer surface and the vascular system in relation to the elliptical scars.
17 I. Transverse section, J. longitudinal section. Blue, central bundles of the petiole; red, elliptical
18 scars; black, rest of bundles; grey, outer part of the specimen. Scale bars = 1 cm.
19
20
21
22
23
24
25
26
27
28
29

30
31 **FIG. 6.** Stems. A. IRSNB b 8425. B-G. Vascular system of specimens IRSNB b 8424 and 8425
32 obtained from segmentation of μ CT-scans. B-C, IRSNB b 8425. B, lateral section. C, transversal
33 section at area where the petiole emerges. D-G. Sequential transversal sections of IRSNB b 8424
34 at possible branching area. Red, central vascular bundles; black, other bundles. Scale bars = 1
35 cm.
36
37
38
39
40
41
42
43
44

45 **FIG. 7.** Models of the vascular system obtained from segmentation. The colouring of the different
46 organs and stem centres is schematic, and was done posteriorly on the models. Red, central
47 vascular strands of the stem; blue, vascular strands of the petiole (P1, oldest petiole in the
48 specimen; P2, youngest petiole in the specimen); green, vascular strands of the rooting organ
49 (R1, oldest rooting organ in the specimen).
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **FIG. 8.** External morphology of Recent fern stems and petioles. A. Scales and tubercles in the
4 bases of petioles in *Angiopteris palmatiformis*. B. Tubercles in the stem and petiole base of
5 *Sphaeropteris intermedia*, there is a possible aerophore or nectary at the base of the petiole. C.
6 Isolated aerophores or lenticels along a line in the petioles of *Sphaeropteris cooperi* (F.Muell.)
7 R.M. Tryon. D. Isolated aerophores or lenticels in an irregular disposition in *Angiopteris evecta*.
8 E. Curved rhizome of *Matonia pectinata* R. Br. (specimen K000407521 from Kew herbarium).
9
10
11
12
13
14
15
16
17
18

19 **FIG. 9.** Comparison of the anatomy between fern species; all shown in transverse section. A,
20 *Weichselia reticulata* (IRSNB b 8425); B, *Cyathea trichiata* Domin drawn after White and Turner
21 (2017); C, *Platycterium aethyopicum* drawn after Allison (1913); D, *Danaea moritziana* Rosenst.
22 drawn after Rolleri (2004); E, *Matonia pectinata* drawn after image 9327 of the Cornell
23 University Plant Anatomy Collection; F, *Osmunda cinnamomea* drawn after image 8798 of the
24 Cornell University Plant Anatomy Collection; G, *Paradoxopteris stromeri* drawn after Edwards
25 (1933). Scale bar represents 10 mm; C-F are shown at two magnifications: image on right is to
26 scale; image on left: C–D, × 2 (scale bar represents 5 mm); E–F, × 5 (scale bar represents 2 mm).
27
28
29
30
31
32
33
34
35
36
37
38

39 **FIG. 10.** A. Central strands of thin section of *Weichselia reticulata* (IRSNB b 0229) B. Anatomy of
40 *Psaronius magnificus* (Herzer) Rothwell and Blicke without the roots (Rothwell and Blicke,
41 1982). Scale bars = 1cm.
42
43
44
45
46
47

48 **FIG. 11.** Vascular system comparison between A, *Weichselia reticulata* and B, *Dracaena*
49 *marginata*, schematic diagram based on Haushahn et al. (2014). Red, central vascular strands of
50 the stem; blue, vascular strands of the petiole in A and vascular strands of the ramification in B.
51
52
53
54
55
56

57 **TABLE 1.** Anatomical, morphological and architectural characters of *Weichselia reticulata* and
58 other fern taxa.
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 2. Calculated height following Niklas (1994) and maximum height from the literature.

TABLE 3. Variables used in the calculations of Corner’s Rules following Brouat et al. (1998).




FIG. 1. A, Representation of all the reconstructions of *Weichselia reticulata* to date. B, fragment of a pedate frond (MCCM-LH 17327) preserved as an impression from the locality of Las Hoyas (upper Barremian of Cuenca, Spain). C, hand specimen from the locality of Tilgate Forest (Valanginian of England) with numerous fragments of secondary pinnae (NHMUK PB V 51405).




FIG. 2. Localization of the locality of Bernissart. A, map of Belgium marking the Mons Basin. B, detailed map of the locality including the different sinkholes. The red line marks the Négresse connecting gallery (coordinates 50°28'60"N 3°38'33,2"E); S1, "puits Négresse" entrance leading to the North sinkhole; S3, "puits Sainte Barbe" entrance leading to the Iguanodon sinkhole. C, simplified log of the different lithological units encountered from south (0m) to north (164 m) in the Négresse connecting gallery. The black line in the lithological unit γ marks the point where the orientation of the layers is changing. The arrow marks the position of the here studied samples.

201x233mm (300 x 300 DPI)




FIG. 3. Sections of stems. A-B, Thin section housed at the RBINS and figured by Alvin (1971) IRSNB b 0229: A, whole section; B-C, enlargement of A x7.5; B, one bundle, C, secretory canal. D-E, Y slices of μ CT-scans of: D, IRSNB b 8424; and E, IRSNB b 8425.




FIG. 4. Scars on the stem. A, Specimen IRSNB b 8426 showing crater-like marks and tubercles. Scale bar = 0.5 cm. B, Specimen IRSNB b 842497. White arrow, petiole insertion with tubercles; black arrow, possible radicular organ insertion. C, Longitudinal and transverse sections of the petiole in specimen IRSNB b 8425. Blue, central strands of the petiole; red, central strands of the stem. D, Longitudinal and transverse sections of the possible radicular organ of specimen IRSNB b 8424. Blue, strands oriented in the opposite direction to the growth of the stem; red, central strands of the stem. B-D scale bars = 1 cm.




FIG. 5. Elliptical scars. A, isolated elliptical scar IRSNB b 8427) left, external view; right, internal view. B-F, Vascular tissue at an elliptical scar in specimen IRSNB b 8424. B, reconstruction of vascular tissue in 3D. C-D, subsequent sections of the frontal view: C, absence of vascular tissue and D, irregular disposition of bundles underneath. E, transverse section. F, longitudinal section, lateral view. G-J. Elliptical scars in specimen IRSNB b 8425. G-H. Photographs of the scars on the specimen. I-J. Diagrams of outer surface and the vascular system in relation to the elliptical scars. I, Transverse section, J, longitudinal section. Blue, central bundles of the petiole; red, elliptical scars; black, rest of bundles; grey, outer part of the specimen. Scale bars = 1 cm.




FIG. 6. Stems. A. IRSNB b 8425. B-G. Vascular system of specimens IRSNB b 8424 and 8425 obtained from segmentation of μ CT-scans. B-C, IRSNB b 8425. B, lateral section. C, transversal section at area where the petiole emerges. D-G. Sequential transversal sections of IRSNB b 8424 at possible branching area. Red, central vascular bundles; black, other bundles. Scale bars = 1 cm.




FIG. 7. Models of the vascular system obtained from segmentation. The colouring of the different organs and stem centres is schematic, and was done posteriorly on the models. Red, central vascular strands of the stem; blue, vascular strands of the petiole (P1, oldest petiole in the specimen; P2, youngest petiole in the specimen); green, vascular strands of the rooting organ (R1, oldest rooting organ in the specimen).




FIG. 8. External morphology of Recent fern stems and petioles. A. Scales and tubercles in the bases of petioles in *Angiopteris palmatifomis*. B. Tubercles in the stem and petiole base of *Sphaeropteris intermedia*, there is a possible aerophore or nectary at the base of the petiole. C. Isolated aerophores or lenticels along a line in the petioles of *Sphaeropteris cooperi* (F.Muell.) R.M. Tryon. D. Isolated aerophores or lenticels in an irregular disposition in *Angiopteris evecta*. E. Curved rhizome of *Matonia pectinata* R. Br. (specimen K000407521 from Kew herbarium).

167x139mm (300 x 300 DPI)




FIG. 9. Comparison of the anatomy between fern species; all shown in transverse section. A, *Weichselia reticulata* (IRSNB b 8425); B, *Cyathea trichiata* Domin drawn after White and Turner (2017); C, *Platycerium aethyopicum* drawn after Allison (1913); D, *Danaea moritziana* Rosenst. drawn after Rolleri (2004); E, *Matonia pectinata* drawn after image 9327 of the Cornell University Plant Anatomy Collection; F, *Osmunda cinnamomea* drawn after image 8798 of the Cornell University Plant Anatomy Collection; G, *Paradoxopteris stromeri* drawn after Edwards (1933). Scale bar represents 10 mm; C-F are shown at two magnifications: image on right is to scale; image on left: C-D, $\times 2$ (scale bar represents 5 mm); E-F, $\times 5$ (scale bar represents 2 mm).

166x200mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




FIG. 10. A. Central strands of thin section of *Weichselia reticulata* (IRSNB b 0229) B. Anatomy of *Psaronius magnificus* (Herzer) Rothwell and Blicke without the roots (Rothwell and Blicke, 1982). Scale bars = 1cm.




FIG. 11. Vascular system comparison between A, *Weichselia reticulata* and B, *Dracaena marginata*, schematic diagram based on Haushahn et al. (2014). Red, central vascular strands of the stem; blue, vascular strands of the petiole in A and vascular strands of the ramification in B.

	<i>Weichselia</i>	<i>Paradoxopteris</i>	<i>Matonia</i>
stem diameter (cm)	7.14	15.83	0.45
bundle size (cm)	0.2	0.53	0.24
bundle / stem	0.03	0.03	0.54
type of stele	Dictyostele	Dictyostele	Solenostele
Polycyclic	yes	yes	Yes
strands to leaf	many	Np	One
accessory strands	?	yes	No
roots from central strand	yes	?	No
secretory canals	?	no	No
endodermis	?	?	yes
Cortex /stem diameter	Np	Np	0.07
Growth habit*	?	?	Terrestrial
position of rhizome	?	?	Creeping
branching	?	Np	yes
Equal branching?	?	Np	yes
Stem curved?	yes	Np	yes
petiole/ stem	0.66	Np	0.75
Leaf	Pedate frond, and bipinnate	Np	Pedate frond, and once pinnate
Leaf area (cm ²)	41902	Np	229
LMA (g/m ²)	352.5*	Np	165
Rhizome length (cm)	857***	2515***	13

* Following Peppe et al. (2014)

** Calculated approximately, from data that are not general or from the same plant (source of data: Roll

***Calculated following Niklas (1994), maximum size if it is upright.

<i>Danaea</i>	<i>Angiopteris</i>	<i>Osmunda</i>	<i>Cyathea</i>	<i>Platyserium</i>
1.4	2	0.41	5.53	1.27
0.15	0.15	0.04	3.42	0.13
0.11	0.08	0.1	0.62	0.1
Dictyostele	Dictyostele	Dictyostele	Dictyostele	Dictyostele
yes	yes	no	no	yes
two	many	one	Many	many
no	no	no	Yes	yes, arising from original dictyostele
Yes	yes	no	no	no
yes	yes	no	no	no
no	no	yes	yes	yes
0.1	Np	Np	0.03	0.03
Terrestrial	Terrestrial	Terrestrial	Arborescent	Epiphytic
Erect or creeping	Erect	Erect	Erect	Creeping
No	no	yes	rare	Lateral bud
No branching	No branching	yes	yes	no
no	no	?	no	?
1.17**	0.25**	0.71	0.62	0.63
Once pinnate	Bipinnate	Bipinnate	Once to many times pinnate	Bifurcately branched
20.9	13.7	81.8	25587	?
50	150	225	158	?
200	100?	?	1000	?

leri, 2002 and Rolleri, 2004).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

<i>Psaronius</i>
43
0.45 - 6.6
0.57
Dictyostele
yes
many
no
rarely
yes
no
0.23
Arborescent
Erect
no
No branching
no
1.2 or 0.2
pinnate
?
?
1500

	Calculated height	Maximum height (literature)
<i>Weichselia</i>	857	?
<i>Danaea</i>	99	200
<i>Cyathea</i>	567	2400
<i>Psaronius</i> min.	345	1500
<i>Psaronius</i> max.	2170	1500

Values in cm. *Danaea*, Rolleri (2004); *Cyathea*, Bower (1926);

Psaronius Morgan (1959).

Psaronius min. and *Psaronius* max., are calculated from the minimum and maximum values from Rothwell and Blicke (1982) respectively.

	Max. stem area	Max. foliage area	Nº leaves obs.	Nº leaves possible
<i>Weichselia</i>	25.57*	41902	?	1 to 2*
<i>Matonia</i>	0.16	211*	1	1 small*
<i>Matonia</i> max.	0.48	633*	1	2*
<i>Cyathea</i>	24.02	39272*	1 to 3	2*
<i>Psaronius</i>	50.27	236527*	5 to 8	7*
<i>Psaronius</i> max.	158.36	671381*	5 to 8	19*

Max. stem area and Max. foliage area in cm². Variables obtained from the formula are marked with an asterisk. Other variables were obtained from the literature: *Weichselia*, from personal observation of specimens published in Blanco-Moreno et al. (2019); *Matonia*, from Table 1 and Kato (1993); *Matonia* max., from the maximum stem diameter in the literature (Kato, 1993); *Cyathea*, from Table 1, Tanner (1983), and Durand and Goldstein (2001); *Psaronius*, most frequent stem diameter measurement in Rothwell and Blicke (1982); and *Psaronius* max., from Table 1 and Rothwell and Blicke (1982).