

AGARICOMYCETES MEDICINAL MUSHROOMS WITH POTENTIAL NEUROPROTECTIVE ACTIVITY GROWING IN ARMENIA

Susanna Badalyan, Sylvie Rapior

▶ To cite this version:

Susanna Badalyan, Sylvie Rapior. AGARICOMYCETES MEDICINAL MUSHROOMS WITH POTENTIAL NEUROPROTECTIVE ACTIVITY GROWING IN ARMENIA. Proceedings of the Yerevan State University - Chemical and Biological Sciences, 2020, 54 (3), 10.46991/PYSU:B/2020.54.3.000. hal-03090953v1

${\rm HAL~Id:~hal\text{-}03090953} \\ {\rm https://hal.umontpellier.fr/hal\text{-}03090953v1}$

Submitted on 30 Dec 2020 (v1), last revised 4 Feb 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chemistry and Biology

2020, **54** (3), p. 000

Biology

AGARICOMYCETES MEDICINAL MUSHROOMS WITH POTENTIAL NEUROPROTECTIVE ACTIVITY GROWING IN ARMENIA

S. M. BADALYAN 1*, S. RAPIOR 2**

- ¹ Laboratory of Fungal Biology and Biotechnology, Institute of Pharmacy, Department of Biomedicine, YSU, Armenia
- ² Laboratoire de Botanique, Phytochimie et Mycologie, Faculté des Sciences Pharmaceutiques et Biologiques, CEFE UMR 5175, CNRS – Université de Montpellier – Université Paul-Valéry Montpellier – EPHE – IRD, Montpellier France

The Agaricomycetes mushrooms (phylum Basidiomycota) are recognised sources of valuable food and medicines. They are producers of bioactive compounds (phenolics, polysaccharides, proteins, steroids, terpenoids, etc.) possessing around 130 therapeutic effects (antimicrobial, anti-inflammatory, antioxidant, immunomodulatory, etc.). Mushrooms are also reported as potential neurotrophic and neuroprotective agents. Seventeen edible and inedible agaricomycetous species from different taxonomic and ecological groups have been reported in Armenia to possess neuroprotective activity. Evaluation resource value and biotechnological potential of Armenian agaricomycetous mushrooms will assist further development of novel myco-pharmaceuticals to prevent andmitigatedifferent disorders, including neurodegenerative.

https://doi.org/10.46991/PYSU:B/2020.54.3.000

Keywords: Agaricomycetes, antioxidant, anti-inflammatory, mushrooms, neurodegenerative, neuroprotective.

Introduction. Agaricomycetes mushrooms (phylum Basidiomycota) are rich bioactive compounds (alkaloids, phenolics, polysaccharides, proteins, ribosomal and non-ribosomal peptides, steroids, terpenoids, etc.) possessing more than 130 therapeutic effects (analgesic, antibacterial, antifungal, anti-inflammatory, antioxidant, cytotoxic, hypocholesterolemic, hepatoprotective, hypoglycemic, hypotensive, immunomodulatory, mitogenic/regenerative, etc.) [1–7]. Theyhave also been reported as neuroprotective and anti-depressive agents [8-14]. New screening strategies based on innovative genetic approaches have identified novel mushroom metabolites-derived products widely applicable in biomedicine [15]. The evaluation of resource value and establishment of specialized culture collections of Agaricomycetes fungi will have an incredible impact on biomedical research that will assist to develop novel myco-pharmaceuticals [6, 16, 17].

* E-mail: s.badalyan@ysu.am

** E-mail: sylvie.rapior@umontpellier.fr

Changes may still occur before final publication.

Biology

Neurodegenerative disorders (NDD) including age-related Alzheimer (AD) and Parkinson diseases (PD), dementia, autism, depression and epilepsy are affecting millions of people worldwide. Stress-less lifestyle and healthy diet prevent development of such pathological conditions [18]. Oxidative stress and neuro-inflammation, protein aggregation and mitochondrial dysfunction are considered to play an important role in the etiopathogenesis of NDD [8, 18, 19]. Available therapies are associated with mild to severe side-effects, therefore, the search for novel approaches and natural resources of bioactive compounds that regulate pathways leading to neuronal death and dysfunction are warranted [6, 8, 20, 24, 33].

Recent studies focus on the pharmacology and feasibility of bioactive compounds of plant and fungal origin as a potential strategy to target a variety of human metabolic and brain disorders [37]. The mechanism and possible synergy of action of these compounds has not been studied yet, however mushrooms-derived products with potential neuroprotective and psychotropic activities can prevent and mitigate development of mental disorders, including depression, anxiety, sleep disturbances and cognitive alterations [1, 7, 8, 12–14, 20, 21–24]. According torecent myco-pharmacological research, Agaricomycetes mushrooms are producers of different neuroprotective compounds [8, 20, 25–27].

The current paper is directed to the evaluation of resource value of Agaricomycetes medicinal mushrooms growing in Armenia with potential neuroprotective activity (NPA) and discusses future perspectives of their usage in biomedicine.

Neuroprotective Compounds of Agaricomycetes Mushrooms. Mushrooms are widely distributed worldwide and have been appreciated in traditional medicine for their nutritional and medicinal properties [6, 7]. Edible and inedible mushrooms are industrially cultivated organisms however their biotechnological potential and perspectives of usage in biomedicine have not been fully exploited [15].

The polysaccharides (β-glucans) are considered one of the major bioactive compounds in mushrooms [6, 7]. The polysaccharides isolated from medicinal mushrooms *Agaricus bisporus*, *Cantharellus cibarius*, *Coprinellus truncorum*, *Coprinus comatus* and *Inonotus obliquus* (Chaga mushroom), distributed also in Armenia showed significant antioxidant, anti-inflammatory, anti-cholinesterase (AChE) and neuroprotective activities which may allow suggest them in the palliative treatment of NDD [11, 28–31].

Mushroom-derived terpenoids, steroids, sterols and phenolics are also known by their diverse pharmacological effects, including anti-inflammatory, antioxidant and neuroprotective [1, 3–5, 32, 33]. The NPA of *Hericium erinaceus* (Lion's mane) has been attributed to cyathane diterpenoids as erinacinesthat can stimulate the production of brain-derived neurotrophic factor [34]. New lanostane triterpenes and aromatic meroterpenoids with antioxidant and neuroprotective activities were isolated from fruiting bodies of *Ganoderma lucidum* (Reishi ou Ling Zhi) [5]. Study of total phenolic content, as well as antioxidant, AChE, tyrosinase, α -amylase and α -glucosidase activities of polypore mushrooms as *Ganoderma applanatum*, *Trametes gibbosa* and *Trametes versicolor* suggest that they may be considered as a source of neuroprotective food supplements [3, 5, 32].

Agaricomycetous Mushrooms as a Source for Neuroprotective Biotech Products. The cultivated edible mushroom, H. erinaceus is an important medicinal fungus with immunomodulatory, anti-mutagenic, antioxidant, anti-inflammatory and antitumor properties. Myco-pharmacological studies have attracted considerable attention on H. erinaceus as a neuroprotector to prevent NDD, including dementia, anxiety or depression [7, 13, 20, 25, 35–38].

Scientific dada have reported antioxidant and neuroprotective activities of *Ganoderma* mushrooms [5, 9, 22, 39]. The ganoderic acid and lucidone A isolated from *G. lucidum* delay AD progression [22]. The inhibition of cholinesterase, tyrosinase, α -amylase and α -glucosidase enzyme activities, as well as antioxidant effect of extracts from medicinal mushrooms of *G. applanatum* and *Ganoderma resinaceum* allows suggesting them as a source to prepare new food supplements and develop new drug formulations with NPA [3].

Oyster mushrooms (*Pleurotus* spp.) possess a high quantity of antioxidants, including ergothioneine, adenosine, and polyphenol derivatives which reduce oxidative stress-related aging. Consumption of edible medicinal mushroom *Pleurotus eryngii* (King Oyster Mushroom) delayed the development of brain atrophy, ameliorated memory deficit in mice and significantly decreased the levels of brain phosphorylated τ -protein, A β plaque deposition and malondialdehyde [26]. The AChE activity was detected in *Pleurotus ostreatus* (Oyster Mushroom), however the effect was weaker than the effect of galantamine [10].

Several white rot polypore *Trametes* (= *Coriolus*) species, such as *Trametes gibbosa*, *T. hirsute*, *T. pubescens* and *T. versicolor* have been used for centuries in traditional medicine [7]. *In vitro* antioxidant, anti-diabetes, anti-dementia and anti-inflammatory activities of *Trametes* species were evaluated. The importance of nutra- and nutriceuticals derived from these polypore mushrooms as neuroprotectores have been considered [19, 32].

The antioxidant power and NPA of edible mushroom *Amanita caesarea* (Caesar's mushroom) have been reported to alleviate the deposition of amyloid beta (Aβ) protein in the brain and improve the central cholinergic system function. *A. caesarea* as a potential food for treatment or prevention of NDD was reported [23]. Muscimol is the main compound found in toxic *Amanita muscaria* (Fly agaric or Fly amanita) with a suppressive effect on essential tremor, without impairing speech and coordination in Parkinson-affected patients. The extract of *A. muscaria* showed significant NPA on *in vitro* neurotoxicity models [40].

Mushrooms with Neuroprotective Potential Distributed in Armenia. The forest regions in Armenia rich with mushroom biodiversity are mainly distributed in North-Eastern (Ijevan and Lori floristic regions) and South-Eastern (Zangezur floristic region) parts of the country [41, 42].

Seventeen edible and inedible agaricoid, polyporoid, hymenochaetoid and cantharelloid Agaricomycetes medicinal species with potential NPA belonging to ecological groups of saprotrophes and xylotrophes have been recorded in Armenia. The antioxidative, anti-inflammatory and neuroprotective effects of these species have been reported (see Table) [1, 3, 5, 8–11, 17, 19, 27, 36, 37, 43, 44].

The Agaricomycetes mushrooms with potential neuroprotective activity growing in Armenia

				F 1 . 1	Di di 1	I
	Species	Order	Edibility	Ecological group	Bioactive compound, medicinal activity	References
1.	Agaricus bisporus (J.E. Lange) Imbach	Agaricales	edible	Soil saprotrophe	Polysaccharide, against Alzheimer's disease	[28]
2.	Amanita muscaria (L.) Lam.	Agaricales	inedible, toxic	Soil saprotrophe	Muscimol, neuroprotective	[40]
3.	Cantharellus cibarius Fr.	Cantharellales	edible	Soil saprotrophe	Polysaccharide, neuroprotective	[11]
4.	Coprinellus truncorum (Scop.) Redhead, Vilgalys & Moncalvo	Psathyrellales	inedible	Soil saprotrophe/ xylotroph	Polysaccharide, antioxidant, anti- cholinesterase	[17, 31]
5.	Coprinus comatus (O.F. Müll.) Pers.	Agaricales	edible	Soil saprotrophe/ xylotroph	Polysaccharide, antioxidant, anti- cholinesterase	[17, 31]
6.	Flammulina velutipes (Curtis) Singer	Agaricales	edible	Xylotrophe	Polysaccharides, neuroprotective, mitigate neurodegeneration	[8, 20]
7.	Ganoderma applana- tum (Pers.) Pat. [=Ganoderma lipsiense (Batsch) G.F.]	Polyporales	inedible	Xylotrophe	Triterpenes, aromatic meroterpenoids, neuroprotective, against Alzheimer's disease	[5]
8.	Ganoderma lucidum (Curtis) P.Karst.	Polyporales	inedible	Xylotrophe	Triterpenes, aromatic meroterpenoids, antioxidant, anti- cholinesterase, neuroprotective, against Alzheimer's disease	[3, 5, 9, 22, 39]
9.	Hericium erinaceus (Bull.: Fr.) Pers.	Russulales	edible	Xylotrophe	Polysaccharides, cyathane diterpenoids, hericenones, erinacines, antioxidant, anti- depressant, memory enhancer, neuro-stimulating,	[4, 13, 21, 25, 30, 34–38]
10.	Inonotus obliquus (Ach. ex Pers.)	Hymeno- chaetales	inedible	Xylotrophe	Polysaccharides, proliferate human neurogliocytoma cells	[29]
11.	Laetiporus sulphureus (Bull.) Murrill	Polyporales	edible	Xylotrophe	Polysaccharides, antioxidant, against Alzheimer and Parkinson diseases	[10]
12.	Phellinus pini (Brot.) A. Ames [=Porodaedalea pini (Brot.) Murrill]	Hymeno- chaetales	inedible	Xylotrophe	Polysaccharides, phenolic compound hispidin, phenolic acids, polysaccharides, anti- cholinesterase, anti-inflammatory	[43]
13.	Pleurotus eryngii (DC.) Quél.	Agaricales	edible	Xylotrophe	Polysaccharides, phenolic compounds, ameliorate memory and learning deficit, against Alzheimer's disease	[14, 26]
14.	Pleurotus ostreatus (Jacq.) P. Kumm.	Agaricales	edible	Xylotrophe	Polysaccharides, antioxidant, against Alzheimer and Parkinson diseases	[10]
15.	Trametes gibbosa (Pers.) Fr.	Polyporales	inedible	Xylotrophe	Polysaccharides, phenolics, anti- oxidative, anti-neurodegenerative	[27]
16.	Trametes pubescens (Schumach.) Pilát	Polyporales	inedible	Xylotrophe	Polysaccharides, phenolics, antioxidative, anti- neurodegenerative	[27, 44]
17.	Trametes versicolor (L.) Lloyd	Polyporales	inedible	Xylotrophe	Polysaccharides, phenolics, antioxidant, anti-dementia, anti- inflammatory	[27, 32]

Conclusion. Given the aging of the population, finding natural resources with a potential effect to prevent or support the treatment of neurodegenerative diseases is a crucial societal issue. Indeed, further evaluation of resource value and biotechnological potential of Armenian medicinal Agaricomycetes mushrooms will assist to use them as potential natural source to formulate novelpotential neuroprotective myco-pharmaceuticals and functional foods without adverse side effects for preventingor mitigating different diseases, including neurodegenerative.

The authors are thankful to Dr. A. Barkhudaryan for a critical review of the manuscript.

This work was supported by the Science Committee of the MESCS RA, in the frames of the research thematic project No. 18T-1F115.

Received 17.07.2020 Reviewed 15.08.2020 Accepted 00.00.2020

REFERENCES

- Palacios I., Lozano M., Moro C., et al. Antioxidant Properties of Phenolic Compounds Occurring in Edible Mushrooms. Food. Chem. 28 (2011), 674–678.
- 2. De Silva D.D., Rapior S., Sudarman E., et al. Bioactive Metabolites from Macrofungi: Ethnopharmacology, Biological Activities, Chemistry. *Fungal Divers.* **62** (2013), 1–40.
- Zengin G., Sarikurkcu C., Gunes E., et al. Two GanodermaSpecies: Profiling of Phenolic Compounds by HPLC-DAD, Antioxidant, Antimicrobial and Inhibitory Activities on Key Enzymes Linked to Diabetes Mellitus, Alzheimer's Disease and Skin Disorders. Food Funct. 6: 8 (2015), 2794–2802.
- Wang X.Y., Zhang D.D., Yin J.Y., et al. Recent Developments in Hericium erinaceus Polysaccharides: Extraction, Purification, Structural Characteristics and Biological Activities. Crit. Rev. Food Sci. Nutr. 59: 1 (2019), S96–S115.
- Wang C., Liu X., Lian C., et al. Triterpenes and Aromatic Meroterpenoids with Antioxidant Activity and Neuroprotective Effects from Ganoderma lucidum. *Molecules* 24: 23 (2019), 4353
- Badalyan S.M., Zambonelli A. Biotechnological exploitation of macrofungi for the production of food, pharmaceuticals and cosmeceuticals. In: Sridhar K.R., Deshmukh S.K. (eds) Advances in Macrofungi: Diversity, Ecology and Biotechnology. CRC Press, Boca Raton (2019), 199-230.
- 7. Badalyan S.M., Barkhudaryan A., Rapior S. Recent progress in research on the pharmacological potential of mushrooms and prospects for their clinical application. In: Agrawal D.C., Dhanasekaran M. (eds) *Medicinal Mushrooms: Recent Progress in Research and Development*. Springer Nature, Singapore (2019), 1–70.
- 8. Sabaratnam V., Phan C.W. Neuroactive components of culinary and medicinal mushrooms with potential to mitigate age-related neurodegenerative diseases. In: Brahmachari G (ed) *Discovery and development of neuroprotective agents from natural products*. Elsevier (2017), 401–413.
- Ćilerdžić J.L., Sofrenić I.V., Tešević V.V., et al. Neuroprotective Potential and Chemical Profile of Alternatively Cultivated Ganoderma lucidumBasidiocarps. *Chem. Biodivers.* 15: 5 (2018), e1800036.
- Ćilerdžić J., Galić M., Vukojević J., Stajic M. Pleurotus ostreatus and Laetiporus sulphureus (Agaricomycetes): Possible Agents against Alzheimer and Parkinson Diseases. *Int. J. Med. Mushrooms* 21: 3 (2019), 275–289.

- Lemieszek M.K., Nunes F.M., Cardoso C., et al. Neuroprotective Properties of Cantharellus cibarius Polysaccharide Fractions in Different in vitro Models of Neurodegeneration. Carbohydr. Polymers 197 (2018), 598–607.
- 12. Bell V., Ferrão J., Chaquisse E., et al. Role of Mushrooms in Autism. *Austin J. Nutri. Food Sci.* **7**: 6 (2019), 1128
- Chong P.S., Fung M.L., Wong K.H., Lim L.W. Therapeutic Potential of Hericium erinaceus for Depressive Disorder. *Int. J. Mol. Sci.* 21: 1 (2019), E163.
- Liang C.-H., Huang P.-C., Mau J.-L., Chiang S.-S. Effect of the King Oyster Culinary-Medicinal Mushroom Pleurotus eryngii (Agaricomycetes) Basidiocarps Powder to Ameliorate Memory and Learning Deficit in Ability in Aβ-Induced Alzheimer's Disease C57BL/6J Mice Model. Int. J. Med. Mushrooms 22: 2 (2020), 145–159.
- Kües U., Badalyan S.M. Making Use of Genomic Information to Explore the Biotechnological Potential of Medicinal Mushrooms. In: Agrawal D.C., et al. (eds) Medicinal Plants and Fungi: Recent Advances in Research and Development, Medicinal and Aromatic Plants of the World, vol 4. Springer, New York (2017), 397–458.
- Badalyan S.M., Gharibyan N.G. Characteristics of Mycelial Structures of Different Fungal Collections. YSU Press, Yerevan, 2017.
- 17. Badalyan S.M. Medicinal Coprinoid Mushrooms (Agaricomycetes) Distributed in Armenia (Review). *Int. J. Med. Mushrooms* **22**: 3 (2020), 257–267.
- 18. Uddin M.S., Ashraf G.M. Introductory Chapter: Alzheimer's Disease The Most Common Cause of Dementia. In: Ashraf G.M., Uddin M.S. (eds) *Advances in Dementia Research*, IntechOpen, 2018.
- 19. Trovato Salinaro A., Pennisi M., Di Paola R., et al. Neuroinflammation and Neurohormesis in the Pathogenesis of Alzheimer's Disease and Alzheimer-Linked Pathologies: Modulation by Nutritional Mushrooms. *Immun. Ageing* **15** (2018), 8.
- Phan C.W., David P., Sabaratnam V. Edible and Medicinal Mushrooms: Emerging Brain Food for the Mitigation of Neurodegenerative Diseases. J. Med. Food 20: 1 (2017), 1–10.
- Rossi P., Cesaroni V., Brandalise F., et al. Dietary Supplementation of Lion's Mane Medicinal Mushroom, Hericium erinaceus (Agaricomycetes), and Spatial Memory in Wild-Type Mice. Int. J. Med. Mushroom 20: 5 (2018), 485–494.
- Lai G., Guo Y., Chen D., et al. Alcohol Extracts from Ganoderma lucidum Delay the Progress of Alzheimer's Disease by Regulating DNA Methylation in Rodents. Front. Pharmacol. 10 (2019), 272.
- Li Z., Chen X., Lu W., et al. Wang D. Anti-Oxidative Stress Activity is Essential for Amanita caesareaMediated Neuroprotection on Glutamate-Induced Apoptotic HT22 Cells and an Alzheimer's Disease Mouse Model. *Int. J. Mol. Sci.* 18: 8 (2017), E1623.
- Ho L.H., Zulkifli N.A., Tan T.C. Edible mushroom: Nutritional properties, potential nutraceutical values, and its utilisation in food product development. In: An introduction to mushroom (2020), IntechOpen.
- Cheng J.H., Tsai C.L., Lien Y.Y., et al. High Molecular Weight of Polysaccharides from Hericium erinaceus against Amyloid beta-Induced Neurotoxicity. BMC Complement Altern. Med. 16 (2016), 170.
- 26. Zhang C.J., Guo J.Y., Cheng H., et al. Protective Effects of the King Oyster Culinary-Medicinal Mushroom, Pleurotus eryngii (Agaricomycetes), Polysaccharides on β-Amyloid-Induced Neurotoxicity in PC12 Cells and Aging Rats, In Vitro and In Vivo Studies. *Int. J. Med. Mushrooms* 22: 4 (2020), 325–333.
- Knežević A., Stajić M., Sofrenić I., et al. Antioxidative, Antifungal, Cytotoxic and Anti-Neurodegenerative Activity of Selected Trametes Species from Serbia. PLoS One 13: 8 (2018), e0203064.
- 28. Mahmoud M.G., Ibrahim A.Y., et al. Therapeutic Potential and Structural Elucidation of a Water-Soluble Polysaccharide of a Wild Edible Mushroom Agaricus bisporus against Neurodegenerative Disease, Alzheimer. *World J. Pharm. Sci.* 2: 10 (2014), 1136–1145.
- Ning X., Luo Q., Li C., et al. Inhibitory Effects of a Polysaccharide Extract from the Chaga Medicinal Mushroom, Inonotus obliquus (Higher Basidiomycetes), on the Proliferation of Human Neurogliocytoma Cells. *Int. J. Med. Mushrooms* 16: 1 (2014), 29–36.

- He X., Wang X., Fang J., et al. Structures, Biological Activities, and Industrial Applications of the Polysaccharides from Hericium erinaceus (Lion's Mane) Mushroom: A Review. *Int. J. Biol. Macromol.* 97 (2017), 228–237.
- 31. Pejin B., Tešanović K., Jakovljević D., et al. The Polysaccharide Extracts from the Fungi Coprinus comatus and Coprinellus truncorum Do Exhibit AChE Inhibitory Activity. *Nat. Prod. Res.* 33: 5 (2018), 750–754.
- Pop R.M., Puia I.C., Puia A., et al. Characterization of Trametes versicolor: Medicinal Mushroom with Important Health Benefits. Not. Bot. Horti. Agrobo. 46: 2 (2018), 343–349.
- 33. Dhakal S., Kushairi N., Phan C.W., et al. Dietary Polyphenols: AMultifactorial Strategy to Target Alzheimer's Disease. *Int. J. Mol Sci.* **20** : 20 (2019), E5090.
- 34. Rupcic Z., Rascher M., Kanaki S., et al. Two New Cyathane Diterpenoids from Mycelial Cultures of the Medicinal Mushroom Hericium erinaceus and the Rare Species, Hericium flagellum. *Int. J. Mol. Sci.* **19**: 3 (2018), E740.
- 35. Phan C.W., Lee G.S., Hong S.L., et al. Hericium erinaceus (Bull.: Fr.) Pers. cultivated under tropical conditions: isolation of hericenones and demonstration of NGF-mediated neurite outgrowth in PC12 cells via MEK/ERK and PI3K-Akt signaling pathways. *Food Funct.* **5**: 12 (2014), 160–169.
- Kushairi N., Phan C.W., Sabaratnam V., et al. Lion's Mane Mushroom, Hericium erinaceus (Bull.: Fr.) Pers. Suppresses H2O2-Induced Oxidative Damage and LPS-Induced Inflammation in HT22 Hippocampal Neurons and BV2 Microglia. *Antioxidants* 8: 8 (2019), 261.
- 37. Saitsu Y., Nishide A., Kikushima K., et al. Improvement of cognitive functions by oral intake of Hericium erinaceus. *Biomed. Res.* **40**: 4 (2019), 125–131.
- 38. Limanaqi F., Biagioni F., Busceti C.L., et al. Potential Antidepressant effects of Scutellaria baicalensis, Hericium erinaceus and Rhodiola rosea. *Antioxidants* 9: 3 (2020), 234.
- Zhao C., Zhang C., Xing Z., et al. Pharmacological Effects of Natural Ganoderma and its Extracts on Neurological Diseases: A Comprehensive Review. *Int. J. Biol. Macromol.* 121 (2019), 1160–1178.
- Kondeva-Burdina M., Voynova M., Shkondrov A., et al. Effects of Amanita muscariaExtract on Different In VitroNeurotoxicity Models at Sub-Cellular and Cellular Levels. Food Chem. Toxicol. 132 (2019), 110687.
- 41. Badalyan S.M., Gharibyan N.G. Macroscopic Fungi from Central Part of Virahayotz Mountains' Forests of Armenia and Their Medicinal Properties. YSU Press, Yerevan (2008)
- 42. Badalyan S.M., Gharibyan N.G. Diversity of Polypore Bracket Mushrooms, Polyporales (Agaricomycetes) Recorded in Armenia and their Medicinal Properties. *Int. J. Med. Mushrooms* **18**: 4 (2016), 347–354.
- 43. Im K.H., Nguyen T.K., Kim J.K., et al. Evaluation of Anticholinesterase and Inflammation Inhibitory Activity of Medicinal Mushroom Phellinus pini (Basidiomycetes) Fruiting Bodies. *Int. J. Med. Mushrooms* 17: 2 (2015), 1011–1022.
- Im K.H., Nguyen T.K., Choi J., Lee T.S. In Vitro Antioxidant, Anti-Diabetes, Anti-Dementia, and Inflammation Inhibitory Effect of Trametes pubescens Fruiting Body Extracts. *Molecules* 21: 5 (2016), E639.

Ս. Մ. ԲԱԴԱԼՅԱՆ, ՍԻԼՎԻ ՌԱՊԻՈՐ

ՀԱՅԱՍՏԱՆՈՒՄ ԱՃՈՂ ԱԳԱՐԻԿՈՄԻՑԵՏԱՅԻՆ ԴԵՂԱՍՆԿԵՐԻ ՆԵՅՐՈՊԱՇՏՊԱՆ ՆԵՐՈՒԺԸ

Ագարիկոմիցետային սնկերը (բաժին Basidiomycota) հայտնի են որպես արժեքավոր սննդի և դեղամիջոցների աղբյուր։ Դրանք կենսաակտիվ միացությունների (ֆենոլներ, պոլիսախարիդներ, ստերոիդներ, տերպենոիդներ և այլն) արտադրողներ են, որոնք ունեն շուրջ 130 բուժականագրեցություն (հակամանրէային, հակաբորբոքային,

հակաօքսիդիչ, իմունոմողուլյատոր և այլն)։ Մնկերը հանդիսանում են նաև որպես նեյրոտրոֆիկ և նեյրոպաշտպան գործոններ։ Հոդվածում բերվում է Հայաստանում աձող տարբեր կարգաբանական և էկոլագիակն խմբերին պատկանող ուտելիևոչ ուտելի ագարիկոմիցետային սնկերի 18 տեսակներ, որոնք օժտված են նյարդապաշտպան ազդեցությամբ։ Նրանց ռեսուրսային արժեքի և կենսատեխնոլոգիական ներուժի գնահատումը կնպաստի սնկային ծագման դեղապատրաստուկների ստացմանն ու կիրառմանը տարբեր նեյրոդեգեներատիվ խանգարումները կանխելու և մեղմելու նպատակով։

С. М. БАДАЛЯН, С. РАПИОР

АГАРИКОМИЦЕТНЫЕ ЛЕКАРСТВЕННЫЕ ГРИБЫ С НЕЙРОРПОТЕКНЫМ ПОТЕНЦИАЛОМ РАСТУЩИХ В АРМЕНИИ

Грибы Agaricomycetes (отдел Basidiomycota) являются источниками ценных пищевых и лекарственных продуктов. Они являются производителями биоактивных соединений (фенолов, полисахаридов, стероидов, терпеноидов и т.д.), обладающих около 130 терапевтическими эффектами (противомикробным, противовоспалительным, антиоксидантным, иммуномодулирующим и др.). Грибы также известны как потенциальные нейротрофические и нейропротекторные агенты. Было обнаружено, что восемнадцать видов съедобных и несъедобных агарикомицетных грибов, растущих в Армении, обладают нейропротекторной активностью. Оценка их ресурсной ценности и биотехнологического потенциала поможет разработке новых микопрепаратов для предотвращения и смягчения симптомов различных нейродегенеративных нарушений.