



HAL
open science

IFN α signaling through PKC- θ is essential for antitumor NK cell function

Natalia R. Comet, Juan Ignacio Aguilo, Moez G. Rathoré, Elena Catalán, Johan Garaude, × Gilles Uzé, Javier Naval, Julián Pardo, Martin Villalba, Alberto Anel

► To cite this version:

Natalia R. Comet, Juan Ignacio Aguilo, Moez G. Rathoré, Elena Catalán, Johan Garaude, et al.. IFN α signaling through PKC- θ is essential for antitumor NK cell function. *OncoImmunology*, 2014, 3 (8), pp.e948705. 10.4161/21624011.2014.948705 . hal-03108492

HAL Id: hal-03108492

<https://hal.umontpellier.fr/hal-03108492>

Submitted on 13 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

IFN α signaling through PKC- θ is essential for antitumor NK cell function

Natalia R Comet^{1,†}, Juan Ignacio Aguiló^{1,†}, Moez G Rathoré², Elena Catalán¹, Johan Garaude², Gilles Uzé³, Javier Naval¹, Julián Pardo^{4,5}, Martín Villalba^{2,6,‡}, and Alberto Anel^{1,*}

¹Apoptosis, Immunity & Cancer Group; Department of Biochemistry and Molecular and Cell Biology; University of Zaragoza and Aragón Health Research Institute (IIS Aragón); Zaragoza, Spain; ²INSERM U1040; Université de Montpellier 1; UFR Médecine; Montpellier, France; ³CNRS UMR 5235; Université de Montpellier II; Place Eugene Bataillon; Montpellier, France; ⁴Immune Effector Cells Group; IIS Aragón; Biomedical Research Center of Aragón (CIBA); Nanoscience Institute of Aragón (INA); Zaragoza, Spain; ⁵Aragón I+D Foundation (ARAID); Zaragoza, Spain; ⁶Institut de Recherche en Biothérapie (IRB); CHU Montpellier; Montpellier, France

[†]These authors contributed equally to this work.

[‡]Shared senior authorship.

Keywords: CXCL10, IFN- α , IL-15, NK cells, PKC- θ

Abbreviations: CDK8, cyclin-dependent kinase 8; CXCL10, (C-X-C motif) ligand 10/*CXCL10*; FCS, fetal calf serum; IFN α , interferon- α ; *IFNA1*; IFN γ , interferon- γ , *IFNG*; IL-15, interleukin-15/*IL15*; mAb, monoclonal antibody; MACS, magnetic cell separation; MEF, murine embryonic fibroblast; MHC-I, major histocompatibility complex class I/*MHC-I*; NK, natural killer; PI3K, phosphatidylinositol-3-kinase; PKC- θ , protein kinase C- θ , *PRKCQ*; PLC, phospholipase-C; Poly I:C, poly-inosinic:cytidilic acid; RT-PCR, real-time polymerase chain reaction; STAT-1, signal transducer and activator of transcription-1/*STAT1*.

We have previously shown that the development of a major histocompatibility complex class I (MHC-I)-deficient tumor was favored in protein kinase C- θ knockout (PKC- $\theta^{-/-}$) mice compared to that occurring in wild-type mice. This phenomenon was associated with scarce recruitment of natural killer (NK) cells to the tumor site, as well as impaired NK cell activation and reduced cytotoxicity *ex vivo*. Poly-inosinic:cytidilic acid (poly I:C) treatment activated PKC- θ in NK cells depending on the presence of a soluble factor produced by a different splenocyte subset. In the present work, we sought to analyze whether interleukin-15 (IL-15) and/or interferon- α (IFN α) mediate PKC- θ -dependent antitumor NK cell function. We found that IL-15 improves NK cell viability, granzyme B expression, degranulation capacity and interferon- γ (IFN γ) secretion independently of PKC- θ . In contrast, we found that IFN α improves the degranulation capability of NK cells against target cancer cells in a PKC- θ -dependent fashion both *ex vivo* and *in vivo*. Furthermore, IFN α induces PKC- θ auto-phosphorylation in NK cells, in a signal transduction pathway involving both phosphatidylinositol-3-kinase (PI3K) and phospholipase-C (PLC) activation. PKC- θ dependence was further implicated in IFN α -induced transcriptional upregulation of chemokine (C-X-C motif) ligand 10 (*CXCL10*), a signal transducer and activator of transcription-1 (STAT-1)-dependent target of IFN α . The absence of PKC- θ did not affect IFN α -induced STAT-1 Tyr701 phosphorylation but affected the increase in STAT-1 phosphorylation on Ser727, attenuating *CXCL10* secretion. This connection between IFN α and PKC- θ in NK cells may be exploited in NK cell-based tumor immunotherapy.

Introduction

The serine threonine specific protein kinase C- θ (PKC- θ) was initially isolated and characterized as a PKC isoenzyme expressed in T cells.¹ In this context, PKC- θ signal transduction downstream of the T cell receptor (TCR) triggers key transcription factors essential for T-cell activation such as activating protein-1 (AP-1, also known as JUN), nuclear factor of κ -light chain polypeptide gene enhancer in B cells (NF- κ B) and nuclear factor of activated T cells (NF-AT).²⁻⁶ However, PKC- θ expression is not exclusive to T cells.⁷

The cancer immunosurveillance hypothesis proposes that the immune system detects and eliminates neoplastic cells undergoing malignant transformation and that the adaptive immune system is

able to maintain small cancer lesions in an equilibrium, i.e., in a growth constricted state.⁸⁻¹² PKC- θ is essential for this immune function at least in the control of leukemia as evinced by prior findings demonstrating that PKC- θ -deficient mice exhibit increased lymphoproliferative disease incidence and onset.¹³ This response presumably involved cytotoxic T lymphocyte (CTL) function, considering that these leukemic cancer cells express major histocompatibility complex class I (MHC-I).

Cells of the innate immune system, including $\gamma\delta$ T cells, natural killer (NK) cells and natural killer T (NKT) cells, can also mediate anticancer immune responses. Specifically, NK cells play an important role in tumor immunosurveillance,¹⁴⁻¹⁶ as they have been observed to control the progression of MHC-I-deficient tumors via

*Correspondence to: Alberto Anel; Email: anel@unizar.es

Submitted: 04/09/2014; Revised: 06/03/2014; Accepted: 06/06/2014
<http://dx.doi.org/10.4161/21624011.2014.948705>

perforin/granzyme- and FasL-mediated cytotoxicity.¹⁷⁻¹⁹ Interestingly, we have previously shown that PKC- θ is required for FasL-expression in T cells, although not for CTL degranulation.^{20,21} Since most cancer cells down-regulate MHC-I expression to escape the CTL-mediated cytotoxicity²² it is important to evaluate further the role of NK cells in mediating antitumor immunity.

We have previously shown that development of a MHC-I-deficient tumor (RMA-S) is favored in PKC- $\theta^{-/-}$ mice as compared to that occurring in wild-type (wt) mice.^{23,24} This pathological condition was found to be associated with a deficient recruitment of NK cells to the tumor site in PKC- $\theta^{-/-}$ mice, correlating with impaired activation of recruited NK cells. Furthermore, *in vitro* testing showed that NK cells isolated from PKC- $\theta^{-/-}$ mice exhibited reduced cytotoxicity against leukemic RMA-S cells after poly I:C treatment *ex vivo*.²³ We also observed that poly I:C treatment increased the relative expression levels and the activation status of PKC- θ in NK cells, both *in vivo* and *in vitro*. In the latter scenario, the presence of the complete splenocyte immune population was apparently needed, suggesting that the effect is presumably partially mediated by macrophages and/or dendritic cells.²³ Our finding is in accord with previous studies demonstrating a NK cell-dependent increase in anticancer activity of NK cells responding to poly I:C stimulation.²⁵ PKC- θ activation in this context is presumably mediated by soluble factor(s), which could be cytokine(s) secreted by macrophages and (or) dendritic cells (DCs) that are known to activate NK cells.²⁴ Thus, we have set out to define factors mediating NK activation and tumoricidal activity through PKC- θ . Since it has been previously reported that interleukin-12 (IL-12) signal transduction is affected in NK cells derived from PKC- $\theta^{-/-}$ mice,²⁶ this cytokine was tested first. Interleukin-15 (IL-15) was also included in the mentioned studies, since it is important in regulating NK cell function and survival,^{27,28} and for efficient antitumor NK cell activity.²⁹ Indeed, we reported that both, IL-12 and IL-15 activated PKC- θ in NK cells, with IL-15 being more potent at inducing PKC- θ phosphorylation. More importantly, in a mixed splenocyte culture stimulated *ex vivo* with poly I:C, neutralizing antibodies against IL-15 substantially reduced NK cell PKC- θ phosphorylation, whereas IL-12 antibody blockade was ineffective.²³ Therefore, IL-15 appeared to be the most feasible candidate to mediate PKC- θ -dependent antitumor NK cell immune function.²⁴ In the present study, we initially set out to test this possibility, testing IL-15 in regards to PKC- θ activation status and NK cell immunophenotypes. Contrary to our expectations, our results implicate interferon- α (IFN α) as the principal cytokine that signals through PKC- θ in NK cells and, as a consequence of downstream transcriptional changes, is primarily responsible for PKC- θ -dependent NK cell anticancer immunity.

Results

PKC- θ in IFN α and IL-15 effect on survival and immune function of NK cells

Our previous studies suggested that IL-15 could be the main cytokine responsible for the PKC- θ -dependent antitumor function of NK cells.²³ In order to evaluate the necessity for PKC-

θ -mediated signal transduction in a particular NK cell biological process, we comparatively analyzed IFN α and IL-15 responses in NK cells derived from PKC- $\theta^{-/-}$ versus wt animals. As shown in Fig. 1A, using an Annexin V externalization assay, we found that IL-15 is crucial for NK cell survival as although the majority (~70%) of isolated murine NK cells were Annexin V positive within the first 24 h in culture, this programmed cell death was almost completely abolished by inclusion of IL-15 in the cultures. However, this effect was found to be independent of PKC- θ , since it was equally achieved in NK cells from wt or PKC- $\theta^{-/-}$ mice. IFN α also seemed to improve survival, although less efficiently than IL-15 and also in a PKC- θ -independent manner. IL-15 also induced interferon- γ (IFN γ) production in purified NK cells in a PKC- θ independent fashion, whereas IFN α had no effect (Fig. 1B).

As shown in Fig. 1C, IL-15 improved NK cell degranulation when co-cultured with YAC-1 target cells as measured by an increase in the percentage of NK cells expressing CD107a, but this effect was again PKC- θ -independent. In sharp contrast, IFN α increased degranulation against YAC-1 cells to a higher magnitude, and this was entirely dependent upon PKC- θ expression, since this immunity-related biological process was abolished in NK cells from PKC- $\theta^{-/-}$ mice (Fig. 1C). Furthermore, although both IL-15 and IFN α modestly increased granzyme B expression in NK cells from wt mice over the already high basal expression level characteristic of spleen NK cells,²³ this increase was dependent on PKC- θ only in the case of IFN α (Fig. 1D). In sum, these experiments show that although IL-15 is important to maintain NK cell viability and in the induction of IFN γ secretion, these immune functions were independent of PKC- θ . On the other hand, our findings are the first to provide evidence that the increase in NK cell cytotoxic potential induced by IFN α is dependent on PKC- θ , with implications in the antitumor function of these molecules.

IFN α -mediated NK cell activation *in vivo* depends on PKC- θ

We next set out to determine the physiological dependence of IFN α -induced increase of NK cell cytotoxic potential by stimulating NK cells with IFN α *in vivo*. To this end, we injected 10,000 IU of IFN α intraperitoneally into wt or PKC- $\theta^{-/-}$ mice and, 24 h later, obtained purified peritoneal or splenic NK cells, and assayed NK cell degranulation (as measured by expression of 107a) against YAC-1 cells and the percentage of NK cells expressing granzyme B. We found that injection of IFN α increased the cytotoxic potential of peritoneal or splenic NK cells against YAC-1 cells (Fig. 2A). This effect was significantly (*p*-value <0.05) reduced in peritoneal NK cells obtained from PKC- $\theta^{-/-}$ mice, confirming the *in vitro* result and implicating PKC- $\theta^{-/-}$ as a key mediator of NK cell immune responses to IFN α . However, despite our findings using NK cells from the peritoneum, no significant difference was observed using splenic NK cells. *In vivo* injection of IFN α also resulted in a net increase in the expression of granzyme B, especially in peritoneal NK cells (Fig. 2B), but this effect was independent of PKC- θ expression, since granzyme B was observed in a similar fraction of NK cells derived from PKC- $\theta^{-/-}$ mice.

IFN α activates PKC- θ in NK cells through a PI3K- and PLC-dependent pathway

As shown in Fig. 3A, IFN α treatment of splenic NK cells from wt mice induced a rapid, efficient and sustained auto-phosphorylation of PKC- θ on Ser676, thus indicating enzymatic activation of the kinase. In contrast, the protein expression level of PKC- θ was not affected (Fig. 3A).

We next sought to identify the signal transduction pathway that connects IFN α to PKC- θ . Interestingly, PKC- θ is the only T-cell expressed PKC isoform that is activated through a PI3K-dependent pathway triggered by TCR ligation.³⁰ Although the main signaling pathway elicited by cytokine receptors is mediated by Janus kinases (JAKs) and signal transducer and activator of transcription (STATs), cytokines such as IFN α , IL-2 and IL-15 also activate the phosphatidylinositol-3-kinase (PI3K) pathway in B and T cells.³¹⁻³⁴ IFN α is also known to activate phospholipase-C (PLC) signaling cascades.³⁵ Hence, we tested the ability of wortmannin, a PI3K inhibitor, and U73122, a PLC inhibitor, or rapamycin, a mammalian target of rapamycin (mTOR) inhibitor, to attenuate PKC- θ activation induced by IFN α in NK cells. As shown in Fig. 3B, both wortmannin and U73122 abrogated the increase in PKC- θ auto phosphorylation induced by IFN α in NK cells, while rapamycin only partially inhibited this process.

The induction of some IFN α -target genes in NK cells is PKC- θ -dependent

We next set out to analyze if transcription of IFN α -responsive genes was affected in NK cells from PKC- $\theta^{-/-}$ mice. We used quantitative reverse transcription polymerase chain reaction (qRT-PCR) to assay the transcriptional levels of interferon-regulatory factor 9 (*IRF-9*) and interferon-stimulated 15 kDa protein (*ISG15*), 2 IFN α -responsive genes that have been previously reported to be dependent upon PKC- θ signaling in T cells,³⁶ as well as the mRNA levels of (C-X-C motif) ligand 10 (*CXCL10*), a candidate IFN α -responsive gene dependent on STAT1.³⁷ We used

interferon- γ (*IFN γ*) as a control gene that is not affected by IFN α in NK cells. As shown in Fig. 4A, IFN α treatment significantly increased the transcription of *IRF9* in NK cells obtained from either wt or heterozygotic PKC- $\theta^{+/-}$ mice, but not in those purified from PKC- $\theta^{-/-}$ mice. *ISG15* transcription was significantly increased in NK cells from all 3 genetic backgrounds, although to a different extent (Fig. 4A). Finally, *CXCL10* transcript levels were substantially increased in NK cells from wt mice, reduced in heterozygotic PKC- $\theta^{+/-}$ mice, and abolished in NK cells from PKC- $\theta^{-/-}$ mice (Fig. 4A). On the contrary, we did not detect a significant increase in the transcription of *IFN γ* in response to treatment with IFN α , demonstrating specificity in transcriptional responses. Fig. 4B shows the increase in the ratio of gene expression of each experimental transcript relative to the transcript levels of Rab




Figure 1. Dependence on PKC- θ for IL-15 and IFN α -induced NK cell survival and immune function. (A–D) Natural killer (NK) cells derived from C57BL/6 mice null for protein kinase C- θ ($PKC-\theta^{-/-}$) vs. wild-type animals ($PKC-\theta^{+/+}$) were functionally characterized for responses to interleukin 15 (IL-15) and interferon- α (IFN α). NK cells were isolated by magnetic cell separation (MACS) technology and cultured in complete medium at 2×10^6 cells/mL in the absence (–) or in the presence of the indicated concentrations of IL-15 or IFN α for 24 h. Afterwards, cells were labeled with an anti-NK1.1 mAb labeled with phycoerythrin (PE) to restrict the cytofluorimetric analysis to NK cells. (A) Cell death was determined by surface staining with Annexin-V labeled with fluorescein isothiocyanate (FITC) and flow cytometry. (B and C) After 24 h cytokine stimulation, NK cells were incubated with YAC-1 cells at a 1:1 ratio during 6 h in the presence of monensin and the production of interferon- γ (IFN γ) (B) and degranulation (C) was tested by flow cytometry. IFN γ was detected by intracellular staining with a specific anti-IFN γ antibody labeled with FITC and degranulation was determined by analysis of CD107a surface expression using an anti-CD107a antibody labeled with FITC. (D) The content of granzyme B was analyzed by permeabilization and intracellular staining with a specific anti-granzyme B antibody detected using a FITC-conjugated secondary and flow cytometry. Data are the mean \pm SD of 2 independent experiments performed in triplicate; statistical analyses were performed by Student's t test: * $P < 0.05$; ** $P < 0.02$.




Figure 2. Dependence on PKC- θ for IFN α -induced NK cell immune function *in vivo*. **(A and B)** Protein kinase C- θ knockout (PKC- $\theta^{-/-}$) or wild-type (PKC- $\theta^{+/+}$) C57BL/6 mice were injected intraperitoneally with 10,000 IU/mL of interferon- α (IFN α) or the same volume of PBS. After 24 h, mice were sacrificed, and natural killer (NK) cells were magnetically isolated by MACS technology either from total spleen cells or cells derived from flushing the peritoneum. NK cells were functionally characterized following a 6 h stimulation with YAC-1 target cells at a 1:1 ratio, immunofluorescence staining and cytofluorimetric analysis. **(A)** The degranulation potential of isolated NK cells against YAC-1 cells was estimated using a fluorophore-conjugated anti-CD107a antibody and flow cytometry. **(B)** The granzyme B content of YAC-1 stimulated NK cells was estimated via immunofluorescence staining and flow cytometry. Data are the mean \pm SD of assaying NK cells derived from 3 different mice in each experimental condition; statistical analyses were performed by Student's t test; * $P < 0.05$; *** $P < 0.01$; ns: difference not statistically significant.

interacting lysosomal protein-like 1 (*RILPL1*, also known as *RLP1*), a stable and abundant mRNA, upon IFN α treatment in the same experiments. The IFN α -dependent increase in the ratio was moderate for *IRF9* and equal in NK cells obtained from either wt or heterozygotic PKC- $\theta^{+/-}$ mice, but decreased by more than 50% in NK cells from PKC- $\theta^{-/-}$ mice, although this difference was not statistically significant (P -value > 0.05). In comparison, the relative increase in the transcript levels of *ISG15* was higher than that of *IRF9*, an increase significantly inhibited in NK cells derived from either PKC- $\theta^{+/-}$ or PKC- $\theta^{-/-}$ mice. Finally, the highest magnitude of IFN α -induced transcriptional increase corresponded to *CXCL10*, a molecular phenotype significantly inhibited in NK cells from either PKC- $\theta^{+/-}$ and PKC- $\theta^{-/-}$ mice. The fact that *IRF9* gene expression was not affected in NK cells from PKC- $\theta^{+/-}$, whereas *ISG15* and *CXCL10* mRNA levels were significantly affected by PKC- θ deficiency, suggests a dosage effect at promoters, some of which require full activation of the transcription factors involved. Our data confirmed that IFN α -mediated *IRF9* and

ISG15 expression is dependent upon PKC- θ in NK cells, as previously reported in T cells.³⁶ However, our data showing a dependency on PKC- θ for the IFN α -mediated induction of *CXCL10* gene expression are new and intriguing.

Reduced IFN α -induced STAT-1 phosphorylation in NK cells derived from PKC- $\theta^{-/-}$ mice

Our observation that IFN α stimulated *CXCL10* transcription is impaired in NK cells from PKC- $\theta^{-/-}$ mice suggested that PKC- θ is involved in STAT1 signaling. IFN α readily induced STAT1 phosphorylation at Tyr701 in NK cells obtained from the spleens of wild-type mice. However, the phospho-STAT1 level was only slightly reduced in splenic NK cells derived from PKC- $\theta^{-/-}$ mice. Considering that STAT1 phosphorylation depends on the direct action of JAK1,³⁸ PKC- θ presumably affects the STAT1 pathway downstream of JAK1-mediated tyrosine phosphorylation. Although tyrosine phosphorylation is indispensable for the transcriptional activity of STAT1, complete transcriptional capacity has been ascribed to additional phosphorylation events, including serine phosphorylation. For example, in macrophages, upon IFN γ treatment, PKC- α has been shown to phosphorylate STAT1 on Ser727, generating a fully functional STAT1 at the transcription level.³⁹ Furthermore, PKC- δ , a close homolog of PKC- θ , has been shown to be able to associate with STAT1 and mediate IFN γ -induced STAT1 phosphorylation in a promyelocytic cell line.⁴⁰ Interestingly, rottlerin, the PKC- δ inhibitor used in that study, also inhibits PKC- θ .^{20,21} Hence, we next tested whether IFN α treatment increased STAT-1 phosphorylation on Ser727 in NK cells. As shown in Fig. 6, 30 min of exposure to IFN α increased STAT1 phosphorylation on Ser727 in NK cells from wt mice. In contrast, NK cells from PKC- $\theta^{-/-}$ mice showed basal Ser727 phosphorylation that did not increase upon IFN α treatment (Fig. 6).

Finally, considering that IFN α induced transcription of *CXCL10* through PKC- θ in NK cells, we next sought to ascertain the PKC- θ -dependent effects of IFN α on *CXCL10* protein levels. We found that IFN α induced a significant increase in

CXCL10 secretion by NK cells, which was statistically decreased in NK cells derived from PKC- $\theta^{-/-}$ mice (Fig. 7). Taken together, these data elucidate a new signal transduction pathway in NK cells, connecting IFN α signaling with STAT-1-regulated genes through PKC- θ enzymatic activity.

Discussion

Our previous studies have shown that PKC- θ is essential for the antitumor function of NK cells.²³ We also previously showed that PKC- θ activation in this context is mediated by soluble factor(s), probably cytokines secreted by macrophages and (or) DCs that are known to activate NK cells.^{23,24} Thus, we prospectively evaluated whether IL-12 or IL-15 could be among the key cytokines activating PKC- θ signaling cascades. In this prior study, we found that although both IL-12 and IL-15 efficiently activated PKC- θ in NK cells, only neutralizing antibodies to IL-15 reduced PKC- θ auto-phosphorylation induced *in vitro* by poly I:C in the context of a mixed splenocyte population.²³ In the present study, we initially tested the possibility that IL-15 could be the main cytokine controlling these responses. Although we confirmed our prior data indicating that IL-15 is essential for NK cell survival and for the secretion of IFN γ , we report here that these functions are actually PKC- θ independent.

Unexpectedly, we have found that the main DC-derived cytokine whose signaling is dependent upon the presence of PKC- θ in NK cells, and further is most likely responsible for PKC- θ -dependent NK-cell mediated antitumor activity is rather IFN α . This interpretation is consistent with the increased sensitivity of IFNAR $^{-/-}$ or PKC- $\theta^{-/-}$ mice to the development of NK cell-sensitive tumors.^{23,41} We have shown that IFN α achieves this immune function by fostering at least 3 different responses in NK cells in a PKC- θ -dependent manner, including: i) augmenting the degranulation capacity of NK cells; ii) contributing to the increase in granzyme B expression in activated NK cells; and iii) inducing CXCL10 secretion from activated NK cells.

The augmentation of the degranulation capacity of NK cells was demonstrated by treating isolated splenic NK cells *in vitro* with IFN α , or alternatively, by activating NK cells *in vivo* by injecting IFN α , followed by assaying their

degranulation responses to YAC-1 cells. In either scenario, IFN α effects were strictly dependent upon PKC- θ , since NK cells from PKC- $\theta^{-/-}$ mice did not increase their basal degranulation potential. In the case of the *in vivo* activation, this was only observed in peritoneal NK cells, probably reflecting the basal activated phenotype of splenic NK cells, especially against YAC-1 cells, target cells that are extremely sensitive to NK cell cytotoxicity.^{17,23}

It has been previously reported that PKC- θ clusters at the NK cell immunological synapse, potentially amplifying effector responses, and possibly regulating NK cell degranulation.⁴² However, in this study the inhibition of degranulation was only partial in NK cells from PKC- $\theta^{-/-}$ mice as compared with wild type mice (41.8% vs. 64.5%), and there were no effects observed on the cytotoxicity against YAC-1 or BaF3 cells transfected with KAR ligands. However, IFN γ secretion was affected,⁴² as has been previously described.⁴³

Our group,²¹ and others,^{44,45} have demonstrated that PKC- θ is dispensable for degranulation induced by the TCR in mature cytotoxic T lymphocyte (CTL) clones and primary T cells, and further, that degranulation is not affected in CTLs from PKC- $\theta^{-/-}$ mice. However, the induction of FasL expression in these cells is strictly dependent on PKC- θ .^{20,21,30}




Figure 3. IFN α -induced PKC- θ activation in NK cells. **(A)** Natural killer (NK) cells were magnetically isolated by MACS technology from spleens of wild-type (wt) C57BL/6 mice and cultured in complete medium at 2×10^6 cells/mL in the absence (–) or presence (+) of the indicated concentrations of interferon- α (IFN α) for 30 min (left panel), or in the presence of 100 IU/mL of IFN α for the indicated periods of time (right panel). **(B and C)** Isolated NK cells from wt C57BL/6 mouse spleens were pre-incubated for 30 min in the absence (control) or in the presence of either 10 μ M U73122 or 100 nM of wortmannin or rapamycin, as indicated. Afterwards, 100 IU/mL of IFN α were added (+) or not (–) and cells were incubated in the presence of the indicated inhibitors for an additional 30 min. **(B)** PKC- θ phosphorylation in Ser676 determined by immunoblot. Total PKC- θ and/or β -actin expression was determined in parallel by immunoblot as loading controls. **(C)** The phospho-PKC- θ / β -actin ratios were determined by densitometry and results expressed as the increase in this ratio. Data shown are the mean \pm SD of at least 2 different experiments for each condition.




Figure 4. PKC-θ-dependent transcriptional changes in NK cells induced by IFNα. **(A and B)** Splens of wild-type (wt), protein kinase C-θ null heterozygotes (PKC-θ^{+/-}) or knockout (PKC-θ^{-/-}) C57BL/6 mice were used to magnetically (MACS) isolate 10 × 10⁶ natural killer (NK) cells. NK cells from each group were cultured in complete medium for 90 min with or without 100 IU/mL of IFNα, as indicated. RNA was extracted, reverse transcribed and cDNAs amplified via real time qPCR. **(A)** mRNA expression level of interferon-γ (*IFNG*) interferon regulatory factor 9 (*IRF-9*), ISG15 ubiquitin-like modifier (*ISG15*) or chemokine (C-X-C motif) ligand 10 (*CXCL10*) as a ratio of the reference *RLP1* mRNA level used for standardization. **(B)** qPCR results from **(A)** shown as the increase in the ratio of the indicated mRNA level expressed in control cells (untreated) versus cells treated with IFNα. Results are the mean ± SD of duplicate determinations in 2 independent experiments; statistical analyses were performed by Student's t test; **P* < 0.05; ***P* < 0.02; ****P* < 0.01; ns: difference not statistically significant.

We have shown previously that basal NK cell degranulation or cytotoxicity against YAC-1 cells was not affected using freshly isolated splenic NK cells from PKC-θ^{-/-} mice,²³ in agreement with previous reports.⁴³ However, YAC-1 cells are extremely sensitive to NK cells and unable to generate tumors in either syngeneic wt or PKC-θ^{-/-} mice. On the other hand, degranulation or cytotoxicity against RMA-S cells has been shown to be reduced in NK cells obtained from PKC-θ^{-/-} mice, in agreement with the increased tumorigenic potential of RMA-S cells in these knockout mice.²³ Moreover, in this system, splenic NK cells




Figure 5. IFNα-induced STAT1 phosphorylation at Tyr701 in NK cells is independent of PKC-θ. **(A and B)** Natural killer (NK) cells were isolated by MACS technology from spleens of wild type (PKC-θ^{+/+}) or protein kinase C-θ (PKC-θ^{-/-}) knockout C57BL/6 mice and cultured in complete medium at 2 × 10⁶ cells/mL in the absence or presence of the indicated concentrations of interferon-α (IFNα) for 30 min. Reactions were stopped in the cold, cells lysed, protein extracts prepared and signal transducer and activator of transcription-1 (STAT1) phosphorylation at residue Tyr701 determined by phospho-STAT1 (P-STAT1) specific immunoblot. β-actin expression was determined in parallel as loading control. **(B)** The phospho-STAT1/β-actin ratios of the experiment shown in **(A)** were determined by densitometry and represented as a bar diagram. Data shown are representative of 2 independent experiments.

must be previously activated by poly I:C to demonstrate cytotoxicity or degranulation potential against RMA-S cells, in contrast to unprimed NK cell activity against YAC-1 cells. Altogether, these data suggest that PKC-θ is a necessary constituent of *in vivo* activation of the degranulation potential of NK cells responding to poly I:C, or alternatively, endogenous tumor danger signals. Further, PKC-θ does not appear to be essential to the degranulation process itself, although it could amplify this response, as previously suggested.⁴² The present data also suggest that one of the PKC-θ dependent mechanisms by which poly I:C or danger signals activate NK cells *in vivo* is via the secretion of IFNα by DC or macrophages.²⁴

Granzyme B is the archetypical effector molecule of CTL and NK cells.⁴⁶ Granzyme B mRNA is abundant in NK cells and cytokine stimulation such as that meted by IL-15 and IFNα affects a net increase in protein content,⁴⁷ a response independently confirmed here. Furthermore, we also found that the intracellular content of granzyme B was lower in IFNα-activated NK cells derived from PKC-θ^{-/-} mice than from those of wt mice. *In vitro*, IFNα action is restricted to NK cells, and in this case the increase in granzyme B is PKC-θ-dependent.




Figure 6. IFN α -mediated STAT-1 phosphorylation on Ser727 is partially dependent upon PKC- θ in NK cells. **(A and B)** Natural killer (NK) cells were isolated by MACS technology from spleens of wt (PKC- $\theta^{+/+}$) or protein kinase C- θ knockout (PKC- $\theta^{-/-}$) C57BL/6 mice and cultured in complete medium at 2×10^6 cells/mL in the absence (-) or in the presence (+) of 100 IU/mL of IFN α for 30 min. **(A)** Reactions were stopped in the cold, cells lysed, protein extracts prepared and STAT1 phosphorylation in Ser727 determined by immunoblot. β -actin expression was determined in parallel by immunoblot as loading control. **(B)** The phospho-STAT1/ β -actin ratios were determined by densitometry and results expressed as the increase in this ratio between control cells and cells treated with IFN α in each case. Data shown are the mean \pm SD of at least 2 different experiments for each condition.




Figure 7. IFN α -induced CXCL10 secretion from NK cells is dependent on PKC- θ . Natural killer (NK) cells were isolated by MACS technology from spleens of wild type (PKC- $\theta^{+/+}$) or protein kinase C- θ knockout (PKC- $\theta^{-/-}$) C57BL/6 mice and cultured in complete medium at 5×10^6 cells/mL in the absence (control) or in the presence (+IFN α) of 100 IU/mL of IFN α for 8 h. After this time, cell culture supernatants were collected and the presence of CXCL10 detected using an ELISA kit. Data are the mean \pm SD of 3 different determinations made in duplicate; statistical analyses were performed by Student's t test; * $P < 0.05$; *** $P < 0.01$; ns: difference not statistically significant.

These differences are not observed *in vivo*, in which IFN α injection readily induced an increase in granzyme B expression, especially in peritoneal NK cells, irrespective of PKC- θ status. This indicates that *in vivo*, other cellular and molecular mechanisms independent of PKC- θ contribute to the observed increase in granzyme B expression. IFN α could induce the secretion of several other cytokines from other cell populations that induce granzyme B expression in NK cells in a PKC- θ -independent fashion. Indeed, it has been demonstrated that Type I IFN induces IL-15 secretion from activated DCs, and that DCs, through cross-presentation and IL-15 signaling, can thus induce NK cell priming.⁴⁸

CXCL10 is a chemokine involved in NK cell recruitment to inflamed tissues and sites of tumor development.⁴⁹⁻⁵¹ In fact, it has recently been demonstrated that the efficacy of adenovirus-engineered dendritic cells as a cancer vaccine correlates with their ability to secrete CXCL10 and CXCL8 and to efficiently recruit antitumor NK cells.⁵¹ CXCL10 secretion has been, so far, primarily attributed to DCs and monocytes/macrophages.^{50,51} Our data indicate that NK cells also secrete CXCL10 upon IFN α treatment, albeit at a lower concentrations than DCs and macrophages. Furthermore, in a recent transcriptomic analysis we found that NK cell activation by target cells in the presence of whole peripheral blood mononuclear cells (PBMCs) induces the expression of CXCL10, as well as its receptor CXCR3, suggesting the presence of an autocrine migration loop occurring during NK cell activation.⁵² The results obtained here suggest that IFN α produced by DC and macrophages/monocytes present in the PBMC population is contributing at least to the upregulation of this chemokine in NK cells. Previously, we demonstrated that recruitment of NK cells to the tumor site was impaired in PKC- $\theta^{-/-}$ mice, thus contributing to their increased sensitivity to tumor development. Hence, this defective response could be partly explained by the defective upregulation of CXCL10 by NK cells that we have uncovered here.

This observation has shed light on a new NK cell signal-transduction pathway connecting the IFN α receptor and PKC- θ activation, and further encompassed both PI3K and PLC activities, similar to that observed in T cells and reported by Srivastava et al.³⁶ PKC- θ is the only T cell expressed PKC isoform known to be activated through a PI3K-dependent pathway after TCR ligation.³⁰ However, in T cells PLC does not seem to play a primary role in PKC- θ activation by TCR ligation.³⁰ Therefore, unlike the route in T cells, it appears as though in IFN α -activated NK cells, both pathways should be activated to guarantee an efficient PKC- θ activation. The partial effect of the mTOR inhibitor rapamycin could be explained since this pathway is downstream of PI3K. These data are also in accord with PLC and PI3K activation in response to IFN α in other cellular contexts as well.^{34,35}

Although the impaired transcription of *ISG15* and *IRF9* in PKC- θ deficient NK cells responding to IFN α is concordant with prior observations in the context of T cells,³⁶ However, the dependence on PKC- θ of IFN- α -mediated *CXCL10* transcription and secretion is novel. *CXCL10* is an archetypical STAT1-dependent IFN-responsive gene.³⁷ In agreement with observations in T cells,³⁶ the main post-translational modification responsible for the activation of the transcriptional activity

of STAT1, specifically JAK1-mediated phosphorylation at Tyr701, is intact in NK cells arising in $PKC-\theta^{-/-}$ mice. Nevertheless, our data show a partially defective IFN α -mediated phosphorylation of STAT1 at Ser727, an alternative regulatory site in these cells. This post-translational modification may be a required signal transduction event for efficient *CXCL10* transcription and secretion.

Recently, it has been reported that the main kinase mediating STAT1 Ser727 phosphorylation in macrophages and murine embryonic fibroblasts (MEFs) is cyclin-dependent kinase 8 (CDK8).⁵³ CDK8 has been implicated as a STAT1 Ser727 phosphorylating enzyme in NK cells as well,⁵⁴ although in this case the enzymatic activity seems rather related with the presence of a basal level of Ser727 STAT1 phosphorylation. With these observations in mind, the basal phosphorylation status of STAT1 in freshly isolated NK cells from both wt and $PKC-\theta^{-/-}$ mice reported here is likely attributable to CDK8.

We further found that this phosphorylation increases in NK cells from wt but not from $PKC-\theta^{-/-}$ mice treated with IFN α . NK cells from mice expressing a mutant STAT1 recalcitrant to phosphorylation in Ser727 have been previously shown to exhibit higher cytotoxicity and granzyme B expression.⁵⁴ Considering that granzyme B expression has been shown to be regulated by a post-transcriptional mechanism in NK cells,⁴⁷ this effect of mutant STAT1 may relate to an unknown, non-transcriptional role of STAT1. In MEFs expressing mutant STAT1 and stimulated with IFN γ the transcription of *CXCL10* is augmented rather than diminished.⁵³ Our data have been obtained in NK cells stimulated with IFN α , and the situation is quite distinct, since CDK8-mediated basal STAT1 phosphorylation is present, and the role of $PKC-\theta$ seems rather to enhance Ser727 phosphorylation, resulting in an increase in *CXCL10* gene transcription. It is likely that phosphorylation on STAT1 Ser727 induces different responses in specific cellular contexts.

In summary, the present study uncovers the connection between IFN α and $PKC-\theta$ in NK cells and point to IFN α and $PKC-\theta$ -dependent signaling pathways as crucial regulators of NK cell function that can be exploited in NK cell-based tumor immunotherapy.

Materials and Methods

Mice

$PKC-\theta^{-/-}$ mice in the C57/BL6 background were a generous gift from Dr. D. Littman⁶ and were housed under the same conditions as the C57/BL6 parental (wt) mice. All experiments involving animals were performed according to the guidelines and regulations of the Center National de la Recherche Scientifique, France. Authors have official degrees for performing animal experimentation delivered by corresponding authorities in Spain or France, respectively.

Tumor cell lines

YAC-1 is a Moloney murine leukemia virus-induced lymphoma of the H-2K^b haplotype that lacks MHC-I expression

and is sensitive to lysis by NK cells.⁵⁵ Cell lines were cultured in RPMI-1640 medium supplemented with 10% heat-inactivated fetal calf serum (FCS), Glutamax and antibiotics (complete medium).

In vitro analysis of the effect of IFN- α and IL-15 on functional parameters in isolated NK cells

NK cells were isolated from the spleens of C57/BL6 wt or $PKC-\theta^{-/-}$ mice by magnetic cell separation (MACS) technology, using magnetic beads coated with anti-CD49b monoclonal antibody (mAb), as indicated previously.²³ NK cells were cultured in complete medium. Briefly, NK cells were isolated by positive selection using LS columns and anti-CD49b magnetic microbeads (Myltenyi Biotec, Madrid) from spleens of wild-type and $PKC-\theta^{-/-}$ C57BL/6 mice. Isolated NK cells were cultured for 24 h in this complete medium, supplemented with or without different concentrations of IL-15 or IFN α , both from Myltenyi Biotec, Madrid. Then, several functional assays were performed via marker staining and cytofluorimetric analyses using a FACScan (BD Biosciences, Madrid) flow cytometer as follows:

- i. Viability was determined by trypan blue staining and labeling with fluorescein isothiocyanate (FITC) conjugated Annexin-V and flow cytometry.
- ii. IFN γ expression was determined by intracellular staining with an anti-IFN γ rat monoclonal antibody (mAb) conjugated to FITC (BD, Madrid), as previously indicated.⁵⁶ In brief, NK cells were mixed at 1:1 ratio for 6 h with activating YAC-1 cells in the presence of monensin (GolgiStop reagent) to inhibit vesicular trafficking, thereby attenuating IFN γ secretion and favoring its intracellular accumulation. At the end of the incubation, cells were washed, fixed with 1% paraformaldehyde in PBS and permeabilized using a solution of 0.1% saponin in PBS with 5% FCS. The last solution contained the anti-IFN γ -FITC and also a mouse anti-NK1.1 mAb labeled with phycoerythrin (PE; BD, Madrid), both at a 1/1000 dilution and cells were incubated for 1 h at room temperature. IFN γ staining was analyzed by flow cytometry via gating within the NK1.1⁺ population.
- iii. Degranulation was analyzed following a similar protocol as indicated for IFN- γ secretion, using also YAC-1 cells as stimulus for NK cells, by analyzing CD107a externalization using a FITC-labeled anti-CD107a rat mAb, as indicated in.²³ Briefly, purified NK cells were incubated with target cells at a 1:1 ratio during 5 hours in the presence of Golgi-Stop (BD Biosciences, Madrid) and FITC-conjugated anti-CD107a mAb (BD Biosciences, Madrid). Then, cells were labeled with PE-conjugated anti-NK1.1 and analyzed via flow cytometry.
- iv. The intracellular expression of the cytotoxic effector granzyme B was analyzed using staining with a specific non-commercial rabbit polyclonal antibody on permeabilized NK cells, as we previously described.^{23,57}

Effect of IFN α on NK cell activation *in vivo*

For *in vivo* characterization of responses, 10,000 IU of IFN α per mouse were injected in wt or PKC- $\theta^{-/-}$ mice and immune cells were obtained 24 h later from dissociated spleen cell preparations or flushed from their peritoneum, essentially as previously described.²³ The NK cells were then purified by MACS technology, as indicated above. Next, their intracellular expression of granzyme B, and their degranulation potential stimulated by YAC-1 cells was analyzed as indicated above.

Analysis of PKC- θ activation in NK cells induced by IFN α

NK cells were isolated from the spleens of wt mice and treated during various times (10 to 120 min) with different doses of IFN α (100 to 1000 IU/mL). The auto phosphorylation of PKC- θ on Ser676 and its level of expression was analyzed by immunoblot using a rabbit pAb (Upstate/Millipore, Madrid) and a mouse mAb (BD, Madrid), respectively. Anti- β -actin antibody (Sigma, Madrid) was used in immunoblots as loading controls. Anti- β -actin reprobing on the same unstrapped membrane in which phospho-PKC- θ had been analyzed was used as a preferred loading control, since stripping of the anti-phospho-PKC- θ antibody resulted in protein loss, and anti-PKC- θ blotting in a separate membrane of a parallel set of samples gave sometimes inconsistent results. The effect of several enzymatic inhibitors on the auto-phosphorylation of PKC- θ at Ser676 following treatment with 100 IU/mL IFN α for 30 min was also analyzed by preincubating isolated NK cells with the indicated inhibitors for 30 min prior to adding IFN α . These included the PLC inhibitor U73122 (Calbiochem/Merck, Madrid) at 10 μ M, the PI3K inhibitor wortmannin (Sigma, Madrid), and the mTOR inhibitor rapamycin (Sigma, Madrid), both at 100 nM. The phospho-PKC- θ / β -actin ratio in each sample was determined using the ImageJ software.

RNA isolation and qRT-PCR of IFN α -responsive genes

ISG15 and *IRF9* mRNA levels were assayed as controls as these have previously been described to be PKC- θ -dependent genes, at least in T cells³⁶; *CXCL10* was tested as a prototype of a STAT1-dependent, IFN-responsive gene³⁷; and *IFNG* was tested as a control gene, not affected by IFN α , at least in NK cells.

Ribonucleic acid (RNA) isolation was performed and reverse transcribed as previously described.⁵⁸ cDNAs (cDNAs) were amplified using the SyberGreen polymerase chain reaction (PCR) Master Mix from Invitrogen. Amplification products were detected by Real Time qPCR using the light cycler 480 (Roche), according to manufacturer specifications. Reactions were carried out for 5 min at 95°C followed by 40 cycles of: 30 sec at 95°C, 30 sec at 64°C and 30 sec at 72°C. qRT-PCR data was analyzed using the Light Cycler 480 Software release 1.5 (Roche), according to the manufacturer specifications. The housekeeping gene β -actin was used to first normalize the mRNA expression levels. RLP13A (RLP1) was then used as a standard reference mRNA and relative mRNA levels were calculated according to this standard. Primers were designed and selected using the Primer3 (v. 0.4.0) program and were the following:

IFN γ : Forward, 5'-GCTCTGAGACAATGAACGCTAC-3',
IFN γ : Reverse, 5'-TCTTCCACATCTATGCCACTTG-3'
CXCL10: Forward, 5'-CCCACGTGTTGAGATCATTG-3',
CXCL10: Reverse, 5'-TCCATCACAGCACCGGG-3',
IRF9: Forward, 5'-CGC TGC TGC TCA CCT TCA T-3',
IRF9: Reverse, 5'-GGC TGT GCA CCT GGG TCT TA-3'
ISG15: Forward, 5'-GATTCCTGGTGTCCGTGAC-3'
ISG15: Reverse, 5'-CTAGCATCACTGTGCTGCTG-3'
 β -actin: Forward, 5'-GAGGGAAATCGTGCGTGACA-3'
 β -actin: Reverse, 5'-AATAGTGATGACCTGGCCGT-3'
RLP13A: Forward, 5'-GGTCCTCAAGACCAACGGACTC-3'
RPL13A: Reverse, 5'-GTGCGCTGTCAGCTCTCTAATG-3'

Analysis of STAT1 phosphorylation induced by IFN α in NK cells

STAT1 phosphorylation in Tyr701³⁸ or Ser727³⁹ was analyzed in control NK cells or in cells treated with IFN α from wt or PKC- $\theta^{-/-}$ mice by immunoblot using anti-STAT1 phospho-specific rabbit pAbs (Cell Signaling, Barcelona). Anti- β -actin reprobing on the same membrane in which phospho-STAT1 was analyzed has been used as a preferred loading control, since stripping of the anti-phospho-STAT1 antibodies resulted in protein loss, and anti-STAT1 blotting in a separate membrane of a parallel set of samples sometimes yielded inconsistent results. In addition, IFN α treatment for short times did not affect to the total STAT1 content (data not shown). The phospho-STAT1/ β -actin ratio in each sample was determined using the ImageJ software.

Analysis of CXCL10 secretion stimulated by IFN α in NK cells

CXCL10 secretion was determined by analyzing the supernatants of purified NK cells incubated for 8 h in complete medium in the presence or absence of 100 IU/mL IFN α . CXCL10 concentrations in the supernatants were measured using a specific quantitative ELISA kit (R&D systems, Abingdon, UK).

Statistical analysis

Results are shown as the average \pm SD and statistical significance was evaluated using the Student's t test. Differences were not considered significant if *P*-values were ≥ 0.05 .

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Funding

This work was supported by the collaborative projects CliNK SOE2/P1/E341 from Sudoe/Interreg IV B (EU) and Grant CTPP5/12 from "Communauté de Travail des Pyrénées"; by grant SAF2010-15341 to AA and grant SAF2011-25390 to JP, both from Ministerio de Economía y Competitividad (Spain); by Fondo Social Europeo and by the association CIEL, the association L'Un pour l'Autre and the federation Ensangle to MV; and by one AOI from the CHU Montpellier and fellowships from the ARC and Higher Education Commission, Pakistan to MGR.

References

- Baier G, Telford D, Giampa KM, Coggeshall KM, Baier-Bitterlich G, Isakov N, Altman A. Molecular cloning and characterization of PKC- η , a human novel member of the protein kinase C (PKC) gene family expressed predominantly in hematopoietic cells. *J Biol Chem* 1993; 268:4997-5004; PMID:8444877
- Altman A, Kaminski S, Busutil V, Droin N, Hu J, Tadevosyan Y, Hipskind RA, Villalba M. Positive feedback regulation of PLC γ 1/Ca²⁺ signaling by PKC η in restimulated T cells via a *tec* kinase-dependent pathway. *Eur J Immunol* 2004; 34:2001-11; PMID:15214048; <http://dx.doi.org/10.1002/eji.200324625>
- Baier-Bitterlich G, Uberall F, Bauer B, Fresser F, Wachter H, Grunicke H, Utermann G, Altman A, Baier G. Protein kinase C- η isoenzyme selective stimulation of the transcription factor complex AP-1 in T lymphocytes. *Mol Cell Biol* 1996; 16:1842-50; PMID:8657160
- Coudronniere N, Villalba M, Englund N, Altman A. NF- κ B activation induced by TCR/CD28 costimulation is mediated by PKC- η . *Proc Natl Acad Sci USA* 2000; 97:3394-9; PMID:10716728
- Pfeifhofer C, Kofler K, Gruber T, Tabrizi NG, Lutz C, Maly K, Leitges M, Baier G. Protein kinase C η affects Ca²⁺ mobilization and NFAT cell activation in primary mouse T cells. *J Exp Med* 2003; 197:1525-35; PMID:12782715; <http://dx.doi.org/10.1084/jem.20020234>
- Sun Z, Arendt CW, Ellmeier W, Schaeffer EM, Sunshine MJ, Gandhi L, Annes J, Petrzilka D, Kupfer A, Schwartzberg PL, et al. PKC- η is required for TCR-induced NF- κ B activation in mature but not in immature T lymphocytes. *Nature* 2000; 404:402-7; PMID:10746729; <http://dx.doi.org/10.1038/3500690>
- Isakov N, Altman A. Protein kinase C η in T cell activation. *Annu Rev Immunol* 2002; 20:761-94; PMID:11861617; <http://dx.doi.org/10.1146/annurev.immunol.20.100301.064807>
- Dunn GP, Bruce AT, Ikeda H, Old LJ, Schreiber RD. Cancer immunoeediting : from immunosurveillance to tumor escape. *Nature Immunol* 2002; 3:991-8; <http://dx.doi.org/10.1038/ni1102-991>
- Dunn GP, Koebel CM, Schreiber RD. Interferons, immunity and cancer immunoeediting. *Nature Rev Immunol* 2006; 6:836-48; <http://dx.doi.org/10.1038/nri1961>
- Koebel CM, Vermi W, Swann JB, Zerafa N, Rodig SJ, Old LJ, Smyth MJ, Schreiber RD. Adaptive immunity maintains occult cancer in an equilibrium state. *Nature* 2007; 450:903-7; PMID:18026089; <http://dx.doi.org/10.1038/nature06309>
- Villalba M, López-Royuela N, Krzywinska E, Rathore M, Hipskind R, Haouas H, Allende-Vega N. Chemical metabolic inhibitors for the treatment of blood-borne cancers. *Anti-Cancer Ag Med Chem* 2014; 14:223-32;
- Villalba M, Rathore MG, López-Royuela N, Krzywinska E, Garaude J, Allende-Vega N. From tumor cell metabolism to tumor immune escape. *Int J Biochem Cell Biol* 2013; 45:106-13; PMID:22568930; <http://dx.doi.org/10.1016/j.biocel.2012.04.024>
- Garaude J, Kaminski S, Charni S, Aguiló JL, Jacquet C, Plays M, Rodriguez F, Hernández J, Hipskind RA, Anel A, et al. Impaired anti-leukemic immune response in PKC η -deficient mice. *Mol Immunol* 2008; 45:3463-9; PMID:18462800; <http://dx.doi.org/10.1016/j.molimm.2008.03.016>
- Ljunggren HG, Karre K. Host resistance directed selectively against H-2-deficient lymphoma variants. Analysis of the mechanism. *J Exp Med* 1985; 162:1745-95; PMID:3877776; <http://dx.doi.org/10.1084/jem.162.6.1745>
- Terme M, Ullrich E, Delahaye NF, Chaput N, Zitvogel L. Natural killer cell-directed therapies: moving from unexpected results to successful strategies. *Nature Immunol* 2008; 9:486-94; <http://dx.doi.org/10.1038/ni1580>
- Vivier E, Tomasello E, Baratin M, Walzer T, Ugolini S. Functions of natural killer cells. *Nature Immunol* 2008; 9:503-10; <http://dx.doi.org/10.1038/ni1582>
- Pardo J, Balkow S, Anel A, Simon MM. Granzymes are critically involved in NK-mediated control of RMA-S tumor growth *in vivo*. *Eur J Immunol* 2002; 32:2881-6; PMID:12355441; [http://dx.doi.org/10.1002/1521-4141\(200210\)32:10<2881::AID-IMMU2881>3.0.CO;2-K](http://dx.doi.org/10.1002/1521-4141(200210)32:10<2881::AID-IMMU2881>3.0.CO;2-K)
- Van den Broek MF, Kägi D, Zinkernagel RM, Hengartner H. Perforin dependence of natural killer cell-mediated tumor control *in vivo*. *Eur J Immunol* 1995; 25:3514-6; PMID:8566046; <http://dx.doi.org/10.1002/eji.1830251246>
- Screpanti V, Wallin RPA, Ljunggren HG, Grandien A. A central role for death receptor-mediated apoptosis in the rejection of tumors by NK cells. *J Immunol* 2001; 167:2068-73; PMID:11489989; <http://dx.doi.org/10.4049/jimmunol.167.4.2068>
- Villalba M, Kasibhatla S, Genestier L, Mahboubi A, Green DR, Altman A. Protein kinase C- η cooperates with calcineurin to induce Fas ligand expression during activation-induced T cell death. *J Immunol* 1999; 163:5813-9; PMID:10570264
- Pardo J, Buferne M, Martínez-Lorenzo MJ, Naval J, Schmitt-Verhulst AM, Boyer C, Anel A. Differential implication of PKC isoforms in TCR/CD3-induced Fas ligand expression and in CTL degranulation. *Int Immunol* 2003; 15:1441-50; PMID:14645153; <http://dx.doi.org/10.1093/intimm/dxg141>
- Garrido F, Algarra I, García-Lora AM. The escape of cancer from T lymphocytes: immunoselection of MHC class I loss variants harboring structural-irreversible "hard" lesions. *Cancer Immunol Immunother* 2010; 59:1601-6; PMID:20625726; <http://dx.doi.org/10.1007/s00262-010-0893-2>
- Aguiló JL, Garaude J, Pardo J, Villalba M, Anel A. Protein kinase C- η is required for NK cell activation and *in vivo* control of tumor progression. *J Immunol* 2009; 182:1972-81; <http://dx.doi.org/10.4049/jimmunol.0801820>
- Anel A, Aguiló JL, Catalán E, Garaude J, Rathore MG, Pardo J, Villalba M. Protein kinase C- η (PKC- η) in natural killer cell function and anti-tumor immunity. *Front Immunol* 2012; 3:187; PMID:22783260; <http://dx.doi.org/10.3389/fimmu.2012.00187>
- Akazawa T, Ebihara T, Okuno M, Okuda Y, Shingai M, Tsujimura K, Takahashi T, Ikawa M, Okabe M, Inoue N, et al. Antitumor NK activation induced by the Toll-like receptor 3-TICAM-1 (TRIF) pathway in myeloid dendritic cells. *Proc Natl Acad Sci USA* 2007; 104:252-7; PMID:17190817; <http://dx.doi.org/10.1073/pnas.0605978104>
- Page KM, Chaudhary D, Goldman SJ, Kasaian MT. Natural killer cells from protein kinase C η mice stimulated with interleukin-12 are deficient in production of interferon- γ . *J Leukoc Biol* 2008; 83:1267-76; PMID:18263766; <http://dx.doi.org/10.1189/jlb.1107745>
- Carson WE, Giri JG, Lindemann MJ, Linett ML, Ahdieh M, Paxton R, Anderson D, Eisenmann J, Grabstein K, Caligiuri MA. Interleukin (IL) 15 is a novel cytokine that activates human natural killer cells via components of the IL-2 receptor. *J Exp Med* 1994; 180:1395-403; PMID:7523571; <http://dx.doi.org/10.1084/jem.180.4.1395>
- Cooper MA, Fehniger TA, VanDeusen JB, Waite RE, Liu Y, Aguila HL, Caligiuri MA. *In vivo* evidence for a dependence on interleukin 15 for survival of natural killer cells. *Blood* 2002; 100:3633-8; PMID:12393617; <http://dx.doi.org/10.1182/blood-2001-12-0293>
- Liu RB, Engels B, Arina A, Schreiber K, Hyjek E, Schietinger A, Binder DC, Butz E, Krausz T, Rowley DA, et al. Densely granulated murine NK cells eradicate large solid tumors. *Cancer Res* 2012; 72:1964-74; PMID:22374983; <http://dx.doi.org/10.1158/0008-5472.CAN-11-3208>
- Villalba M, Bi K, Hu J, Altman Y, Bushway P, Reits E, Neeffes J, Baier G, Abraham RT, Altman A. Translocation of PKC η in T cells is mediated by a nonconventional, PI3-K- and Vav-dependent pathway, but does not absolutely require phospholipase C. *J Cell Biol* 2002; 157:253-63; PMID:11956228; <http://dx.doi.org/10.1083/jcb.200201097>
- Ben Ahmed M, Belhadj Hmdia N, Moes N, Buyse S, Abledadhim M, Louzir H, Cerf-Bensussan N. IL-15 renders conventional lymphocytes resistant to suppressive functions of regulatory T cells through activation of the phosphatidylinositol 3-kinase pathway. *J Immunol* 2009; 182:6763-70; PMID:19454671; <http://dx.doi.org/10.4049/jimmunol.0801792>
- Brennan P, Babage JW, Burgering BMT, Gromer B, Reif K, D. A. C. Phosphatidylinositol 3-kinase couples the interleukin-2 receptor to the cell cycle regulator E2F. *Immunity* 1997; 7:679-89; PMID:9390691; [http://dx.doi.org/10.1016/S1074-7613\(00\)80388-X](http://dx.doi.org/10.1016/S1074-7613(00)80388-X)
- Truitt KE, Mills GB, Turck CW, Imboden JB. SH2-dependent association of phosphatidylinositol 3'-kinase 85-kDa regulatory subunit with the interleukin-2 receptor β chain. *J Biol Chem* 1994; 269:5937-43; PMID:7509794
- Uddin S, Fish E, Sher D, Gardziola C, White M, Platanias L. Activation of the phosphatidylinositol 3-kinase serine kinase by IFN- α . *J Immunol* 1997; 158:2390-7; PMID:9036989
- Premecz G, Markovits A, Bagi G, Farkas T, Földes I. Phospholipase C and phospholipase A₂ are involved in the antiviral activity of human interferon- α . *FEBS Lett* 1989; 249:257-60; PMID:2544450; [http://dx.doi.org/10.1016/0014-5793\(89\)80635-0](http://dx.doi.org/10.1016/0014-5793(89)80635-0)
- Srivastava KK, Batra S, Sassano A, Li Y, Majchrzak B, Kiyokawa H, Altman A, Fish EN, Platanias LC. Engagement of protein kinase C- η in interferon signaling in T-cells. *J Biol Chem* 2004; 279:29911-20; PMID:15150272; <http://dx.doi.org/10.1074/jbc.M401997200>
- Wang J, Campbell IL. Innate STAT1-dependent genomic response of neurons to the antiviral cytokine alpha interferon. *J Virol* 2005; 79:8295-302; PMID:15956575; <http://dx.doi.org/10.1128/JVI.79.13.8295-8302.2005>
- Fu YY. A transcription factor with SH2 and SH3 domains is directly activated by an interferon alpha-induced cytoplasmic protein tyrosine kinase(s). *Cell* 1992; 70:323-35.
- Hardy PO, Diallo TO, Matte C, Descoteaux A. Roles of phosphatidylinositol 3-kinase and p38 mitogen-activated protein kinase in the regulation of protein kinase C- α activation in interferon- γ -stimulated macrophages. *Immunology* 2009; 128:e652-e60; PMID:19740326; <http://dx.doi.org/10.1111/j.1365-2567.2009.03055.x>
- Deb D, Sassano A, Lekmine F, Majchrzak B, Verma A, Kambhampati S, Uddin S, Rahman A, Fish E, Platanias L. Activation of protein kinase C δ by IFN- γ . *J Immunol* 2003; 171:267-73; PMID:12817007; <http://dx.doi.org/10.4049/jimmunol.171.1.267>
- Swann JB, Hayakawa Y, Zerafa N, Sheelman KCF, Scott B, Schreiber RD, Hertzog P, Smyth MJ. Type I IFN contributes to NK cell homeostasis, activation, and antitumor function. *J Immunol* 2007; 178:7540-9; PMID:17548588; <http://dx.doi.org/10.4049/jimmunol.178.12.7540>
- Merino E, Abeyweera TP, Firth MA, Zawislak CL, Basu R, Liu X, Sun JC, Huse M. Protein kinase C- η clustering at immunological synapses amplifies effector responses in NK cells. *J Immunol* 2012; 189:4859-69; PMID:23077238; <http://dx.doi.org/10.4049/jimmunol.1200825>

43. Tassi I, Cella M, Presti R, Colucci A, Gilfillan S, Littman DR, Colonna M. NK cell activating receptors require PKC ζ for sustained signaling, transcriptional activation and IFN- γ secretion. *Blood* 2008; 112:4109-16; PMID:18784374; <http://dx.doi.org/10.1182/blood-2008-02-139527>
44. Grybko MJ, Pores-Fernando AT, Wurth GA, Zweifach A. Protein kinase C activity is required for cytotoxic T cell lytic granule exocytosis, but the ζ isoform does not play a preferential role. *J Leukoc Biol* 2007; 81:509-19; PMID:17077164; <http://dx.doi.org/10.1189/jlb.0206109>
45. Puente LG, He JS, Ostergaard HL. A novel PKC regulates ERK activation and degranulation of cytotoxic T lymphocytes: Plasticity in PKC regulation of ERK. *Eur J Immunol* 2006; 36:1009-18; PMID:16552708; <http://dx.doi.org/10.1002/eji.200535277>
46. Pardo J, Aguiló J, Anel A, Martin P, Joeckel L, Borner C, Wallich R, Müllbacher A, Froelich C, Simon M. The biology of cytotoxic cell granule exocytosis pathway: granzymes have evolved to induce cell death and inflammation. *Microbes & Infection* 2009; 11:452-9; PMID:19249384; <http://dx.doi.org/10.1016/j.micinf.2009.02.004>
47. Fehniger TA, Cai SF, Cao X, Bredemeyer AJ, Presti RM, French AR, Ley TJ. Acquisition of murine NK cell cytotoxicity requires the translation of a pre-existing pool of granzyme B and perforin mRNAs. *Immunity* 2007; 26:798-811; PMID:17540585; <http://dx.doi.org/10.1016/j.immuni.2007.04.010>
48. Lucas M, Schachterle W, Oberle K, Aichele P, Dieffenbach A. Dendritic cells prime natural killer cells by trans-presenting interleukin 15. *Immunity* 2007; 26:503-17; PMID:17398124; <http://dx.doi.org/10.1016/j.immuni.2007.03.006>
49. Gustafsson K, Ingelsten M, Bergqvist L, Nyström J, Andersson B, Karlsson-Parra A. Recruitment and activation of natural killer cells in vitro by a human dendritic cell vaccine. *Cancer Res* 2008; 68:5965-71; PMID:18632652; <http://dx.doi.org/10.1158/0008-5472.CAN-07-6494>
50. Robertson M. Role of chemokines in the biology of natural killer cells. *J Leukoc Biol* 2002; 71:173-83; PMID:11818437
51. Vujanovic L, Ballard W, Thorne SH, Vujanovic NL, Butterfield LH. Adenovirus-engineered human dendritic cells induce natural killer cell chemotaxis via CXCL8/IL-8 and CXCL10/IP-10. *Oncol Immunol* 2012; 1:448-57; <http://dx.doi.org/10.4161/onci.19788>
52. Sánchez-Martínez D, Krzywinska E, Rathore MG, Saument A, Cornilon A, López-Royuela N, Martínez-Lostao L, Ramírez-Labrada A, Lu Z, Rossi J, et al. All-trans retinoic acid (ATRA) induces miR-23a expression, decreases CTSC expression and granzyme B activity leading to impaired NK cell cytotoxicity. *Int J Biochem Cell Biol* 2014; 49:42-52; <http://dx.doi.org/10.1016/j.biocel.2014.01.003>
53. Bancerek J, Poss Z, Steinparzer I, Sedlyarov V, Pfaffenwimmer T, Mikulic I, Dölken L, Strobl B, Müller M, Taatjes D, et al. CDK8 kinase phosphorylates transcription factor STAT1 to selectively regulate the interferon response. *Immunity* 2013; 38:250-62; PMID:23352233; <http://dx.doi.org/10.1016/j.immuni.2012.10.017>
54. Putz E, Gotthardt D, Hoemann G, Csiszar A, Wirth S, Berger A, Straka E, Rigler D, Wallner B, Jamieson A, et al. CDK8-mediated STAT1-S727 phosphorylation restrains NK cell cytotoxicity and tumor surveillance. *Cell Rep* 2013; 4:437-44; PMID:23933255; <http://dx.doi.org/10.1016/j.celrep.2013.07.012>
55. Kiessling R, Klein E, Wigzell H. "Natural" killer cells in the mouse. I. Cytotoxic cells with specificity for Moloney leukemia cells. Specificity and distribution according to genotype. *Eur J Immunol* 1975; 5:112-7; PMID:1234049; <http://dx.doi.org/10.1002/eji.1830050208>
56. Murali-Krishna K, Altman J, Suresh M, Sourdive D, Zajac A, Miller J, Slansky J, Ahmed R. Counting antigen-specific CD8⁺ T cells: a re-evaluation of bystander activation during viral infection. *Immunity* 1998; 8:177-87; PMID:9491999; [http://dx.doi.org/10.1016/S1074-7613\(00\)80470-7](http://dx.doi.org/10.1016/S1074-7613(00)80470-7)
57. Pardo J, Wallich R, Ebnet K, Iden S, Zentgraf H, Martin P, Ekiciler A, Prins A, Müllbacher A, Huber M, et al. Granzyme B is expressed in mouse mast cells in vivo and in vitro and causes delayed cell death independent of perforin. *Cell Death Differ* 2007; 14:1768-79; PMID:17599099; <http://dx.doi.org/10.1038/sj.cdd.4402183>
58. Charni S, de Bettignies G, Rathore MG, Aguiló JJ, van den Elsen PJ, Haouzi D, Hipskind RA, Enriquez JA, Sanchez-Beato M, Pardo J, et al. Oxidative phosphorylation induces de novo expression of the MHC class I in tumor cells through the ERK5 pathway. *J Immunol* 2010; 185:3498-503; PMID:20729331; <http://dx.doi.org/10.4049/jimmunol.1001250>