



HAL
open science

In-depth characterization of bacterial and archaeal communities present in the abandoned Kettara pyrrhotite mine tailings (Morocco)

Odile Bruneel, N. Mghazli, R. Hakkou, I. Dahmani, A. Filali Maltouf, L. Sbabou

► To cite this version:

Odile Bruneel, N. Mghazli, R. Hakkou, I. Dahmani, A. Filali Maltouf, et al.. In-depth characterization of bacterial and archaeal communities present in the abandoned Kettara pyrrhotite mine tailings (Morocco). *Extremophiles*, 2017, 21 (4), pp.671-685. 10.1007/s00792-017-0933-3 . hal-03143128

HAL Id: hal-03143128

<https://hal.umontpellier.fr/hal-03143128>

Submitted on 8 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

In-depth characterization of bacterial and archaeal communities present in the abandoned Kettara pyrrhotite mine tailings (Morocco)

Odile Bruneel^{1,2} · N. Mghazli¹ · R. Hakkou³ · I. Dahmani¹ · A. Filali Maltouf¹ · L. Sbabou¹

Abstract In Morocco, pollution caused by closed mines continues to be a serious threat to the environment, like the generation of acid mine drainage. Mine drainage is produced by environmental and microbial oxidation of sulfur minerals originating from mine wastes. The fundamental role of microbial communities is well known, like implication of Fe-oxidizing and to a lesser extent S-oxidizing microorganism in bioleaching. However, the structure of the microbial communities varies a lot from one site to another, like diversity depends on many factors such as mineralogy, concentration of metals and metalloids or pH, etc. In this study, prokaryotic communities in the pyrrhotite-rich tailings of Kettara mine were characterized using the Illumina sequencing. In-depth phylogenetic analysis revealed a total of 12 phyla of bacteria and 1 phyla of Archaea. The majority of sequences belonged to the phylum of *Proteobacteria* and *Firmicutes* with a predominance

of *Bacillus*, *Pseudomonas* or *Corynebacterium* genera. Many microbial populations are implicated in the iron, sulfur and arsenic cycles, like *Acidiferrobacter*, *Leptospirillum*, or *Alicyclobacillus* in Fe; *Acidiferrobacter* and *Sulfobacillus* in S; and *Bacillus* or *Pseudomonas* in As. This is one of the first description of prokaryotic communities in pyrrhotite-rich mine tailings using high-throughput sequencing.

Keywords Microbial diversity · Pyrrhotite-rich tailings · Intermittent acid mine drainage · Illumina sequencing · Kettara mine

Introduction

The mining of metallic sulfide ores produces huge quantities of sulfide-rich mine wastes that generate acid mine drainage (AMD) when exposed to oxygen and water (Hallberg 2010). Although this process occurs naturally, mining operations like excavation and milling accelerate the process by increasing the exposure of sulfide minerals to water, oxygen and microorganisms (Simate and Ndlovu 2014). AMD generally contains high concentrations of sulfates, iron, metals (Pb, Cd, Zn), and metalloids like arsenic, with acid pH, that represent a threat to the environment and to human health with high environmental and socio-economic costs (Akcil and Koldas 2006; Hallberg 2010).

The overall reaction of the oxidation of pyrite (FeS₂, the most abundant sulfide mineral on Earth) can be written as: $\text{FeS}_2 + 14\text{Fe}^{3+} + 8\text{H}_2\text{O} \rightarrow 15\text{Fe}^{2+} + 2\text{SO}_4^{2-} + 16\text{H}^+$. In contrast to FeS₂ oxidation, acid-soluble metal sulfides like pyrrhotite (Fe_{1-x}S with x = 0–0.125) can be dissolved by the combined action of Fe(III) oxidative attacks (the most widespread oxidant in acidic environments) and proton

✉ Odile Bruneel
odile.bruneel@ird.fr

¹ Laboratoire de Microbiologie et Biologie Moléculaire, LMBM, Faculté des Sciences, Université Mohammed V, Av Ibn Batouta, BP1014, Rabat, Morocco

² Laboratoire HydroSciences Montpellier, UMR5569 (CNRS/IRD/UM), Université de Montpellier, CC0057 (MSE), 16, rue Auguste Broussonet, 34090 Montpellier, France

³ Laboratoire de Chimie des Matériaux et de l'Environnement (LCME), Faculté des Sciences et Technique Guéliz, Université de Cadi Ayyad, Avenue Abdelkarim Elkhattabi, Guéliz, P.O. Box 549, Marrakech, Morocco

attacks (Vera et al. 2013). The main role of microorganisms in leaching consists in the regeneration of Fe(III) ions (Schippers and Sand 1999), by microorganisms such as *Acidithiobacillus ferrooxidans* capable of increasing the iron oxidation rate one hundred to one million fold (Singer and Stumm 1970). Elemental sulfur can be also biologically oxidized to sulfuric acid thus contributing to the dissolution of pyrrhotite. Furthermore, S-oxidizing microorganisms can minimize the accumulation of elemental sulfur on the surface of the metal sulfide, which is known to reduce metal sulfide dissolution (reviewed in Vera et al. 2013).

Microbial populations in mine tailings have attracted considerable interest in the last decade. Indeed, knowledge of bioleaching microorganisms can be of great importance for the long-term prediction of mine waste and the development and monitoring of countermeasures to limit the pollution (Rawlings and Johnson 2007; Schippers et al. 2010; Chen et al. 2013). The microorganisms involved in sulfidic mine waste generating AMD are similar to those used in commercial leaching operations to recover precious and base metals, so understanding how they function could have a strong economic impact (Schippers et al. 2010). Concerning the bioleaching of metal sulfides, the best studied bacterial strains are *Acidithiobacillus ferrooxidans* (the first isolated) or *Leptospirillum ferrooxidans*. However, in recent decades research has revealed a much wider diversity of leaching bacteria than previously thought, with bacteria distributed among different phyla like *Proteobacteria* (including *Acidithiobacillus*, *Acidiferrobacter*, *Ferrovum*); *Nitrospirae* (*Leptospirillum*); *Firmicutes* (*Alicyclobacillus*, *Sulfobacillus*) or *Actinobacteria* (*Acidimicrobium*, *Ferrimicrobium* or *Ferritrix*, Vera et al. 2013). Many iron- and sulfur-oxidizing Archaea strains have also been found in the order of *Sulfolobales*, such as *Sulfolobus*, *Acidianus*, *Metallosphaera*, *Sulfurisphaera*, etc., or in the *Thermoplasmales* like *Ferroplasma* (Schippers et al. 2010; Vera et al. 2013).

Although the chemistry of AMD generation is well known, their characteristics vary greatly from one region to another and the production of AMD is a function of the geology of the mining site, the type of microorganisms present, the concentration of metals and metalloids, the temperature, the pH, and the presence of water and oxygen (Schippers et al. 2007; Chen et al. 2013; Simate and Ndlovu 2014). In the last decade, rapid advances in sequencing technology have revolutionized microbial ecology, from sequencing hundreds of 16S rRNA gene fragments per study with clone libraries to millions of fragments using next-generation sequencing (NGS) technologies like 454 or Illumina. This has greatly improved our understanding of the composition and diversity of microbial communities in the environment (Caporaso et al. 2011; Kozich et al. 2013).

The abandoned Kettara mine is a problematic mine site located near the city of Marrakech (Morocco), where inhabitants are directly exposed (Lghoul et al. 2014). The sulfide deposit is composed of major and minor lenses of massive pyrrhotite, with small amounts of sphalerite, galena, chalcopyrite, pyrite, arsenopyrite and glaucodot (Khalil et al. 2013). The substrate is composed of fractured and altered shale, which facilitates AMD infiltration into the groundwater. The groundwater table is located at a depth of only 10–20 m, and is contaminated by AMD with high concentration of sulfates and high conductivity (Lghoul et al. 2012; Khalil et al. 2013).

The aim of the present work was to characterize the diversity and taxonomic composition of the microbial communities (archaea and bacteria) using a MiSeq approach targeting the 16S rRNA gene and to investigate their spatial evolution in relation to the physicochemical structure of the tailings. Our results provide insights into the potential role of microorganisms in the formation of AMD at the abandoned Kettara pyrrhotite ore mine, which is located in a semi-arid environment.

Experimental procedures

Study area

Kettara pyrrhotite mine is located 30 km north-west of Marrakech (Morocco) and was exploited for the production of sulfuric acid from 1965 to 1982 (Hakkou et al. 2008a). The mine produced more than 5.2 million metric tons of pyrrhotite concentrate containing an average of 1.6–14.5 wt% sulfur, mainly as sulfide minerals (e.g., pyrrhotite, pyrite, chalcopyrite, galena, and sphalerite, Hakkou et al. 2008a; Khalil et al. 2013). Three million tons of mine residues were directly stored on a shale and sandstone fractured bedrock distributed over an area of approximately 16 ha, without any concern for the environment (Khalil et al. 2013). The tailings materials can be divided into two broad classes: coarse tailings (fine gravel) deposited on the 15-m-high dyke or on 1-m-high piles, and fine mine residues (silt) stored in the tailings pond (Hakkou et al. 2008a). The climate is semi-arid with mean annual rainfall of approximately 250 mm, characterized by short but intensive rainfall events (Lghoul et al. 2014). Sulfide minerals can be easily oxidized due to the high hydraulic permeability of the tailings, generating AMD after each rainfall event characterized by acid pH (ranging from 2.4 to 3.4) and very high concentrations of sulfate (8000–49600 mg l⁻¹), Fe (40–1650 mg l⁻¹), and Al (730–4170 mg l⁻¹, Lghoul et al. 2014). Results of laboratory experiments confirmed field observations indicating that fine and coarse tailings at Kettara are highly acid generating (Hakkou et al. 2008a, b).

The mining village of Kettara (with around 2000 inhabitants) is located downstream from the tailings pond. Groundwater, sampled from wells in Kettara village, is contaminated by AMD (Lghoul et al. 2012; Toughzaoui et al. 2015). Furthermore, windblown dust and sulfur emanations represent a real threat for the village inhabitants and farms in the vicinity (Babi et al. 2015).

Sample collection

Nine tailings samples were collected from the field site in September 2014, after a dry and hot summer, and were analyzed in the laboratory to determine their chemical composition. Sampling locations are shown in Fig. 1 and Table S1.


Three trenches were excavated to a depth of 60 and 120 cm in the tailings pond. In each trench, significant variations in color were observed, unoxidized tailings being grey and oxidized tailings yellowish-brown (Table S1). In trench 1, samples of fresh, unaltered fine tailings (T1-RF) were taken at a depth of 60–70 cm and samples of oxidized fine tailings (T1-RO) at a depth of 4–5 cm. In trench 2, unaltered fine tailings (T2-RF) were sampled at a depth of 40–50 cm, and samples of oxidized

fine tailings (T2-RO) at a depth of 4–5 cm. Trench 3 was excavated in the main dyke, corresponding to the coarse tailings whose oxidized zone was thicker than those observed in the fine tailings. Samples of unaltered coarse tailings (T3-S1F) were taken at a depth of 110–120 cm and samples of oxidized coarse tailings (T3-S1O) at a depth of 4–5 cm. Secondary precipitates (called D8) were sampled in the dyke and correspond to clear grey efflorescent salts. Samples K1 and K2 were taken at a depth of 4–5 cm in the 1-m-high piles of coarse tailings. Samples for chemical analysis were preserved at 4 °C until processing, and samples for DNA extraction were preserved on the field at 4 °C then stored at –80 °C until analysis.

Characterization of tailing samples

The chemical composition of the tailings was determined with a thermo scientific NITON X-ray fluorescence (XRF) analyzer (Thermo Scientific NITON 2008). The quality of the data was assessed using duplicate sample analyses and measurement accuracy was estimated at $\pm 5\%$ for all the elements analyzed. pH was analyzed according to the method of Mathieu and Pieltain (2003).

Fig. 1 Map of the Kettara mining site and location of the 9 sampling stations, sampled in September 2014. 1 T1-RF, 2 T1-RO, 3 T2-RF, 4 T2-RO, 5 T3-S1F, 6 T3-S1O, 7 D8, 8 K1 and 9 K2



DNA extraction, PCR amplification and Illumina MiSeq sequencing

Genomic DNA was extracted in triplicate from sediments using the PowerSoil DNA Isolation Kit and analyzed following the manufacturer's recommendations (MoBio Laboratories Inc., Carlsbad, CA, USA). All extracted genomic DNA samples were stored at -20°C until further analysis.

The genes coding for 16S rRNA were amplified by PCR for multiplexed Illumina using barcoded primers. The universal primer set 515F (5'-GTGCCAGCMGCCGCGG-TAA-3') and 806R (5'-GGACTACVSGGGTATCTAAT-3') designed to target nearly all bacterial and archaeal taxa (Bates et al. 2011) was used to amplify a 291-bp region targeting the V4 region of 16S rRNA genes.

A PCR reaction was performed on all three extracted DNA samples using a different barcode for each sample. The reaction mixture was used in a 28-cycle PCR using the HotStarTaq Plus Master Mix Kit (Qiagen, USA) with the following program: 3 min at 94°C , followed by 28 cycles at 94°C for 30 s, 53°C for 40 s and 72°C for 1 min, and a final elongation step at 72°C for 5 min. After amplification, the PCR products were checked in 2% agarose gel to determine the success of amplification and the relative intensity of the bands. Samples were pooled and purified using calibrated Ampure XP beads. Then the pooled and purified PCR product was used to prepare a DNA library following the Illumina TruSeq DNA library preparation protocol. Sequencing was performed at MR DNA (<http://www.mrdnab.com>, Shallowater, TX, USA) on an Illumina MiSeq 2x300 bp, following the manufacturer's guidelines.

Processing and taxonomic classification

Sequence data were processed (paired end reads were merged and fasta and qual files produced) using MR DNA analysis pipeline (MR DNA, Shallowater, TX, USA). Bioinformatics analyses were then processed using the mothur platform, version 1.35.1 (Schloss et al. 2009). The raw reads obtained from sequencing were quality trimmed (with a minimum Phred average quality score of 35 over a 50-bp window), with exact barcode and primer matching. Reads containing ambiguous characters and less than 150 nucleotides were removed. Identical sequences were grouped, and representative sequences were aligned against the SILVA bacterial and archaeal reference database using the Needleman–Wunsch algorithm (Needleman and Wunsch 1970). Only sequences aligned in the same area were kept. The resulting sequences were denoised using a pre-clustering method that allowed one mismatch (Huse et al. 2010; Schloss et al. 2011). Chimeric sequences were detected and removed using chimera uchime (Edgar et al. 2011) and sequences that did not classify to the level of kingdom or

that classified as Eukaryota, chloroplasts, or mitochondria were deleted. The remaining high-quality sequences were used to generate a distance matrix and were clustered into operational taxonomic units (OTUs) defined with a cutoff of 97% using the average neighbor algorithm. Singletons were culled. The OTUs were phylogenetically classified to genus level using the naive Bayesian classifier (80% confidence threshold) against a manually curated database derived from the RDP reference file (<http://rdp.cme.msu.edu>) and implemented in mothur. To obtain comparable data and to equalize the depth of sequencing, samples were downsized to 17557 reads by random re-sampling in triplicate (i.e., 52671 per site).

Estimation of diversity and statistical analysis

Diversity indices

Nonparametric Chao1 and Shannon alpha diversity estimates as well as the coverage and rarefaction curves for each sampling site were calculated using mothur software at 97% sequence similarity.

Principal component analyses

Statistical analyses were performed with R 3.3.1 (2016-06-21) (R Development Core Team 2012) using the ade4 and Vegan packages. Principal component analysis (PCA) was used to explore variations in the chemical composition of our set of environmental variables.

Non Metric Multidimensional Scaling (NMDS) was then used to graphically depict differences between microbial communities. The significance of the observed clustering of samples on the ordination plot was assessed by an Analysis OF SIMilarity (ANOSIM, 999 permutations). Environmental variables were fitted to the NMDS ordinations as vectors with the 'envfit' function of the 'vegan' package.

Results

Physicochemical characteristics of the tailing samples

The main physicochemical characteristics of the tailings samples collected from the different stations are listed in Table S2. The Kettara mine tailings were characterized by low pH (from 1.83 to 4.81) and by extremely high concentrations of Fe (up to 403561 mg kg^{-1}), sulfur (up to 109013 mg kg^{-1}), Cu (up to 2794 mg kg^{-1}), As (up to 344 mg kg^{-1}), or Cr (up to 436 mg kg^{-1}). These values are higher than many of the geochemical background values established for Kettara mine by Khalil

et al. (2013) ($\text{Cu} \approx 43.8 \text{ mg kg}^{-1}$, $\text{Pb} \approx 21.8 \text{ mg kg}^{-1}$, $\text{Zn} \approx 102.6 \text{ mg kg}^{-1}$, $\text{As} \approx 13.9 \text{ mg kg}^{-1}$ and $\text{Fe} \approx 56.978 \text{ mg kg}^{-1}$).

Table S2 shows that in the trenches, the pH was always lower at the surface than at depth which, logically, would be linked with oxidation of pyrrhotite in the surface tailings impoundment that is more subject to weathering than deeper layers. The concentrations of Fe and S varied with depth and the values of samples taken in the oxidized zone near the surface were always slightly higher than those of the samples from the deeper unoxidized zone. The highest concentration of Fe, S, and Cr were measured in the oxidized zone of trench T1. Sampling site D8 contained the highest concentration of Si, As and high concentrations of S, but also the lowest concentrations of Fe, Zn, Cu Cr, Mn, Al and Mg associated with the lowest pH (1.83) compared to the other sampling sites. The chemical composition of the coarse tailings in the dyke (T3-S1F and T3-S1O) and the piles (K1–K2) was quite similar.

Principal component analysis, describing the variations of the chemical parameters of the 9 tailings mine samplings, have identified around four groups: the secondary precipitates (D8), the 2 oxidized (T1-RO and T2-RO, in surface) and unoxidized (T1-RF and T2-RF, in depth) parts of the fine tailings and the coarse tailings (T3-S1O, T3S1F, K1 and K2, Fig. 2). The first PCA axis, explaining around 53% of the variations, separated the tailing sample D8 and the oxidized (T1-RO and T2-RO) and unoxidized (T1-RF and T2-RF) parts of the fine tailings. The second PCA axis, explaining around 25% of the variations, separated the tailing sample D8 and the oxidized and unoxidized parts of

the fine tailings. The chemical composition of the oxidized parts of the fine tailings (T1-RO and T2-RO) seems positively correlated with S, and the chemical composition of the unoxidized parts of the fine tailings (T1-RF and T2-RF) seems positively correlated with Zn, Cu, Mn, Mg and pH. The chemical composition of the sample D8 seems positively correlated with As and Si and negatively correlated with Fe and Cr. T1-RF and T2-RF contained the most important concentration of Al (60130 and 75557 mg kg^{-1} , respectively) compared to T1-RO, T2-RO and D8 (around 10000 mg kg^{-1}).

Sequencing results and diversity indices

A total of 1584954 good-quality reads with a mean length of 272 bp were obtained from the nine samples through MiSeq sequencing analysis. The number of sequences per sample was then made equal by random resampling (17557 sequences in triplicate, i.e., 52671 sequences per site) giving 474039 good-quality reads (Table 1). The clustering of these sequences allowed the identification of 430 OTUs defined with 97% identity.

The rarefaction curves in Figure S1 tend to reach an asymptote for all of the samples, except for T3-S1O, suggesting that the majority of the microbial phylotypes present in each sample were identified. This was confirmed by the very high Good's coverage (100%) for all samples (Table 1), which provides an estimate of sampling completeness. However, the total number of OTUs for each sample estimated by the Chao1 estimator, indicated that the richness of the microbial communities was a little higher

Fig. 2 Principal component analysis (PCA) analysis of the Kettara mine tailings and associated environmental variables




Table 1 Estimated OTU richness, diversity indices and estimated sample coverage for the nine 16S rRNA gene libraries

Sampling stations	No. of normalized sequences	No. of OTUs ^a	Good's coverage ^b	Shannon diversity ^c	Chao1 richness
T1-RF	52671	137	100	3.32 (3.31; 3.32)	183 (158; 241)
T1-RO	52671	147	100	2.86 (2.85; 2.87)	203 (174; 264)
T2-RF	52671	127	100	3.57 (3.56; 3.58)	148 (136; 180)
T2-RO	52671	143	100	3.23 (3.23; 3.24)	205 (173; 272)
T3-S1F	52671	139	100	3.24 (3.23; 3.25)	178 (157; 225)
T3-S1O	52671	229	100	2.28 (2.27; 2.30)	286 (258; 340)
D8	52671	166	100	3.18 (3.17; 3.19)	201 (182; 240)
K1	52671	148	100	2.81 (2.80; 2.83)	178 (161; 217)
K2	52671	132	100	1.49 (1.47; 1.51)	163 (145; 209)

Results are presented for the normalized data, randomly resampled to have an equal sample size

Values in brackets are 95% confidence intervals

^a OTUs were defined at 97% cutoff

^b Sum of probabilities of observed classes calculated as $(1 - (n/N))$, where n is the number of singleton sequences and N is the total number of sequences

^c Takes into account the number and evenness of species

than the observed number of OTUs, ranging from 148 for T2-RF to 286 for T3-S1O, suggesting the presence of a few more endemic taxa. The Shannon index, which measures the diversity of the microbial communities, varied slightly in the tailings (Table 1). The least microbial diversity was observed at K2 (1.49) and the highest at T2-RF (3.57).

Taxonomic composition at Kettara pyrrhotite mine and spatial variation among the sampling sites

In this study, the sequences of bacteria represented around 97% of the total sequences compared to those of Archaea (only 12646 sequences, Table 2). For the bacteria, 313825 of the 461393 qualified reads (68%) were assigned to a known phylum of bacteria (Table 2) at a confidence threshold of 80%. For Archaea, only 47% of reads could be assigned to a known phylum. The largest proportion of unassigned sequences was mainly found in the coarse tailings and represented 85% of the unclassified sequences of bacteria (around 10% for T3-S1F, 28% for T3-S1O, 19% for K1 and 29% for K2) and around 99% of Archaea (mainly limited to T3-S1F sampling site). In all the samples, a total of 12 phyla of bacteria and one phylum of Archaea were identified. Sixty-seven percent of sequences of bacteria clustered in 4 phyla: *Proteobacteria* (29%) and *Firmicutes* (25%), followed by *Actinobacteria* (10%) and *Bacteroidetes* (3%, Table 2). The eight additional phyla were each represented by less than 1% of the sequences and belonged to *Nitrospirae*, *Planctomycetes*, *Fusobacteria*, *Chloroflexi*, *Acidobacteria*, *Verrucomicrobia*, *Deinococcus-Thermus* and *Gemmatimonadetes*. All the sequences of Archaea were assigned to the phylum of *Euryarchaeota*.

The OTUs were assigned to 87 genera of bacteria and two genera of Archaea (Table 3). The unclassified sequences of bacteria represented 43% of the total sequences of bacteria. Only 18 genera had sequences representing more than 1% of the total number of sequences of bacteria. The dominant bacteria genera were represented by sequences of *Bacillus* (representing around 8% of the total sequences of bacteria), followed by the genus *Pseudomonas* (5%) and *Corynebacterium* (5%). The unclassified sequences of Archaea represented 53% of the total sequences of Archaea (Table 3). The two genera of Archaea, *Methanobacterium* and *Methanosaeta* represented, respectively, around 46 and 1% of the total sequences of Archaea.

Nonmetric MultiDimensional Scaling (NMDS) analysis was used to show the relationship between diversity patterns of microbial communities and environmental parameters (Fig. 3). The NMDS ordination showed a separation of the microbial communities along the first ordination axis, between the fine tailing (T1 and T2) and the secondary precipitate (D8), opposed to the coarse tailing (T3, K1 and K2). This analysis showed also that the main parameter discriminating our microbial communities was the concentration of Al ($r^2 = 0.606$, $P = 0.040$, Fig. 3; Table S2). *Bacillus* seemed more associated with the site T1-RO while *Enterococcus* and *Burkholderia* with the site D8 and *Staphylococcus* with the samples T2 (T2-RF and T2-RO).

Discussion

The aim of this study was to perform an in-depth characterization of the bacterial and archaeal diversity and their spatial distribution in nine different tailings of the Kettara

Table 2 Distribution of normalized sequences of bacterial and archaeal 16S rRNA genes from each sample assigned to different phyla

Phylum	Total	T1-RF	T1-RO	T2-RF	T2-RO	T3-SIF	T3-SIO	D8	K1	K2
Bacteria										
Unclassified	147568 (32.0)	4206 (2.9)	2678 (1.8)	744 (0.5)	7944 (5.4)	15227 (10.3)	41389 (28.0)	4722 (3.2)	28204 (19.1)	42454 (28.8)
<i>Proteobacteria</i>	132792 (28.8)	23359 (17.6)	15619 (11.8)	20185 (15.2)	18628 (14.0)	13455 (10.1)	6462 (4.9)	15882 (12.0)	13085 (9.9)	6117 (4.6)
<i>Firmicutes</i>	115664 (25.1)	14198 (12.3)	29009 (25.1)	13884 (12.0)	19320 (16.7)	5730 (5.0)	2974 (2.6)	20352 (17.6)	7899 (6.8)	2298 (2.0)
<i>Actinobacteria</i>	46547 (10.1)	7462 (16.0)	5187 (11.1)	10677 (22.9)	3474 (7.5)	8905 (19.1)	1270 (2.7)	6207 (13.3)	1990 (4.3)	1375 (3.0)
<i>Bacteroidetes</i>	14761 (3.2)	3439 (23.3)	165 (1.1)	4322 (29.3)	3295 (22.3)	8 (0.1)	74 (0.5)	2220 (15.0)	861 (5.8)	377 (2.6)
<i>Nitrospirae</i>	1641 (0.4)	3 (0.2)	2 (0.1)	1 (0.1)	0 (0.0)	1633 (99.5)	0 (0.0)	1 (0.1)	1 (0.1)	0 (0.0)
<i>Planctomycetes</i>	1210 (0.3)	0 (0.0)	1 (0.1)	1 (0.1)	6 (0.5)	777 (64.2)	420 (34.7)	2 (0.2)	3 (0.2)	0 (0.0)
<i>Fusobacteria</i>	736 (0.2)	0 (0.0)	1 (0.1)	731 (99.3)	1 (0.1)	0 (0.0)	0 (0.0)	2 (0.3)	1 (0.1)	0 (0.0)
<i>Chloroflexi</i>	388 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	232 (59.8)	45 (11.6)	1 (0.3)	109 (28.1)	1 (0.3)
<i>Acidobacteria</i>	66 (<0.1)	3 (4.5)	2 (3.0)	1 (1.5)	1 (1.5)	1 (1.5)	10 (15.2)	2 (3.0)	0 (0.0)	46 (69.7)
<i>Verrucomicrobia</i>	13 (<0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	9 (69.2)	4 (30.8)	0 (0.0)	0 (0.0)
<i>Deinococcus-Thermus</i>	6 (<0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	6 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
<i>Gemmatimonadetes</i>	1 (<0.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
Archaea										
Unclassified	6719 (53.1)	1 (<0.1)	3 (<0.1)	2 (<0.1)	2 (<0.1)	6699 (99.7)	7 (0.1)	2 (<0.1)	1 (<0.1)	2 (<0.1)
<i>Euryarchaeota</i>	5927 (46.9)	0 (0.0)	4 (0.1)	2123 (35.8)	0 (0.0)	4 (0.1)	4 (0.1)	3274 (55.2)	517 (8.7)	1 (<0.1)

The number in brackets represents the relative abundance (in %)

Table 3 Distribution of the relative abundance (in %) of bacterial and archaeal 16S rRNA genes from each samples assigned to different genus

Genus	Total	T1-RF	T1-RO	T2-RF	T2-RO	T3-S1F	T3-S1O	D8	K1	K2
Bacteria										
Unclassified	43.3	8.3	6.9	4.1	7.7	10.9	21.9	2.9	15.6	21.6
<i>Bacillus</i>	7.8	4.9	49.5	11.4	9.6	13.9	0.3	1.9	5.8	2.7
<i>Pseudomonas</i>	5.3	17.4	14.1	9.8	13.1	4.8	1.9	11.0	16.4	11.4
<i>Corynebacterium</i>	4.7	15.9	18.2	24.8	0.2	16.7	0.2	17.5	1.7	5.0
<i>Staphylococcus</i>	3.5	21.4	0.4	39.1	26.0	2.2	0.1	7.6	3.1	0.1
<i>Enterococcus</i>	2.8	0.4	0.5	0.2	18.1	0.2	0.6	64.2	12.1	3.7
<i>Saccharibacillus</i>	2.4	29.6	3.7	9.9	22.4	2.6	0.2	25.7	5.6	0.2
<i>Burkholderia</i>	2.0	0.2	0.1	11.0	2.0	0.1	0.2	74.5	10.0	1.9
<i>Cloacibacterium</i>	2.0	22.1	0.1	34.1	20.9	0.0	0.5	15.1	6.4	0.8
<i>Stenotrophomonas</i>	1.9	0.1	0.2	8.0	65.0	1.8	0.0	6.1	18.8	0.0
<i>Acidiferrobacter</i>	1.8	0.2	0.3	0.2	0.2	66.2	29.8	0.3	0.2	2.6
<i>Alcaligenes</i>	1.8	54.2	0.2	9.0	0.1	2.7	0.1	20.1	13.6	0.0
<i>Agrobacterium</i>	1.7	14.2	36.2	21.8	0.2	23.8	0.1	0.1	2.6	0.9
<i>Variovorax</i>	1.6	27.0	21.2	5.7	9.9	8.0	2.5	5.5	19.6	0.8
<i>Micrococcus</i>	1.5	10.9	15.5	24.2	34.3	0.1	0.1	0.2	13.8	0.9
<i>Phyllobacterium</i>	1.3	0.2	9.1	50.1	25.5	8.4	2.1	0.1	0.2	4.3
<i>Acidiphilium</i>	1.1	0.2	0.2	0.3	0.3	0.3	49.3	0.2	23.8	25.4
<i>Thermicanus</i>	1.0	8.4	35.4	0.2	17.5	0.0	0.3	33.6	4.2	0.4
<i>Geobacillus</i>	1.0	1.3	0.2	0.2	83.2	0.0	0.3	14.6	0.1	0.1
<i>Enterobacter</i>	0.9	2.4	72.5	0.2	16.6	2.8	3.1	0.3	2.1	0.0
<i>Herbaspirillum</i>	0.7	0.2	6.7	0.2	60.2	12.1	0.2	0.1	19.0	1.4
<i>Brevibacterium</i>	0.6	0.1	0.0	50.1	0.1	0.2	0.1	46.4	2.8	0.1
<i>Alicyclobacillus</i>	0.6	0.3	0.5	0.4	0.3	0.1	58.1	0.4	27.5	12.4
<i>Ruminococcus</i>	0.5	0.2	0.0	0.0	0.1	0.0	0.0	95.6	4.0	0.0
<i>Lactobacillus</i>	0.4	99.7	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0
<i>Moraxella</i>	0.4	0.3	0.2	22.6	0.1	0.1	1.4	66.5	8.7	0.0
<i>Acinetobacter</i>	0.4	0.3	13.8	39.1	0.3	33.6	1.7	0.3	6.7	4.3
<i>Alcanivorax</i>	0.4	76.9	0.1	19.2	0.1	0.2	0.3	1.9	1.3	0.1
<i>Escherichia</i>	0.4	7.2	29.8	60.9	0.2	0.1	0.2	0.0	0.6	1.1
<i>Sphingobacterium</i>	0.4	55.6	0.1	0.1	0.1	0.1	0.7	22.6	4.2	16.6
<i>Actinoallomurus</i>	0.4	0.1	0.1	0.1	0.1	99.2	0.1	0.1	0.2	0.1
<i>Streptococcus</i>	0.4	57.7	0.3	0.9	36.2	0.1	1.4	0.8	0.2	2.5
<i>Bifidobacterium</i>	0.4	0.2	0.1	0.1	35.4	0.0	0.1	55.7	0.2	8.2
<i>Leptospirillum</i>	0.4	0.2	0.1	0.1	0.0	99.5	0.0	0.1	0.1	0.0
<i>Brevundimonas</i>	0.3	0.7	0.0	98.1	0.0	0.0	0.0	1.2	0.0	0.0
<i>Aeromonas</i>	0.3	0.1	0.1	0.0	11.9	0.1	0.0	73.1	0.0	14.7
<i>Weissella</i>	0.3	0.0	0.0	0.1	0.1	0.1	0.1	99.5	0.1	0.0
<i>Legionella</i>	0.3	0.2	0.0	99.3	0.1	0.0	0.2	0.1	0.2	0.1
<i>Novosphingobium</i>	0.3	59.1	0.2	0.1	0.2	0.1	0.0	0.2	40.0	0.0
<i>Cereibacter</i>	0.2	99.3	0.4	0.2	0.0	0.1	0.0	0.1	0.0	0.0
<i>Tepidimonas</i>	0.2	32.6	0.2	61.2	0.1	0.1	0.6	5.2	0.0	0.0
<i>Anaerococcus</i>	0.2	0.3	0.4	63.7	0.1	0.0	0.2	28.0	7.2	0.0
<i>Tumebacillus</i>	0.2	0.1	0.0	0.0	0.1	0.1	0.5	99.2	0.0	0.0
<i>Paracoccus</i>	0.2	69.7	0.2	29.7	0.1	0.0	0.0	0.1	0.0	0.1
<i>Sulfobacillus</i>	0.2	0.1	0.4	0.0	0.3	1.3	96.4	0.3	0.5	0.9
<i>Achromobacter</i>	0.2	0.6	16.5	67.5	0.0	0.3	15.0	0.0	0.0	0.1
<i>Massilia</i>	0.2	0.1	91.9	0.0	0.0	0.3	0.1	0.0	0.1	7.4
Archaea										
Unclassified	53.1	0.0	0.1	16.8	0.0	53.0	0.1	25.9	4.1	0.0

Table 3 continued

Genus	Total	T1-RF	T1-RO	T2-RF	T2-RO	T3-S1F	T3-S1O	D8	K1	K2
<i>Methanobacterium</i>	46.3	0.0	0.1	36.2	0.0	0.1	0.1	54.8	8.8	0.0
<i>Methanosaeta</i>	0.5	0.0	0.0	1.5	0.0	0.0	0.0	98.5	0.0	0.0

Only genera comprising more than 700 sequences are presented




Fig. 3 Non-metric multidimensional scaling (NMDS) ordination plot derived from pairwise matrix distances of microbial communities for each sampling site. The stress values were <0.2 which indicates that these data were well-represented by the two dimensional representation. The angle and length of the vector indicate the direction and strength of the variable. *Arrows* are the projections of possible explanation variables obtained by vector fitting. Only correlations with $p < 0.05$ were indicated. The r^2 correlation coefficient and the p values are presented in Table S2. The 10 most abundant OTUs were also presented

pyrrhotite mine producing AMD, to better understand the potential role of microorganisms in the formation of mine drainage. Many studies have investigated the microbial and community structure of sulfidic mine waste (Schippers et al. 2010; Chen et al. 2013; Liu et al. 2014), but most of them mainly sampled pyrite tailings and little is known about the microbial communities inhabiting pyrrhotite mine tailings (Schippers et al. 2007; Kock and Schippers 2008; Korehi et al. 2014). To our knowledge, this is the first investigation of the microbial structure of pyrrhotite-rich mine tailings using next-generation sequencing.

Microbial diversity and taxonomic composition of the sediments

This work has revealed a higher microbial diversity in acid generating mine tailings than those reported in studies based on clone library analyses (Diaby et al. 2007; Tan et al. 2008; Huang et al. 2011) but a lower diversity than those using next-generation sequencing (Chen et al. 2013; Liu et al. 2014). The rarefaction curves and the coverage suggested an identification of the majority of the microbial communities. The high level of sequences of bacteria

and Archaea that could not be classified at the phylum level suggested that many novel microorganisms were also detected.

More than 69 genera of bacteria had a number of sequences representing less than 1% of the total number of sequences of bacteria, i.e., corresponding to the “rare biosphere”. The ecological and functional roles of this rare biosphere, encountered in quite large proportions in most studies since the arrival of next-generation sequencing, are still not well understood (Galand et al. 2009). It is hypothesized that such organisms could become dominant under changing environmental conditions and that their functional redundancy could allow biogeochemical processes to be maintained (Bachy and Worden 2014). Other studies question the accuracy of OTU richness estimates resulting from high-throughput sequencing, which could correspond to sequencing errors or could be generated by the use of inappropriate algorithms (Huse et al. 2010).

The dominant bacteria genera were related to *Bacillus* (representing around 8% of the total sequences of bacteria), followed by the genus *Pseudomonas* (5%), *Corynebacterium* (5%) and *Staphylococcus* (4%). These genera have already been detected in mine dumps or heaps, sometimes as dominant populations (Schippers et al. 2010; Chen et al. 2013).

Ecological role and potential impact on the environment

Bioleaching of metal sulfides is performed by a diverse group of microorganisms, but metal sulfide dissolution is mainly mediated by Fe(III) and to a certain extent by sulfur-oxidizing communities thanks to their capacity to reduce the accumulation of sulfur on the surface of metal sulfides and through the generation of protons that dissolve acid-soluble metal sulfides like pyrrhotite (Vera et al. 2013).

Concerning the biogeochemical cycling of iron in the Kettara mine, two microorganisms well known for their leading role in the bioleaching and formation of AMD, were identified: *Acidiferrobacter* and *Leptospirillum* representing, respectively, 2% and less than 0.4% of the total bacterial sequences. *Acidiferrobacter*, formerly called “*Thiobacillus ferrooxidans* m-1”, is an acidophilic iron-oxidizing bacterium (Hallberg et al. 2011). Although the majority of *Alicyclobacillus* species mainly utilize organic

compounds, different species are also able to oxidize ferrous iron for growth, including *Alicyclobacillus ferrooxidans*, *A. disulfidooxidans* and *A. tolerans* (Karavaiko et al. 2005; Jiang et al. 2008). This genus, forming endospores under adverse environmental or nutritional conditions, representing around 1% of the total sequences of bacteria identified in Kettara mine, is known as leaching bacteria (Vera et al. 2013). Another acidophilic and spore forming genus, known for its active role in the oxidation of sulfide minerals is the genus *Sulfobacillus* containing many isolates identified as iron oxidizers (Norris et al. 1996; Justice et al. 2014). This genus represented 0.2% of the sequences identified in Kettara mine. Some species of *Pseudomonas*, like *P. stutzeri* (Weber et al. 2006; Muehe et al. 2009) or *Pseudomonas* sp. GE-1 forming ferrihydrite (Xiu et al. 2015), are able to oxidize iron. Other genera like *Paracoccus*, representing 0.2% of the total bacteria identified in Kettara mine such as *Paracoccus ferrooxidans*, was also shown to oxidize iron (Kumaraswamy et al. 2006).

Fe(III)-reducing microorganisms are known to dissolve Fe(III)(hydr)oxides, thereby releasing adsorbed or precipitated metals (Schippers et al., 2010). *Acidiferrobacter* is known to grow by ferric iron respiration (Hallberg et al., 2011) like some strains of *Sulfobacillus* (*Sulfobacillus thermosulfidooxidans*, *Sulfobacillus acidophilus*, Bridge and Johnson 1998; Justice et al. 2014). Many other genera are involved in the reduction of Fe(III) including *Alicyclobacillus* (Weber et al. 2006; *A. tolerans*, Karavaiko et al. 2005); *Bacillus* (Schippers et al. 2010 including *Bacillus infernus*, Weber et al. 2006); *Pseudomonas* (Schippers et al. 2010, *Pseudomonas stutzeri*, Wang et al. 2014a); *Acidiphilium* (an acidophilic genus quite common in AMD that represented 1% of the total sequence of bacteria in Kettara mine, like *Acidiphilium acidophilum*, Johnson and Bridge 2002; Weber et al. 2006) or *Aeromonas* (*Aeromonas hydrophila*, Weber et al. 2006; Ventura et al. 2015). This species, which represented 0.3% of the sequences of bacteria identified at Kettara mine, is able to use different Fe(III)-oxyhydroxides, some weathering compounds commonly retrieved at mine sites like ferrihydrite, or jarosite as sole source of iron (Garcia Balboa et al. 2010).

Numerous microorganisms are known to be implicated in the redox transformation of sulfur, using elemental sulfur or inorganic sulfur compounds as electron donors or acceptors (Dopson and Johnson 2012). In the Kettara tailings, different populations able to oxidize the reduced inorganic sulfur compounds were retrieved, including *Acidiferrobacter* (Hallberg et al. 2011); *Sulfobacillus* (Norris et al. 1996; Justice et al. 2014); *Alicyclobacillus* (*A. disulfidooxidans* or *A. ferrooxidans*, Karavaiko et al. 2005; Jiang et al. 2008); *Pseudomonas* (Turner 1954, *P. stutzeri*, Sorokin et al. 1999; Mahmood et al. 2009; *Pseudomonas* sp. C27, Xu et al. 2016) or *Paracoccus* (*P. denitrificans*, Friedrich and

Mitrenga 1981; *P. versutus* (formerly *Thiobacillus versutus*, Katayama et al. 1995), *P. ferrooxidans* (Kumaraswamy et al. 2006) or *P. pantotrophus*, Vikromvarasiri et al. 2015). Chemolithotrophic growth with elemental sulfur or thiosulphate as energy source had also been observed in the genus *Acidiphilium* but differs from species to species (Okamura et al. 2015). Species of the genus *Alcaligenes* (representing 2% in the Kettara tailings) can also grow autotrophically by using sulfur compounds as sole energy sources (Anandham et al. 2008; Luo et al. 2013). Other species, less encountered in mine environments, can also oxidize reduced inorganic sulfur compounds like *Bacillus* (*Bacillus pumilus*, *Bacillus subtilis*, *Bacillus megaterium*) and *Micrococcus* spp. (Behera et al. 2014) or *Corynebacterium* (Omori et al. 1992). Genus of archaea like *Methanobacterium*, possess also the capacity to grow with sulfide or elemental sulfur (Zeikus and Wolfe 1972; Bhatnagar et al. 1984).

Microorganisms like *Clostridium* (representing less than 0.1% of total bacteria at Kettara mine) appear also able to reduce polysulfide to hydrogen sulfide (Takahashi et al. 2010), such as *Clostridium sulfidigenes*, a mesophilic, thiosulfate and sulfur reducing bacterium (Sallam and Steinbüchel 2009).

Among all the toxic elements present at the Kettara site, arsenic is of particular concern due to the serious environmental and health impact of this metalloid on living organisms (Mandal and Suzuki 2002; Abdul et al. 2015). Along with abiotic processes, the activity of As transforming microorganisms has a major influence on the behavior of this element because different As species present differences in their toxicity, solubility, mobility, and bioavailability depending on its state of oxidation (Macur et al. 2004; Sarkar et al. 2014). For example, As(III) can be oxidized in the less soluble and less toxic form As(V), thus redox transformations of As are of great importance to predict its fate in the environment (Oremland and Stolz 2003; Yamamura and Amachi 2014). In the pyrrhotite-rich Kettara tailings, containing high concentration of arsenic (up to 344 mg kg⁻¹), we have identified many genera, which contained species known to oxidize As. It is the case of the two predominant populations in Kettara, *Bacillus* (Fisher and Hollibaugh 2008, *Bacillus firmus*, Bachate et al. 2013) and *Pseudomonas* (Turner 1954, *P. stutzeri*, *P. arsenitoxidans* or *P. arsenoxydans*, etc., Das et al. 2014; Bahar et al. 2013a). It is also the case of different species, representing around 2% of sequences of bacteria identified at Kettara mine, such as *Stenotrophomonas* (Bahar et al. 2012, 2013a); *Agrobacterium* (Santini et al. 2000, *Agrobacterium tumefaciens*, Macur et al. 2004); *Variovorax* (*Variovorax paradoxus*, Macur et al. 2004; Bahar et al. 2013b); *Burkholderia* (*Burkholderia cepacia*, Campos et al. 2009) or *Alcaligenes* (Osborne and Ehrlich (1976),

including *Alcaligenes faecalis*, Bahar et al. 2013a). Minor species, representing less than 0.4% at Kettara mine, had also this capacity, like *Acinetobacter* (Chang et al. 2011; Das et al. 2014); *Achromobacter* (Turner 1954; Bachate et al. 2012); *Arthrobacter* (Duan et al. 2013) or *Brevibacillus* (*Brevibacillus brevis*, Banerjee et al. 2013).

A wide variety of microorganisms are also able to reduce As, which could have implications for the remobilization of As, including *Bacillus*, able to respire this metalloid like *Bacillus arsenicoselenatis* or *Bacillus selenitireducens* (Blum et al. 1998); *Agrobacterium* (*Agrobacterium tumefaciens*, Macur et al. 2004); *Enterobacter* sp. (representing around 3% of the total bacteria identified at Kettara mine, Liao et al. 2011); *Acinetobacter* (Paul et al. 2014); *Escherichia* (representing around 0.4% of the total bacteria identified at Kettara, Bahar et al. 2013a) or *Clostridium* sp. (Stolz et al. 2006).

Some species are also able to methylate As or to produce arsine, like the archaea *Methanobacterium bryantii* or *Methanobacterium formicium* producing methylated arsines and arsine, while *Clostridium collagenovorans* produces trimethyl arsine (Stolz et al. 2006). According to Wang et al. (2014b), species of genus like *Escherichia* and *Corynebacterium* sp., produce dimethylarsine; *Enterobacter* sp., *Achromobacter* sp. or *Aeromonas* sp. produce mono and dimethylarsine; *Alcaligenes* sp. produce mono-, dimethylarsine and arsine and *Pseudomonas* sp. produce mono-, di- and trimethylarsine and arsine. Microbial volatilization of arsines in the environment could play an important role in the biogeochemical cycling of As, and could be an interesting method for arsenic bioremediation, like bio-volatilization (Wang et al. 2014b).

Autotrophic and heterotrophic microorganisms are also important in these extreme environments. Indeed, autotrophic iron- and sulfur-oxidizing microorganisms that fix inorganic carbon provide organic materials used as growth substrates by heterotrophic populations (Johnson and Hallberg 2008) like *Acidiferrobacter*, an obligate autotroph and facultative anaerobe that fixes inorganic carbon and could also assimilate dinitrogen for growth (Hallberg et al. 2011). Heterotrophic bacteria that degrade organic compounds are also important in this environment, because they enable the continued growth of autotrophic Fe(II) and sulfur-oxidizing bacteria, which are often sensitive to the presence of this type of compound (Schippers et al. 2010; Hallberg 2010). Many microorganisms including *Bacillus*, *Pseudomonas*, *Acidiphilium*, *Enterobacter*, *Alicyclobacillus*, *Acinetobacter* or *Sulfobacillus* are known for their ability to oxidize organic carbon and are found in mine tailings and dumps (Schippers et al. 2010; Sanchez Andrea et al. 2011).

Spatial variation in microbial communities presents in the mine tailings of Kettara

According to Hakkou et al. (2008a), the main sulfide minerals were pyrrhotite and pyrite in both unweathered and oxidized Kettara tailings. The most abundant gangue minerals are chlorite-serpentine, quartz, talc, and goethite (most likely secondary phase). Muscovite and albite are present in trace proportions and carbonates (calcite) occur in very negligible quantities. Trace amounts of jarosite were also observed.

Principal component analysis revealed substantial differences in tailings microbial community composition with four groups identified: the secondary precipitates; the oxidized parts of the fine tailings; the unoxidized parts of the fine tailings and the coarse tailings. The alteration process was more intense in the coarse tailings of the dyke than in the tailings pond with an extensive oxidation front through the upper 75 cm in the coarse tailings compared to 5–15 cm in the tailings dam. Indeed, coarse tailings facilitate oxygen access and increase oxidation of sulfides due to their large particle size, contrary to the tailings pond where alteration was limited, due to the presence of a hardpan layer formed at the surface of tailings pond (Hakkou et al. 2008a). The principal Fe(II) and sulfur-oxidizing bacteria, well known for metal sulfide oxidation were mainly present in the coarse tailings, like *Acidiferrobacter* representing, in the dyke trench T3, 66% in T3-S1F and 30% in T3-S1O; *Alicyclobacillus* (58% in T3-S1O, 27% in K1 and 12% in K2; *Leptospirillum* (more than 99% in T3-S1F) or *Sulfobacillus* (96% in T3-S1O).

Our results highlighted also a quite different microbial community, compared to other sulfuric mine tailings producing AMD with our 4 most abundant populations similar to *Bacillus*, *Pseudomonas*, *Corynebacterium* and *Staphylococcus* and not to the well-known iron-oxidizing populations. Studies of microbial population in acidic mine tailings are not numerous, but many of them have identified the predominance of typical leaching bacteria such as *Acidithiobacillus*, *Leptospirillum*, *Sulfobacillus* or *Ferroplasma*. It is the case for different mine tailings containing mainly pyrite; a lead and zinc mine in China (Huang et al. 2011; Chen et al. 2013); a massive copper tailing in China (Liu et al. 2014); a copper and molybdenum porphyry mine in Chile (Diaby et al. 2007) or an abandoned semi-arid lead–zinc mine tailing site in USA (Mendez et al. 2008). Korehi et al. (2014) obtained similar results in 3 sulfidic mine tailings: a Pb, Zn and Ag mine containing pyrite and arsenopyrite in a temperate zone in Germany; a Zn and Cu mine dominated by pyrite in a cold and humid climate in Sweden and also in a former Ni, Cu, Zn and Co mine in Botswana containing mainly pyrrhotite in a hornblende and feldspar as gangue minerals in semi-arid climate, thus quite similar to our conditions

(Schipper et al. 2007; Kock and Schippers 2008). However, three of our four dominant bacterial population (*Pseudomonas*, *Corynebacterium* and *Staphylococcus*), were also the dominant genera in the extremely acidic part of a mine tailings of the lead and zinc mine in China, present in a warm and humid subtropical climate (Huang et al. 2011; Chen et al. 2013). The metallic minerals in this mine are pyrite, sphalerite, galena, chalcopyrite, tetrahedrite and magnetic pyrite and the major gangue minerals are calcite, dolomite and quartz (Han et al. 2012).

pH seemed to be often the most important factor shaping the microbial communities in mine tailings (Korehi et al. 2014; Chen et al. 2013; Liu et al. 2014). However, in our study, NMDS showed that Al was the main parameter discriminating our microbial communities (Fig. 3). In the Kettara mine, concentration of Al is very important in mine tailing (from 10407 to 74557 mg kg⁻¹) and in AMD (from 730 to 4170 mg l⁻¹). This element is known to be toxic for some microorganisms when present in high concentrations, competing with metabolism of iron and magnesium, and binding to DNA, membranes or cell walls (Pina and Cervantes 1996; Auger et al. 2013).

The generation, mobility, and attenuation of AMD involve complex processes governed by a combination of physical, chemical, but also biological factors (Simate and Ndlovu 2014). Acid mine drainage (AMD) is considered as one of the main causes of water pollution in many countries that have historic or current mining activities and this phenomenon is particularly problematic in semi-arid areas with water shortage. In Kettara mine, sequences related to *Bacillus*, *Pseudomonas*, *Corynebacterium* and *Staphylococcus* were identified as the dominant genera. However, many species are also related to genera known to play a key role in Fe and S cycle implicated in the oxidation of mineral sulfides but also in As and C cycles, helping us to gain a deeper understanding of the functioning of this ecosystem that produce intermittent AMD. To our knowledge, this is one of the first descriptions of microbial communities in a pyrrhotite mine using high-throughput sequencing, and the first description of the microbial community at Kettara mine. Such knowledge of microbial populations is useful to secure reactive mine tailings and will also help for the further development of waste management and bioremediation strategies.

Acknowledgements The authors thank the International Research Chairs Initiative funded by the International Development Research Centre (IDRC), Canada, and by the Canada Research Chairs program and the IRD (Institut de Recherche pour le Développement) for supporting this research.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Abdul SM, Jayasinghe SS, Chandana EPS, Jayasumana C et al (2015) Arsenic and human health effects: a review. *Environ Toxicol Pharmacol* 40:828–846
- Akcil A, Koldas S (2006) Acid mine drainage (AMD): causes, treatment and case studies. *J Clean Prod* 14:1139–1145
- Anandham R, Indiragandhi P, Madhaiyan M, Ryu KY et al (2008) Chemolithoautotrophic oxidation of thiosulfate and phylogenetic distribution of sulfur oxidation gene (*soxB*) in rhizobacteria isolated from crop plants. *Res Microbiol* 159:579–589
- Auger C, Han S, Appanna VP, Thomas SC, Ulibarri G, Appanna VD (2013) Metabolic reengineering invoked by microbial systems to decontaminate aluminum: implications for bioremediation technologies. *Biotechnol Adv* 31:266–273
- Babi K, Asselin H, Benzaazoua M (2015) Stakeholders' perceptions of sustainable mining in Morocco: a case study of the abandoned Kettara mine. *Extr Ind Soc* 1:185–192
- Bachate SP, Khapare RM, Kodam KM (2012) Oxidation of arsenite by two β -proteobacteria isolated from soil. *Appl Microbiol Biotechnol* 93:2135–2145
- Bachate SP, Nandre VS, Ghatpande NS, Kodam KM (2013) Simultaneous reduction of Cr(VI) and oxidation of As(III) by *Bacillus firmus* TE7 isolated from tannery effluent. *Chemosphere* 90:2273–2278
- Bachy C, Worden AZ (2014) Microbial ecology: finding structure in the rare biosphere. *Curr Biol* 24:R315–R317
- Bahar MM, Megharaj M, Naidu R (2012) Arsenic bioremediation potential of new arsenic oxidizing bacterium *Stenotrophomonas* sp. MM-7 isolated from soil. *Biodegradation* 23:803–812
- Bahar MM, Megharaj M, Naidu R (2013a) Bioremediation of arsenic-contaminated water: recent advances and future prospects. *Water Air Soil Pollut* 224:1722
- Bahar MM, Megharaj M, Naidu R (2013b) Kinetics of arsenite oxidation by *Variovorax* sp. MM-1 isolated from a soil containing low arsenic and identification of arsenite oxidase gene. *J Hazard Mater* 262:997–1003
- Banerjee S, Majumdar J, Samal AC, Bhattachariya P et al (2013) Biotransformation and bioaccumulation of arsenic by *Brevibacillus brevis* isolated from arsenic contaminated region of West Bengal. *IOSR J Environ Sci Toxicol Food Technol* 3:1–10
- Bates ST, Berg-Lyons D, Caporaso JG et al (2011) Examining the global distribution of dominant archaeal populations in soil. *ISME J* 5:908–917
- Behera BC, Patra M, Dutta SK, Thatoi HN (2014) Isolation and characterisation of sulphur oxidising bacteria from mangrove soil of Mahanadi River Delta and their sulphur oxidising ability. *J Appl Environ Microbiol* 2:1–5
- Bhatnagar L, Henriquet M, Zeikus JG, Aubert JP (1984) Utilization of mercapto-2-ethanol as a medium reductant for determination of the metabolic response of methanogens towards inorganic sulfur compounds. *FEMS Microbiol Lett* 22:155–158
- Blum JS, Bindi AB, Buzelli J, Stolz JF, Oremland RS (1998) *Bacillus arsenicoselenatis*, sp. nov., and *Bacillus selenitireducens*, sp. nov.: two haloalkaliphiles from Mono Lake, California that respire oxyanions. *Arch Microbiol* 171:19–30
- Bridge TAM, Johnson DB (1998) Reduction of soluble iron and reductive dissolution of ferric iron-containing minerals by moderately thermophilic iron-oxidizing bacteria. *Appl Environ Microbiol* 64:2181–2186
- Campos VL, Escalante G, Yañez J, Zaror CA et al (2009) Isolation of arsenite-oxidizing bacteria from a natural biofilm associated to volcanic rocks of Atacama Desert, Chile. *J Basic Microbiol* 49:S93–S97

- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D et al (2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci* 108:4516–4522
- Chang J-S, Lee J-H, Kim I-S (2011) Bacterial aox genotype from arsenic contaminated mine to adjacent coastal sediment: evidences for potential biogeochemical arsenic oxidation. *J Hazard Mater* 193:233–242
- Chen LX, Li JT, Chen YT, Huang LN et al (2013) Shifts in microbial community composition and function in the acidification of a lead/zinc mine tailings. *Environ Microbiol* 15:2431–2444
- Das S, Jean J-S, Kar S, Chou M-L et al (2014) Screening of plant growth-promoting traits in arsenic-resistant bacteria isolated from agricultural soil and their potential implication for arsenic bioremediation. *J Hazard Mater* 272:112–120
- Diaby N, Dold B, Pfeifer HR, Holliger C, Johnson DB, Hallberg KB (2007) Microbial communities in a porphyry copper tailings impoundment and their impact on the geochemical dynamics of the mine waste. *Environ Microbiol* 9:298–307
- Dopson M, Johnson DB (2012) Biodiversity, metabolism and applications of acidophilic sulfur-metabolizing microorganisms. *Environ Microbiol* 14:2620–2631
- Duan M, Wang Y, Xie X, Su C et al (2013) Arsenite oxidizing bacterium isolated from high arsenic groundwater aquifers from Datong Basin, Northern China. *Proc Earth Planet Sci* 7:232–235
- Edgar RC, Haas BJ, Clemente JC, Quince C et al (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–2200
- Fisher J, Hollibaugh JT (2008) Selenate-dependent anaerobic arsenite oxidation by a bacterium from Mono Lake, California. *Appl Environ Microbiol* 74:2588–2594
- Friedrich CG, Mitrenga G (1981) Oxidation of thiosulfate by *Paracoccus denitrificans* and other hydrogen bacteria. *FEMS Microbiol Lett* 10:209–212
- Galand PE, Casamayor EO, Kirchman DL, Lovejoy C (2009) Ecology of the rare microbial biosphere of the Arctic Ocean. *Proc Natl Acad Sci USA* 106:22427–22432
- García Balboa C, Bedoya IC, González F, Blázquez ML, Muñoz JA, Ballester A (2010) Bio-reduction of Fe(III) ores using three pure strains of *Aeromonas hydrophila*, *Serratia fonticola* and *Clostridium celerecrescens* and a natural consortium. *Biores Technol* 101:7864–7871
- Hakkou R, Benzaazoua M, Bussière B (2008a) Acid mine drainage at the abandoned Kettara mine (Morocco): 1. Environmental characterization. *Mine Water Environ* 27:145–159
- Hakkou R, Benzaazoua M, Bussière B (2008b) Acid mine drainage at the abandoned Kettara mine (Morocco): 2. Mine waste geochemical behavior. *Mine Water Environ* 27:160–170
- Hallberg KB (2010) New perspectives in acid mine drainage microbiology. *Hydrometallurgy* 104:448–453
- Hallberg KB, Hedrich S, Johnson DB (2011) *Acidiferrobacter thiooxydans*, gen. nov. sp. nov.; an acidophilic, thermo-tolerant, facultatively anaerobic iron-and sulfur-oxidizer of the family *Ectothiorhodospiraceae*. *Extremophiles* 15:271–279
- Han Y, Wang JB, Zhu XY, Guo NN (2012) Stable isotope and REE geological and geochemical characteristics of the calcite in the Fankou Zinc-Lead deposit, Guangdong Province, China. In *Adv Mater Res* 524:205–212
- Huang LN, Zhou WH, Hallberg KB, Wan CY, Li J, Shu WS (2011) Spatial and temporal analysis of the microbial community in the tailings of a Pb-Zn mine generating acidic drainage. *Appl Environ Microbiol* 77:5540–5544
- Huse SM, Welch DM, Morrison HG, Sogin ML (2010) Ironing out the wrinkles in the rare biosphere through improved OTU clustering. *Environ Microbiol* 12:1889–1898
- Jiang CY, Liu Y, Liu YY, You XY, Guo X, Liu SJ (2008) *Alicyclobacillus ferrooxydans* sp. nov., a ferrous-oxidizing bacterium from solfataric soil. *Int J Syst Evol Microbiol* 58:2898–2903
- Johnson DB, Bridge TAM (2002) Reduction of ferric iron by acidophilic heterotrophic bacteria: evidence for constitutive and inducible enzyme systems in *Acidiphilium* spp. *J Appl Microbiol* 92:315–321
- Johnson DB, Hallberg KB (2008) Carbon, iron and sulfur metabolism in acidophilic micro-organisms. *Adv Microb Physiol* 54:201–255
- Justice NB, Norman A, Brown CT, Singh A, Thomas BC, Banfield JF (2014) Comparison of environmental and isolate *Sulfobacillus* genomes reveals diverse carbon, sulfur, nitrogen, and hydrogen metabolisms. *BMC Genomics* 15:1107
- Karavaiko GI, Bogdanova TI, Tourova TP, Kondrat'eva TF et al (2005) Reclassification of 'Sulfobacillus thermosulfidooxidans subsp. thermotolerans' strain K1 as *Alicyclobacillus tolerans* sp. nov. and *Sulfobacillus disulfidooxidans* Dufresne et al 1996 as *Alicyclobacillus disulfidooxidans* comb. nov., and emended description of the genus *Alicyclobacillus*. *Int J Syst Evol Microbiol* 55:941–947
- Katayama Y, Hiraishi A, Kuraishi H (1995) *Paracoccus thiocyanatus* sp. nov., a new species of thiocyanate-utilizing facultative chemolithotroph, and transfer of *Thiobacillus versutus* to the genus *Paracoccus* as *Paracoccus versutus* comb. nov. with emendation of the genus. *Microbiology* 141:1469–1477
- Khalil K, Hanich L, Bannari A, Zouhri L et al (2013) Assessment of soil contamination around an abandoned mine in a semi-arid environment using geochemistry and geostatistics: pre-work of geochemical process modeling with numerical models. *J Geochem Explor* 125:117–129
- Kock D, Schippers A (2008) Quantitative microbial community analysis of three different sulfidic mine tailing dumps generating acid mine drainage. *Appl Environ Microbiol* 74:5211–5219
- Korehi H, Blöthe M, Schippers A (2014) Microbial diversity at the moderate acidic stage in three different sulfidic mine tailings dumps generating acid mine drainage. *Res Microbiol* 165:713–718
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD (2013) Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 79:5112–5120
- Kumaraswamy R, Sjollem K, Kuenen G, Van Loosdrecht M, Muyzer G (2006) Nitrate-dependent [Fe(II) EDTA]²⁻ oxidation by *Paracoccus ferrooxydans* sp. nov., isolated from a denitrifying bioreactor. *Syst Appl Microbiol* 29:276–286
- Lghoul M, Teixidó T, Peña JA, Hakkou R, Kchikach A, Guérin R et al (2012) Electrical and seismic tomography used to image the structure of a tailings pond at the abandoned Kettara mine, Morocco. *Mine Water Environ* 31:53–61
- Lghoul M, Maqsoud A, Hakkou R, Kchikach A (2014) Hydrogeochemical behavior around the abandoned Kettara mine site, Morocco. *J Geochem Explor* 144:456–467
- Liao VHC, Chu YJ, Su YC, Hsiao SY, Wei CC, Liu CW et al (2011) Arsenite-oxidizing and arsenate-reducing bacteria associated with arsenic-rich groundwater in Taiwan. *J Contam Hydrol* 123:20–29
- Liu J, Hua ZS, Chen LX, Kuang JL, Li SJ, Shu WS, Huang LN (2014) Correlating microbial diversity patterns with geochemistry in an extreme and heterogeneous environment of mine tailings. *Appl Environ Microbiol* 80:3677–3686
- Luo J, Tian G, Lin W (2013) Enrichment, isolation and identification of sulfur-oxidizing bacteria from sulfide removing bioreactor. *J Environ Sci* 25:1393–1399

- Macur RE, Jackson CR, Botero LM, Mcdermott TR, Inskip WP (2004) Bacterial populations associated with the oxidation and reduction of arsenic in an unsaturated soil. *Environ Sci Technol* 38:104–111
- Mahmood Q, Zheng P, Hu B, Jilani G, Azim MR, Wu D, Liu D (2009) Isolation and characterization of *Pseudomonas stutzeri* QZ1 from an anoxic sulfide-oxidizing bioreactor. *Anaerobe* 15:108–115
- Mandal BK, Suzuki KT (2002) Arsenic round the world: a review. *Talanta* 58:201–235
- Mathieu C, Pieltain F (2003) Analyse chimique des sols. In: Méthode choisies. Ed Lavoisier, Tec and Doc
- Mendez MO, Neilson JW, Maier RM (2008) Characterization of a bacterial community in an abandoned semiarid lead-zinc mine tailing site. *Appl Environ Microbiol* 74:3899–3907
- Muehe EM, Gerhardt S, Schink B, Kappler A (2009) Ecophysiology and the energetic benefit of mixotrophic Fe(II) oxidation by various strains of nitrate-reducing bacteria. *FEMS Microbiol Ecol* 70:335–343
- Needleman SB, Wunsch CD (1970) A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J Mol Biol* 48:443–453
- Norris PR, Clark DA, Owen JP, Waterhouse S (1996) Characteristics of *Sulfobacillus acidophilus* sp. nov. and other moderately thermophilic mineral-sulphide-oxidizing bacteria. *Microbiology* 142:775–783
- Okamura K, Kawai A, Wakao N, Yamada T, Hiraishi A (2015) *Acidiphilium iwatense* sp. nov., isolated from an acid mine drainage treatment plant, and emendation of the genus *Acidiphilium*. *Int J Syst Evol Microbiol* 65:42–48
- Omori T, Monna LISA, Saiki Y, Kodama T (1992) Desulfurization of dibenzothiophene by *Corynebacterium* sp. strain SY1. *Appl Environ Microbiol* 58:911–915
- Oremland RS, Stolz JF (2003) The ecology of arsenic. *Science* 300:939–944
- Osborne FH, Ehrlich HL (1976) Oxidation of arsenite by a soil isolate of *Alcaligenes*. *J Appl Microbiol* 41:295–305
- Paul D, Poddar S, Sar P (2014) Characterization of arsenite-oxidizing bacteria isolated from arsenic-contaminated groundwater of West Bengal. *J Environ Sci Health Part A-Toxic/Hazard Subst Environ Eng* 49:1481–1492
- Pina RG, Cervantes C (1996) Microbial interactions with aluminium. *BioMetals* 9:311–316
- R Development Core Team (2012) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org>
- Rawlings DE, Johnson DB (2007) The microbiology of biomining: development and optimization of mineral-oxidizing microbial consortia. *Microbiology* 153:315–324
- Sallam A, Steinbüchel A (2009) *Clostridium sulfidigenes* sp. nov., a mesophilic, proteolytic, thiosulfate- and sulfur-reducing bacterium isolated from pond sediment. *Int J Syst Evol Microbiol* 59:1661–1665
- Sanchez Andrea I, Rodriguez N, Amils R, Sanz JL (2011) Microbial diversity in anaerobic sediments at Rio Tinto, a naturally acidic environment with a high heavy metal content. *Appl Environ Microbiol* 77:6085–6093
- Santini JM, Sly LI, Schnagl RD, Macy JM (2000) A new chemolithoautotrophic arsenite-oxidizing bacterium isolated from a gold mine: phylogenetic, physiological, and preliminary biochemical studies. *Appl Environ Microbiol* 66:92–97
- Sarkar A, Kazy KS, Sar P (2014) Studies on arsenic transforming groundwater bacteria and their role in arsenic release from subsurface sediment. *Environ Sci Pollut Res* 21:8645–8662
- Schippers A, Sand W (1999) Bacterial leaching of metal sulfides proceeds by two indirect mechanisms via thiosulfate or via polysulfides and sulfur. *Appl Environ Microbiol* 65:319–321
- Schippers A, Kock D, Schwartz M, Böttcher ME, Vogel H, Hagger M (2007) Geomicrobiological and geochemical investigation of a pyrrhotite-containing mine waste tailings dam near Selebi-Phikwe in Botswana. *J Geochem Explor* 92:151–158
- Schippers A, Breuker A, Blazejak A, Bosecker K, Kock D, Wright TL (2010) The biogeochemistry and microbiology of sulfidic mine waste and bioleaching dumps and heaps, and novel Fe(II)-oxidizing bacteria. *Hydrometallurgy* 104:342–350
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB et al (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75:7537–7541
- Schloss PD, Gevers D, Westcott SL (2011) Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS One* 6:e27310
- Simate GS, Ndlovu S (2014) Acid mine drainage: challenges and opportunities. *J Environ Chem Eng* 2:1785–1803
- Singer PC, Stumm W (1970) Acid mine drainage: the rate-determining step. *Science* 167:1121–1123
- Sorokin DY, Teske A, Robertson LA, Kuenen JG (1999) Anaerobic oxidation of thiosulphate to tetrathionate by obligately heterotrophic bacteria, belonging to the *Pseudomonas stutzeri* complex. *FEMS Microbiol Ecol* 30:113–123
- Stolz JF, Basu P, Santini JM, Oremland RS (2006) Arsenic and selenium in microbial metabolism. *Annu Rev Microbiol* 60:107–130
- Takahashi Y, Suto K, Inoue C (2010) Polysulfide reduction by *Clostridium* relatives isolated from sulfate-reducing enrichment cultures. *J Biosci Bioeng* 109:372–380
- Tan GL, Shu WS, Hallberg KB, Li F, Lan CY, Zhou WH, Huang LN (2008) Culturable and molecular phylogenetic diversity of microorganisms in an open-dumped, extremely acidic Pb/Zn mine tailings. *Extremophiles* 12:657–664
- Thermo Scientific NITON (2008) Thermo Scientific NITON® XL3t 900 Series Product specifications. <http://www.thermo.com/niton> (last accessed 3 Nov 2014)
- Toughzaoui S, El Amari K, Benkaddour A, Hibti M, Essarraj S (2015) Hydrogeochemical and isotopic studies of the Kettara mine watershed, Morocco. *Mine Water Environ* 34:308–319
- Turner AW (1954) Bacterial oxidation of arsenite. I. Description of bacteria isolated from arsenical cattle-dipping fluids. *Aust J Biol Sci* 7:452–476
- Ventura BA, González F, Ballester A, Blázquez ML, Muñoz JA (2015) Bioreduction of iron compounds by *Aeromonas hydrophila*. *Int Biodeterior Biodegrad* 103:69–76
- Vera M, Schippers A, Sand W (2013) Progress in bioleaching: fundamentals and mechanisms of bacterial metal sulfide oxidation. Part A. *Appl Microbiol Biotechnol* 97:7529–7541
- Vikromvarasiri N, Boonyawanich S, Pisutpaisal N (2015) Optimizing sulfur oxidizing performance of *Paracoccus pantotrophus* isolated from leather industry wastewater. *Energy Proc* 79:629–633
- Wang GW, Chen TH, Yue ZB, Zhou YF, Wang J (2014a) Isolation and characterization of *Pseudomonas stutzeri* capable of reducing Fe(III) and nitrate from skarn-type copper mine tailings. *Geomicrobiol J* 31:509–518
- Wang P, Sun G, Jia Y, Meharg AA, Zhu Y (2014b) A review on completing arsenic biogeochemical cycle: microbial volatilization of arsines in environment. *J Environ Sci* 26:371–381
- Weber KA, Achenbach LA, Coates JD (2006) Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat Rev Microbiol* 4:752–764
- Xiu W, Guo H, Liu Q, Liu Z, Zhang B (2015) Arsenic removal and transformation by *Pseudomonas* sp. strain GE-1-induced ferrihydrite: co-precipitation versus adsorption. *Water Air Soil Pollut* 226:1–14

Xu XJ, Chen C, Guo HL, Wang AJ, Ren NQ, Lee DJ (2016) Characterization of a newly isolated strain *Pseudomonas* sp. C27 for sulfide oxidation: reaction kinetics and stoichiometry. *Sci Rep* 6:21032. doi:[10.1038/srep21032](https://doi.org/10.1038/srep21032)

Yamamura S, Amachi S (2014) Microbiology of inorganic arsenic: from metabolism to bioremediation. *J Biosci Bioeng* 118:1–9

Zeikus JG, Wolfe RS (1972) *Methanobacterium thermoautotrophicum* sp. nov., an anaerobic, autotrophic, extreme thermophile. *J Bacteriol* 109:707–713