



**HAL**  
open science

## The origin of pre-plasmablastic cells

Jerome Moreaux

► **To cite this version:**

Jerome Moreaux. The origin of pre-plasmablastic cells. Blood, American Society of Hematology, 2021, 137 (9), pp.1134-1135. 10.1182/blood.2020009746 . hal-03243758

**HAL Id: hal-03243758**

**<https://hal.umontpellier.fr/hal-03243758>**

Submitted on 31 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **The origin of pre-plasmablastic cells**

Tracking no: BLD-2020-009746-C

Jerome Moreaux (Institute of Human Genetics, France)

### **Abstract:**

**Conflict of interest:** No COI declared

**COI notes:** The author has no conflict of interest to declare.

**Preprint server:** No;

**Author contributions and disclosures:** JM wrote the commentary.

**Non-author contributions and disclosures:** No;

**Agreement to Share Publication-Related Data and Data Sharing Statement:**

**Clinical trial registration information (if any):**

## The origin of pre-plasmablastic cells

Jerome MOREAUX<sup>1,2,3,4</sup>

<sup>1</sup> Department of Biological Hematology, CHU Montpellier, Montpellier, France;

<sup>2</sup> Institute of Human Genetics, UMR 9002 CNRS-UM, Montpellier, France;

<sup>3</sup> University of Montpellier, UFR Medicine, Montpellier, France;

<sup>4</sup> Institut Universitaire de France (IUF)

In this issue of Blood, Pignarre A et al. characterize the genomic events involved in the cell fate decision between activated B cells and plasmablast<sup>1</sup>. Plasma cells play an important role in humoral immunity by synthesizing and secreting antibodies<sup>2</sup>. Understanding the biological processes that control the production of plasma cells is critical both to ensure efficient immune response without autoimmunity or immune deficiency and prevent tumorigenesis. The production of IL-4 by follicular helper T cells drives B cell amplification and maturation<sup>3</sup>. However, the full molecular mechanisms behind these functions are not fully understood. Pignarre A et al. reported new biological events driving normal B to plasma cell differentiation. Using an *in vitro* model, naïve B cells are cultured in a two-step process which results in differentiation into plasmablasts<sup>4</sup>, they demonstrated that cells that are destined to differentiate into plasma cells if there is an early response to IL-4, which results in downregulation of the CD23 cell surface protein and IL-4/STAT6 signaling. However, B cells maintaining IL-4 signaling did not differentiate. Furthermore, the differentiation of CD23 negative cells is associated with CBLb E3 ubiquitin ligase downregulation, coinciding with IRF4 induction and with specific chromatin and transcriptional **modifications**. The changes were identified by ATAC-sequencing and hydroxymethylation profiling. However, no major changes in expression of epigenetic factors was noted. CBLb is known to prevent premature germinal center (GC) exit

promoting IRF4 degradation in light zone B cells<sup>5</sup>. Pignarre et al. reported potential STAT6 binding sites in the CBLb promoter suggesting potential direct regulation, hence the interest in characterizing STAT6 targets using chromatin immunoprecipitation. CD23 negative B cells, post activation, have the characteristics of preplasmablasts with a significant increase in chromatin accessibility at immunoglobulin heavy chains coding loci. Full transcriptomic characterization of the proposed model at a single-cell level would be particularly useful in deciphering the heterogeneity and transcriptional trajectories during B to plasma cell differentiation. The major transcriptional and epigenetic changes reported by Pignarre et al. may be associated with changes in nuclear organization during terminal B cell differentiation. Gene regulation depends on the 3-dimensional chromatin organization and its regulatory elements. Recent data obtained in a mouse model revealed that B to plasma cell differentiation is associated with major changes in chromosome topology that could be driven by plasma cell biology or reflect enhancer-induced modifications in chromatin organization<sup>6</sup>. B to plasma cell transition is associated with compartmentalization changes along with **gain in genomic interactions across the Prdm1 locus increasing genomic interactions between promoter regions and regulatory elements, concurrent with transcriptional induction. In contrast, early B-cell factor 1 (Ebf1) locus repositions to peri-centromeric heterochromatin in association with transcriptional repression.** Furthermore, the inter-chromosomal hubs reported during B to plasma cell maturation are associated with histone marks that define transcriptionally active or repressive hubs. The epigenetic landscape characterization together with nuclear architecture study of the human B to PC differentiation model developed by Pignarre et al. may provide important findings to progress in the understanding of the molecular mechanisms driving B to plasma cell fate.

IL-4 production by T follicular helper and STAT6 mutations activate and drive the IL4/STAT6 axis in follicular lymphoma<sup>7</sup>. Recently, single-cell characterization of germinal center B cells generated a new single-cell cell of origin classification that identified distinct prognostic subgroups within the GCB and ABC diffuse large B cell lymphoma subgroups<sup>8</sup>. The analysis of these single-cell transcriptomic resources derived from germinal center purified B cells, in light of the new data provided by Pignarre et al. may be of particular interest. The results provided by Pignarre et al. provide new insights in the molecular mechanisms driving follicular lymphoma biology.

## References

1. Pignarre A, Chatonnet F, Caron G, Haas M, Desmots-Loyer F, Fest T. Plasmablasts derive from CD23-negative activated B cells after the extinction of IL-4/STAT6 signaling and IRF4 induction. *Blood*. 2020.
2. Shapiro-Shelef M, Calame K. Regulation of plasma-cell development. *Nat Rev Immunol*. 2005;5(3):230-242.
3. Crotty S. T follicular helper cell differentiation, function, and roles in disease. *Immunity*. 2014;41(4):529-542.
4. Le Gallou S, Caron G, Delaloy C, Rossille D, Tarte K, Fest T. IL-2 requirement for human plasma cell generation: coupling differentiation and proliferation by enhancing MAPK-ERK signaling. *J Immunol*. 2012;189(1):161-173.
5. Li X, Gadzinsky A, Gong L, et al. Cbl Ubiquitin Ligases Control B Cell Exit from the Germinal-Center Reaction. *Immunity*. 2018;48(3):530-541 e536.
6. Bortnick A, He Z, Aubrey M, et al. Plasma Cell Fate Is Orchestrated by Elaborate Changes in Genome Compartmentalization and Inter-chromosomal Hubs. *Cell Rep*. 2020;31(1):107470.
7. Ame-Thomas P, Hoeller S, Artchounin C, et al. CD10 delineates a subset of human IL-4 producing follicular helper T cells involved in the survival of follicular lymphoma B cells. *Blood*. 2015;125(15):2381-2385.
8. Holmes AB, Corinaldesi C, Shen Q, et al. Single-cell analysis of germinal-center B cells informs on lymphoma cell of origin and outcome. *J Exp Med*. 2020;217(10).

Figure legend

After activation, B cells that are committed to differentiate into plasma cells downregulate the CD23 cell surface protein, IL-4/STAT6 signaling and CBLb activity concomitantly with IRF4 induction. B cells that maintain the IL-4 signaling will not differentiate.

