



**HAL**  
open science

## Frugivory-related traits promote speciation of tropical palms

Renske Onstein, William Baker, Thomas Couvreur, Søren Faurby,  
Jens-Christian Svenning, W. Daniel Kissling

► **To cite this version:**

Renske Onstein, William Baker, Thomas Couvreur, Søren Faurby, Jens-Christian Svenning, et al.. Frugivory-related traits promote speciation of tropical palms. *Nature Ecology & Evolution*, 2017, 1 (12), pp.1903-1911. 10.1038/s41559-017-0348-7. hal-03346431

**HAL Id: hal-03346431**

**<https://hal.umontpellier.fr/hal-03346431v1>**

Submitted on 16 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Frugivory-related traits promote speciation of tropical palms**

2

3

4 Renske E. Onstein<sup>1,\*</sup>, William J. Baker<sup>2</sup>, Thomas L. P. Couvreur<sup>3</sup>, Søren Faurby<sup>4</sup>, Jens-Christian  
5 Svenning<sup>5</sup> & W. Daniel Kissling<sup>1,\*</sup>

6

7 <sup>1</sup>Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, Amsterdam,  
8 The Netherlands

9 <sup>2</sup>Royal Botanic Gardens, Kew, Richmond, Surrey, UK

10 <sup>3</sup>Institut de Recherche pour le Développement, UMR-DIADE, Montpellier, France

11 <sup>4</sup>Department of Biological and Environmental Sciences, University of Gothenburg, Box 461, SE 405  
12 30, Göteborg, Sweden

13 <sup>5</sup>Section for Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University, Aarhus,  
14 Denmark

15

16 \*corresponding authors e-mails: [onsteinre@gmail.com](mailto:onsteinre@gmail.com); [wdkissling@gmail.com](mailto:wdkissling@gmail.com)

17

18 Running title: Frugivory and palm speciation

19

20 Total word count (Abstract, Introduction, Results, Discussion, Methods, References, and  
21 Acknowledgements): 7300 words

22

23 Abstract (148 out of 150 words)

24 Main text (3137 out of 3500 words)

25 Introduction (812 words)

26 Methods: (2005 out of 3000 words)

27 References: 59 (excl. method references)

28

29 Number of figures: 4 (Figs. 1, 2, 3, 4 in colour)

30 Tables: 1

31 Supporting information: Yes

32 **Abstract** [148 words]

33

34 **Animal-mediated seed dispersal by frugivorous birds and mammals is central to the ecology and**  
35 **functioning of tropical ecosystems, but whether and how frugivory-related traits have affected**  
36 **plant speciation remains little explored. Fruit size is directly linked to plant dispersal capacity**  
37 **and therefore influences gene flow and genetic divergence of plant populations. Using a global**  
38 **species-level phylogeny with comprehensive data on fruit sizes and plant species distributions we**  
39 **test whether fruit size has affected speciation rates of palms (Arecaceae), a characteristic**  
40 **tropical plant family. Globally, results reveal that palms with small fruit sizes have elevated**  
41 **speciation rates compared to those with large (megafauna) fruits. Speciation of small-fruited**  
42 **palms is particularly high in the understory of tropical rainforests in the New World, and on**  
43 **islands in the Old World. This suggests that frugivory-related traits in combination with**  
44 **geography and the movement behaviour of frugivores can influence speciation of tropical plants.**

45

46 The dispersal of seeds by fruit-eating animals such as birds and mammals is a key plant-animal  
47 interaction, especially in the tropics<sup>1,2</sup>. Frugivores constantly move around the seeds of animal-  
48 dispersed plants and thereby affect dispersal, gene flow, and genetic structure of plant populations<sup>3</sup>.  
49 This ultimately can influence plant speciation<sup>2</sup>. Of particular importance for speciation are the  
50 frequencies of both restricted and long-distance dispersal because the degree of genetic divergence of  
51 plant populations depends on them<sup>2,4</sup>. To date, few case studies have examined the relationship  
52 between seed dispersal and speciation in animal-dispersed plants<sup>2,4</sup>, and large-scale studies testing the  
53 generality of this remain scarce.

54 Fruit size is a key trait in plant-frugivore interactions (Table 1). Fruit size sets a limit to the  
55 ingestion of fruits by relatively small-sized seed dispersers and therefore tends to be positively  
56 correlated with body sizes and gape widths of consumers<sup>5-8</sup>. Large fruits such as megafaunal fruits (> 4  
57 cm in size<sup>9,10</sup>) are predominantly dispersed by large-bodied, non-flying mammalian frugivores (e.g.  
58 elephants, a number of extinct proboscideans, tapirs, large primates, ground sloths) which have large  
59 home ranges<sup>11,12</sup>. This leads to frequent dispersal across large distances, high gene flow among plant  
60 populations and a low speciation probability<sup>2</sup>. In contrast, small fruits are predominantly dispersed by  
61 small- and medium-sized frugivores, including frugivorous birds, bats, scatter-hoarding rodents or  
62 other small-bodied non-flying mammals. Compared to mammalian megafauna, these frugivores  
63 generally have smaller home ranges and less frequent dispersal across large distances<sup>2</sup>, and island  
64 colonization is possible (e.g. via birds and bats)<sup>13</sup>. The more ‘restricted dispersal’ of non-megafaunal  
65 frugivores combined with occasional long-distance dispersal, e.g. as typically suggested from fat-  
66 tailed seed dispersal kernels<sup>3</sup>, can promote the divergence of isolated plant populations and hence  
67 increase the probability of speciation<sup>2</sup>. Consequently, a higher speciation rate can be predicted for  
68 plant lineages with small fruits compared to those with large, megafaunal fruits (‘fruit-size  
69 hypothesis’, H1 in Table 1).

70 Beyond fruit size, plants with animal-dispersed fruits in the understory of tropical rainforests  
71 have been associated with high speciation rates<sup>4</sup>, especially when compared to taller plant growth  
72 forms (e.g. canopy trees) (Table 1). This could be caused by the sedentary nature of small- and  
73 medium-sized seed dispersers in the forest understory because their spatially restricted dispersal will  
74 result in low gene flow among plant populations<sup>14,15</sup> and thus a higher probability of allopatric plant  
75 speciation<sup>4</sup>. Animal-dispersed understory plants should therefore diversify more extensively than other  
76 plants<sup>4</sup>, especially in regions where small-bodied understory birds are abundant and species-rich<sup>16</sup>.  
77 This predicts a higher speciation rate of understory lineages compared to other growth forms  
78 (‘understory habitat hypothesis’, H2 in Table 1). Only few studies have tested the understory habitat  
79 hypothesis<sup>4,17</sup> and it remains unclear how general and widely applicable it is across taxa and regions.

80 Oceanic barriers make seed dispersal to islands challenging. The isolation of islands can  
81 restrict colonization and limit gene flow among plant populations (Table 1). Long-distance dispersal of  
82 animal-dispersed plants to remote oceanic islands usually depends on frugivores that are strong fliers

83 and hence able to cross large stretches of open water, including birds such as hornbills, macaws and  
84 fruit pigeons, and volant mammals such as fruit bats<sup>13,18</sup>. Seed dispersal to remote islands is therefore  
85 generally rare, leading to increased possibilities for genetic differentiation by isolation and allopatric  
86 plant speciation ('island colonization hypothesis', H3 in Table 1). Especially on islands that have been  
87 isolated for millions of years (i.e. volcanic islands or atolls), plant speciation rates can be expected to  
88 be higher compared to the mainland or continental islands because the latter have experienced more  
89 connectivity (and hence gene flow) through geological time. Although adaptive radiations on islands  
90 have been studied extensively<sup>19</sup>, tests of the island colonization hypothesis for animal-dispersed plant  
91 taxa remain sparse.

92 Here, we quantify speciation rates in relation to fruit size, understory habitat and island  
93 colonization in palms (Arecaceae), a species-rich animal-dispersed plant family typical for tropical  
94 rainforests<sup>20,21</sup>. Among angiosperm families, the palm family is one of the major food plant groups for  
95 vertebrate frugivores in the tropics<sup>2</sup> and a large number of both avian and mammalian frugivores have  
96 been observed to feed on them<sup>22</sup>. Together with data on fruit sizes, growth forms and species  
97 distributions we estimate speciation rates from a species-level phylogeny of the palm family<sup>23</sup> globally  
98 as well as separately for the New World (including South, Central and North America and the  
99 Caribbean) and the Old World (including Australia, Indomalaya, Oceania, Pacific, Africa, Madagascar  
100 and surrounding islands)<sup>24</sup>. More specifically, we tested the following three hypotheses (Table 1): (H1)  
101 palms with small fruit sizes (< 4 cm) have higher speciation rates than palms with large fruit sizes (i.e.  
102 megafaunal fruits  $\geq 4$  cm) ('fruit size hypothesis'); (H2) understory palms (especially those with small  
103 fruits) show higher speciation rates than canopy palms ('understory habitat hypothesis'); and (H3)  
104 dispersal to oceanic islands has increased speciation rates compared to speciation on the mainland and  
105 continental islands ('island colonization hypothesis').

## 106 **Results**

107 The majority of palms have relatively small fruits (< 4 cm in length;  $n = 1607$  species), but about 12%  
108 of all sampled palm species have large, megafauna fruits ( $\geq 4$  cm in length;  $n = 229$  species) (Fig. 1).  
109 Overall, fruit sizes of animal-dispersed palms vary widely from small 0.4–0.5 cm fruits in some *Areca*,  
110 *Bactris*, *Calamus*, *Chamaedorea*, *Coccothrinax*, *Dyopsis*, *Geonoma*, *Licuala* and *Pinanga* species to  
111 large > 10 cm fruits in genera such as *Borassus*, *Metroxylon* and *Phytelephas* (Supplementary Table  
112 1). Some palm species with particularly large fruits such as the coconut (*Cocos nucifera*, 22.5 cm fruit  
113 size), the nipa palm (*Nypa fruticans*, 11.5 cm fruit size) and the double coconut (*Lodoicea maldivica*,  
114 45 cm fruit size) are not dispersed by animals (Supplementary Table 1). Apart from these few  
115 exceptions, all other palm species have vertebrate-dispersed fruit types (drupes and berries) and both  
116 birds and mammals are their main seed dispersers<sup>22</sup>.

117

118 **Fruit size-dependent speciation.** Using information on fruit sizes of 1836 palm species together with  
119 the Binary State Speciation and Extinction (BiSSE) model<sup>25,26</sup> we tested whether small fruits are  
120 associated with high speciation rates (H1, Table 1). The best fitting BiSSE model showed that  
121 speciation rates of small-fruited palm lineages are higher than those of large-fruited palm lineages  
122 (H1; Fig. 2). This supported the fruit size hypothesis both globally (Fig. 2a) and in the Old World (Fig.  
123 2c), but interestingly not in the New World (Fig. 2b). The global BiSSE model indicated that small-  
124 fruited palms have a 1.6-fold higher speciation rate than large-fruited palms (median  $\lambda_{\text{small}} = 0.22$ ,  $\lambda_{\text{large}}$   
125  $= 0.14$ ). A 3.4-fold increase in speciation rate due to small fruit sizes was inferred for the Old World  
126 (median  $\lambda_{\text{small}} = 0.48$ ,  $\lambda_{\text{large}} = 0.13$ ). A potential confounding factor between speciation rate and fruit  
127 size could be the allometric relationship between fruit size and overall plant size (Supplementary Fig.  
128 1). Using data on palm stem heights as a measure of overall plant size showed that the high speciation  
129 rate of small-fruited palm lineages persisted when overall plant size was accounted for (for details see  
130 Supplementary Note 1). Hence, there was strong evidence for the hypothesized increase of speciation  
131 rates in small-fruited vertebrate-dispersed palm clades relative to large, megafaunal-fruited palms, at  
132 least for the Old World.

133  
134 **Understory habitat.** About 39% of all sampled palm species ( $n = 802$  species) occur in the forest  
135 understory, and palms with small fruits are more common in the understory than palms with large  
136 fruits (91% and 9%, respectively). In general, understory palms also have smaller fruits than canopy  
137 palms (median<sub>understory</sub> = 1.25 cm vs. median<sub>canopy</sub> = 1.7 cm, Supplementary Fig. 2). To quantify the  
138 relative importance of fruit size and understory habitat for speciation rates in palms, we used fruit  
139 sizes and additional data on understory habitat in a Multiple State Speciation and Extinction model  
140 (MuSSE multistate)<sup>27</sup> to test the understory habitat hypothesis (H2, Table 1). We compared the  
141 additive and interaction effects of these two binary traits (small fruits and understory habitat) to a  
142 baseline MuSSE model that estimated speciation rates when both traits were absent, i.e. relative to  
143 palm lineages that have large fruits and a canopy habit. Globally, the best-fitting MuSSE model  
144 indicated that both small fruits and understory habitat had a positive effect on speciation rates relative  
145 to the baseline (H2; Fig. 3a, compare yellow vs. grey box-and-whisker plots). In addition, there was a  
146 positive interaction effect (red box-and-whisker plot, Fig. 3a), indicating that fruit size and understory  
147 habitat acted synergistically to increase speciation rates. Hence, understory palms with small fruits had  
148 the highest speciation rates at a global scale. Since understory palm species are more common in the  
149 New World than the Old World relative to canopy species (45% and 36%, respectively), we further  
150 tested whether this interaction effect differed among these regions. In the New World, the best-fitting  
151 MuSSE model confirmed the global analysis, i.e. both additive and interactive effects of small fruit  
152 size and understory habitat were detected (Fig. 3b). However, in the Old World only positive additive  
153 effects were supported, but no interaction term (Fig. 3c).

154

155 **Island colonization.** About 13% of all included palm species ( $n = 331$  species) are restricted in their  
156 occurrence to oceanic islands, 80% are restricted to the mainland or to continental islands ( $n = 2036$   
157 species), and 7% occur both on oceanic islands and mainland or continental islands ( $n = 190$  species).  
158 Oceanic island-distributed palms have on average slightly larger fruits than palms that are distributed  
159 on mainland or continental islands ( $\text{median}_{\text{island}} = 1.6$  cm vs.  $\text{median}_{\text{mainland}} = 1.5$  cm, Supplementary  
160 Fig. 2). Using MuSSE models, we tested whether small fruit size in combination with oceanic island  
161 colonization has an effect on speciation rates (H3, Table 1). At a global scale, the best-fitting MuSSE  
162 model for fruit size and island colonization indicated that small fruits and island colonization have  
163 both positive additive effects (Fig. 4a, compare yellow vs. grey box-and-whisker plots) as well as  
164 positive interactive effects (red box-and-whisker plot, Fig. 4a), compared to large-fruited mainland-  
165 distributed palm lineages. Interestingly, in the New World the best-fitting MuSSE model only  
166 supported positive additive effects of small fruit size and island colonization, but no interaction effect  
167 (Fig. 4b). Moreover, the posterior distributions of the speciation rates resulting from the Bayesian  
168 analysis strongly overlapped, suggesting only a slight increase in speciation rates due to small fruits  
169 and island colonization relative to large-fruited mainland palms (Fig. 4b). However, in the Old World  
170 the results from the global analysis were confirmed (Fig. 4c), suggesting that small-fruited palm  
171 lineages on oceanic islands have particularly high speciation rates.

172

## 173 **Discussion**

174 Using trait-dependent diversification models and time-calibrated species-level phylogenies of palms  
175 we show that dispersal-relevant traits are important drivers of palm radiations. Speciation rates were  
176 higher for palm lineages with small fruits ( $< 4$  cm in length) compared to large-fruited, megafauna-  
177 adapted lineages ( $\geq 4$  cm in length), especially in the understory of New World tropical forests as well  
178 as on Old World oceanic islands. These results suggest that plant speciation is enhanced by the  
179 evolution of small fruit sizes in conjunction with understory habitat and island colonization. This  
180 directly relates to the dispersal and movement behaviour of particular frugivores, e.g. the spatially-  
181 restricted seed dispersal of small-bodied frugivores in the understory of rainforests or the seed  
182 dispersal to isolated islands by strong-flying frugivores that can cross oceanic barriers. These results  
183 suggest that frugivory-related traits are important drivers of speciation in vertebrate-dispersed tropical  
184 plants, and hence provide trait-based insights into how frugivory might influence biodiversity in the  
185 tropics<sup>28</sup>.

186

187 **Fruit size-dependent speciation.** We hypothesized that speciation of small-fruited palms is higher  
188 than speciation of large-fruited palms (H1, Table 1). This was supported by our results in the global  
189 and Old World analyses (Fig. 2). In general, vertebrate-dispersed plants with small fruit sizes tend to  
190 be dispersed more frequently by small-bodied frugivores than by large-bodied frugivores<sup>6-8</sup>. Since  
191 vertebrate body size scales with home range area, small-bodied frugivores on average have a more

192 restricted space use than large-bodied frugivores<sup>11,12,29</sup>. As a consequence, small-fruited plants  
193 typically show frequent short-distance and rare long-distance dispersal events<sup>3,30</sup>. Large-bodied  
194 frugivores often show large-scale movements<sup>18,31</sup>, which increases the frequency of long-distance  
195 dispersal events, particularly in large-fruited plants<sup>2</sup>. Dispersal distances of small-fruited vertebrate-  
196 dispersed plants therefore tend to be shorter than those of large-fruited plants, which results in lower  
197 gene flow among plant populations and therefore an increased probability of genetic differentiation  
198 and allopatric speciation<sup>2</sup>. Biogeographic comparisons of fruit sizes<sup>5</sup> and studies of disperser loss in  
199 tropical forest fragments<sup>32</sup> also support the idea that seed disperser body size imposes a strong  
200 selective pressure on fruit size. Our results provide macroevolutionary evidence that fruit size can have  
201 a strong influence on diversification dynamics of vertebrate-dispersed plants.

202  
203 **Understory habitat.** Some vertebrate-dispersed plant clades are particularly species-rich in the  
204 understory of tropical rainforests<sup>4,17</sup>. This has been used to hypothesize that understory habitat  
205 generally promotes plant speciation in the tropics (H2, Table 1). Our results (Fig. 3) show that small  
206 fruit size in combination with understory habitat leads to exceptionally high speciation rates in palms,  
207 but only in the New World and not in the Old World. Indeed, several Neotropical understory palm  
208 genera with small fruits (e.g. *Chamaedorea* and *Geonoma*) have been mentioned in support of the  
209 understory habitat hypothesis<sup>4</sup>. These genera along with some other palm genera (i.e. the node leading  
210 to *Desmoncus*, *Bactris* and *Astrocaryum*) represent a diversification rate shift across the palm  
211 phylogenetic tree<sup>33</sup>. Radiations of these palm genera could be, at least partly, driven by the spatially-  
212 restricted movements of the many small-bodied understory frugivores in Neotropical forests<sup>4,16</sup>. For  
213 instance, many Neotropical understory birds show higher genetic differentiation than canopy birds,  
214 indicating low dispersal across biogeographic barriers such as rivers<sup>14,34</sup>. This limited dispersal of  
215 understory frugivores reduces gene flow and ultimately promotes speciation<sup>2,15</sup>. In the Old World,  
216 speciation rates of understory palms were also higher than those of large-fruited canopy palms, but no  
217 additional increase in speciation rates due to small fruit sizes was supported. This may be explained by  
218 the paucity of small-bodied, sedentary understory frugivores in the Old World<sup>16,35</sup> and radiations of  
219 large-bodied, ground-living avian frugivores comparable to, for example, the New World cracids  
220 (Cracidae) and trumpeters (Psophiidae), are relatively rare in the Old World<sup>35,36</sup>. Furthermore, these  
221 results conform to the (relative) scarcity of Old World understory palms at present (36% vs. 45% in  
222 the Old World and New World, respectively). Beyond frugivory, differences in speciation of small-  
223 fruited understory palms between the New World and the Old World may also be explained by the  
224 taller status of the Old World rainforests<sup>37</sup>, potentially due to competition-driven selection for larger  
225 growth forms, or due to historical climate stresses, notably in Africa<sup>38</sup>.

226  
227 **Island colonization.** Many islands show spectacular palm radiations<sup>38-41</sup>, suggesting that island  
228 colonization could be a major driver of palm speciation. Oceanic islands and island-like environments



229 are characterized by restricted colonization and limited gene flow<sup>4,42-45</sup> that can increase speciation  
230 rates relative to the mainland or continental islands (H3, Table 1). Our results (Fig. 4) supported this  
231 hypothesis by demonstrating a particularly high speciation rate for small-fruited palm lineages on Old  
232 World islands. This result is primarily driven by palm diversification in Southeast Asia (95% of the  
233 Old World palm species occur in Indomalaya, Australasia, Pacific and the Western Indian Ocean)  
234 rather than diversification on the relatively species-poor African continent. This was supported by a  
235 similar result when removing the Afrotropical species ( $n = 56$ ) from the analysis (results not shown).  
236 The high diversity of palms on Old World islands<sup>39,46</sup> coincides with a high species richness of large-  
237 bodied, strong-flying avian frugivores in this region, especially the predominance of fruit pigeons  
238 (Columbidae) in Australasia and frugivorous hornbills (Bucerotidae) in Indomalaya<sup>1</sup>. These birds<sup>42</sup> as  
239 well as fruit bats (Pteropodidae)<sup>13</sup> successfully colonize remote islands and thereby contribute to long-  
240 distance seed dispersal of vertebrate-dispersed plants across oceanic barriers. Further empirical studies  
241 provide evidence of frequent long-distance seed dispersal of large-bodied birds at landscape and  
242 biogeographic scales<sup>18,31,47</sup> as well as of dispersal of palm fruits to remote islands<sup>22,48</sup>. Hence, the  
243 diversity of these frugivorous birds may have facilitated island colonisation by palm lineages with  
244 relatively small fruit sizes (i.e. < 4 cm, small enough to be swallowed by ocean-crossing frugivorous  
245 birds and bats) in the Old World. The lack of a relationship between speciation rates, fruit size and  
246 island colonization in the New World may be due to fewer oceanic islands in this part of the world,  
247 congruent with the occurrence of relatively few island-distributed palm species in the New World  
248 compared to the Old World (8% vs. 28% in the New World and Old World, respectively).  
249 Furthermore, many bird families that are widespread in Neotropical rainforests often lack  
250 representative species on oceanic islands<sup>34</sup>, which may have constrained the overall dispersal and  
251 subsequent radiation of Neotropical palms on oceanic islands.

252

253 **Other potential drivers of palm radiations.** Remarkable evolutionary radiations and exceptionally  
254 high diversification rates have been previously identified for various palm genera (e.g.<sup>20,33,38</sup>). Such  
255 fast diversification might not only be driven by interactions with frugivorous vertebrates, but also by  
256 heterogeneity in topography, soils and microenvironments<sup>49-51</sup>, long-term climate and biome  
257 stability<sup>38</sup>, or other types of biotic interactions such as those with herbivores and pathogens<sup>52</sup>. A  
258 potentially confounding factor in terms of correlated evolution between fruit size and plant size<sup>53</sup> was  
259 not supported in our analyses because the negative association between fruit size and speciation  
260 remained after correcting for plant height (see Supplementary Note 1). Beyond fruit sizes, other fruit  
261 traits (e.g. fruit colour, softness, odour, and exposure) or defence traits (e.g. spines) could also  
262 influence the diversification of vertebrate-dispersed plants<sup>54-56</sup>. Moreover, the former presence of a  
263 rich, now largely extinct megafauna (e.g. extinct proboscideans, pilosans, cingulates and  
264 notoungulates) in the Neotropics would have influenced long-distance seed dispersal and gene flow of

265 large-fruited palms<sup>9,10</sup>, and may explain some of the observed differences in diversification between  
266 New World and Old World megafauna-fruited palms (Figs 2-4 and Supplementary Fig. 4).

## 267 **Conclusions**

268 Although several lines of evidence have previously been used to infer a potential role of frugivores in  
269 the diversification of vertebrate-dispersed plants (e.g.<sup>2,4,17,57</sup>), rigorous quantitative tests of specific  
270 hypotheses in a phylogenetic framework have been limited, especially beyond sister clade  
271 comparisons<sup>2</sup>. Using a species-level phylogeny of palms combined with extensive trait datasets, we  
272 demonstrated that speciation rates of palms are highest in small-fruited palm lineages (< 4 cm fruit  
273 size), especially in the understory of New World rainforests as well as in insular environments of  
274 Southeast Asia and the Pacific. In both systems, comparably small fruits probably promote the  
275 establishment of isolated populations through their interaction with particular frugivores. Considering  
276 the evolution of intrinsic traits (e.g. fruit size, growth form) in interaction with geography (e.g. oceanic  
277 islands) and the biotic environment (e.g. frugivores and their movement behaviours) is therefore  
278 essential for understanding plant radiations<sup>58,59</sup>. The combination of time-calibrated phylogenies with  
279 ecological, interaction-relevant traits is thus particularly useful for gaining a deeper understanding of  
280 how biotic interactions have constrained or mediated the evolutionary radiations of tropical plants.

281

## 282 **Methods**

283 **Phylogeny.** We used an all-evidence species-level supertree of palms which includes almost all  
284 accepted palm species ( $n = 2539$ )<sup>23</sup>. This time-calibrated, phylogenetic tree is based on a backbone  
285 generated from nine plastid and four nuclear markers as well as morphological data<sup>60</sup>, and additional  
286 molecular and morphological data for several genera<sup>23</sup>. The phylogenetic tree was dated using five  
287 calibration points<sup>21</sup>. A Bayesian modeling approach was used to place species without genetic or  
288 morphological data in the phylogeny, based on taxonomy (for details see ref.<sup>23</sup>). As this leads to  
289 uncertainty in the exact placement of a species within the phylogeny, all analyses were performed on a  
290 set of 100 randomly sampled palm phylogenetic trees available from ref.<sup>23</sup>.

291

292 **Data on fruit sizes.** Information on fruit sizes was collected for a total of 1836 palm species (ca. 70%  
293 of all palm species) from various sources, including primary literature, monographs, herbaria and palm  
294 websites (all sources are listed in Supplementary data sources). Specifically, we calculated the average  
295 fruit length for each species (based on multiple records per species if available), because fruit length is  
296 the most commonly reported fruit size trait in monographs and species descriptions. For the analyses,  
297 we classified species into two groups: small-fruited palms (fruits < 4 cm in length) and large-fruited  
298 palms (fruits  $\geq$  4 cm in length). Since palm fruits are usually single-seeded<sup>20</sup>, the large-fruited palms  
299 represent species with ‘megafaunal’ fruits<sup>9,10</sup>. Across the palm family, at least 229 palm species have

300 megafaunal fruits (Fig. 1), and about one third of the palm genera have at least one species with such  
301 fruits (Supplementary Table 1). We used the binary state of fruit size (small/large) rather than a  
302 continuous variable because the implemented diversification models (see below) deal with binary data,  
303 and because species with large, megafaunal fruit sizes are dependent on seed dispersal by large-bodied  
304 mammalian frugivores (megafauna), thereby excluding volant frugivores (birds, bats) and small- and  
305 medium-sized frugivores. In contrast, species with small fruit sizes are predominantly ingested by  
306 birds and small- and medium-sized mammalian seed dispersers<sup>2</sup>.

307 **Data on understory habitat.** To quantify affiliation with the forest understory, we compiled species-  
308 level data on maximum stem height for 2073 palm species (ca. 81% of all palm species) from the same  
309 sources as used for the fruit size data (see Supplementary data sources). For all palm species, we  
310 additionally determined their main growth form (climber, acaulescence, erect shrub/tree). From these  
311 data, we estimated whether palms present their fruits in the understory. This included short-stemmed  
312 palms (maximum stem height  $\leq 5$  m) as well as all acaulescent species (i.e. having no or only a very  
313 short stem concealed in the ground). Palms with a stem height  $> 5$  m were considered to be non-  
314 understory plants, i.e. tall-stemmed or medium-sized palms and most climbers (referred to as  
315 ‘canopy’).

316  
317 **Data on island colonization.** The palm family has a pantropical distribution (i.e. it occurs in all  
318 tropical regions). To quantify species distributions on islands, we compiled global presence-absence  
319 data for all palm species from the world checklist of palms<sup>61</sup>. This exhaustive, authoritative checklist  
320 records palm species occurrence within level 3 geographic units as defined by the International  
321 Working Group on Taxonomic Databases (TDWG) (referred to as ‘botanical countries’)<sup>61</sup>. These  
322 generally correspond to countries although larger countries such as the United States are normally  
323 broken down into smaller political units. Palm occurrence data are freely available from the  
324 continuously updated World Checklist of Monocotyledons (<http://apps.kew.org/wcsp>), and we here  
325 used a database version downloaded on July 2015. For our analyses, we defined a binary state  
326 (island/mainland) describing whether a species occurs on volcanic and atoll islands (referred to as  
327 ‘island’), or on the mainland or on continental islands (referred to as ‘mainland’), following the  
328 classification from ref<sup>62</sup>. This classification follows geology as a surrogate for isolation, in which  
329 oceanic and atoll islands have arisen as newly formed land, whereas continental islands are either part  
330 of the continental shelf or were once connected to continental landmasses (e.g. Madagascar). The  
331 occurrence of palm species on oceanic and atoll islands consequently must have resulted from  
332 colonization and speciation in isolation, whereas palm lineages on continental islands have  
333 experienced less isolation. This classification closely matches the classification based on the  
334 connectivity of islands to the mainland during the last glacial maximum, as quantified by ref.<sup>63</sup>.

335

336 **Performance of trait-based models.** In this study, we tested the impact of specific traits on the  
337 diversification of lineages using the maximum likelihood based ‘state speciation and extinction’ or  
338 ‘SSE’ models<sup>25,26</sup>. These methods calculate the probability that a lineage evolved as observed given a  
339 model of character evolution. However, SSE models have recently been criticized for high type I error  
340 rates<sup>64</sup>, suggesting that a significant effect of a trait on speciation rates can be detected even if it is not  
341 truly present. We evaluated this bias by performing simulations in which neutral binary traits evolved  
342 on 100 empirical palm phylogenies<sup>23</sup> under several transition rate scenarios, as recommended by ref.<sup>64</sup>.

343 These simulated (neutral) binary traits are expected to be neutral with respect to speciation  
344 rates. We evaluated the Bayesian credible intervals in speciation rates between these simulated binary  
345 traits after running a Bayesian Markov Chain Monte Carlo (MCMC) for 10,000 generations on the 100  
346 palm phylogenies. Our results showed a strong overlap of Bayesian credible intervals between  
347 character states on speciation rates under all transition rate scenarios (Supplementary Fig. 3),  
348 supporting the expectation that these neutral traits do not affect speciation rates in palms. This  
349 suggests that the empirical results can be reliably obtained from the ‘SSE’ models. In addition, our  
350 dataset meets the other requirements for applying ‘SSE’ models, such as sufficient replication events  
351 (e.g. the independent evolution of small fruit sizes)<sup>65</sup>, > 300 species and balanced character state  
352 distributions<sup>66</sup> (for more details on these simulations see Supplementary Note 2).

353  
354 **Fruit size-dependent speciation.** We used the Binary State Speciation and Extinction (BiSSE)  
355 model<sup>25,26</sup> implemented in the ‘diversitree’ package<sup>27</sup> in R<sup>67</sup> to model speciation ( $\lambda$ ), extinction ( $\mu$ )  
356 and transition ( $q$ ) rates of small-fruited vs. large-fruited palm lineages (H1 in Table 1). The BiSSE  
357 model jointly estimates speciation, extinction and transition rates of a binary trait by using dated  
358 phylogenetic trees, and trait states assigned to the species at the tips of the trees. We focus on  
359 speciation rates because our hypotheses (Table 1) directly refer to speciation rather than extinction or  
360 net diversification. Nevertheless, the joint-estimation of these rates is desirable as trait changes may  
361 not be independent from speciation and extinction rates<sup>26</sup>. We report and discuss all evolutionary rates  
362 other than speciation in Supplementary Note 3 and Supplementary Fig. 4, and provide an overview of  
363 the model selection globally as well as for the New World and Old World, respectively  
364 (Supplementary Tables 2–4). We fitted eight BiSSE models with decreasing complexity (parameters)  
365 and selected the best-fitting models based on likelihood-ratio tests under a Chi-square distribution and  
366 the Akaike Information Criterion (AIC) (Supplementary Tables 2–4). These models included  
367 constraints on speciation, extinction and transition rates between trait states<sup>25</sup>. Maximum likelihood  
368 was used to optimize full and constrained models. BiSSE enables correcting for species and their traits  
369 not sampled in the datasets by indicating a sampling fraction, i.e. 32% of small-fruited and 18% of  
370 large-fruited palm species were not sampled in the global dataset. This fraction was based on imputed  
371 trait values from the PhyloPars algorithm<sup>68</sup> for those species sampled in the phylogenetic tree but  
372 lacking fruit size data (these imputed trait values were only used to calculate sampling fractions, not in

373 the actual analyses). A MCMC was run for the best-fitting model for 10,000 generations on 100  
374 randomly sampled palm phylogenies. We evaluated the posterior distribution of these Bayesian rates,  
375 and in case the 95% Bayesian credibility intervals between parameter states did not overlap, we  
376 considered them significantly different from each other<sup>25</sup>.

377 All analyses were run globally as well as separately for the New World (including South,  
378 Central and North America as well as the Caribbean) and the Old World (including Australia,  
379 Indomalaya, Oceania, Pacific, Africa, Madagascar and surrounding islands). This geographic division  
380 was used because most palm species (as well as genera) are endemic to one of these regions<sup>38</sup>. Hence,  
381 these regions are characterized by distinct historical differences in terms of palm diversification<sup>33,69</sup>,  
382 frugivore communities<sup>1</sup>, and representation of understory palm species (45% and 36% in the New  
383 World and Old World, respectively) and island-distributed palm species (8% and 28% in the New  
384 World and in the Old World, respectively).

385 As results may be biased by the allometric relationship between fruit size and plant size, we  
386 repeated the analyses after accounting for the correlation between palm maximum stem heights and  
387 fruit sizes. The effect of residual fruit sizes (after correcting for maximum plant height) on speciation  
388 rates was assessed with BiSSE (for details on the approach see Supplementary Note 1, for model  
389 selection see Supplementary Table 5 and for results see Supplementary Figs. 1 and 4). However, as  
390 these residuals do not represent ‘true’ small and large fruits, we also investigated the effect of residual  
391 fruit sizes on speciation rates using the Quantitative Speciation and Extinction model (QuaSSE)<sup>70</sup>.  
392 QuaSSE can be used to test the effect of a continuous trait on speciation rates by testing the fit of  
393 models describing the distribution of the response (i.e. speciation rate) to the trait (e.g. constant, linear  
394 or sigmoidal). Details on the methods and results of this analysis are provided in Supplementary Note  
395 1, Supplementary Table 6 and Supplementary Fig. 5.

396 Furthermore, to test whether our binary classification of fruit size biased the results, we  
397 additionally tested the effect of fruit size as a continuous trait on speciation rates. We first estimated  
398 speciation rate heterogeneity across the phylogeny with a Bayesian Analysis of Macroevolutionary  
399 Mixtures (BAMM)<sup>71</sup> and then compared the observed difference in speciation rate between palms that  
400 exhibit different fruit sizes to a background speciation rate through randomizing the estimated tip  
401 speciation rates from the BAMM outputs. These additional analyses also confirmed the high  
402 speciation rate of small-fruited palm lineages. Details on the methods and results of this analysis are  
403 provided in the Supplementary Note 4 and Supplementary Fig. 6.

404

405 **Speciation rates due to understory habitat and island colonization.** The effects of a trait (e.g. fruit  
406 size) on speciation rates may be enhanced by an interaction effect with another trait. For example,  
407 palms in the understory or on islands may have particularly high speciation rates if they have also  
408 small fruits. To disentangle such effects we implemented the Multiple State Speciation and Extinction  
409 model (MuSSE multistate)<sup>27</sup>. The MuSSE model can be used to quantify the additive and interactive

410 effects of two binary traits (e.g. small fruit size and understory growth form, or small fruit size and  
411 island colonization) on speciation, extinction and transition rates. The model intercept of the MuSSE  
412 model (the ‘base model’) estimates speciation rates when both traits are absent (e.g. palm lineages  
413 with large fruits that do not grow in the understory). The interaction term (when both traits are present)  
414 will indicate whether these traits may interact in either a positive way (i.e. both traits increase  
415 speciation rates) or a negative way (i.e. both traits decrease speciation rates).

416 To quantify trait-dependent diversification for both binary trait combinations (H2: small/large  
417 fruit size and understory/canopy habitat; H3: small/large fruit size and island/mainland distribution),  
418 we compared the likelihood of a total of sixteen models with increasing complexity (Supplementary  
419 Table 7). We used stepwise AIC model selection, globally as well as separately for the New World  
420 and Old World, and selected the models with the lowest AIC (Supplementary Tables 8–9). A MCMC  
421 for the best-fitting model (based on AIC) was run for 10,000 generations on 100 palm phylogenies.  
422 We tested for the additive and interactive effects of small fruits and understory habitat (compared to  
423 large-fruited canopy palms) on speciation rates (Supplementary Table 8), and between small fruits and  
424 oceanic island colonization (compared to large-fruited mainland/continental island-distributed palms)  
425 on speciation rates (Supplementary Table 9). We report and discuss all evolutionary rates other than  
426 speciation rates in Supplementary Note 5 and Supplementary Fig. 7.

427

428 **Data availability.** The phylogenetic data that support the findings of this study are available from  
429 ref.<sup>23</sup>. The palm species distribution data are available from the World Checklist of Selected Plant  
430 Families (<http://apps.kew.org/wcsp>). All scripts to perform the analyses in this study are available  
431 upon request from the first author [REO]. The palm trait data that support the findings of this study are  
432 available upon request from the last author [WDK].

433 **References**

434

- 435 1 Kissling, W. D., Böhning-Gaese, K. & Jetz, W. The global distribution of frugivory in birds.  
 436 *Global Ecology and Biogeography* **18**, 150-162, doi:10.1111/j.1466-8238.2008.00431.x  
 437 (2009).
- 438 2 Fleming, T. H. & Kress, W. J. *The ornaments of life: coevolution and conservation in the*  
 439 *tropics*. (Chicago University Press, 2013).
- 440 3 Nathan, R. & Muller-Landau, H. C. Spatial patterns of seed dispersal, their determinants and  
 441 consequences for recruitment. *Trends in Ecology & Evolution* **15**, 278-285,  
 442 doi:[http://dx.doi.org/10.1016/S0169-5347\(00\)01874-7](http://dx.doi.org/10.1016/S0169-5347(00)01874-7) (2000).
- 443 4 Givnish, T. J. Ecology of plant speciation. *Taxon* **59**, 1326-1366 (2010).
- 444 5 Lord, J. M. Frugivore gape size and the evolution of fruit size and shape in southern  
 445 hemisphere floras. *Austral Ecology* **29**, 430-436 (2004).
- 446 6 Wheelwright, N. T. Fruit-Size, gape width, and the diets of fruit-eating birds. *Ecology* **66**,  
 447 808-818, doi:10.2307/1940542 (1985).
- 448 7 Jordano, P. in *Seeds: the ecology of regeneration in plant communities 2* 125-166 (2000).
- 449 8 Herrera, C. M. in *Plant-animal interactions: an evolutionary approach* (eds Carlos M.  
 450 Herrera & Olle Pellmyr) 185-208 (John Wiley & Sons, 2002).
- 451 9 Guimarães, P. R., Jr., Galetti, M. & Jordano, P. Seed dispersal anachronisms: rethinking the  
 452 fruits extinct megafauna ate. *PLoS ONE* **3**, e1745, doi:10.1371/journal.pone.0001745 (2008).
- 453 10 Janzen, D. H. & Martin, P. S. Neotropical anachronisms: the fruits the gomphotheres ate.  
 454 *Science* **215**, 19-27, doi:10.1126/science.215.4528.19 (1982).
- 455 11 Haskell, J. P., Ritchie, M. E. & Olff, H. Fractal geometry predicts varying body size scaling  
 456 relationships for mammal and bird home ranges. *Nature* **418**, 527-530 (2002).
- 457 12 Milton, K. & May, M. L. Body weight, diet and home range area in primates. *Nature* **259**,  
 458 459-462 (1976).
- 459 13 Shanahan, M., Harrison, R. D., Yamuna, R., Boen, W. & Thornton, I. W. B. Colonization of  
 460 an island volcano, Long Island, Papua New Guinea, and an emergent island, Motmot, in its  
 461 caldera lake. V. colonization by figs (*Ficus* spp.), their dispersers and pollinators. *Journal of*  
 462 *Biogeography* **28**, 1365-1377 (2001).
- 463 14 Burney, C. W. & Brumfield, R. T. Ecology predicts levels of genetic differentiation in  
 464 neotropical birds. *The American Naturalist* **174**, 358-368, doi:10.1086/603613 (2009).
- 465 15 Salisbury, C. L., Seddon, N., Cooney, C. R. & Tobias, J. A. The latitudinal gradient in  
 466 dispersal constraints: ecological specialisation drives diversification in tropical birds. *Ecology*  
 467 *Letters* **15**, 847-855, doi:10.1111/j.1461-0248.2012.01806.x (2012).
- 468 16 Karr, J. R. Geographical variation in the avifaunas of tropical forest undergrowth. *The Auk* **97**,  
 469 283-298 (1980).
- 470 17 Smith, J. F. High species diversity in fleshy-fruited tropical understory plants. *The American*  
 471 *Naturalist* **157**, 646-653, doi:10.1086/320625 (2001).
- 472 18 Holbrook, K. M., Smith, T. B. & Hardesty, B. D. Implications of long-distance movements of  
 473 frugivorous rain forest hornbills. *Ecography* **25**, 745-749, doi:10.1034/j.1600-  
 474 0587.2002.250610.x (2002).
- 475 19 Losos, J. B. & Ricklefs, R. E. Adaptation and diversification on islands. *Nature* **457**, 830-836  
 476 (2009).
- 477 20 Dransfield, J. *et al.* *Genera Palmarum: The evolution and classification of palms*. (Kew  
 478 Publishing, 2008).
- 479 21 Couvreur, T. L. P., Forest, F. & Baker, W. J. Origin and global diversification patterns of  
 480 tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC*  
 481 *Biology* **9**, 1-12, doi:10.1186/1741-7007-9-44 (2011).
- 482 22 Zona, S. & Henderson, A. A review of animal-mediated seed dispersal of palms. *Selbyana* **11**,  
 483 6-21 (1989).
- 484 23 Faurby, S., Eiserhardt, W. L., Baker, W. J. & Svenning, J.-C. An all-evidence species-level  
 485 supertree for the palms (Arecaceae). *Molecular Phylogenetics and Evolution* **100**, 57-69,  
 486 doi:<http://dx.doi.org/10.1016/j.ympev.2016.03.002> (2016).

- 487 24 Brummitt, R. K., Pando, F., Hollis, S. & Brummitt, N. *World geographical scheme for*  
488 *recording plant distributions*. (International Working Group on Taxonomic Databases for  
489 Plant Sciences (TDWG), 2001).
- 490 25 FitzJohn, R. G., Maddison, W. P. & Otto, S. P. Estimating trait-dependent speciation and  
491 extinction rates from incompletely resolved phylogenies. *Systematic Biology* **58**, 595-611,  
492 doi:10.1093/sysbio/syp067 (2009).
- 493 26 Maddison, W. P., Midford, P. E. & Otto, S. P. Estimating a binary character's effect on  
494 speciation and extinction. *Systematic Biology* **56**, 701-710, doi:10.1080/10635150701607033  
495 (2007).
- 496 27 FitzJohn, R. G. Diversitree: comparative phylogenetic analyses of diversification in R.  
497 *Methods in Ecology and Evolution* **3**, 1084-1092, doi:10.1111/j.2041-210X.2012.00234.x  
498 (2012).
- 499 28 Richardson, J. E. & Pennington, R. T. Editorial: Origin of tropical diversity: from clades to  
500 communities. *Frontiers in Genetics* **7**, doi:10.3389/fgene.2016.00186 (2016).
- 501 29 Jetz, W., Carbone, C., Fulford, J. & Brown, J. H. The scaling of animal space use. *Science*  
502 **306**, 266-268, doi:10.1126/science.1102138 (2004).
- 503 30 Jordano, P., García, C., Godoy, J. A. & García-Castaño, J. L. Differential contribution of  
504 frugivores to complex seed dispersal patterns. *Proceedings of the National Academy of*  
505 *Sciences* **104**, 3278-3282, doi:10.1073/pnas.0606793104 (2007).
- 506 31 Lenz, J. *et al.* Seed-dispersal distributions by trumpeter hornbills in fragmented landscapes.  
507 *Proceedings of the Royal Society B: Biological Sciences* **278**, 2257-2264,  
508 doi:10.1098/rspb.2010.2383 (2011).
- 509 32 Galetti, M. *et al.* Functional extinction of birds drives rapid evolutionary changes in seed size.  
510 *Science* **340**, 1086-1090, doi:10.1126/science.1233774 (2013).
- 511 33 Baker, W. J. & Couvreur, T. L. P. Global biogeography and diversification of palms sheds  
512 light on the evolution of tropical lineages. II. Diversification history and origin of regional  
513 assemblages. *Journal of Biogeography* **40**, 286-298, doi:10.1111/j.1365-2699.2012.02794.x  
514 (2013).
- 515 34 Moore, R. P., Robinson, W. D., Lovette, I. J. & Robinson, T. R. Experimental evidence for  
516 extreme dispersal limitation in tropical forest birds. *Ecology Letters* **11**, 960-968,  
517 doi:10.1111/j.1461-0248.2008.01196.x (2008).
- 518 35 Corlett, R. T. Frugivory and seed dispersal by vertebrates in tropical and subtropical Asia: An  
519 update. *Global Ecology and Conservation* **11**, 1-22,  
520 doi:<https://doi.org/10.1016/j.gecco.2017.04.007> (2017).
- 521 36 Fleming, T. H., Breitwisch, R. & Whitesides, G. H. Patterns of tropical vertebrate frugivore  
522 diversity. *Annual Review of Ecology and Systematics* **18**, 91-109,  
523 doi:10.1146/annurev.es.18.110187.000515 (1987).
- 524 37 Banin, L. *et al.* What controls tropical forest architecture? Testing environmental, structural  
525 and floristic drivers. *Global Ecology and Biogeography* **21**, 1179-1190, doi:10.1111/j.1466-  
526 8238.2012.00778.x (2012).
- 527 38 Kissling, W. D. *et al.* Cenozoic imprints on the phylogenetic structure of palm species  
528 assemblages worldwide. *Proceedings of the National Academy of Sciences of the United*  
529 *States of America* **109**, 7379-7384, doi:10.1073/pnas.1120467109 (2012).
- 530 39 Baker, W. J. & Couvreur, T. L. P. in *Biotic evolution and environmental change in Southeast*  
531 *Asia* (eds D. Gower *et al.*) 164-190 (Cambridge University Press, 2012).
- 532 40 Bacon, C. D., Baker, W. J. & Simmons, M. P. Miocene dispersal drives island radiations in  
533 the palm tribe Trachycarpeae (Arecaceae). *Systematic Biology* **61**, 426-442,  
534 doi:10.1093/sysbio/syr123 (2012).
- 535 41 Dransfield, J. & Beentje, H. *The palms of Madagascar*. (Royal Botanic Gardens Kew and The  
536 International Palm Society, 1995).
- 537 42 Diamond, J. M., Gilpin, M. E. & Mayr, E. Species-distance relation for birds of the Solomon  
538 Archipelago, and the paradox of the great speciators. *Proceedings of the National Academy of*  
539 *Sciences of the United States of America* **73**, 2160-2164 (1976).
- 540 43 Moyle, R. G., Filardi, C. E., Smith, C. E. & Diamond, J. Explosive Pleistocene diversification  
541 and hemispheric expansion of a "great speciator". *Proceedings of the National Academy of*  
542 *Sciences* **106**, 1863-1868, doi:10.1073/pnas.0809861105 (2009).



- 543 44 Baldwin, B. G. & Sanderson, M. J. Age and rate of diversification of the Hawaiian  
544 silversword alliance (Compositae). *Proceedings of the National Academy of Sciences* **95**,  
545 9402-9406 (1998).
- 546 45 Hughes, C. & Eastwood, R. Island radiation on a continental scale: Exceptional rates of plant  
547 diversification after uplift of the Andes. *Proceedings of the National Academy of Sciences*  
548 **103**, 10334-10339, doi:10.1073/pnas.0601928103 (2006).
- 549 46 Kissling, W. D. *et al.* Quaternary and pre-Quaternary historical legacies in the global  
550 distribution of a major tropical plant lineage. *Global Ecology and Biogeography* **21**, 909-921,  
551 doi:10.1111/j.1466-8238.2011.00728.x (2012).
- 552 47 Kitamura, S. Frugivory and seed dispersal by hornbills (Bucerotidae) in tropical forests. *Acta*  
553 *Oecologica* **37**, 531-541 (2011).
- 554 48 Morici, C. in *Ecologia Insular* (eds JM Fernandez-Palacios & C Morici) 81-122  
555 (Asociacion Española de Ecología Terrestre, 2004).
- 556 49 Pintaud, J.-C. & Jaffre, T. Patterns of diversity and endemism in palms on ultramafic rocks in  
557 New Caledonia. *South African Journal of Science* **97**, 548-550 (2001).
- 558 50 Sanín, M. J. *et al.* The Neogene rise of the tropical Andes facilitated diversification of wax  
559 palms (Ceroxylon: Arecaceae) through geographical colonization and climatic niche  
560 separation. *Botanical Journal of the Linnean Society* **182**, 303-317, doi:10.1111/boj.12419  
561 (2016).
- 562 51 Svenning, J.-C. On the role of microenvironmental heterogeneity in the ecology and  
563 diversification of neotropical rain-forest palms (Arecaceae). *The Botanical Review* **67**, 1-53,  
564 doi:10.1007/BF02857848 (2001).
- 565 52 Eiserhardt, W. L., Svenning, J.-C., Kissling, W. D. & Balslev, H. Geographical ecology of the  
566 palms (Arecaceae): determinants of diversity and distributions across spatial scales. *Annals of*  
567 *Botany* **108**, 1391-1416, doi:10.1093/aob/mcr146 (2011).
- 568 53 Herrera, C. M. Interspecific Variation in Fruit Shape: Allometry, Phylogeny, and Adaptation  
569 to Dispersal Agents. *Ecology* **73**, 1832-1841, doi:10.2307/1940034 (1992).
- 570 54 Lomáscolo, S. B., Levey, D. J., Kimball, R. T., Bolker, B. M. & Alborn, H. T. Dispersers  
571 shape fruit diversity in *Ficus* (Moraceae). *Proceedings of the National Academy of Sciences*  
572 **107**, 14668-14672, doi:10.1073/pnas.1008773107 (2010).
- 573 55 Voigt, F. A. *et al.* A comparison of morphological and chemical fruit traits between two sites  
574 with different frugivore assemblages. *Oecologia* **141**, 94-104, doi:10.1007/s00442-004-1654-8  
575 (2004).
- 576 56 Dominy, N. J., Svenning, J. C. & Li, W. H. Historical contingency in the evolution of primate  
577 color vision. *Journal of Human Evolution* **44**, 25-45, doi:10.1016/s0047-2484(02)00167-7  
578 (2003).
- 579 57 Ricklefs, R. E. & Renner, S. S. Species richness within families of flowering plants. *Evolution*  
580 **48**, 1619-1636, doi:10.2307/2410252 (1994).
- 581 58 Bouchenak-Khelladi, Y., Onstein, R. E., Xing, Y., Schwery, O. & Linder, H. P. On the  
582 complexity of triggering evolutionary radiations. *New Phytologist* **207**, 313-326,  
583 doi:10.1111/nph.13331 (2015).
- 584 59 Onstein, R. E. & Linder, H. P. Beyond climate: convergence in fast evolving sclerophylls in  
585 Cape and Australian Rhamnaceae predates the mediterranean climate. *Journal of Ecology* **104**,  
586 665-677, doi:10.1111/1365-2745.12538 (2016).
- 587 60 Baker, W. J. *et al.* Complete generic-level phylogenetic analyses of palms (Arecaceae) with  
588 comparisons of supertree and supermatrix approaches. *Syst Biol* **58**, 240-256 (2009).
- 589 61 Govaerts, R. & Dransfield, J. *World checklist of palms.*, (Royal Botanic Gardens Kew, 2005).
- 590 62 Kreft, H., Jetz, W., Mutke, J., Kier, G. & Barthlott, W. Global diversity of island floras from a  
591 macroecological perspective. *Ecology Letters* **11**, 116-127, doi:10.1111/j.1461-  
592 0248.2007.01129.x (2008).
- 593 63 Weigelt, P., Jetz, W. & Kreft, H. Bioclimatic and physical characterization of the world's  
594 islands. *Proceedings of the National Academy of Sciences* **110**, 15307-15312,  
595 doi:10.1073/pnas.1306309110 (2013).
- 596 64 Rabosky, D. L. & Goldberg, E. E. Model inadequacy and mistaken inferences of trait-  
597 dependent speciation. *Systematic Biology* **64**, 340-355, doi:10.1093/sysbio/syu131 (2015).

- 598 65 Maddison, W. P. & FitzJohn, R. G. The unsolved challenge to phylogenetic correlation tests  
599 for categorical characters. *Systematic Biology* **64**, 127-136, doi:10.1093/sysbio/syu070 (2015).
- 600 66 Davis, M. P., Midford, P. E. & Maddison, W. Exploring power and parameter estimation of  
601 the BiSSE method for analyzing species diversification. *BMC Evolutionary Biology* **13**, 1-11,  
602 doi:10.1186/1471-2148-13-38 (2013).
- 603 67 R Development Core Team. *R: a language and environment for statistical computing, version*  
604 *3.1.0 (2014-04-10)*, <<http://www.R-project.org>> (2014).
- 605 68 Bruggeman, J., Heringa, J. & Brandt, B. W. PhyloPars: estimation of missing parameter  
606 values using phylogeny. *Nucleic Acids Research* **37**, W179-W184, doi:10.1093/nar/gkp370  
607 (2009).
- 608 69 Baker, W. J. & Couvreur, T. L. P. Global biogeography and diversification of palms sheds  
609 light on the evolution of tropical lineages. I. Historical biogeography. *Journal of*  
610 *Biogeography* **40**, 274-285, doi:10.1111/j.1365-2699.2012.02795.x (2013).
- 611 70 FitzJohn, R. G. Quantitative Traits and Diversification. *Systematic Biology* **59**, 619-633,  
612 doi:10.1093/sysbio/syq053 (2010).
- 613 71 Rabosky, D. L. Automatic Detection of Key Innovations, Rate Shifts, and Diversity-  
614 Dependence on Phylogenetic Trees. *PLOS ONE* **9**, e89543, doi:10.1371/journal.pone.0089543  
615 (2014).
- 616

## 617 **Acknowledgements**

618 We thank Henrik Balslev, Anders Barfod, Anne Blach-Overgaard, Finn Borchsenius, John Dransfield,  
619 Wolf Eiserhardt and María José Sanín for discussions about palm biology, John Dransfield and  
620 Andrew J. Henderson for the use of pictures for figure 1. W.D.K. was supported by the University of  
621 Amsterdam (starting grant), the Danish Council for Independent Research–Natural Sciences (grant 11-  
622 106163), and the Netherlands Organisation for Scientific Research (grant 824.15.007). W.J.B. was  
623 supported by a grant from the Garfield Weston Foundation to the Global Tree Seed Bank Project at the  
624 Royal Botanic Gardens, Kew. J.C.S. was supported by the European Research Council (ERC-2012-  
625 StG-310886-HISTFUNC), and also considers this work a contribution to his VILLUM Investigator  
626 project “Biodiversity Dynamics in a Changing World” funded by VILLUM FONDEN.

627

## 628 **Author contributions**

629 W.D.K. conceived the idea, W.D.K. and R.E.O. designed the study; W.D.K. and R.E.O. collected  
630 data; R.E.O. analysed the data; R.E.O. and W.D.K. wrote the manuscript; all authors discussed the  
631 results and commented on the manuscript.

## 632 **Additional information**

633 **Supplementary information** is available for this paper.

634 **Reprints and permissions** information is available at [www.nature.com/reprints](http://www.nature.com/reprints).

635 **Correspondence and requests for materials** should be addressed to R.E.O and W.D.K.

636 **How to cite this article:** TO BE ADDED.

637 **Competing interests**

638 The authors declare no competing financial interests.

639  
640**Table 1 Key hypotheses of how frugivorous vertebrates influence plant speciation.**

Hypotheses	Explanation	Case studies and lines of evidence	References
<b>H1: Fruit size</b> Speciation of plants with small fruits is higher than speciation of large-fruited plants	Dispersal of large-fruited plants by large-bodied frugivores leads to high gene flow among plant populations and therefore reduces the probability of genetic differentiation and allopatric speciation <sup>2</sup>	<ol style="list-style-type: none"> <li>1. Fruit size of vertebrate-dispersed plants correlates with body size (or gape width) of their frugivorous consumers so that large-fruited plants are dispersed by large-bodied frugivores</li> <li>2. Large-bodied birds and mammals have larger home ranges and movement distances than small-bodied species</li> <li>3. Fruit sizes of species in trans-regional plant genera are smaller in regions where large-bodied mammalian frugivores have been absent (e.g. New Zealand) compared to regions where plants have evolved in the presence of such frugivores (e.g. Australia)</li> <li>4. Fruit sizes of vertebrate-dispersed plants become smaller at sites where large-bodied frugivores have become functionally extinct</li> </ol>	6,8 11,12 5 32
<b>H2: Understorey habitat</b> Plants with small, vertebrate-dispersed fruits in the understorey of tropical moist and wet forests have higher speciation rates than taller plants	Vertebrate-dispersed understorey plants have low gene flow among populations because dispersal distance by sedentary understorey frugivores is spatially restricted. This increases the probability of genetic differentiation and allopatric speciation <sup>4</sup>	<ol style="list-style-type: none"> <li>1. Avian seed dispersers in the understorey show spatially restricted dispersal and therefore higher genetic divergence and more subspecies than canopy birds</li> <li>2. Several vertebrate-dispersed plant genera in tropical forest understoreies show an extraordinary species richness</li> <li>3. Clades of Neotropical vertebrate-dispersed understorey plants are more species rich than understorey sister clades with dry fruits</li> </ol>	14,15 4 17
<b>H3: Island colonization</b> Vertebrate-dispersed plants on oceanic islands have higher speciation rates than plants on the mainland or on continental islands	Oceanic barriers lead to restricted colonization and limited gene flow among plant populations which increases the probability of genetic differentiation and allopatric speciation <sup>4</sup>	<ol style="list-style-type: none"> <li>1. Avian seed dispersers show higher speciation rates on oceanic island archipelagos than on the mainland, possibly because limited dispersal leads to high genetic differentiation</li> <li>2. Rapid radiations of plants have been described for oceanic island archipelagos as well as island-like mountainous habitats</li> <li>3. Long-distance dispersal of vertebrate-dispersed plants to oceanic islands or fragmented habitat patches is dependent on occasional movement of large-bodied volant frugivores (e.g. hornbills, fruit pigeons and fruit bats)</li> </ol>	42,43 44, 45 13,18

641

642 **Figures**


643

644 **Figure 1 | Global variation in palm fruit size.** (a) Fruit sizes of all species in the analysis ( $n = 1836$ 645 extant palm species). The classification of small ( $< 4$  cm) and large ( $\geq 4$  cm) fruits as used in the646 models in this study is indicated, as well as the median ( $m$ ) and sample size ( $n$ ) for each of these

647 groups. (b–i) Pictures illustrating the diversity of vertebrate-dispersed fruits in palms, representing (b–

648 e) small-fruited and (f–i) large-fruited palms. (b) *Iguanura elegans* (John Dransfield); (c) *Pinanga*649 *disticha* (John Dransfield); (d) *Calamus erioacanthus* (John Dransfield); (e) *Ravenea dransfieldii*650 (John Dransfield); (f) *Manicaria saccifera* (John Dransfield); (g) *Mauritia flexuosa* (Andrew J.651 Henderson); (h) *Pholidocarpus sumatranus* (John Dransfield); (i) *Metroxylon sagu* (William J. Baker).

652




653

654

655

656

657 **Figure 2 | Speciation rate estimates for palm lineages with small (< 4 cm) and large ( $\geq$  4 cm)**  
 658 **fruits.** Rates are inferred (a) globally as well as separately for (b) the New World (the Americas) and  
 659 (c) the Old World (Africa, Asia and Australia) using Binary State Speciation and Extinction (BiSSE)  
 660 models with 100 palm phylogenies. Box-and-whiskers indicate the 95% Bayesian credibility intervals  
 661 of the speciation rates as estimated through Bayesian Markov Chain Monte Carlo methods. Small-  
 662 fruited palm lineages show higher speciation rates than large-fruited palms globally and in the Old  
 663 World, but not in the New World (where the best model suggested an equal rate of both large and  
 664 small-fruited palms, see Supplementary Table 3).  
 665




666

667

668


669 **Figure 3 | Understory habitat and its effect on speciation rates for palm lineages with small (<4**  
 670 **cm) fruits.** Rates are inferred (a) globally as well as separately for (b) the New World (the Americas)  
 671 and (c) the Old World (Africa, Asia and Australia) using Multiple State Speciation and Extinction  
 672 (MuSSE) models with 100 palm phylogenies. Box-and-whiskers indicate the 95% Bayesian credibility  
 673 intervals of the speciation rates as estimated through Bayesian Markov Chain Monte Carlo methods.  
 674 The base model indicates rates of large-fruited ( $\geq 4$  cm) canopy palms. Small fruits and understory  
 675 habitat both add positively to the speciation rate as compared to the base model. In the global and New  
 676 World analyses an interaction term was supported, suggesting the highest rates for small-fruited  
 677 understory palms. No value is given for the interaction for the Old World since a model without it was  
 678 preferred by AIC (indicated by *n.a.* [not applicable]).  
 679



680

681

682 **Figure 4 | Island colonization and its effect on speciation rates for palm lineages with small (<4**  
 683 **cm) fruits.** Rates are inferred (a) globally as well as separately for (b) the New World (the Americas)  
 684 and (c) the Old World (Africa, Asia and Australia) using Multiple State Speciation and Extinction  
 685 (MuSSE) models with 100 palm phylogenies. Box-and-whiskers indicate the 95% Bayesian credibility  
 686 intervals of the speciation rates as estimated through Bayesian Markov Chain Monte Carlo methods.  
 687 The base model indicates rates of large-fruited ( $\geq 4$  cm) mainland or continental island-distributed  
 688 palms. Small fruits and island distribution both add positively to the speciation rate as compared to the  
 689 base model. In the global and Old World analyses an interaction term was supported, suggesting the  
 690 highest rates for small-fruited island-distributed palms. No value is given for the interaction for the  
 691 New World since a model without it was preferred by AIC (indicated by *n.a.* [not applicable]).  
 692



693



## SUPPLEMENTARY INFORMATION

### **Frugivory-related traits promote speciation of tropical palms**

Renske E. Onstein<sup>1,\*</sup>, William J. Baker<sup>2</sup>, Thomas L. P. Couvreur<sup>3</sup>, Søren Faurby<sup>4,5</sup>, Jens-Christian Svenning<sup>6</sup> & W. Daniel Kissling<sup>1,\*</sup>

<sup>1</sup>Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, Amsterdam, The Netherlands

<sup>2</sup>Royal Botanic Gardens, Kew, Richmond, Surrey, UK

<sup>3</sup>Institut de Recherche pour le Développement, UMR-DIADE, Montpellier, France

<sup>4</sup>Department of Biological and Environmental Sciences, University of Gothenburg, Box 461, SE 405 30, Göteborg, Sweden

<sup>5</sup>Gothenburg Global Biodiversity Centre, Box 461, SE-405 30 Göteborg, Sweden

<sup>6</sup>Section for Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University, Aarhus, Denmark

\*corresponding authors e-mails: [onsteinre@gmail.com](mailto:onsteinre@gmail.com); [wdkissling@gmail.com](mailto:wdkissling@gmail.com)

# CONTENT

Supplementary Notes .....	3
Supplementary Note 1. Fruit size-dependent diversification rates corrected for plant size.....	3
Supplementary Note 2. Performance of trait-based models.....	4
Supplementary Note 3. Fruit size-dependent diversification rates.....	5
Supplementary Note 4. Fruit size as a continuous trait, and its effect on speciation rates.....	6
Supplementary Note 5. Understory and island-dependent diversification rates.....	7
Supplementary Tables .....	9
Supplementary Table 1 Summary statistics of fruit sizes for each palm genus.....	9
Supplementary Table 2 Model selection for large/small fruits globally (BiSSE).....	14
Supplementary Table 3 Model selection for large/small fruits in the New World (BiSSE).....	15
Supplementary Table 4 Model selection for large/small fruits in the Old World (BiSSE).....	16
Supplementary Table 5 Model selection after correcting for plant size (BiSSE).....	17
Supplementary Table 6 Model selection after correcting for plant size (QuaSSE).....	18
Supplementary Table 7 Models tested in the Multiple State Speciation and Extinction (MuSSE) framework.....	19
Supplementary Table 8 Model selection for fruit size and understory habitat (MuSSE).....	20
Supplementary Table 9 Model selection for fruit size and island colonization (MuSSE).....	21
Supplementary Figures.....	22
Supplementary Figure 1   Correlation between fruit size and plant size in palms.....	22
Supplementary Figure 2   Fruit size frequency distributions of palms.....	23
Supplementary Figure 3   Diversification rates under simulated binary traits (BiSSE).....	24
Supplementary Figure 4   Diversification rates of small- and large-fruited palms (BiSSE).....	25
Supplementary Figure 5   Speciation rate in response to fruit size (QuaSSE).....	26
Supplementary Figure 6   Phylorate plot of speciation rates in palms (BAMM).....	27
Supplementary Figure 7   Trait interaction effects on diversification rates (MuSSE).....	28
Supplementary data sources .....	29
Supplementary references .....	33

## Supplementary Notes

### **Supplementary Note 1. Fruit size-dependent diversification rates corrected for plant size.**

**Methods.** As a positive effect of small fruit size on speciation rates could be driven by a confounding effect of plant size on fruit size (i.e. smaller plants generally have smaller fruits and may have shorter generation times), we corrected for this by performing trait-dependent diversification analyses on the residuals of a linear regression model with log (fruit size) as the response variable and log (maximum stem height) as the explanatory variable (see Supplementary Fig. 1). The traits were log-transformed to approximate normality and to accommodate linear model assumptions such as normality in the residuals.

The linear regression model indicated a significant positive effect of log (maximum stem height) on log (fruit size) ( $df = 1683$ ,  $F = 115.98$ ,  $p < 0.001$ , model estimate = 0.142, standard error = 0.232), although the adjusted  $R^2$  was only 0.06, indicating that ca. 6% of the variation in fruit size can be explained by palm maximum stem height. Model residuals (i.e. the non-explained variation in fruit size) were reassessed for their effect on speciation, extinction and transition rates with the BiSSE model (large fruit residuals  $\geq 4$  cm), repeating the analyses as described in the main text (related to H1) (see Supplementary Table 5 for model selection). However, as these residuals do not represent ‘true’ small and large fruits, we also investigated the effect of residual log-transformed fruit size on speciation rates using the Quantitative Speciation and Extinction model (QuaSSE)<sup>1</sup>. QuaSSE can be used to test the effect of a continuous trait on speciation rates by testing the fit of models describing the distribution of the response (i.e. speciation rate) to the trait (e.g. constant, linear or sigmoidal) (see Supplementary Table 6 for model selection). The best model was selected using likelihood-ratio tests (nested models) and the Akaike Information Criterion (AIC) (non-nested models). The relationship between residual fruit size and speciation rate resulting from the best model can be plotted to visualize the results (Supplementary Fig. 5).

**Results.** Our results indicate that after correcting for a plant size effect, speciation rates between small and large fruited palm lineages remain significantly different. Specifically, small-fruited palms show higher speciation rates than large-fruited palms (see Supplementary Fig. 4: compare residuals to global, New World and Old World distributions of Bayesian posterior rates for small- and large-fruited palms). Furthermore, the effect of palm residual fruit size (cm) on speciation rates under the best-fitting QuaSSE model indicates that larger fruit sizes have lower rates of speciation than smaller fruit sizes, following a sigmoidal relationship (Supplementary Fig. 5).

## Supplementary Note 2. Performance of trait-based models.

The class of ‘SSE’ models has recently been criticized for high type I error rates<sup>2</sup>, suggesting that a significant effect of a trait on speciation or extinction rates can be detected even if it is not truly present. This error may be driven by the phylogenetic tree shape (i.e. distribution of branch lengths in the tree)<sup>2</sup>. We evaluated this bias by performing simulations as recommended by ref.<sup>2</sup>. First, we randomly evolved a neutral binary trait on 100 empirical palm phylogenies under three transition rate scenarios ( $q = 0.01$ ,  $q = 0.1$  and  $q = 1$ )<sup>2</sup>, providing a gradient from rare to frequent character state changes. This simulated trait is expected to be neutral with respect to speciation and extinction rates, as it evolved under a simple ‘Markov discrete’ (Mk)<sup>3</sup> model of evolution. We then evaluated the Bayesian credible intervals in speciation and extinction rates between these simulated trait states after running the Bayesian Markov Chain Monte Carlo chain for 10,000 generations on 100 palm phylogenies (Supplementary Fig. 3). We repeated this procedure using our observed transition rates (i.e.  $q_{\text{large to small}} = 0.017$ ;  $q_{\text{small to large}} = 0.006$ ). Importantly, strongly overlapping Bayesian credible intervals of speciation and extinction rates of the simulated binary traits would suggest that our empirical palm tree shapes unlikely biased our results, and type I error rates should be neglectable. Our results show that this is the case for all transition rate scenarios (Supplementary Fig. 3), suggesting that we can rely on the empirical results obtained from the ‘SSE’ models.

A second criticism on the ‘SSE’ models is that datasets should consist of sufficient species at the tips of the tree ( $> 300$  species), a balanced distribution of trait states at the tips of the tree so that at least 10% of the species has a certain trait state<sup>4,5</sup>, and the trait should evolve multiple times to have sufficient independent events to evaluate correlations between trait states and speciation rates<sup>6</sup>. These criteria should be met to have the statistical power to reliably estimate the parameters in the SSE models. Our dataset comprises 1774 species, large fruits are found in 12%, understory habitats in 36%, and island distributions in 20% of all sampled palm species. Large fruits are found within 63, understory habitats within 73, and island distributions within 83 (out of 183) palm genera. Although this does not directly proof the repeated independent evolution of these ‘traits’ (i.e. these genera may form a clade), it does strongly suggest that at least several independent events have occurred. We thus have the statistical power to detect correlates between speciation rates and traits without erroneously relying on unbalanced character state distributions across the tree<sup>5</sup>, or pseudo-replication<sup>6</sup>.

### **Supplementary Note 3. Fruit size-dependent diversification rates.**

**Methods.** The Binary State Speciation and Extinction (BiSSE) model used to test the effect of fruit size on speciation rates (H1, see main text), simultaneously estimates speciation, extinction and transition rates, as these rates do not evolve independently from each other. The net diversification rate can be calculated by subtracting the extinction from the speciation rate. Although not directly related to the hypotheses presented in the main text, we here report the extinction, transition and net diversification rate estimates based on the best-fitting models for global, New World and Old World palms (see Supplementary Tables 2-4 for model selections).

**Results.** The Bayesian rates indicate that large-fruited Old World palm lineages have lower extinction rates than small-fruited Old World palm lineages (Supplementary Table 4, Supplementary Fig. 4). For the global and New World datasets we did not detect support for a different extinction rate for small- and large-fruited palms (Supplementary Tables 2 and 3). Globally, this has resulted in a higher net diversification rate (speciation rate minus extinction rate) for small-fruited compared to large-fruited palm lineages, but this difference is not recovered in the New World or Old World (overlapping 95% Bayesian posterior densities, see Supplementary Fig. 4). Last, transition rates from small to large fruits have been lower than from large to small fruits in all analyses (globally, New World and Old World). These results suggest that although small fruits positively affect speciation rates globally and in the Old World (Fig. 2), the net diversification of small-fruited palms is only evident in the global analysis, as Old World small-fruited palms show high speciation rates, but also high extinction rates (resulting in high turnover rates). The transition rate results suggest that the evolution of smaller fruits from larger fruits has been more frequently observed during the diversification of palms than vice versa. All results are visualized in Supplementary Fig. 4.

#### **Supplementary Note 4. Fruit size as a continuous trait, and its effect on speciation rates.**

**Methods.** In this study we compared the effect of small and large (megafaunal) fruits on speciation rates (main text). To do so, we defined megafaunal fruits as those with length  $\geq 4$  cm. However, fruit size in palms is a continuous trait, and we therefore also tested the effect of fruit size as a continuous trait on speciation rates, using the Bayesian Analysis of Macroevolutionary Mixtures approach implemented in BAMM version 2.5.0<sup>7</sup>. BAMM explicitly accounts for diversification rate variation through time and uses a reversible jump MCMC algorithm to explore numerous candidate models of lineage diversification. These models thereby identify probabilities of diversification rate shifts (i.e. increases or slowdowns in diversification) on phylogenetic trees.

We ran BAMM on the palm Maximum Clade Credibility (MCC) tree<sup>8</sup>. Priors were estimated with the BAMMTools R package<sup>9</sup> using the function “setBAMMpriors”. A compound Poisson process for the prior probability of a rate shift along any branch was implemented, and we ran a MCMC for 10 million generations sampling event data every 5000 steps. We checked for convergence by plotting the log-likelihood trace of the MCMC output file and checked that the effective sample sizes of the runs exceeded 200. BAMMTools was used to generate the mean phylorate plot, which represents the mean speciation rate sampled from the posterior at any point in time along any branch of the phylogenetic tree<sup>9</sup>.

To evaluate whether speciation rates are correlated to fruit size, we ran SStructured Rate Permutations on Phylogenies (STRAPP)<sup>10</sup>, implemented in BAMMTools. The traitDependentBAMM function was used to compare the observed difference in speciation rate between palms that exhibit different fruit sizes to a background speciation rate through randomizing the estimated tip speciation rates from the BAMM outputs<sup>10</sup>.

**Results.** Palms show substantial speciation rate heterogeneity, congruent with the BAMM analysis performed previously on the genus-level phylogeny<sup>11</sup> (Supplementary Fig. 6). We found that the speciation rate in palms correlates negatively with fruit size (one-tailed Pearson correlation,  $r = -0.3$ ,  $p < 0.05$ ) assessed by 1000 permutations. This indicates that the observed correlation from the posterior samples is more negative (i.e. negative correlation between fruit size and speciation rate) than the correlations calculated with permuted rates, in more than 95% of the simulations. These results indicate that fruit size in palms is generally associated with low speciation rates, congruent with results from the binary classification of fruit size we made in the main text (Fig. 2).

## Supplementary Note 5. Understory and island-dependent diversification rates.

**Methods.** To disentangle the effect of two binary traits on speciation, extinction and transition rates, we implemented the Multiple State Speciation and Extinction model (MuSSE multistate)<sup>12</sup>. The model intercept of the MuSSE model (the ‘base model’) estimates speciation, extinction and transition rates when both traits are absent (illustrated with code 000, in which the numbers refer to speciation, extinction and transition respectively, see Supplementary Table 7). 0 refers to absence of the traits, 1 to an additive effect of the traits, and 2 to an interaction effect of the traits. 000 therefore refers to, for example, palm lineages with large fruits that do not grow in the understory (absence of small fruits and understory habitat). Significant support for the additive effects of the traits on for example speciation rates (illustrated with code 100) or extinction rates (illustrated with code 010) or both (illustrated with code 110) would indicate that the individual traits affect speciation or extinction rates or both (speciation and extinction rates), respectively. Support for an interaction term (when both traits are present) will indicate whether these traits may interact in either a positive way (i.e. both traits increase speciation rates) or a negative way (i.e. both traits decrease speciation rates). This would be illustrated with code 210 for a significant interaction effect on speciation rates, and 120 for a significant interaction effect on extinction rates. Similarly, when additive effects on transition rates are detected (but not on speciation or extinction rates), this would be illustrated with code 001.

To quantify trait-dependent diversification for both binary trait combinations (H2: small/large fruit size and understory yes/no; H3: small/large fruit size and island colonization yes/no), we compared the likelihood of a total of sixteen models with increasing complexity, using a stepwise approach. We started with the base model (code 000) and added parameters for additive effects on speciation (code 100), extinction (code 010) or transition rates (code 001), or combinations of these (codes 110, 011, 101). Then we evaluated the support for including an interaction term for speciation (code 211) and extinction (code 121) rates to the model. The support for more complex models was evaluated using a likelihood ratio test (nested models) and the Akaike Information Criterion (AIC) (non-nested models). In case a more complex model was rejected, we accepted the best-fitting model given the fewest number of parameters. All models are described in Supplementary Table 7. A Bayesian Markov Chain Monte Carlo for the best-fitting model was run for 10,000 generations on 100 palm phylogenies for the global, New World and Old World palms separately (Supplementary Fig. 7).

**Results.** Here, we report the extinction, transition and net diversification rates as estimated with the Multiple State Speciation and Extinction (MuSSE) model for H2: small/large fruit size and understory yes/no (Supplementary Fig. 7a); and H3: small/large fruit size and island colonization yes/no (Supplementary Fig. 7b), globally, in New World and in Old World palm lineages. For speciation rate results see the main text.

The Bayesian rates indicate that small fruit size and understory habitat do not have an interactive effect on extinction rates (Supplementary Table 8). However, small fruit sizes do have a positive effect on extinction rates compared to large-fruited canopy palms in the global, New World and Old World analyses, congruent with the results obtained from the global and Old World BiSSE analyses (see Supplementary Note 3 and Supplementary Fig. 4). Transition rates from understory habitat to canopy habitat were higher than vice versa in all analyses (global, New World and Old World), and transition rates from large to small fruits

were higher than from small to large fruits, congruent with the results obtained from the BiSSE analyses (see Supplementary Note 3 and Supplementary Fig. 4). The highest diversification rates were detected for small-fruited understory palms in the global and New World analyses, but not in the Old World (Supplementary Fig. 7a). This suggests that fruit size and understory habit act in synergy to increase diversification rates in global and New World palms.

Furthermore, the Bayesian rates indicate that small fruit size and island colonization do not have an interactive effect on extinction rates (Supplementary Table 9). However, small fruit sizes do have a positive effect on extinction rates globally and in the Old World compared to large-fruited mainland-distributed palms, congruent with the results obtained from the global and Old World BiSSE analyses (see Supplementary Note 3 and Supplementary Fig. 4). Transition rates from island to mainland were not significantly different from mainland to island (overlapping posterior densities in Supplementary Fig. 7b). The highest diversification rates were detected for small-fruited island-distributed palms in the global and Old World analyses, but not in the New World (Supplementary Fig. 7b). This suggests that fruit size and island colonization act in synergy to increase diversification rates in global and Old World palms.



## Supplementary Tables

### **Supplementary Table 1 Summary statistics of fruit sizes for each palm genus.**

Fruit size characteristics and species richness for each palm genus are summarized as follows: mean = average fruit size based on all species within the genus, sd = standard deviation, var = variance, #obs = number of species observations used for fruit measures, se = standard error, min = minimum, max = maximum, range = maximum minus minimum, rich = species richness, NA = not applicable.

\*these genera are not dispersed by animals.

<i>Genus</i>	mean	sd	var	#obs	se	min	max	range	rich
<i>Acanthophoenix</i>	1.22	0.70	0.49	3	0.40	0.65	2	1.35	3
<i>Acoelorrhaphe</i>	0.70	NA	NA	1	NA	0.70	0.70	0	1
<i>Acrocomia</i>	2.68	1.48	2.20	3	0.86	1.30	4.25	2.95	8
<i>Actinokentia</i>	2.60	NA	NA	1	NA	2.60	2.60	0	2
<i>Actinorhysis</i>	10	NA	NA	1	NA	10	10	0	1
<i>Adonidia</i>	2.50	NA	NA	1	NA	2.50	2.50	0	2
<i>Aiphanes</i>	1.50	1.09	1.19	23	0.23	0.50	4.70	4.20	29
<i>Allagoptera</i>	2.32	1.19	1.40	5	0.53	1.35	4.25	2.90	5
<i>Ammandra</i>	10	NA	NA	1	NA	10	10	0	1
<i>Aphandra</i>	35	NA	NA	1	NA	35	35	0	1
<i>Archontophoenix</i>	1.60	0.48	0.23	6	0.20	1.10	2.30	1.20	6
<i>Areca</i>	3.21	1.66	2.76	32	0.29	0.50	7.50	7	45
<i>Arenga</i>	2.71	2.19	4.78	16	0.55	0.70	7	6.30	24
<i>Asterogyne</i>	1.75	1.06	1.13	2	0.75	1	2.50	1.50	5
<i>Astrocaryum</i>	4.92	1.70	2.90	31	0.31	1.20	8	6.80	37
<i>Attalea</i>	7.58	2.37	5.60	39	0.38	3.75	12.50	8.75	67
<i>Bactris</i>	1.73	0.88	0.77	75	0.10	0.50	5	4.50	77
<i>Balaka</i>	2.41	0.95	0.91	5	0.43	1.60	4	2.40	9
<i>Barcella</i>	3.50	NA	NA	1	NA	3.50	3.50	0	1
<i>Basselinia</i>	0.80	0.24	0.06	11	0.07	0.40	1.20	0.80	14
<i>Beccariophoenix</i>	2.95	0.78	0.61	2	0.55	2.40	3.50	1.10	2
<i>Bentinckia</i>	1.50	0	0	2	0	1.50	1.50	0	2
<i>Bismarckia</i>	4.40	NA	NA	1	NA	4.40	4.40	0	1
<i>Borassodendron</i>	10.50	2.12	4.50	2	1.50	9	12	3	2
<i>Borassus</i>	18.10	7.30	53.30	5	3.26	12	30	18	5
<i>Brahea</i>	1.68	0.65	0.43	11	0.20	0.65	3	2.35	11
<i>Brassiophoenix</i>	3.25	0.07	0.01	2	0.05	3.20	3.30	0.10	2
<i>BurretioKentia</i>	3.88	4.55	20.67	5	2.03	1.60	12	10.40	5
<i>Butia</i>	2.23	0.72	0.51	12	0.21	1.20	3.60	2.40	20
<i>Calamus</i>	1.48	0.71	0.50	193	0.05	0.50	5	4.50	379
<i>Calyptrocalyx</i>	1.51	0.89	0.79	26	0.17	0.40	5	4.60	26
<i>Calyptrogyne</i>	1.23	0.51	0.26	8	0.18	0.21	1.75	1.54	17

<i>Calyptronoma</i>	1.45	0.97	0.94	3	0.56	0.58	2.50	1.92	3
<i>Carpentaria</i>	2	NA	NA	1	NA	2	2	0	1
<i>Carpoxydon</i>	6	NA	NA	1	NA	6	6	0	1
<i>Caryota</i>	2.45	0.72	0.51	11	0.22	1.40	3.50	2.10	14
<i>Ceratolobus</i>	1.59	0.42	0.18	6	0.17	1.10	2.10	1	6
<i>Ceroxydon</i>	1.76	0.16	0.03	10	0.05	1.50	2	0.50	12
<i>Chamaedorea</i>	1.02	0.32	0.10	97	0.03	0.40	2.15	1.75	106
<i>Chamaerops</i>	1	NA	NA	1	NA	1	1	0	1
<i>Chambeyronia</i>	3.75	1.06	1.13	2	0.75	3	4.50	1.50	2
<i>Chelyocarpus</i>	1.99	0.34	0.12	4	0.17	1.75	2.50	0.75	4
<i>Chuniophoenix</i>	2.05	0.64	0.41	2	0.45	1.60	2.50	0.90	3
<i>Clinosperma</i>	2.15	1.18	1.40	4	0.59	1	3.75	2.75	4
<i>Clinostigma</i>	1.23	0.54	0.29	8	0.19	0.50	2.20	1.70	11
<i>Coccothrinax</i>	1.21	1.23	1.51	20	0.27	0.50	6.30	5.80	51
<i>Cocos*</i>	22.50	NA	NA	1	NA	22.50	22.50	0	1
<i>Colpotherinax</i>	1.62	0.32	0.10	3	0.19	1.25	1.85	0.60	3
<i>Copernicia</i>	1.94	0.42	0.17	13	0.12	1.35	2.75	1.40	22
<i>Corypha</i>	4.50	2.12	4.50	4	1.06	2.50	7.50	5	5
<i>Cryosophila</i>	1.87	0.63	0.40	9	0.21	1.25	3.35	2.10	10
<i>Cyphokentia</i>	1.23	0.04	0	2	0.03	1.20	1.25	0.05	2
<i>Cyphophoenix</i>	1.89	0.48	0.23	4	0.24	1.20	2.30	1.10	4
<i>Cyphosperma</i>	1.33	0.48	0.23	3	0.28	0.90	1.85	0.95	5
<i>Cyrtostachys</i>	1.16	0.21	0.05	4	0.11	1	1.45	0.45	7
<i>Daemonorops</i>	1.78	0.67	0.45	72	0.08	0.65	4.50	3.85	107
<i>Deckenia</i>	1.20	NA	NA	1	NA	1.20	1.20	0	1
<i>Desmoncus</i>	1.54	0.57	0.33	23	0.12	1	3.62	2.62	24
<i>Dictyocaryum</i>	3.05	0.35	0.12	3	0.20	2.65	3.25	0.60	3
<i>Dictyosperma</i>	1.25	NA	NA	1	NA	1.25	1.25	0	1
<i>Dransfieldia</i>	1.55	NA	NA	1	NA	1.55	1.55	0	1
<i>Drymophloeus</i>	1.13	0.81	0.66	2	0.58	0.55	1.70	1.15	3
<i>Dypsis</i>	1.36	0.51	0.27	99	0.05	0.50	2.75	2.25	162
<i>Elaeis</i>	3.50	1.06	1.13	2	0.75	2.75	4.25	1.50	2
<i>Eleiodoxa</i>	2.50	NA	NA	1	NA	2.50	2.50	0	1
<i>Eremospatha</i>	2.51	0.44	0.20	8	0.16	1.75	3.25	1.50	11
<i>Eugeissona</i>	6.90	1.67	2.80	5	0.75	5	9	4	6
<i>Euterpe</i>	1.31	0.34	0.12	7	0.13	1.05	2	0.95	7
<i>Gaussia</i>	1.26	0.26	0.07	5	0.12	1	1.55	0.55	5
<i>Geonoma</i>	0.85	0.27	0.07	63	0.03	0.53	1.89	1.36	68
<i>Guihaia</i>	0.80	0.28	0.08	2	0.20	0.60	1	0.40	2
<i>Hedysepe</i>	4	NA	NA	1	NA	4	4	0	1
<i>Hemithrinax</i>	0.90	NA	NA	1	NA	0.90	0.90	0	3
<i>Heterospatha</i>	1.22	0.58	0.33	27	0.11	0.50	3.25	2.75	41
<i>Howea</i>	3.75	0.35	0.13	2	0.25	3.50	4	0.50	2
<i>Hydriastele</i>	1.13	0.32	0.10	25	0.06	0.60	1.80	1.20	49

<i>Hyophorbe</i>	2.53	0.90	0.81	4	0.45	1.80	3.80	2	5
<i>Hyospathe</i>	1.08	0.16	0.03	3	0.09	0.90	1.20	0.30	4
<i>Hyphaene</i>	6.29	1.32	1.74	7	0.50	4.50	8.50	4	8
<i>Iguanura</i>	1.30	0.54	0.29	17	0.13	0.60	2.40	1.80	32
<i>Iriartea</i>	2.35	NA	NA	1	NA	2.35	2.35	0	1
<i>Iriartella</i>	1.38	0.25	0.06	2	0.18	1.20	1.55	0.35	2
<i>Itaya</i>	2.25	NA	NA	1	NA	2.25	2.25	0	1
<i>Johannesteijsmannia</i>	4.20	1.13	1.28	2	0.80	3.40	5	1.60	4
<i>Juania</i>	1.65	NA	NA	1	NA	1.65	1.65	0	1
<i>Jubaea</i>	3.75	NA	NA	1	NA	3.75	3.75	0	1
<i>Jubaeopsis</i>	3	NA	NA	1	NA	3	3	0	1
<i>Kentiopsis</i>	1.95	0.21	0.04	4	0.10	1.70	2.20	0.50	4
<i>Kerriodoxa</i>	4	NA	NA	1	NA	4	4	0	1
<i>Korthalsia</i>	1.64	0.68	0.47	23	0.14	0.80	3.50	2.70	27
<i>Laccospadix</i>	1.35	NA	NA	1	NA	1.35	1.35	0	1
<i>Laccosperma</i>	1.63	0.30	0.09	5	0.13	1.35	2	0.65	6
<i>Lanonia</i>	0.79	0.11	0.01	7	0.04	0.65	1	0.35	8
<i>Latania</i>	5	1.32	1.75	3	0.76	4	6.50	2.50	3
<i>Lemurophoenix</i>	0.50	NA	NA	1	NA	0.50	0.50	0	1
<i>Leopoldinia</i>	3.32	0.68	0.46	2	0.48	2.84	3.80	0.96	2
<i>Lepidocaryum</i>	2.25	NA	NA	1	NA	2.25	2.25	0	1
<i>Lepidorrhachis</i>	1.20	NA	NA	1	NA	1.20	1.20	0	1
<i>Leucothrinax</i>	0.70	NA	NA	1	NA	0.70	0.70	0	1
<i>Licuala</i>	1.19	0.56	0.31	99	0.06	0.45	4	3.55	162
<i>Linospadix</i>	1.31	0.47	0.22	6	0.19	0.70	2.15	1.45	7
<i>Livistona</i>	1.61	0.67	0.45	26	0.13	0.80	3.25	2.45	28
<i>Lodoicea*</i>	45	NA	NA	1	NA	45	45	0	1
<i>Loxococcus</i>	2.50	NA	NA	1	NA	2.50	2.50	0	1
<i>Lytocaryum</i>	2.58	0.81	0.66	2	0.58	2	3.15	1.15	4
<i>Manicaria</i>	35	NA	NA	1	NA	35	35	0	2
<i>Marojejya</i>	2.10	0.21	0.05	2	0.15	1.95	2.25	0.30	2
<i>Masoala</i>	2.85	0.57	0.32	2	0.40	2.45	3.25	0.80	2
<i>Mauritia</i>	6.38	0.88	0.78	2	0.63	5.75	7	1.25	2
<i>Mauritiella</i>	3.25	1.15	1.31	3	0.66	2.25	4.50	2.25	4
<i>Maxburretia</i>	0.98	0.04	0	2	0.03	0.95	1	0.05	3
<i>Medemia</i>	4.50	NA	NA	1	NA	4.50	4.50	0	1
<i>Metroxylon</i>	10.10	2.25	5.05	5	1	7	12.50	5.50	7
<i>Myrialepis</i>	2.50	NA	NA	1	NA	2.50	2.50	0	1
<i>Nenga</i>	3.70	1.96	3.86	5	0.88	1.40	6.50	5.10	5
<i>Neonicholsonia</i>	0.95	NA	NA	1	NA	0.95	0.95	0	1
<i>Neoveitchia</i>	5.50	0.71	0.50	2	0.50	5	6	1	2
<i>Nephrosperma</i>	1.20	NA	NA	1	NA	1.20	1.20	0	1
<i>Normanbya</i>	4.25	NA	NA	1	NA	4.25	4.25	0	1
<i>Nypa*</i>	11.50	NA	NA	1	NA	11.50	11.50	0	1

<i>Oenocarpus</i>	2.14	0.65	0.42	9	0.22	1.40	3.50	2.10	9
<i>Oncocalamus</i>	1.97	0.06	0	3	0.03	1.90	2	0.10	5
<i>Oncosperma</i>	1.28	0.30	0.09	4	0.15	1	1.70	0.70	5
<i>Orania</i>	3.76	1.77	3.13	16	0.44	0.60	6.50	5.90	28
<i>Oraniopsis</i>	3.40	NA	NA	1	NA	3.40	3.40	0	1
<i>Parajubaea</i>	4.50	0.90	0.81	3	0.52	3.50	5.25	1.75	3
<i>Pelagodoxa</i>	12.50	NA	NA	1	NA	12.50	12.50	0	1
<i>Phoenicophorium</i>	0.80	NA	NA	1	NA	0.80	0.80	0	1
<i>Phoenix</i>	2.20	1.41	2	14	0.38	1.30	7	5.70	14
<i>Pholidocarpus</i>	7.10	3.03	9.18	5	1.35	4	11.50	7.50	6
<i>Pholidostachys</i>	1.95	0.55	0.31	7	0.21	1.22	2.86	1.64	8
<i>Physokentia</i>	2.01	0.55	0.31	6	0.23	1.20	2.55	1.35	7
<i>Phytelephas</i>	9	2.68	7.17	4	1.34	7.50	13	5.50	6
<i>Pigafetta</i>	0.90	0	0	2	0	0.90	0.90	0	2
<i>Pinanga</i>	1.59	0.57	0.33	101	0.06	0.40	4.50	4.10	138
<i>Plectocomia</i>	2.28	0.46	0.21	10	0.15	1.50	3	1.50	15
<i>Plectocomiopsis</i>	2.59	0.12	0.02	5	0.06	2.50	2.75	0.25	6
<i>Podococcus</i>	2.50	NA	NA	1	NA	2.50	2.50	0	2
<i>Pogonotium</i>	1.47	0.21	0.04	3	0.12	1.30	1.70	0.40	3
<i>Ponapea</i>	3.08	0.88	0.77	3	0.51	2.25	4	1.75	4
<i>Prestoea</i>	1	0.19	0.04	9	0.06	0.85	1.40	0.55	10
<i>Pritchardia</i>	3.09	1.47	2.15	27	0.28	0.60	7	6.40	28
<i>Pseudophoenix</i>	1.97	0.82	0.67	4	0.41	1.30	3.07	1.77	4
<i>Ptychococcus</i>	3.10	2.69	7.22	2	1.90	1.20	5	3.80	2
<i>Ptychosperma</i>	1.50	0.25	0.06	24	0.05	1	2	1	30
<i>Raphia</i>	7.05	1.61	2.60	19	0.37	5	11.50	6.50	20
<i>Ravenea</i>	1.57	0.60	0.35	16	0.15	0.75	2.85	2.10	20
<i>Reinhardtia</i>	1.58	0.28	0.08	6	0.11	1.20	2	0.80	6
<i>Retispatha</i>	2	NA	NA	1	NA	2	2	0	1
<i>Rhapidophyllum</i>	2	NA	NA	1	NA	2	2	0	1
<i>Rhapis</i>	1.17	0.78	0.60	9	0.26	0.70	2.95	2.25	11
<i>Rhopaloblaste</i>	2.17	0.77	0.60	6	0.32	1.10	3.25	2.15	6
<i>Rhopalostylis</i>	1.35	0.07	0.01	2	0.05	1.30	1.40	0.10	2
<i>Roscheria</i>	1	NA	NA	1	NA	1	1	0	1
<i>Roystonea</i>	1.29	0.15	0.02	10	0.05	1	1.50	0.50	10
<i>Sabal</i>	1.34	0.37	0.14	14	0.10	0.80	1.85	1.05	14
<i>Salacca</i>	5.24	2.15	4.63	12	0.62	1.80	8	6.20	22
<i>Saribus</i>	1.72	1.11	1.23	9	0.37	0.39	4	3.61	9
<i>Satakentia</i>	1.30	NA	NA	1	NA	1.30	1.30	0	1
<i>Satranala</i>	5.60	NA	NA	1	NA	5.60	5.60	0	1
<i>Schippia</i>	2.50	NA	NA	1	NA	2.50	2.50	0	1
<i>Sclerosperma</i>	3.67	0.76	0.58	3	0.44	3	4.50	1.50	3
<i>Serenoa</i>	2.05	NA	NA	1	NA	2.05	2.05	0	1
<i>Socratea</i>	3.50	0.71	0.50	4	0.35	3	4.50	1.50	5

<i>Sommieria</i>	1.20	NA	NA	1	NA	1.20	1.20	0	1
<i>Syagrus</i>	3.39	1.24	1.53	49	0.18	2	7	5	53
<i>Synechanthus</i>	2.05	0.42	0.18	2	0.30	1.75	2.35	0.60	2
<i>Tahina</i>	3.20	NA	NA	1	NA	3.20	3.20	0	1
<i>Tectiphiala</i>	1.10	NA	NA	1	NA	1.10	1.10	0	1
<i>Thrinax</i>	0.80	0.13	0.02	3	0.08	0.70	0.95	0.25	3
<i>Trachycarpus</i>	1.05	0.32	0.10	8	0.11	0.70	1.70	1	9
<i>Trithrinax</i>	0.98	0.16	0.03	3	0.09	0.80	1.10	0.30	4
<i>Veitchia</i>	2.71	1.32	1.74	7	0.50	1.50	5	3.50	11
<i>Verschaffeltia</i>	2.50	NA	NA	1	NA	2.50	2.50	0	1
<i>Voanioala</i>	7.50	NA	NA	1	NA	7.50	7.50	0	1
<i>Wallichia</i>	1.56	0.42	0.18	5	0.19	1.10	2.20	1.10	8
<i>Washingtonia</i>	0.80	0.28	0.08	2	0.20	0.60	1	0.40	2
<i>Welfia</i>	3.21	NA	NA	1	NA	3.21	3.21	0	1
<i>Wendlandiella</i>	0.90	NA	NA	1	NA	0.90	0.90	0	1
<i>Wettinia</i>	2.79	0.80	0.64	20	0.18	1.70	4.50	2.80	21
<i>Wodyetia</i>	5.75	NA	NA	1	NA	5.75	5.75	0	1
<i>Zombia</i>	1.75	NA	NA	1	NA	1.75	1.75	0	1

**Supplementary Table 2 Model selection for large/small fruits globally (BiSSE).**

Eight Binary State Speciation and Extinction (BiSSE) models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree to compare speciation, extinction and transition rates of small- vs. large-fruited palm lineages. The best-fitting model given the fewest number of parameters (i.e. 5 Df) is indicated in bold (\*). This model indicates that, globally, small- and large-fruited palm lineages have different speciation rates, but similar extinction rates.

<b>Model constraints</b>	<b>Df</b>	<b>LnLik</b>	<b>AIC</b>	<b>ChiSq</b>	<b>P</b>
Full (no constrain)	6	-5978.8	11970		
$\lambda_{\text{small}} \sim \lambda_{\text{large}}$	5	-5990.1	11990	22.610	2e-06 ***
<b>*<math>\mu_{\text{small}} \sim \mu_{\text{large}}</math></b>	<b>5</b>	<b>-5979.4</b>	<b>11969</b>	<b>1.144</b>	<b>0.285</b>
$q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	5	-5986.1	11982	14.590	0.0001 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}$	4	-6026.9	12062	96.045	< 2.2e-16 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-5990.7	11989	23.621	7.4e-06 ***
$\mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-5988.7	11985	19.732	5.2e-05 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	3	-6033.1	12072	108.491	< 2.2e-16 ***

Df = degrees of freedom, LnLik = log likelihood, AIC = Akaike Information Criterion, ChiSq = Chi-square, P = significance of the model compared to the full model,  $\lambda$  = speciation rate,  $\mu$  = extinction rate, q = transition rate,  $\sim$  = equal to (constrain). Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

**Supplementary Table 3 Model selection for large/small fruits in the New World (BiSSE).**

Eight Binary State Speciation and Extinction (BiSSE) models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree to compare speciation, extinction and transition rates of small- vs. large-fruited New World palm lineages. The best-fitting model given the fewest number of parameters (i.e. 4 Df) is indicated in bold (\*). This model indicates that New World small- and large-fruited palm lineages have similar speciation and extinction rates.

<b>Model constraints</b>	<b>Df</b>	<b>LnLik</b>	<b>AIC</b>	<b>ChiSq</b>	<b>P</b>
Full (no constrain)	6	-2218.3	4448.6		
$\lambda_{\text{small}} \sim \lambda_{\text{large}}$	5	-2219.1	4448.2	1.5633	0.211
$\mu_{\text{small}} \sim \mu_{\text{large}}$	5	-2218.6	4447.2	0.5500	0.458
$q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	5	-2224.0	4458.0	11.4470	0.0007 ***
<b>* <math>\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}</math></b>	<b>4</b>	<b>-2219.9</b>	<b>4447.8</b>	<b>3.2443</b>	<b>0.197</b>
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-2224.6	4457.2	12.5629	0.002 **
$\mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-2226.7	4461.4	16.7662	0.0002 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	3	-2233.3	4472.7	30.0633	1.3e-06 ***

Df = degrees of freedom, LnLik = log likelihood, AIC = Akaike Information Criterion, ChiSq = Chi-square, P = significance of the model compared to the full model,  $\lambda$  = speciation rate,  $\mu$  = extinction rate,  $q$  = transition rate,  $\sim$  = equal to (constrain). Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

**Supplementary Table 4 Model selection for large/small fruits in the Old World (BiSSE).**

Eight Binary State Speciation and Extinction (BiSSE) models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree to compare speciation, extinction and transition rates of small- vs. large-fruited Old World palm lineages. The best-fitting model given the fewest number of parameters (i.e. 6 Df) is indicated in bold (\*). This model indicates that Old World small- and large-fruited palm lineages have different speciation and extinction rates.

Model constraints	Df	LnLik	AIC	ChiSq	P
<b>* Full (no constrain)</b>	<b>6</b>	<b>-3754.9</b>	<b>7521.8</b>		
$\lambda_{\text{small}} \sim \lambda_{\text{large}}$	5	-3780.9	7571.7	51.903	5.8e-13 ***
$\mu_{\text{small}} \sim \mu_{\text{large}}$	5	-3767.8	7545.6	25.739	3.9e-07 ***
$q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	5	-3761.9	7533.8	13.940	0.0002 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}$	4	-3822.6	7653.2	135.351	< 2.2e-16 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-3782.1	7572.3	54.474	1.5e-12 ***
$\mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-3767.8	7543.6	25.737	2.6e-06 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	3	-3823.0	7652.1	136.278	< 2.2e-16 ***

Df = degrees of freedom, LnLik = log likelihood, AIC = Akaike Information Criterion, ChiSq = Chi-square, P = significance of the model compared to the full model,  $\lambda$  = speciation rate,  $\mu$  = extinction rate, q = transition rate,  $\sim$  = equal to (constrain). Signif. codes: 0 '\*\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1



**Supplementary Table 5 Model selection after correcting for plant size (BiSSE).**

Trait-dependent speciation, extinction and transition rate model selection under the Binary State Speciation and Extinction (BiSSE) model for all palms, after correcting for the effect of maximum palm stem height on fruit size. Eight BiSSE models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree. The best-fitting model given the fewest number of parameters is indicated in bold (\*). This model indicates that speciation, extinction and transition rates are different between small- and large-fruited palm lineages.

Model constraints	Df	LnLik	AIC	ChiSq	P
<b>*Full (no constrain)</b>	<b>6</b>	<b>-5501.5</b>	<b>11015</b>		
$\lambda_{\text{small}} \sim \lambda_{\text{large}}$	5	-5511.1	11032	19.355	1.1e-05 ***
$\mu_{\text{small}} \sim \mu_{\text{large}}$	5	-5506.1	11022	9.200	0.002 **
$q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	5	-5532.5	11075	62.126	3.2e-15 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}$	4	-5525.3	11059	47.701	4.4e-11 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-5536.1	11080	69.274	8.9e-16 ***
$\mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	4	-5544.7	11098	86.566	< 2.2e-16 ***
$\lambda_{\text{small}} \sim \lambda_{\text{large}}, \mu_{\text{small}} \sim \mu_{\text{large}}, q_{\text{small} \rightarrow \text{large}} \sim q_{\text{large} \rightarrow \text{small}}$	3	-5557.4	11121	111.906	< 2.2e-16 ***

Df = degrees of freedom, LnLik = log likelihood, AIC = Akaike Information Criterion, ChiSq = Chi-square, P = significance of the model compared to the full model,  $\lambda$  = speciation rate,  $\mu$  = extinction rate, q = transition rate,  $\sim$  = equal to (constrain). Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

**Supplementary Table 6 Model selection after correcting for plant size (QuaSSE).**

Trait-dependent speciation rate model selection under the Quantitative Speciation and Extinction (QuaSSE) model for all palms, after correcting for the effect of maximum stem height on fruit size. Residuals of log (fruit length) against log (maximum stem height) as a continuous trait were tested for their effect on speciation rates on the palm Maximum Clade Credibility (MCC) phylogenetic tree. QuaSSE can be used to test whether the speciation rate follows a constant, a linear or a sigmoidal response curve to (residual) log-transformed fruit length. Values range from -1 to 1, indicating residual sizes from ca. 0.1 cm to 10 cm. The best model is indicated in bold (\*). This model indicates that speciation rates follow a sigmoidal relationship in response to residual fruit sizes.

<b>Model</b>	<b>Df</b>	<b>lnLik</b>	<b>AIC</b>	<b>ΔAIC</b>
Constant $\lambda$	3	-4582.0	9169.9	159.9
Linear $\lambda$	4	-4507.6	9023.2	13.2
Sigmoidal $\lambda$	6	-4500.4	9012.8	2.8
Linear $\lambda$ + drift	5	-4506.8	9023.6	13.5
<b>*Sigmoidal <math>\lambda</math> + drift</b>	<b>7</b>	<b>-4498.0</b>	<b>9010.0</b>	<b>0</b>

Sampling fraction: 1685 / 2539 species = 0.66, Df = degrees of freedom, lnLik = log-likelihood, AIC = Akaike Information Criterion, ΔAIC = different in AIC compared to the best model.

**Supplementary Table 7 Models tested in the Multiple State Speciation and Extinction (MuSSE) framework.**

The sixteen models describe the effect of two binary traits on speciation, extinction and transition rates. These models increase in complexity from the base model in which traits do not affect speciation, extinction and transition rates (model 0) to the most complex model in which speciation, extinction and transition rates are affected by additive and interaction effects of the two binary traits (model 15). \*\*0: no additive or interaction effects; 1: additive effects; 2: additive and interaction effects on rates.

<b>Model</b>	<b>Model description</b>	<b>Model parameters (<math>\lambda</math> <math>\mu</math> <math>q</math>)**</b>	<b>Df</b>
0	Base model	000	6
1	Additive trait effects on $\lambda$	100	8
2	Additive trait effects on $\mu$	010	8
3	Additive and interaction trait effects on $\lambda$	200	9
4	Additive and interaction trait effects on $\mu$	020	9
5	Additive trait effects on $q$	001	10
6	Additive trait effects on $\lambda$ and $\mu$	110	10
7	Additive and interaction trait effects on $\lambda$ , additive effect on $\mu$	210	11
8	Additive and interaction trait effects on $\mu$ , additive effect on $\lambda$	120	11
9	Additive trait effects on $\lambda$ and $q$	101	12
10	Additive trait effects on $\mu$ and $q$	011	12
11	Additive and interaction trait effects on $\lambda$ and $\mu$	220	12
12	Additive trait effects on $\lambda$ , $\mu$ and $q$	111	14
13	Additive and interaction trait effects on $\lambda$ , additive effects on $\mu$ and $q$	211	15
14	Additive and interaction trait effects on $\mu$ , additive effects on $\lambda$ and $q$	121	15
15	Additive and interaction trait effects on $\lambda$ and $\mu$ , additive effects on $q$	221	16

Df = degrees of freedom,  $\lambda$  = speciation rate,  $\mu$  = extinction rate,  $q$  = transition rate.

**Supplementary Table 8 Model selection for fruit size and understory habitat (MuSSE).**

Sixteen Multiple State Speciation and Extinction (MuSSE) models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree for global, New World and Old World palms. These models include the additive and interaction effects of small fruits and understory habitat on speciation, extinction and transition rates, as compared to large-fruited canopy palms (the base model). The best-fitting model given the fewest number of parameters is indicated in bold (\*). As a stepwise model selection approach was applied, not all models were tested for each dataset. For details on the models see Supplementary Table 7. \*\*0: no additive or interaction effects; 1: additive effects; 2: additive and interaction effects on rates.

Model	Model parameters ( $\lambda \mu q$ )**	Df	AIC
<b>Global</b>			
0	000	6	15730
1	100	8	15528
2	010	8	15549
5	001	10	15715
6	110	10	15518
7	111	14	15504
<b>13*</b>	<b>211</b>	<b>15</b>	<b>15491</b>
14	121	15	15506
<b>New World</b>			
0	000	6	5686.9
1	100	8	5590.3
2	010	8	5582.2
5	001	10	5655.2
6	110	10	5588.4
12	111	14	5561.3
<b>13*</b>	<b>211</b>	<b>15</b>	<b>5545.4</b>
14	121	15	5552.5
15	221	16	5546.3
<b>Old World</b>			
0	000	6	10094.2
1	100	8	9916
2	010	8	9935.9
5	001	10	10093
<b>6*</b>	<b>110</b>	<b>10</b>	<b>9899.2</b>
7	210	11	10104.2
8	120	11	10105
12	111	14	9903.1

Df = degrees of freedom, AIC = Akaike Information Criterion,  $\lambda$  = speciation rate,  $\mu$  = extinction rate,  $q$  = transition rate.

**Supplementary Table 9 Model selection for fruit size and island colonization (MuSSE).**

Sixteen Multiple State Speciation and Extinction (MuSSE) models were fitted to the palm Maximum Clade Credibility (MCC) phylogenetic tree for global, New World and Old World palms. These models include the additive and interaction effects of small fruits and oceanic island distributions on speciation, extinction and transition rates, as compared to large-fruited mainland or continental island-distributed palms (the base model). The best-fitting model given the fewest number of parameters is indicated in bold (\*). As a stepwise model selection approach was applied, not all models were tested for each dataset. For details on the models see Supplementary Table 7. \*\*0: no additive or interaction effects; 1: additive effects; 2: additive and interaction effects on rates.


Model	Model parameters ( $\lambda$ $\mu$ $q$ )**	Df	AIC
<b>Global</b>			
0	000	6	13757
1	100	8	13668
2	010	8	13661
5	001	10	13746
6	110	10	13656
12	111	14	13658
<b>7*</b>	<b>210</b>	<b>11</b>	<b>13647</b>
8	120	11	13653
11	220	12	13651
<b>New World</b>			
0	0	6	4670.4
<b>1*</b>	<b>100</b>	<b>8</b>	<b>4659.4</b>
<b>2*</b>	<b>010</b>	<b>8</b>	<b>4658.9</b>
3	200	9	4660.4
4	020	9	4658.9
5	001	10	4665.4
6	110	10	4661.1
9	101	12	4661
10	011	12	4659
<b>Old World</b>			
0	0	6	8998.3
1	100	8	8867.1
2	010	8	8905.2
5	001	10	8996
6	110	10	8857.9
12	111	14	8852.4
<b>13*</b>	<b>211</b>	<b>15</b>	<b>8832.5</b>
14	121	15	8839.5
15	221	16	8834.5

Df = degrees of freedom, AIC = Akaike Information Criterion,  $\lambda$  = speciation rate,  $\mu$  = extinction rate,  $q$  = transition rate.

## Supplementary Figures


### **Supplementary Figure 1 | Correlation between fruit size and plant size in palms.**

The figure shows the relationship between log (fruit size) (cm) and log (maximum stem height) (m), with larger palms having on average larger fruits. The smoothed line was generated in the ggplot R package. The grey box indicates the confidence interval around the smoothed line.




**Supplementary Figure 2 | Fruit size frequency distributions of palms.**

Fruit size frequencies for small- and large-fruited palms, indicated for all palms, understory, canopy, island-distributed and mainland- or continental island- distributed palm species. Understory palms generally have smaller fruits than canopy palms, both for the small fruit (< 4 cm) and large fruit ( $\geq 4$  cm) groups. Mainland palms generally have slightly smaller fruits than island-distributed palms, but only for the small fruit (< 4 cm) group.  $m$  = median,  $n$  = sample size.



### Supplementary Figure 3 | Diversification rates under simulated binary traits (BiSSE).

Bayesian posterior densities resulting from the Binary State Speciation and Extinction (BiSSE) model for speciation ( $\lambda$ ), extinction ( $\mu$ ), transition ( $q$ ), and net diversification ( $r$ ) rates. These rates were estimated by Bayesian Markov Chain Monte Carlo methods for 10,000 generations for palm lineages after simulating binary traits on 100 randomly sampled, empirical palm phylogenetic trees. These binary traits were simulated under different transition rate scenarios. As 95% posterior densities between trait states (indicated with 0 and 1 on the x-axes) within each scenario strongly overlap (compare same colored box-and-whiskers in each plot), we can reject the hypothesis that the distribution of branch lengths in our empirical data causes our observed results (compare to Fig. 2 and Supplementary Fig. 4).




Transition rate used for neutral trait evolution:  
■ 0.01  
■ 0.1  
■ 1  
■ observed (empirical):  $q$  large to small fruits = 0.017;  $q$  small to large fruits = 0.006



### Supplementary Figure 4 | Diversification rates of small- and large-fruited palms (BiSSE).

Bayesian posterior densities resulting from the Binary State Speciation and Extinction model (BiSSE) of speciation ( $\lambda$ ), extinction ( $\mu$ ), transition ( $q$ ) and net diversification ( $r$ ) rates as estimated by Bayesian Markov Chain Monte Carlo methods on 100 randomly sampled palm phylogenetic trees for 10,000 generations, for global, New World (NW), Old World (OW) and residual fruit size (after correcting for maximum stem height) small and large-fruited palm lineages. ‘0’ refers to small-fruited palm lineages, ‘1’ to large-fruited palm lineages, NW = New World, OW = Old World.



Speciation ( $\lambda$ ), extinction ( $\mu$ ) and net diversification ( $r$ ) rate:


- Large fruits  $\geq 4$  cm
- Small fruits  $< 4$  cm

Transition rate ( $q$ )

- Large to small fruits
- Small to large fruits


**Supplementary Figure 5 | Speciation rate in response to fruit size (QuaSSE).**

The effect of palm residual fruit size (cm) on speciation rates under the best-fitting Quantitative Speciation and Extinction (QuaSSE) model (sigmoidal + drift, Supplementary Table 6). This indicates that palms with large fruit sizes have lower rates of speciation than palms with small fruit sizes, following a sigmoidal relationship.




**Supplementary Figure 6 | Phylorate plot of speciation rates in palms (BAMM).**

Phylorate plot showing speciation rates as estimated from Bayesian Analysis of Macroevolutionary Mixtures (BAMM) on the palm Maximum Clade Credibility (MCC) phylogenetic tree. The speciation rates at the tips of the phylogenetic tree (i.e. the species) were regressed against fruit sizes of palms and compared to 1000 permutations. These results indicated that a significant negative relationship between fruit size and speciation rates exists in palms (i.e. palms with larger fruits have on average lower speciation rates than palms with smaller fruits).



**Supplementary Figure 7 | Trait interaction effects on diversification rates (MuSSE).** Bayesian posterior densities resulting from the Multiple State Speciation and Extinction (MuSSE) model for extinction, transition and net diversification rates. These rates were estimated by Bayesian Markov Chain Monte Carlo methods on 100 randomly sampled palm phylogenetic trees for 10,000 generations, for global, New World (NW) and Old World (OW) palms. (a) Rates as estimated for small-fruited palm lineages in the understory, compared to a base model of large-fruited canopy palm lineages. (b) Rates as estimated for small-fruited palm lineages distributed on oceanic islands, compared to a base model of large-fruited mainland- or continental island-distributed palm lineages.



## Supplementary data sources

Trait data sources used to obtain fruit sizes and maximum stem heights of palms, as used in this study.

### **Herbaria:**

Aarhus University Herbarium

Kew Royal Botanic Gardens Herbarium

### **Palm websites:**

Palmweb ([www.palmweb.org](http://www.palmweb.org))

Palmpedia ([www.palmpedia.net](http://www.palmpedia.net))

### **Literature:**

Bacon, C. D. & Baker, W. J. (2011): *Saribus* resurrected. *Palms* 55: 109-116.

Barfod, A. S. (1991): A monographic study of the subfamily Phytelephantoideae (Arecaceae). *Opera Botanica* 105: 1-73

Barrow S. C. (1998): A monograph of *Phoenix* L. (Palmae: Coryphoideae). *Kew Bulletin* 53: 513-575

Bergmann, Birgitte (pers. comm.)

Bernal, R., Galeano, G. & Hodel, D.R. (2004): A new species of *Chamaedorea* from Columbia. *Palms* 48: 27-29.

Blicher-Mathiesen, U. Unpubl.: The subtribe Attaleinae (Palmae) in Ecuador

Borchsenius, F. & Bernal, R. (1996): *Aiphanes* (Palmae). *Flora Neotropica* 70: 1-94

Borchsenius, F., Pedersen, H. B. & Balslev, H. (1998): Manual to the palms of Ecuador. AAU Reports 37, Aarhus University Press, Aarhus.

Cascante, A. (2000): Additions to the Genus *Batris* (Arecaceae) of Mesoamerica. *Palms* 44: 146-153.

de Guzman, E. D. & Fernando, E. S. (1986): Guide to Philippine flora and fauna. Natural Resources Management Center, Ministry of Natural Resources: University of the Philippines.

Dowe J. L. & Hodel D. R. (1994): A revision of *Archontophoenix* H.Wendl. & Drude (Arecaceae). *Austrobaileya* 4: 227-44.

Dowe, J.L. & Ferrero, M.D. (2000): A new species of rheophytic palm from New Guinea. *Palms* 44: 194-197.

Dowe, J. L. & Ferrero, M. D. (2001): Revision of *Calyptrocalyx* and the New Guinea species of *Linospadix* (Linospadicinae: Arecoideae: Arecaceae). *Blumea* 46: 207-251.

Dowe, J. L. (2010): Australian palms: biogeography, ecology and systematics. CSIRO Publishing, Collingwood, Australia.

Dowe, J. L. & Irvine, A. K. (1997): A revision of *Linospadix* in Australia, with the description of a new species. *Principes* 41: 192-197, 211-217.

Dransfield (1979): A manual of the rattans of the Malay Peninsula. *Malayan Forest Records* 29.

Dransfield, J. (1972a): The genus *Borassodendron* in Malesia. *Reinwardtia* 8: 351-363.

Dransfield, J. (1972b): The genus *Johannesteijsmannia* H.E. Moore Jr. *Gard Bull. Singapore* 26: 63-83.

- Dransfield, J. (1978b): The growth forms of rain forest palms. Pages 247–268 in Tomlinson, P.B. and M.H. Zimmermann (eds). *Tropical trees as living systems*. Cambridge University Press. New York.
- Dransfield, J. (1986): *Palmae*. In: *Flora of tropical East Africa* (ed. Polhill, R. M.), Balkema, Rotterdam.
- Dransfield, J. & Beentje, H. (1995): *Palms of Madagascar*. Royal Botanical Gardens, Kew.
- Dransfield, J. (1984): The rattans of Sabah. *Sabah forest record* 13 (1984).
- Dransfield, J., Uhl, N. W., Asmussen, C. B., Baker, W. J., Harley, M. M. & Lewis, C. E. (2008): *Genera Palmarum - the evolution and classification of palms*. Royal Botanic Gardens, Kew, London.
- Essig F. B. (1978): A revision of the genus *Ptychosperma* Labill. (*Arecaceae*). *Allertonia* 1: 415-78.
- Essig, F. B. & Hernandez, N. (2002) A systematic histological study of palm fruits. V. Subtribe *Archontophoenicinae* (*Arecaceae*). *Brittonia*, 54, 65-71.
- Essig, F. B. & Litten, L. (2004) A systematic histological analysis of palm fruits VII. The *Cyrtostachydinae* (*Arecaceae*). *Brittonia*, 56, 375-379.
- Essig, F. B. (2002) A systematic histological study of palm fruits. VI. Subtribe *Linospadicinae* (*Arecaceae*). *Brittonia*, 54, 196-201.
- Essig, F. B. (2008) A systematic histological study of palm fruits. VIII. Subtribe *Dypsidinae* (*Arecaceae*). *Brittonia*, 60, 82-92.
- Essig, F. B., Bussard, L. & Hernandez, N. (2001) A systematic histological study of palm fruits. IV. Subtribe *Oncospermatinae* (*Arecaceae*). *Brittonia*, 53, 466-471.
- Essig, F. B., Manka, T. J. & Bussard, L. (1999) A systematic histological study of palm fruits. III. Subtribe *Iguanurinae* (*Arecaceae*). *Brittonia*, 51, 307-325.
- Evans (2001): The Indochinese rattan *Calamus acanthophyllus*: a fire-loving palm. *Palms* 45. (1): 25-28.
- Fernando, E. S. (1983): A revision of the genus *Nenga*. *Principes* 27: 55-70.
- Fong, F. (1986): Studies on the population structure, growth dynamics and resource importance of nipa palm (*Nypa fruticans* Wurmb.). Ph.D. diss., University of Malaya.
- Galeano, G. & Bernal, R. (2010): *Palmas de Colombia: guia de campo*. Panamericana Formas e Impresos S.A., Bogota.
- Gibbons, M. & Spanner, T. W. (1996): *Medemia argun* lives! *Principes* 40: 65–74.
- Glassman S.F. 1987: Revisions of the palm genus *Syagrus* Mart. And other selected genera in the Cocos alliance. *Illinois Biological Monographs* 59: 1-414.
- Grayum, M.H. (1998 publ. 1999): Nomenclatural and taxonomic notes on Costa Rican palms (*Arecaceae*), with five new species. *Phytologia* 84: 307-327.
- Hahn, W. (1993): *Biosystematics and evolution of the genus Caryota* (*Palmae: Arecoideae*). PhD Thesis, University of Wisconsin, Madison, Wisconsin.
- Henderson A. (2002) *Evolution and ecology of palms*. The New York Botanical Garden Press, Bronx.
- Henderson (2007): A revision of *Wallichia* (*Palmae*). *Taiwania* 52: 1-11.
- Henderson A. (2009): *Palms of Southern Asia*. Princeton University Press, Princeton.
- Henderson A. (2011): A revision of *Geonoma* (*Arecaceae*). *Phytotaxa* 17: 1–271.
- Henderson A., Galeano G. & Bernal R. (1995): *Field guide to the palms of the Americas*. Princeton University Press, Princeton, NJ.

- Henderson, A. (1990): *Arecaceae* part I. Introduction and the *Iriarteinae*. *Flora Neotropica* 53: 1-101.
- Henderson, A. (2000): *Bactris* (Palmae). *Flora Neotropica Monograph* 79: 1-181.
- Henderson, A. (2005): A multivariate study of *Calypstrogyne* (Palmae). *Systematic Botany* 30: 60-83.
- Henderson A. J. & Bacon C. D. (2011): *Lanonia* (Arecaceae: Palmae), a new genus from Asia, with a revision of the species. *Systematic Botany* 36: 883-95.
- Hodel, D. R. (1992a): *Chamaedorea* palms. The International Palm Society, Allen Press, Lawrence, Kansas.
- Hodel, D. R. (1992b): Additions to *Chamaedorea* palms: new species from Mexico and Guatemala and miscellaneous notes. *Principes* 36: 188-202.
- Hodel, D. R. (1995): Three new species of *Chamaedorea* from Panama. *Principes* 39: 14-20.
- Hodel, D. R. (1996): Two new species of *Chamaedorea* from Costa Rica. *Principes* 40: 212-216.
- Hodel, D. R. (1997): Two new species of *Chamaedorea* (Arecaceae). *Novon* 7: 35-37.
- Hodel, D. R. (1998): The palms of Thailand. Allen Press, Lawrence, Kansas.
- Hodel, D. R. (2007): A review of the genus *Pritchardia*. *Palms* 51: S1-S52.
- Hodel, D. R., Herrera, G. & Cascante, A. (1997): A remarkable new species and additional novelties of *Chamaedorea* from Costa Rica and Panama. *Palm Journal* 137: 32-44.
- Hodel, D. R., Mont, J. J. C. & Zuniga, R. (1995): Two new species of *Chamaedorea* from Honduras. *Principes* 39: 183-189.
- Hodel, D. R. & Pintaud, J.-C. (1998): The palms of New Caledonia. Pattaya, Thailand: Kampon Tansacha, Nong Nooch Tropical Garden.
- Jones, D. L. (1995): *Palms throughout the world*. Smithsonian Institution Press, Washington, D.C.
- Keat, L.C. (1998): Notes on recent palm species and records from Peninsular Thailand. *Principes* 42: 110-119.
- Kiew, R. 1977. Taxonomy, ecology and biology of sago palms in Malaysia and Sarawak. In *Sago-76*. Proc. 1st International Sago Symposium, Kuching, Malaysia 5-7 July 1976 (K. Tan, ed.), pp. 147-154.
- Lorenzi, H. (2010): *Brazilian Flora Arecaceae (Palms)*. Nova Odessa, Instituto Plantarum.
- Mogea, J. P. (1991): Indonesia: palm utilization and conservation. In: D. Johnson (ed.), *Palms for human needs in Asia*, A.A. Balkema, Rotterdam, pp. 37-73.
- Moore, H. E. & Guého, L. J. (1984): Palmiers. In: Bosser, J., Cadet, T., Guého, J. & Marais, W. (Eds.). *Flore des Mascareignes. Famille 189*. The Sugar Industry Research Institute, Mauritius, l'Office de la Recherche Scientifique Outre-Mer, Paris, France & Royal Botanic Gardens, Kew, Richmond, United Kingdom. 34 pp.
- Mónica M. R. & Henderson A. (1990): The genus *parajubaea* (Palmae). *Brittonia* 42: 92-9.
- Rauwerdink (1986): Rauwerdink, Jan B. "An essay on *Metroxylon*, the sago palm." *Principes (USA)* (1986).
- Read, R. & Hodel, D. R. (1990): *Arecaceae*. Palm Family. In: W. L. Wagner, D. R. Herbst, and S. H. Sohmer (eds.), *Manual of the Flowering Plants of Hawai'i*. Bishop Museum Special Publication 83. B.P. Bishop Museum, HI, pp. 1360-1375.
- Rodd, A. N. (1998): Revision of *Livistona* (Arecaceae) in Australia. *Telopea* 8: 49-153.
- Russell, T. A. (1968): Palmae. In: *Flora of west tropical Africa* (ed. Hepper, F.N.), 2nd ed. Whitefriars Press, London.

Saw, L. G., Dransfield, J. and Keith-Lucas, D. M. (2003): Morphological diversity of the genus *Licuala* (Palmae). *Telopea* 10: 187-206

Schönfeldter R., unpubl.

Skov (1994): *Geonoma polyandra* (Arecaceae), a new species from Ecuador. *Nordic journal of botany* 14.1: 39-41.

Skov, F. (1989): HyperTaxonomy - a new tool for revisional work and a revision of *Geonoma* (Palmae) in Ecuador, unpublished

Skov and Balslev (1989): Skov, Flemming, and Henrik Balslev. "A revision of *Hyospathe* (Arecaceae)." *Nordic journal of botany* 9.2 (1989): 189-202.

Stauffer F. W., Asmussen C. B., Henderson A. & Endress P. K. (2003): A revision of *Asterogyne* (Arecaceae: Arecoideae: Geonomeae). *Brittonia* 55: 326-56.

Sunderland (2007): Sunderland, Terry CH. *Field guide to the rattans of Africa*. Royal Botanic Gardens, 2007.

Tuley, P. (1995): *The palms of Africa*. The Trendrine Press, Zennor.

Zona, S. (1990): A monograph of *Sabal* (Arecaceae: Coryphoideae). *Aliso* 12: 583-666.



## Supplementary references

- 1 FitzJohn, R. G. Quantitative Traits and Diversification. *Systematic Biology* **59**, 619-633, doi:10.1093/sysbio/syq053 (2010).
- 2 Rabosky, D. L. & Goldberg, E. E. Model inadequacy and mistaken inferences of trait-dependent speciation. *Systematic Biology* **64**, 340-355, doi:10.1093/sysbio/syu131 (2015).
- 3 Lewis, P. O. A likelihood approach to estimating phylogeny from discrete morphological character data. *Systematic Biology* **50**, 913-925 (2001).
- 4 Gamisch, A. Notes on the Statistical Power of the Binary State Speciation and Extinction (BiSSE) Model. *Evolutionary Bioinformatics Online* **12**, 165-174, doi:10.4137/EBO.S39732 (2016).
- 5 Davis, M. P., Midford, P. E. & Maddison, W. Exploring power and parameter estimation of the BiSSE method for analyzing species diversification. *BMC Evolutionary Biology* **13**, 1-11, doi:10.1186/1471-2148-13-38 (2013).
- 6 Maddison, W. P. & FitzJohn, R. G. The unsolved challenge to phylogenetic correlation tests for categorical characters. *Systematic Biology* **64**, 127-136, doi:10.1093/sysbio/syu070 (2015).
- 7 Rabosky, D. L. Automatic Detection of Key Innovations, Rate Shifts, and Diversity-Dependence on Phylogenetic Trees. *PLOS ONE* **9**, e89543, doi:10.1371/journal.pone.0089543 (2014).
- 8 Faurby, S., Eiserhardt, W. L., Baker, W. J. & Svenning, J.-C. An all-evidence species-level supertree for the palms (Arecaceae). *Molecular Phylogenetics and Evolution* **100**, 57-69, doi:<http://dx.doi.org/10.1016/j.ympev.2016.03.002> (2016).
- 9 Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of evolutionary dynamics on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701-707, doi:10.1111/2041-210X.12199 (2014).
- 10 Rabosky, D. L. & Huang, H. A robust semi-parametric test for detecting trait-dependent diversification. *Systematic Biology* **65**, 181-193, doi:10.1093/sysbio/syv066 (2016).
- 11 Couvreur, T. L. P. *et al.* Global diversification of a tropical plant growth form: environmental correlates and historical contingencies in climbing palms. *Frontiers in Genetics* **5**, 452, doi:10.3389/fgene.2014.00452 (2014).
- 12 FitzJohn, R. G. Diversitree: comparative phylogenetic analyses of diversification in R. *Methods in Ecology and Evolution* **3**, 1084-1092, doi:10.1111/j.2041-210X.2012.00234.x (2012).