



HAL
open science

Complete plastome sequences of 14 African yam species (*Dioscorea* spp.)

Jacqueline Magwé-Tindo, Jan J. Wieringa, Bonaventure Sonké, Louis Zapfack, yves Vigouroux, Thomas Couvreur, Nora Scarcelli

► To cite this version:

Jacqueline Magwé-Tindo, Jan J. Wieringa, Bonaventure Sonké, Louis Zapfack, yves Vigouroux, et al.. Complete plastome sequences of 14 African yam species (*Dioscorea* spp.). *Mitochondrial DNA Part B Resources*, Taylor & Francis Online, 2019, 4 (1), pp.74-76. 10.1080/23802359.2018.1536466 . hal-03364209

HAL Id: hal-03364209

<https://hal.umontpellier.fr/hal-03364209>

Submitted on 4 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

Complete plastome sequences of 14 African yam species (*Dioscorea* spp.)

Jacqueline Magwé-Tindo, Jan J. Wieringa, Bonaventure Sonké, Louis Zapfack, Yves Vigouroux, Thomas L. P. Couvreur & Nora Scarcelli

To cite this article: Jacqueline Magwé-Tindo, Jan J. Wieringa, Bonaventure Sonké, Louis Zapfack, Yves Vigouroux, Thomas L. P. Couvreur & Nora Scarcelli (2019) Complete plastome sequences of 14 African yam species (*Dioscorea* spp.), *Mitochondrial DNA Part B*, 4:1, 74-76, DOI: [10.1080/23802359.2018.1536466](https://doi.org/10.1080/23802359.2018.1536466)

To link to this article: <https://doi.org/10.1080/23802359.2018.1536466>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 04 Jan 2019.



Submit your article to this journal [↗](#)



View Crossmark data [↗](#)

Complete plastome sequences of 14 African yam species (*Dioscorea* spp.)

Jacqueline Magwé-Tindo^a, Jan J. Wieringa^b, Bonaventure Sonké^c, Louis Zapfack^a, Yves Vigouroux^d, Thomas L. P. Couvreur^{b,d} and Nora Scarcelli^d

^aDepartment of Plant Biology, Faculty of Science, The University of Yaoundé I, Yaoundé, Cameroon; ^bNational Herbarium of The Netherlands, Naturalis Biodiversity Centre, Leiden, The Netherlands; ^cDepartment of Biology, Higher Teachers' Training College, The University of Yaoundé I, Yaoundé, Cameroon; ^dIRD, University Montpellier, Montpellier, France

ABSTRACT

Complete plastomes of 14 African yam species were reconstructed from whole genome sequencing. These plastomes sizes varied from 151,908 to 155,155 bp. We predicted 130 genes, including 18 duplicated genes located in the two inverted regions. Phylogenetic analysis obtained using the maximum likelihood procedure revealed that each species was distinct with high support. This resource will be useful for phylogenetic and diversity analysis of *Dioscorea* species.

ARTICLE HISTORY

Received 7 September 2018
Accepted 20 September 2018

KEYWORDS

Africa; chloroplast;
Dioscorea; NGS; yam

The genus *Dioscorea* L. (Dioscoreaceae) plays a considerable role in tropical Africa because of its important staple crop *D. rotundata*, a vine cultivated for its starchy tuber. *Dioscorea* is a very species-rich and morphologically complex genus. Therefore, the taxonomy of this genus is difficult, as proven by the variation in the number of accepted species names (Caddick et al. 2002; Wilkin et al. 2005; Govaerts et al. 2017). Molecular phylogenetic analyses focusing on the genus are also poorly resolved (Terauchi et al. 1992; Ramser et al. 1997; Chair et al. 2005). Here we reconstructed complete plastomes of 14 West-African species.

Fourteen *Dioscorea* leaf samples, (*D. abyssinica*, *D. baya*, *D. bulbifera*, *D. burkilliana*, *D. cayenensis*, *D. dumetorum*, *D. hirtiflora*, *D. praehensilis*, *D. preussii*, *D. quartiniana*, *D. sagittifolia*, *D. sansibarensis*, *D. schimperiana*, and *D. togoensis*), were collected from WAG and G herbaria (Table 1). DNAs were extracted from dried leaves according to Scarcelli et al. (2006) and whole genome sequencing was performed as described in Mariac et al. (2014) on an Illumina MiSeq (Illumina, San Diego, CA). For each sample, the whole plastome was reconstructed using MITObim 1.7 (Hahn et al. 2013) as explained in Magwé-Tindo et al. (2018) with *D. rotundata* plastome as a reference (GenBank NC_024170.1). Annotations were recovered from the alignment of the target plastome to NC_024170.1 using Geneious Pro 4.8.5 with Mauve 2.2.0 plugin (Darling et al. 2010), then manually checked for start and stop codons (protein-coding genes) and by blasting the tRNA and rRNA sequences to GenBank nucleotide database (nr/nt, Altschul et al. 1990). Annotated

plastomes were deposited into the GenBank database (Table 1).


Plastome lengths varied from 151,908 bp (*D. burkilliana*) to 155,155 bp (*D. baya*) and exhibited a typical quadripartite structure with a LSC (Large Single Copy) and a SSC (Short Single Copy) separated by two IR (Inverted Region). The mean GC content was 37% and varied from 31% within the SSC to 43% within the IR. The 14 species presented the same genetic structure as the reference (NC_024170.1): 130 genes corresponding to 84 protein-coding genes, 38 tRNA and 8 rRNA. Among these genes, 18 were duplicated in the IR: 6 protein-coding genes, 8 tRNA and 4 rRNA. The only exception was *D. quartiniana* which missed the *matK* gene due to a mutation of the second codon into a stop codon. All species lost the *rps16* gene.

A phylogenetic study was conducted using these 14 complete plastomes, with *D. rotundata* (NC_024170.1) as reference and *D. elephantipes* (NC_009601.1) as outgroup species. The 16 plastomes were aligned using MAFFT 7 (Katoh et al. 2002) and the alignment was cleaned using Gblocks 0.91b (Talavera and Castresana 2007). The phylogenetic tree was constructed using RAxML 8.2.9 (Stamatakis 2014) with 500 bootstraps. The phylogenetic tree obtained was well-resolved and revealed that each species was clearly distinct with strong support (Figure 1), except within the cultivated species complex.

Resources reconstructed here will help to understand the relationships between cultivated yams and their wild relatives and provide useful information for further taxonomic study.

Table 1. Characterization of the samples used for sequencing.

Species	Herbarium voucher accession	Collection locality	Longitude	Latitude	Plastome GenBank ID
<i>D. abyssinica</i>	WAG.1781186	Cameroon	13.9	10.9	MG805602
<i>D. baya</i>	WAG.1781135	Cameroon	10.6	2.9	MG805603
<i>D. bulbifera</i>	WAG.1783436	Liberia	-8.6	6.3	MG805604
<i>D. burkilliana</i>	WAG.1783323	Cameroon	11.2	2.8	MG805605
<i>D. cayenensis</i>	WAG.1783287	Cameroon	11.5	3.9	MG805606
<i>D. dumetorum</i>	WAG.1943398	Gabon	10.7	-3.5	MG805607
<i>D. hirtiflora</i>	WAG.1783636	Senegal	-16.7	12.9	MG805608
<i>D. praehensilis</i>	G00420196	Côte d'Ivoire	-5.1	6.3	MG805609
<i>D. preussii</i>	WAG.1786123	Gabon	11.8	-0.1	MG805610
<i>D. quartiniana</i>	WAG.1786396	Malawi	35.5	-15.9	MG805611
<i>D. sagittifolia</i>	WAG.1786430	Cameroon	13.8	7	MG805612
<i>D. sansibarensis</i>	WAG.1786456	Gabon	11.6	-0.2	MG805613
<i>D. schimperiana</i>	WAG.1786876	Cameroon	11.5	3.9	MG805614
<i>D. togoensis</i>	WAG0283548	Togo	1.6	6.7	MG805615

**Figure 1.** Phylogenetic tree of 16 African *Dioscorea* species constructed using maximum likelihood procedure and whole plastome sequences. *Dioscorea elephantipes* was included as outgroup species. Bootstrap values are indicated for each node. Each species is followed by its GenBank accession ID.

Acknowledgements

The authors thank the Conservatoire et Jardin botaniques de la ville de Genève (G) for the *D. praehensilis* sample.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This project is supported by Agropolis Fondation under the reference ID 1502-206 through the « Investissements d'avenir » program (Labex Agro:ANR-10-LABX-0001-01), under the frame of I-SITE MUSE (ANR-16-IDEX-0006); by ARCAD and FEDER; and by Agence Nationale de la Recherche under Grant ANR-13-BSV7-0017. This work was done in collaboration with the GeT core facility, Toulouse, France (ANR-10-INBS-09).

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol.* 215:403–410.
- Caddick LR, Rudall PJ, Wilkin P, Hedderson TAJ, Chase MW. 2002. Phylogenetics of Dioscoreales based on combined analyses of morphological and molecular data. *Bot J Linn Soc.* 138: 123–144.
- Chair H, Perrier X, Agbangla C, Marchand JL, Daïnou O, Noyer JL. 2005. Use of cpSSRs for the characterisation of yam phylogeny in Benin. *Genome.* 48:674–684.
- Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One.* 5: e11147.
- Govaerts R, Wilkin P, Raz L, Tellez-Valdes O. 2017. World checklist of *Dioscoreales*. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <http://wmsp.science.kew.org> (accessed 30 June 2017).
- Hahn C, Bachmann L, Chevreux B. 2013. Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads—a baiting and iterative mapping approach. *Nucleic Acids Res.* 41:e129.
- Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30:3059–3066.
- Magwé-Tindo J, Wieringa JJ, Sonké B, Zapfack L, Vigouroux Y, Couvreur TLP, Scarcelli N. Forthcoming 2018. Guinea yam (*Dioscorea* spp.) wild relatives identified using whole plastome phylogenetic analyses. *Taxon.* 67(5):905–915.
- Mariac C, Scarcelli N, Pouzadou J, Barnaud A, Billot C, Faye A, Kougbéadjou A, Maillol V, Martin G, Sabot F, et al. 2014. Cost-effective enrichment hybridization capture of chloroplast genomes at deep multiplexing levels for population genetics and phylogeography studies. *Mol Ecol Resour.* 14:1103–1113.
- Ramser J, Weising K, Lopez-Peralta C, Terhalle W, Terauchi R, Kahl G. 1997. Molecular marker based taxonomy and phylogeny of Guinea yam (*Dioscorea rotundata* - *D. cayenensis*). *Genome.* 40:903–915.
- Scarcelli N, Tostain S, Mariac C, Agbangla C, Da O, Berthaud J, Pham J-L. 2006. Genetic nature of yams (*Dioscorea* sp.) domesticated by farmers in Benin (West Africa). *Genet Resour Crop Evol.* 53:121–130.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 30:1312–1313.
- Talavera G, Castresana J. 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol.* 56:564–577.
- Terauchi R, Chikaleke VA, Thottappilly G, Hahn SK. 1992. Origin and phylogeny of Guinea yams as revealed by RFLP analysis of chloroplast DNA and nuclear ribosomal DNA. *Theor Appl Genet.* 83:743–751.
- Wilkin P, Schols P, Chase MW, Chayamarit K, Furness CA, Huysmans S, Rakotonasolo F, Smets E, Thapayai C. 2005. A plastid gene phylogeny of the yam genus, *Dioscorea*: roots, fruits and Madagascar. *Syst Botany.* 30:736–749.