



**HAL**  
open science

## **Cabbage and fermented vegetables: From death rate heterogeneity in countries to candidates for mitigation strategies of severe COVID-19**

Jean Bousquet, Josep Anto, Wienczyslawa Czarlewski, Tari Haahtela, Susana Fonseca, Guido Iaccarino, Hubert Blain, Alain Vidal, Aziz Sheikh, Cezmi Akdis, et al.

### ► To cite this version:

Jean Bousquet, Josep Anto, Wienczyslawa Czarlewski, Tari Haahtela, Susana Fonseca, et al.. Cabbage and fermented vegetables: From death rate heterogeneity in countries to candidates for mitigation strategies of severe COVID-19. *Allergy*, 2021, 76 (3), pp.735-750. 10.1111/all.14549 . hal-03367962

**HAL Id: hal-03367962**

**<https://hal.umontpellier.fr/hal-03367962v1>**

Submitted on 19 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

2 DR. TARI HAAHTELA (Orcid ID : 0000-0003-4757-2156)

3 PROF. CEZMI AKDIS (Orcid ID : 0000-0001-8020-019X)

4 PROF. TORSTEN ZUBERBIER (Orcid ID : 0000-0002-1466-8875)

5

6

7 Article type : Review

8

9

10 **Cabbage and fermented vegetables: from death rate heterogeneity in countries**  
11 **to candidates for mitigation strategies of severe COVID-19**

12

13 Jean Bousquet(1,2), Josep M Anto (3-6), WiencyslawaCzarlewski (7,8), TariHaahtela(9), Susana C Fonseca  
14 (10), Guido Iaccarino(11), HubertBlain(12), Alain Vidal (13), Aziz Sheikh (14), Cezmi A Akdis(15),  
15 TorstenZuberbier(1), and the ARIA group

16

17 1. Charité, Universitätsmedizin Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health,  
18 Comprehensive Allergy Center, Department of Dermatology and Allergy, Berlin, Germany.

19 2. MACVIA-France and CHU, Montpellier, France.

20 3. ISGlobAL, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain.

21 4. IMIM (Hospital del Mar Research Institute), Barcelona, Spain.

22 5. Universitat Pompeu Fabra (UPF), Barcelona, Spain.

23 6. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain.

24 7. MASK-air, Montpellier, France.

25 8. Medical Consulting Czarlewski, Levallois, France.

26 9. Skin and Allergy Hospital, Helsinki University Hospital, and University of Helsinki, Finland.

27 10. GreenUPorto - Sustainable Agrifood Production Research Centre, DGAOT, Faculty of Sciences, University of  
28 Porto, Campus de Vairão, Porto, Portugal.

29 11. Federico II University, Department of Advanced Biomedical Sciences, Napoli, Italy.

30 12. Department of Geriatrics, Montpellier University hospital and MUSE, Montpellier, France.

31 13. World Business Council for Sustainable Development (WBCSD), Maison de la Paix, Geneva, Switzerland &

32 AgroParisTech - Paris Institute of Technology for Life, Food and Environmental Sciences, Paris, France.

33 14. Usher Institute, University of Edinburgh, Scotland, UK.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14549](https://doi.org/10.1111/ALL.14549)

This article is protected by copyright. All rights reserved

34 15. Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland.

35

36

37

38 **Short title: Mitigation of COVID-19 by diet**

39

40 **Address for correspondence**

41

42 Professor Jean Bousquet

43 273 avenue d'Occitanie, 34090 Montpellier, France Tel +33 611 42 88 47, Fax +33 467 41 67 01

44 jean.bousquet@orange.fr

45

## 46 **Abstract**

47

48 Large differences in COVID-19 death rates exist between countries and between regions of the same  
49 country. Some very low death rate countries such as Eastern Asia, Central Europe or the Balkans have a  
50 common feature of eating large quantities of fermented foods. Although biases exist when examining  
51 ecological studies, fermented vegetables or cabbage were associated with low death rates in European  
52 countries. SARS-CoV-2 binds to its receptor, the angiotensin converting enzyme 2 (ACE2). As a result of  
53 SARS-Cov-2 binding, ACE2 downregulation enhances the angiotensin II receptor type 1 (AT<sub>1</sub>R) axis  
54 associated with oxidative stress. This leads to insulin resistance as well as lung and endothelial damage, two  
55 severe outcomes of COVID-19. The nuclear factor (erythroid-derived 2)-like 2 (Nrf2) is the most potent  
56 antioxidant in humans and can block the AT<sub>1</sub>R axis. Cabbage contains precursors of sulforaphane, the most  
57 active natural activator of Nrf2. Fermented vegetables contain many lactobacilli, which are also potent Nrf2  
58 activators. Three examples are given: Kimchi in Korea, westernized foods and the slum paradox. It is  
59 proposed that fermented cabbage is a proof-of-concept of dietary manipulations that may enhance Nrf2-  
60 associated antioxidant effects helpful in mitigating COVID-19 severity.

61 **Key words:** COVID-19, diet, sulforaphane, Lactobacillus, Angiotensin converting enzyme 2, kimchi,  
62 cabbage, fermented vegetable

## 63 **Abbreviations**

64 ACE: Angiotensin converting enzyme

65 Ang II: Angiotensin II

66 AT<sub>1</sub>R: Angiotensin II receptor type 1

67 COVID-19: Coronavirus 19 disease

68 GI: Gastro-intestinal

69 LAB: Lactic acid bacilli

70 NF-κB: Nuclear factor kappa B

71 Nrf2: Nuclear factor (erythroid-derived 2)-like 2

72 PEDV: Porcine epidemic diarrhea virus

73 ROS: Reactive oxygen species

74 SARS: Severe acute respiratory syndrome

75 SARS-Cov-2: Severe acute respiratory syndrome coronavirus 2

76 TGEV: Transmissible Gastroenteritis Coronavirus Infection

## 80 Introduction

81 A COVID-19 epidemic started in China and then disseminated to other Asian countries before becoming a  
82 pandemic. There is a large variability across countries in both incidence and mortality, and most of the  
83 current debates on COVID-19 focus on the differences between countries. Several intertwined factors can  
84 be proposed: social distancing, health system capacity, age of the population, social lifestyle (gathering of  
85 family/friends, social behavior), testing capacity and/or timing and intensity of the first outbreak. German  
86 fatalities are strikingly low as compared to many European countries. Among the several explanations  
87 proposed, an early and large testing of the population was put forward <sup>1</sup> as well as social distancing.  
88 However, little attention has been given to regional within-country differences that may propose new  
89 hypotheses.

90 It would appear that the pandemic has so far resulted in proportionately fewer deaths in some Central  
91 European countries, the Balkans, China, in most Eastern Asian countries as well as in many Sub-Saharan  
92 African countries. Several reasons can explain this picture. One of them may be the type of diet in these  
93 low mortality countries. <sup>2,3</sup>

94 Diet has been proposed to mitigate COVID-19.<sup>4,5</sup> Some foods or supplements may have a benefit on the  
95 immune response to respiratory viruses. However, to date, there are no specific data available to confirm  
96 the putative benefits of diet supplementation, probiotics, and nutraceuticals in the current COVID-19  
97 pandemic. <sup>6</sup> News and social media platforms have implicated dietary supplements in the treatment and  
98 prevention of COVID-19 without evidence.<sup>7</sup>

99 In this paper, we discuss country and regional differences in COVID-19 deaths. We attempt to find  
100 potential links between foods and differences at the national or regional levels in the aim to propose a  
101 common mechanism focusing on oxidative stress that may be relevant in COVID-19 mitigation strategies.  
102 We used cabbage and fermented vegetable as a proof-of-concept.

### 103 1- Biases to be considered

104 According to the Johns Hopkins coronavirus resource center (<https://coronavirus.jhu.edu>), one of the most  
105 important ways of measuring the burden of COVID-19 is mortality. However, death rates are assessed  
106 differently between countries and there are many biases that are almost impossible to assess. Using the  
107 rates of COVID-19 confirmed cases is subject to limitations that are similar to or even worse than the  
108 differences in the use of COVID-19 testing.

109 Differences in the mortality rates depend on health care systems, the reporting method and many unknown  
110 factors. Countries throughout the world have reported very different case fatality ratios - the number of  
111 deaths divided by the number of confirmed cases - but these numbers cannot be compared easily due to  
112 biases. On the other hand, for many countries, the methodology used to report death rates in the different  
113 regions is standardized across the country.

114 We used mortality per number of inhabitants to assess death rates, as proposed by the European Center for  
115 Disease Prevention and Control (ecdc, <https://www.ecdc.europa.eu/en>), and to report trends with cutoffs at  
116 25, 50, 100 and 250 per million.

117 Our hypothesis is mostly based on ecological data that are hypothesis-generating and that require  
118 confirmation by proper studies.

## 119 **2- Multifactorial origin of the COVID-19 epidemic**

120 Like most diseases, COVID-19 exhibits large geographical variations which frequently remain unexplained  
121 <sup>8</sup>. The COVID-19 epidemic is multifactorial, and factors like climate, population density, age, phenotype  
122 and prevalence of non-communicable diseases are also associated to increased incidence and mortality <sup>9</sup>.  
123 Diet represents only one of the possible causes of the COVID-19 epidemic and its importance needs to be  
124 better assessed. Some risk factors for the COVID-19 epidemics are proposed at the individual and country  
125 levels in Table 1.

## 126 **3- Ecological data on COVID-19 death rates**

127 When comparing death rates, large differences exist between and within countries and the evolution of the  
128 pandemic differs largely between countries (Figure 1). Although there are many pitfalls in analyzing death  
129 rates for COVID-19,<sup>3</sup> the evolution of death rates between May 20 and July 18 shows a dramatic increase  
130 in Latin America and only some increase in European countries, certain African countries, the Middle East,  
131 India, Pakistan and some of the South East Asian countries. However, there is no change in the very low  
132 death rates of Cambodia, China, Japan, Korea, Lao, Malaysia, Taiwan, Vietnam and of many Sub-Saharan  
133 African countries, Australia and New Zealand. This geographical pattern is very unlikely to be totally due  
134 to reporting differences between countries.

135 In some high death-rate countries such as Italy (Figure 2), variations are extremely large from 50 per  
136 million in Calabria to over 1,600 in Lombardia. In Switzerland, the French- and Italian-speaking cantons  
137 have a far higher death rate than the German-speaking ones (*Office fédéral de la santé publique*,  
138 Switzerland) (Figure 3). It may be proposed that the high-death rate cantons were contaminated by French

139 and Italian people. However, the Mulhouse airport serves the region of Basel (Switzerland), the Haut-Rhin  
140 department (France) and the region of Freiburg (Germany). There was a COVID-19 outbreak in the Haut-  
141 Rhin department, in particular in Mulhouse and Colmar. The death rate for COVID-19 (May 20, 2020) was  
142 935 per million inhabitants in France but only 10 to 25 in Switzerland and 7 in Germany. It is important to  
143 consider these regional differences since reporting of deaths is similar within the country and many factors  
144 may be considered.

145 In many Western countries, large cities (e.g. London, Madrid, Milan, New York, Paris) have been the most  
146 affected. This seems to be true also for many countries in which the rural areas have much fewer cases.

147 The number of deaths is relatively low in Sub-Saharan Africa compared to other regions, and the low  
148 population density (which applies in rural areas but not in megacities such as Cairo or Lagos) or the  
149 differences in health infrastructure are unlikely to be the only explanation. <sup>10</sup> It has been proposed that hot  
150 temperature may reduce COVID-19, but, in Latin American countries, death rates are high (e.g. Brazil,  
151 Ecuador, Peru and Mexico).

#### 152 **4- Is diet partly responsible for differences between and within countries?**

153 Nutrition may play a role in the immune defense against COVID-19 and may explain some of the  
154 differences seen in COVID-19 between and within countries <sup>3</sup>. In this concept paper, raw and fermented  
155 cabbage were proposed to be candidates.

156 To test the potential role of fermented foods in the COVID-19 mortality in Europe, an ecological study, the  
157 European Food Safety Authority (EFSA) Comprehensive European Food Consumption Database, was used  
158 to study the country consumption of fermented vegetables, pickled/marinated vegetables, fermented milk,  
159 yoghurt and fermented sour milk. <sup>11</sup> Of all the variables considered, including confounders, only fermented  
160 vegetables reached statistical significance with the COVID-19 death rate per country. For each g/day  
161 increase in consumption of fermented vegetables of the country, the mortality risk for COVID-19 was  
162 found to decrease by 35.4% (Figure 4).

163 A second ecological study has analyzed cruciferous vegetables (broccoli, cauliflower, head cabbage (white,  
164 red and savoy cabbage), leafy brassica) and compared them with spinach, cucumber, courgette, lettuce and  
165 tomato <sup>12</sup>. Only head cabbage and cucumber reached statistical significance with the COVID-19 death rate  
166 per country. For each g/day increase in the average national consumption of some of the vegetables (head  
167 cabbage and cucumber), the mortality risk for COVID-19 decreased by a factor of 11, to 13.6 %. The  
168 negative ecological association between COVID-19 mortality and consumption of cabbage and cucumber

169 supports the *a priori* hypothesis previously reported. However, these are ecological studies that need to be  
170 further tested.

171 Another diet component potentially relevant in COVID-19 mortality may be the food supply chain and  
172 traditional groceries.<sup>13</sup> The impact of the long supply chain of food on health is measurable by an increase  
173 in metabolic syndrome and insulin resistance.<sup>14</sup> Therefore, areas that are more prone to short supply food  
174 and traditional groceries may have been able to better tolerate COVID-19 with a lower death toll. These  
175 considerations may be partly involved in the lower death rates of Southern Italy compared to the Northern  
176 part (Figure 2).

## 177 **5- Fermented foods, microbiome and lactobacilli**

178 The fermentation process, born as a preservation method in the Neolithic age, enabled humans to eat not-  
179 so-fresh food and to survive.<sup>15</sup> Indigenous fermented foods such as bread, cheese, vegetables and alcoholic  
180 beverages have been prepared and consumed for thousands of years, are strongly linked to culture and  
181 tradition, especially in rural households and village communities, and are consumed by hundreds of  
182 millions of people.<sup>16</sup> Fermented foods are “foods or beverages made via controlled microbial growth  
183 (including lactic acid bacteria (LAB)) and enzymatic conversions of food components.”<sup>17</sup> Not all fermented  
184 foods contain live cultures, as some undergo further processing after fermentation: pasteurization, smoking,  
185 baking, or filtration. These processes kill or remove the live microorganisms in foods such as soy sauces,  
186 bread, most beers and wines as well as chocolate. Live cultures can be found in fermented vegetables and  
187 fermented milk (fermented sour milk, yoghurt, probiotics, etc.).

188 Most traditional foods with live bacteria in the low-death rate countries are based on LAB fermentation  
189 <sup>18</sup>. A number of bacteria are involved in the fermentation of kimchi and other Korean traditional fermented  
190 foods, but LAB - including *Lactobacillus*- are the dominant species in the fermentation process<sup>19,20</sup>.  
191 *Lactobacillus* is also an essential species in the fermentation of sauerkraut, Taiwanese <sup>21</sup>, Chinese <sup>22</sup> or  
192 other fermented foods<sup>23</sup>. Lactobacilli are among the most common microorganisms found in kefir, a  
193 traditional fermented milk beverage <sup>24</sup>, milk and milk products <sup>25,26</sup>. During fermentation, LAB synthesize  
194 vitamins and minerals, and produce biologically-active peptides with anti-oxidant activity <sup>17,27-31</sup>.

195 Humans possess two protective layers of biodiversity, and the microbiome has been proposed as an  
196 important actor of COVID-19 <sup>32</sup>. The environment (outer layer) affects our lifestyle, shaping the  
197 microbiome (inner layer). <sup>33</sup> Many fermented foods contain living microorganisms and modulate the  
198 intestinal microbiome. <sup>17,31,34-36</sup>



199 The composition of microbiomes varies in different regions of the world.<sup>37</sup> Gut microbiota has an inter-  
200 individual variability due to genetic predisposition and diet.<sup>38</sup> As part of the gut  
201 microbiome, *Lactobacillus* spp. contributes to its diversity and modulates oxidative stress in the GI tract.  
202 Some foods like cabbage can be fermented by the gut microbiota.<sup>39</sup>

203 Westernized foods usually lack fermented vegetables and milk-derived products have less biodiversity than  
204 traditional ones. Urbanization in western countries was associated with changes in the gut microbiome and  
205 with intestinal diversity reduction.<sup>38,40-43</sup> Westernized food in Japan led to changes in the microbiome and  
206 in insulin resistance.<sup>44</sup> The gut microbiome of westernized urban Saudis had a lower biodiversity than that  
207 of the traditional Bedouin population.<sup>45</sup> Fast food consumption was characterized by reduced Lactobacilli in  
208 the microbiome.<sup>46</sup>

209 The links between gut microbiome, inflammation, obesity and insulin resistance are being observed but  
210 further large studies are needed for a definite conclusion.<sup>47-49</sup>

211 Some COVID-19 patients have intestinal microbial dysbiosis<sup>50</sup> with decreased probiotics such  
212 as *Lactobacillus* and *Bifidobacterium*<sup>51</sup>. Many bacteria are involved in the fermentation of vegetables but  
213 most traditional foods with live bacteria in the low-death rate countries are based on LAB fermentation.<sup>18-</sup>  
214 <sup>20,23,30</sup> Lactobacilli are among the most common microorganisms found in milk and milk products<sup>24-26</sup>.

## 215 **6- Angiotensin-converting enzyme 2 (ACE2) and COVID-19**

216 COVID-19 is more severe in older adults and/or patients with comorbidities, such as diabetes, obesity or  
217 hypertension, suggesting a role for insulin resistance.<sup>52</sup> Although differences exist between countries, the  
218 same risk factors for severity were found globally, suggesting common mechanisms. A strong relationship  
219 between hyperglycemia, impaired insulin pathway, and cardiovascular disease in type 2 diabetes is linked  
220 to oxidative stress and inflammation.<sup>53</sup> Lipid metabolism has an important role to play in obesity, in  
221 diabetes and its multi-morbidities, and in ageing.<sup>54</sup> The increased severity of COVID-19 in diabetes,  
222 hypertension, obese or elderly individuals may be related to insulin resistance, with oxidative stress as a  
223 common pathway.<sup>55</sup> Moreover, the severe outcomes of COVID-19 - including lung damage, cytokine storm  
224 or endothelial damage - appear to exist globally, again suggesting common mechanisms.

225 The angiotensin-converting enzyme 2 (ACE2) receptor is part of the dual system -renin-angiotensin-  
226 system (RAS) - consisting of an ACE-Angiotensin-II-AT<sub>1</sub>R axis and an ACE-2-Angiotensin-(1-7)-Mas  
227 axis. AT<sub>1</sub>R is involved in most of the effects of Ang II, including oxidative stress generation,<sup>56</sup> which in  
228 turn upregulates AT<sub>1</sub>R<sup>57</sup>. In metabolic disorders and with older age, there is an upregulation of the AT<sub>1</sub>R  
229 axis leading to pro-inflammatory, pro-fibrotic effects in the respiratory system, and to insulin resistance.<sup>58</sup>

230 SARS-CoV-2 binds to its receptor ACE2 and exploits it for entry into the cell. The ACE2 downregulation,  
231 as a result of SARS-CoV-2 binding, enhances the AT<sub>1</sub>R axis<sup>59</sup> likely to be associated with insulin  
232 resistance<sup>60,61</sup> but also to severe outcomes of COVID-19 (Figure 5A).

## 233 **7- Anti-oxidant activities of foods linked with COVID-19**

234 Many foods have an antioxidant activity<sup>62-64</sup> and the role of nutrition has been proposed to mitigate  
235 COVID-19<sup>65</sup>. Many antioxidant mechanisms have been proposed, and several foods can interact with  
236 transcription factors related to antioxidant effects such as the Nuclear factor (erythroid-derived 2)-like 2  
237 (Nrf2).<sup>4</sup> Some processes like fermentation increase the antioxidant activity of milk, cereals, fruit,  
238 vegetables, meat and fish.<sup>29</sup>

### 239 **7-1- Nrf2, a central antioxidant system**

240 Reactive oxygen species (ROS), such as hydrogen peroxide and superoxide anion, exert beneficial and  
241 toxic effects on cellular functions. Nrf2 is a pleiotropic transcription factor at the centre of a complex  
242 regulatory network that protects against oxidative stress and the expression of a wide array of genes  
243 involved in immunity and inflammation, including antiviral actions.<sup>66</sup> Nrf2 activity in response to chemical  
244 insults is regulated by a thiol-rich protein named KEAP1 (Kelch-like ECH-associated protein 1). The  
245 KEAP1-Nrf2 system is the body's dominant defense mechanism against ROS.<sup>67</sup> Induction of the antioxidant  
246 responsive element and the ROS mediated pathway by Nrf2 reduces the activity of nuclear factor kappa B  
247 (NF-κB),<sup>68</sup> whereas NF-κB can modulate Nrf2 transcription and activity, having both positive and negative  
248 effects on the target gene expression<sup>69</sup>.

249 Natural compounds derived from plants, vegetables, fungi and micronutrients (e.g. curcumin, sulforaphane,  
250 resveratrol and vitamin D) or physical exercise can activate Nrf2.<sup>70,71</sup> However, sulforaphane is the most  
251 potent activator of Nrf2.<sup>3,34</sup> "Ancient foods", and particularly those containing *Lactobacillus*, activate  
252 Nrf2.<sup>72</sup>

253 Nrf2 may be involved in diseases associated with insulin-resistance.<sup>60,73-75</sup> Nrf2 activity declines with age,  
254 making the elderly more susceptible to oxidative stress-mediated diseases.<sup>76</sup> Nrf2 is involved in the  
255 protection against lung<sup>77</sup> or endothelial damage.<sup>78</sup> Nrf2 activating compounds downregulate ACE2 mRNA  
256 expression in human liver-derived HepG2 cells.<sup>79</sup> Genes encoding cytokines including IL-6 and many  
257 others specifically identified in the "cytokine storm" have been observed in fatal cases of COVID-19. ACE2  
258 can inhibit NF-κB and activate Nrf2.<sup>80</sup>

### 259 **7-2- Sulforaphane, the most potent Nrf2 natural activator**

260 Isothiocyanates are stress-response chemicals formed from glucosinolates in plants often belonging to the  
261 cruciferous family, and more broadly to the Brassica genus including broccoli, watercress, kale, cabbage,  
262 collard greens, Brussels sprouts, bok choy, mustard greens and cauliflower.<sup>81</sup>The formation of  
263 isothiocyanates from glucosinolates depends on plant-intrinsic factors and extrinsic postharvest factors  
264 such as industrial processing, domestic preparation, mastication, and digestion.<sup>82</sup>

265 Sulforaphane [1-isothiocyanato-4-(methylsulfinyl)butane] is an isothiocyanate occurring in a stored form  
266 such as glucoraphanin in cruciferous vegetables<sup>83,84</sup>. Sulphoraphanes are also found in fermented cabbage  
267<sup>31,85</sup>. Present in the plant as its precursor, glucoraphanin, sulforaphane is formed through the actions of  
268 myrosinase, a  $\beta$ -thioglucosidase present in either the plant tissue or the mammalian microbiome<sup>86,87</sup>.

269 Sulforaphane is a clinically relevant nutraceutical compound used for the prevention and treatment of  
270 chronic diseases and may be involved in ageing.<sup>88</sup> Along with other natural nutrients, sulforaphane has  
271 been suggested to have a therapeutic value for the treatment of coronavirus disease 2019 (COVID-19).<sup>89</sup>

272 One of the key mechanisms of action of sulforaphane involves the activation of the Nrf2-Keap1 signaling  
273 pathway.<sup>90</sup>Sulforaphane is the most effective natural activator of the Nrf2 pathway, and Nrf2 expression  
274 and function is vital for sulforaphane-mediated action.<sup>91,92</sup>Sulforaphanes were suggested to be effective in  
275 diseases associated with insulin resistance<sup>1,93-95</sup>It has been proposed that SARS-CoV-2 downregulates  
276 ACE2 and that there is an increased insulin resistance associated with oxidative stress through the AT<sub>1</sub>R  
277 pathway. Fermented vegetables and Brassica vegetables release glucoraphanin, converted by the plant or by  
278 the gut microbiome into sulforaphane, which activates Nrf2 and subsequently reduces insulin intolerance  
279 (Figure 5B).

### 280 **7-3- Lactic acid bacteria**

#### 281 *Antioxidant activity of Lactobacillus*

282 The gastrointestinal (GI) tract is challenged with oxidative stress induced by a wide array of factors, such  
283 as exogenous pathogenic microorganisms and dietary aspects. Redox signaling plays a critical role in the  
284 physiology and pathophysiology of the GI tract<sup>96</sup>. The redox mechanisms of *Lactobacillus* spp. are  
285 involved in the downregulation of ROS-forming enzymes,<sup>97,98</sup>and redox stress resistance proteins or genes  
286 differ largely between LAB species. In addition, Nrf-2 and NF- $\kappa$ B are two common transcription factors,  
287 through which *Lactobacillus* spp. also modulates oxidative stress.<sup>99</sup>

#### 288 *Do lactobacilli prevent insulin resistance?*

289 Hundreds of studies have attempted to find an efficacy of LAB on insulin resistance-associated diseases.  
290 However, most of them are underpowered or have some methodological flaws. Moreover, not all LAB  
291 strains have the same action on insulin resistance <sup>100</sup> and new better designed studies with the appropriate  
292 LAB are required. A large meta-analysis found that the intake of probiotics resulted in minor but consistent  
293 improvements in several metabolic risk factors in subjects with metabolic diseases, and particularly in  
294 insulin resistance <sup>101</sup>. Another recent meta-analysis found that an oral supplementation with probiotics or  
295 synbiotics has a small effect in reducing waist circumference but no effect on body weight or body mass  
296 index (BMI) <sup>102</sup>. Kefir, a fermented milk product, was not found to be more effective than yoghurt in the  
297 glycemic control of obesity, possibly because there are insufficient differences between both <sup>103</sup>.

#### 298 *Lactobacillus and Nrf2*

299 Nrf2 may be involved in diseases associated with insulinresistance <sup>73-75</sup>. “Ancient foods”, and particularly  
300 those containing *Lactobacillus*, activate Nrf2<sup>72</sup>. The microbiome is highly related to insulin resistance. In  
301 mice, several strains of *Lactobacillus* were found to regulate Nrf2 in models of ageing <sup>104</sup>, in  
302 cardioprotective effects <sup>105</sup>, and in non-alcoholic fatty acid liver disease <sup>106</sup>. *Lactobacillus*  
303 *plantarum* CQPC11 - isolated from Sichuan pickled cabbages - antagonizes oxidation and ageing in mice  
304 <sup>107</sup>. *Lactobacillus* protects against ulcerative colitis by modulation of the gut microbiota and Nrf2/Ho-1  
305 pathway <sup>108</sup>. The sugary kefir strain, *Lactobacillus mali* APS1, ameliorates hepatic steatosis by regulation  
306 of Nrf2 and the gut microbiota in rats <sup>109</sup>. *In vitro* studies have also found an effect of *Lactobacilli*  
307 mediated by Nrf2 <sup>110-112</sup>. Interestingly, the symbiotic combination of prebiotic grape pomace extract and  
308 probiotic *Lactobacillus* sp. reduces intestinal inflammatory markers.<sup>113</sup>

#### 309 *Coronavirus disease in animals and lactic acid bacteria.*

310 The porcine epidemic diarrhea virus (PEDV) and the Transmissible Gastroenteritis Coronavirus Infection  
311 (TGEV) are worldwide-distributed coronaviruses. Low levels of *Lactobacillus* were found in the intestine  
312 of piglets infected by TGEV <sup>114</sup> or PEDV. *Lactobacillus* inhibits PEDV or TGEV effects *in vitro*<sup>115,116</sup>.

#### 313 **7-4-Nrf2 and COVID-19**

314 A putative mechanism may be proposed (Figure 5). SARS-CoV-2 downregulates ACE2 inducing an  
315 increased insulin resistance associated with oxidative stress through the AT<sub>1</sub>R pathway. This may explain  
316 risk factors for severe COVID-19.

317 Fermented vegetables are often made from cruciferous (Brassica) vegetables that release glucoraphanin  
318 converted by the plant or by the gut microbiome into sulforaphane which activates Nrf2 and subsequently

319 reduces insulin intolerance by its potent antioxidant activities. Fermented vegetables contain a high content  
320 of *Lactobacillus* that can activate Nrf2 and impact on the microbiome. <sup>117</sup>Sulforaphane and LAB both  
321 therefore have the ability to reduce insulin resistance.

322 Other putative actions on COVID-19 severity may be postulated. The down-regulation of ACE2 reduces  
323 the Ang-1,7 anti-oxidant activity that was found to activate Nrf2. <sup>118,119</sup> Nrf2 protects against hallmarks of  
324 severe COVID-19. It has anti-fibrotic effects on various organs including the lungs, <sup>120</sup> protects against  
325 lung injury and acute respiratory distress syndrome, <sup>121</sup> and endothelial damage<sup>78</sup> . Finally, Nrf2 can block  
326 IL-6 in different models of inflammation <sup>122</sup> and might play a role in the COVID-19 cytokine storm.

327 These different mechanisms may explain the importance of fermented cabbage in preventing the severity of  
328 COVID-19. It is clear that other nutrients, vitamin D<sup>123</sup> and many different foods act on NRF2 and that  
329 mechanisms other than Nrf2 may be operative.

330 It is not yet known whether sulforaphane and/or LAB may act on the infectivity of SARS-CoV-2. Disulfide  
331 bonds can be formed under oxidizing conditions and play an important role in the folding and stability of  
332 some proteins. The receptor-binding domain of the viral spike proteins and ACE2 have several cysteine  
333 residues. Using molecular dynamics simulations, the binding affinity was significantly impaired when all  
334 of the disulfide bonds of both ACE2 and SARS-CoV/CoV-2 spike proteins were reduced to thiol groups.  
335 This computational finding possibly provides a molecular basis for the differential COVID-19 cellular  
336 recognition due to the oxidative stress.<sup>124</sup>

337 It is likely that foods with anti-oxidant activity can interact with COVID-19 and that fermented or  
338 cruciferous vegetables represent one of the possible foods involved. If some foods are found to be  
339 associated with a prevention of COVID-19 prevalence or severity, it may be of interest to study their LAB  
340 and/or sulforaphane composition in order to eventually find some common mechanisms and targets for  
341 therapy.

## 342 **8- May dietary modifications change the course of COVID-19?**

### 343 **8-1- Fermented vegetables and Kimchi**

344 It is tempting to propose that countries where traditional LAB-fermented vegetables are largely consumed  
345 are those showing lower COVID-19 death rates and that fermented vegetables represent one possible  
346 preventive approach. Other nutrients are found in these products that may enhance their effect (e.g. vitamin  
347 K <sup>125</sup>). Kimchi fermented from many vegetables including cabbage has several effects on insulin resistance  
348 associated diseases: anti-diabetic properties,<sup>126,127</sup> cardiovascular diseases,<sup>28</sup> dyslipidemia <sup>128</sup> or

349 ageing.<sup>129</sup>Kimchi, when fermented for a long time, reduces insulin intolerance to a greater extent than fresh  
350 kimchi,<sup>126</sup> indicating that newly formed products during fermentation are important. In particular, Kimchi  
351 from cabbage and Chinese cabbage contains several glucosinolates<sup>130-132</sup> that can be transformed in  
352 sulforaphanes either in the plant itself or by the human microbiome.<sup>60</sup> In central European countries, raw  
353 and fermented cabbage is commonly consumed.

354 In Sub-Saharan Africa, people commonly eat fermented foods, mainly cereal-based foods like sorghum,  
355 millet and maize, roots such as cassava, fruits and vegetables. Fermented cassava products  
356 (like *gari* and *fufu*) are a major component of the diet of over 800 million people and, in some areas, these  
357 products constitute over 50% of the diet.<sup>16</sup>

358 It is clear that sauerkraut is consumed in Alsace (France) where a COVID-19 outbreak has been identified,  
359 but it is not a regular meal.

## 360 **8-2- Westernized diet**

361 Westernized diets contain a reduced amount of fermented vegetables<sup>43,133</sup> and may be prone to increasing  
362 insulin resistance<sup>44,134</sup> and diseases associated with it,<sup>135</sup> and thereby severe COVID-19.

363 In the Mediterranean diet, well known for reducing insulin resistance,<sup>136</sup> Nrf2 appears to play an important  
364 role.<sup>71,137</sup> The COVID-19 death rate differences in Italian (Figure 2) and Spanish<sup>3</sup> regions suggest a role for  
365 Mediterranean diet and short chain food supply. This also indicates that many foods can have an effect and  
366 that cabbage and fermented foods represent a proof-of-concept. Nrf2 is also involved in the Okinawan-  
367 based diet<sup>71</sup>, active on insulin intolerance.<sup>138</sup> Taken altogether, it is possible that diet is partly involved in  
368 the COVID-19 death clusters found in large Western cities where traditional diet is often replaced by long  
369 chain food supply.

370 It is clear that diet is not the only risk factor and should be considered in the context of COVID-19 in a  
371 given setting. For example, Nordic/central European people socialize less than the Mediterraneans and  
372 simultaneously may consume more fermented vegetables.

## 373 **8.3. The COVID-19 slum paradox**

374 It was expected that the COVID-19 pandemic will be catastrophic if it reached deprived areas of low- and  
375 middle-income countries, in particular informal settlements (slum areas) where social distancing and  
376 lockdown are almost impossible to set up.<sup>139</sup>

377 In the US, highly populated, regional air hub areas, minorities and poverty had an increased risk of  
378 COVID-19 related mortality. <sup>140</sup> It was proposed that the inequality might be due to the workforce of  
379 essential services, poverty, access to care or air pollution <sup>141</sup>. These are common risk factors in mortality  
380 observed in deprived areas of the US. <sup>142</sup> Moreover, in the US and the UK, there are unique health issues  
381 facing black, Asian and minority ethnic communities. <sup>143,144</sup> This greater risk of hospitalizations in these  
382 populations was not explained by socio-economic or behavioural factors. <sup>145</sup> Social distancing is an  
383 important factor to be considered <sup>146</sup> but diet may also be involved.

384 On the other hand, a recent report of the Municipal Corporation of Greater Mumbai (Public Relation  
385 Department, 28-07-2020) found that 57% of subjects tested in the slum area had antibodies against SAR-  
386 CoV-2 but only 16% in the non-slum areas. The fatality rate in slum areas was very low (0.05-0.1%). <sup>147</sup>  
387 Although precise data are lacking, in Brazilian favelas the spread of COVID-19 is not noticed. <sup>148</sup>  
388 Temperature does not seem to be an important factor to contain the pandemic. Fermented foods are popular  
389 throughout the world and in many regions they represent a widespread tradition as well as they make a  
390 significant contribution to the diet of millions of individuals.<sup>16</sup> This is the case in slum areas and it is  
391 possible that fermented foods explain, at least partly, the paradox.

## 392 **Conclusion**

393 Cabbage contains precursors of sulforaphane, the most active natural activator of Nrf2. Fermented  
394 vegetables contain many lactobacilli, also potent Nrf2 activators. It is proposed that fermented cabbage is a  
395 proof-of-concept of dietary manipulations that may enhance Nrf2-associated antioxidant effects helpful in  
396 mitigate COVID-19 severity.

397 Mainstream COVID-19 control strategies including social distancing, confinement and intensive case  
398 finding, testing, tracing and isolating are so far not enough to provide a SARS-CoV-2-free environment and  
399 restore a safe social life. There are hopes for a safe and effective vaccine, but this is unlikely to become  
400 rapidly available. So, there is a need to explore other potentially useful strategies. An area that has not been  
401 sufficiently considered is diet, both as a preventive and/or therapeutically useful intervention, encouraging  
402 people to eat more traditional foods containing fermented vegetables (Figure 6). We have suggested that  
403 fermented vegetables could be associated with a lower COVID-19 mortality due to their potent antioxidant  
404 effect among which sulforaphane and LAB are important. However, many other foods may have a similar  
405 activity. It should be noted that dietary supplements that over-activate Nrf2 may have side-effects.<sup>149</sup>

406

407 Robust evidence from observational studies would be helpful to formally investigate associations between  
408 fermented foods and clinical outcomes in COVID-19. State-of-the-art methods, including the use of DAGs

409 (Directed Acyclic Graphs), may be needed to help assess whether the associations seen are likely to  
410 represent causal relationship <sup>150</sup>. A faster approach would be to develop large clinical trials in the  
411 appropriate populations. Interventions based on diets with a high intake of fermented foods like Kimchi or  
412 other fermented foods are unlikely to present ethical difficulties. Furthermore, the fact that a precise  
413 mechanism has been proposed would facilitate adding reliable biomarkers to the relevant clinical outcomes.  
414 Moreover, new drugs based on the components of these fermented foods may be of interest.

415 If the hypothesis is proved, COVID-19 will be the first infectious disease outbreak associated with a loss of  
416 “nature” <sup>151</sup> and to be ascribed as a disease of the Anthropocene <sup>152</sup>. Imbalance in the gut microbiota is  
417 responsible for the pathogenesis of various disease types including allergy, asthma, rheumatoid arthritis,  
418 different types of cancer, diabetes mellitus, obesity and cardiovascular disease <sup>153</sup>. Fermentation was  
419 introduced during the Neolithic age and was essential for the survival of human kind. When modern life led  
420 to eating reduced amounts of fermented foods, the microbiome drastically changed <sup>154</sup>, allowing SARS-  
421 CoV-2 to spread or to be more severe <sup>155</sup>. It is time for mitigation <sup>156</sup>.

422



423 **ARIA group :**

424

- 425 Amir Hamzah Abdul Latiff, Baharudin Abdullah, Werner Aberer, Nancy Abusada, Ian Adcock, Alejandro Afani,  
426 IoanaAgache, XenofonAggelidis, Jenifer Agustin, Cezmi A Akdis, MübeccelAkdis, Mona Al-Ahmad, Abou Al-Zahab  
427 Bassam, Hussam Alburdan, Oscar Aldrey-Palacios, Emilio Alvarez Cuesta, Hiba Alwan Salman, Ashraf Alzaabi, Salma  
428 Amade, Gene Ambrocio, Rosana Angles, Isabella Annesi-Maesano, Ignacio J Ansotegui, JosepAnto, Paula Ara  
429 Bardajo, Stefania Arasi, Hasan Arshad, Maria Cristina Artesani, Estrella Asayag, Francesca Avolio, KhuzamaAzhari,  
430 Claus Bachert, Diego Bagnasco, Ilaria Baiardini, NisseraBajrović, Petros Bakakos, Sergio BakeyalaMongono, Christine  
431 Balotro-Torres, Sergio Barba, Cristina Barbara, Elsa Barbosa, Bruno Barreto, Joan Bartra, Eric D Bateman,  
432 LkhagvaaBattur, Anna Bedbrook, Martín Bedolla Barajas, Bianca Beghé, Antra Bekere, Elizabeth Bel, Ali Ben Kheder,  
433 Mikael Benson, Emilia Camelia Berghea, Karl-Christian Bergmann, Roberto Bernardini, David Bernstein, Mike Bewick,  
434 Slawomir Bialek, Artur Białoszewski, Thomas Bieber, Nils E Billo, Maria Beatrice Bilo, Carsten Bindslev-Jensen, Leif  
435 Bjermer, Hubert Blain, Irina Bobolea, Malgorzata Bochenska Marciniak, Christine Bond, Attilio Boner, Matteo Bonini,  
436 Sergio Bonini, SinthiaBosnic-Anticevich, Isabelle Bosse, Sofia Botskariova, Jacques Bouchard, Louis-Philippe Boulet,  
437 Rodolphe Bourret, Philippe Bousquet, FulvioBraidó, Andrew Briggs, Christopher E Brightling, Jan Brozek, Luisa  
438 Brussino, Roland Buhl, Roxana Bumbacea, Rosalva Buquicchio, María-Teresa Burguete Cabañas, Andrew Bush,  
439 William W Busse, Jeroen Buters, Fernan Caballero-Fonseca, Moïses A Calderon, Mario Calvo, Paulo Camargos,  
440 Thierry Camuzat, FR Canevari, Antonio Cano, G Walter Canonica, Arnaldo Capriles-Hulett, Luis Caraballo, Vicky  
441 Cardona, Kai-Hakon Carlsen, Jonas Carmon Pirez, Jorge Caro, Warner Carr, Pedro Carreiro-Martins, Fredelita  
442 Carreon-Asuncion, Ana-Maria Carriazo, Thomas Casale, Mary-Ann Castor, Elizabeth Castro, A.G. Caviglia, Lorenzo  
443 Cecchi, Alfonso Cepeda Sarabia, Ramanathan Chandrasekharan, Yoon-Seok Chang, Victoria Chato-Andeza,  
444 LidaChatzi, Christina Chatzidaki, Niels H Chavannes, Claudia Chaves Loureiro, Marta Chelninska, Yuzhi Chen, Lei  
445 Cheng, Sharon Chinthrajah, Tomas Chivato, EkaterineChkhartishvili, George Christoff, Henry Chrystyn, Derek K Chu,  
446 Antonio Chua, Alexander Chuchalin, Kian Fan Chung, Alberto Cicerán, CemalCingi, Giorgio Ciprandi, IevaCirule, Ana  
447 Carla Coelho, Enrico Compalati, JannisConstantinidis, Jaime Correia de Sousa, Elisio Manuel Costa, David Costa,  
448 María del Carmen Costa Domínguez, André Coste, M. Cottini, Linda Cox, Carlos Crisci, Maria AngiolaCrivellaro, Alvaro  
449 A Cruz, John Cullen, Adnan Custovic, BiljanaCvetkovski, WienczyslawaCzarlewski, Gennaro D'Amato, Jane da Silva,  
450 Ronald Dahl, Sven-Erik Dahlen, Vasilis Daniilidis, LoueiDarjaziniNahas, Ulf Darsow, Janet Davies, Frédéric de Blay,  
451 Giulia De Feo, Eloisa De Guia, Chato de los Santos, Esteban De Manuel Keenoy, Govert De Vries, Diana Deleanu,  
452 Pascal Demoly, Judah Denburg, Philippe Devillier, Alain Didier, SanjaDimicJanjic, Maria Dimou, Anh Tuan Dinh-Xuan,  
453 Ratko Djukanovic, Maria Do CeuTexeira, DejanDokic, Margarita Gabriela, Domínguez Silva, Habib Douagui, Nikolaos  
454 Douladiris, Maria Doulaptsi, Gérard Dray, RutaDubakiene, Eve Dupas, Stephen Durham, Marzia Duse, Mark  
455 Dykewicz, Didier Ebo, NatalijaEdelbaher, Thomas Eiwegger, Eklund, Yehia El-Gamal, Zeinab A. El-Sayed, Shereen S.  
456 El-Sayed, Magda El-Seify, Regina Emuzyte, Lourdes Enecilla, Marina Erhola, Heidilita Espinoza, Jesús Guillermo  
457 Espinoza Contreras, John Farrell, Lenora Fernandez, Antje Fink Wagner, Alessandro Fiocchi, Wytske J Fokkens,

- 458 LeniaFolletti, Joao A Fonseca, Jean-François Fontaine, Francesco Forastiere, Jose Miguel Fuentes Pèrez, Emily  
459 Gaerlan–Resureccion, Mina Gaga, José Luis Gálvez Romero, AmiranGamkrelidze, Alexis Garcia, Cecilia Yvonne García  
460 Cobas, María de la Luz Hortensia García Cruz, Jacques Gayraud, Matteo Gelardi, BilunGemicioglu, Dimitra  
461 Gennimata, Sonya Genova, José Gereda, Roy Gerth van Wijk, Antonio Giuliano, Maximiliano Gomez, Sandra  
462 González Diaz, Maia Gotua, Christos Grigoreas, InetaGrisle, Leo Gualteiro, Marta Guidacci, Nick Guldemon, Zdenek  
463 Gutter, Antonieta Guzmán, Tari Haahtela, RamsaHalloum, David Halpin, Eckard Hamelmann, Suleiman Hammadi,  
464 Richard Harvey, Enrico Heffler, Joachim Heinrich, Adnan Hejjaoui, BirtheHellquist-Dahl, Luiana Hernández Velázquez,  
465 Mark Hew, Elham Hossny, Peter Howarth, Martin Hrubíško, YunuenRocío Huerta Villalobos, Marc Humbert, Husain  
466 Salina, Michael Hyland, Guido Iaccarino, Moustafa Ibrahim, Natalia Ilina, Maddalena Illario, Cristoforo Incorvaia,  
467 Antonio Infantino, Carla Irani, ZhanatIspayeva, Juan-Carlos Ivancevich, Edgardo EJ Jares, Deborah Jarvis, Ewa Jassem,  
468 KlemenJenko, Rubén Darío JiménezracruzUscanga, Sebastian L Johnston, Guy Joos, Maja Jošt, KajaJulge, Ki-Suck Jung,  
469 Jocelyne Just, Marek Jutel, Igor Kaidashev, Omer Kalayci, FuatKalyoncu, Jeni Kapsali, PrzemyslawKardas,  
470 JussiKarjalainen, Carmela A. Kasala, Michael Katotomichelakis, Loreta Kavaliukaite, Bennoor S Kazi, Thomas Keil, Paul  
471 Keith, Musa Khaitov, Nikolai Khaltaev, You-Young Kim, Bruce Kirenga, JorgKleine-Tebbe, Ludger Klimek, Bernard  
472 KoffiN’Goran, EvangeliaKompoti, Peter Kopač, Gerard Koppelman, Anja KorenJeverica, Seppo Koskinen, MitjaKošnik,  
473 Kosta V. Kostov, Marek L Kowalski, Tanya Kralimarkova, Karmen Kramer Vrščaj, Helga Kraxner, SamoKreft, Vicky  
474 Kritikos, Dmitry Kudlay, Mikael Kuitunen, Inger Kull, Piotr Kuna, MaciejKupczyk, Violeta Kvedariene,  
475 MarialenaKyriakakou, Nika Lalek, Massimo Landi, Stephen Lane, DésireeLarenas-Linnemann, Susanne Lau, Daniel  
476 Laune, Jorge Lavrut, Lan Le, Martina Lenzenhuber, Marcus Lessa, Michael Levin, Jing Li, Philip Lieberman, Giuseppe  
477 Liotta, Brian Lipworth, Xuandao Liu, Rommel Lobo, Karin C Lodrup Carlsen, Carlo Lombardi, Renaud Louis, Stelios  
478 Loukidis, Olga Lourenço, Jorge A. Luna Pech, BojanMadjar, Enrico Maggi, Antoine Magnan, Bassam Mahboub,  
479 AlpanaMair, Yassin Mais, Anke-Hilse Maitland van der Zee, Mika Makela, Michael Makris, Hans-Jorgen Malling,  
480 Mariana Mandajieva, Patrick Manning, ManolisManousakis, PavlosMaragoudakis, Gianluigi Marseglia, Gailen  
481 Marshall, Mohammad Reza Masjedi, Jorge F. Máspero, Juan José Matta Campos, Marcus Maurer, Sandra Mavale-  
482 Manuel, CemMeço, Erik Melén, Giovanni Melioli, Elisabete Melo-Gomes, Eli O Meltzer, EnricaMenditto, Andrew  
483 Menzies-Gow, Hans Merk, Jean-Pierre Michel, Yann Micheli, Neven Miculinic, Luís Midão, Florin Mihaltan, Nikolaos  
484 Mikos, Manlio Milanese, BranislavaMilenkovic, DimitriosMitsias, Bassem Moalla, Giuliana Moda, María Dolores  
485 Mogica Martínez, Youssef Mohammad, Mostafa Moin, Mathieu Molimard, Isabelle Momas, Monique Mommers,  
486 Alessandro Monaco, Steve Montefort, Dory Mora, Mario Morais-Almeida, Ralph Mösges, BadrEldin Mostafa,  
487 Joaquim Mullol, Lars Münter, Antonella Muraro, Ruth Murray, Antonio Musarra, TihomirMustakov, Robert Naclerio,  
488 Kari C. Nadeau, Rachel Nadif, AllaNakonechna, Leyla Namazova-Baranova, Gretchen Navarro-Locsin, Hugo Neffen,  
489 Kristof Nekam, AngelosNeou, Eustachio Nettis, Daniel Neuberger, Laurent Nicod, Stefania Nicola, Verena  
490 Niederberger-Leppin, Marek Niedoszytko, Antonio Nieto, Ettore Novellino, Elisabete Nunes, DieudonnéNyembue,  
491 Robyn O’Hehir, CvetankaOdjakova, Ken Ohta, Yoshitaka Okamoto, Kimi Okubo, Brian Oliver, Gabrielle L Onorato,  
492 Maria Pia Orru, Solange Ouédraogo, KampadilembaOuoba, Pier Luigi Paggiaro, Aris Pagkalos, Giovanni Pajno, Gianni  
493 Pala, SP Palaniappan, Isabella Pali-Schöll, Susanna Palkonen, Stephen Palmer, Carmen PanaitescuBunu, Petr Panzner,

- 494 Nikos G Papadopoulos, Vasilis Papanikolaou, Alberto Papi, BojidarParalchev, Giannis Paraskevopoulos, Hae-Sim Park,  
495 Giovanni Passalacqua, Vincenzo Patella, Ian Pavord, Ruby Pawankar, Soren Pedersen, SusetePeleve, Simona  
496 Pellegino, Ana Pereira, Tamara Pérez, Andrea Perna, Diego Peroni, Oliver Pfaar, Nhân Pham-Thi, Bernard Pigearias,  
497 Isabelle Pin, Konstantina Piskou, ConstantinosPitsios, Davor Plavec, Dagmar Poethig, Wolfgang Pohl,  
498 AntonijaPoplasSusic, Todor A. Popov, Fabienne Portejoie, Paul Potter, Lars Poulsen, Alexandra Prados-Torres,  
499 FotisPrarros, David Price, Emmanuel Prokopakis, Francesca Puggioni, Elisa Puig-Domenech, Robert Puy, Klaus Rabe,  
500 Filip Raciborski, Josephine Ramos, Marysia T. Recto, Shereen M. Reda, Frederico S Regateiro, Norbert Reider,  
501 SietzeReitsma, Susana Repka-Ramirez, Erminia Ridolo, Janet Rimmer, Daniela Rivero Yeverino, José Angelo Rizzo,  
502 Carlos Robalo-Cordeiro, Graham Roberts, Nicolas Roche, Mónica Rodríguez González, Eréndira Rodríguez Zagal,  
503 Giovanni Rolla, Christine Rolland, Regina Roller-Wirnsberger, Miguel Roman Rodriguez, Antonino Romano, Jan  
504 Romantowski, Philippe Rombaux, Joel Romualdez, Jose Rosado-Pinto, Nelson Rosario, Lanny Rosenwasser, Oliviero  
505 Rossi, Menachem Rottem, Philip Rouadi, NikoletaRovina, Irma RozmanSinur, Mauricio Ruiz, Lucy Tania Ruiz Segura,  
506 Dermot Ryan, Hironori Sagara, Daiki Sakai, Daiju Sakurai, Wafaa Saleh, Johanna Salimaki, Konstantinos Samitas,  
507 Boleslaw Samolinski, María Guadalupe Sánchez Coronel, Mario Sanchez-Borges, Jaime Sanchez-Lopez,  
508 CodrutSarafoleanu, FaradibaSarquisSerpa, Joaquin Sastre-Dominguez, Eleonora Savi, BisherSawaf, Glenis K Scadding,  
509 Sophie Scheire, Peter Schmid-Grendelmeier, Juan Francisco Schuhl, Holger Schunemann, Maria Schvalbová, Jorgen  
510 Schwarze, Nicola Scichilone, Gianenrico Senna, Cecilia Sepúlveda, Elie Serrano, Aziz Sheikh, Mike Shields, Vasil  
511 Shishkov, Nikos Siafakas, Alexander Simeonov, Estelle FER Simons, Juan Carlos Sisul, BrigitaSitkauskiene,  
512 IngelbjorgSkrindo, Tanja SokličKošak, DirceuSolé, TalantSooronbaev, Manuel Soto-Martinez, Manuel Soto-Quiros,  
513 Barnaro Sousa Pinto, Milan Sova, Michael Soyka, Krzysztof Specjalski, Otto Spranger, Sofia Stamataki, Lina Stefanaki,  
514 Cristiana Stellato, Rafael Stelmach, Timo Strandberg, Petra Stute, Abirami Subramaniam, Charlotte SuppliUlrik,  
515 Michael Sutherland, Silvia Sylvestre, AikateriniSyrigou, Luis TabordaBarata, NadejdaTakovska, Rachel Tan, Frances  
516 Tan, Vincent Tan, Ing Ping Tang, Masami Taniguchi, Line Tannert, PongsakornTantilipikorn, Jessica Tattersall, Filippo  
517 Tesi, Carel Thijs, Mike Thomas, Teresa To, Ana Maria Todo-Bom, AlkisTogias, Peter-Valentin Tomazic, Vesna Tomic-  
518 Spiric, SannaToppila-Salmi, Elina Toskala, Massimo Triggiani, Nadja Triller, Katja Triller, IoannaTsiligianni, M. Uberti,  
519 RuxandraUlmeanu, Jure Urbancic, Marilyn Urrutia Pereira, Martina Vachova, Felipe Valdés, Rudolf Valenta, Marylin  
520 Valentin Rostan, Antonio Valero, ArunasValiulis, Mina Vallianatou, ErkkaValovirta, Michiel Van Eerd, Eric Van Ganse,  
521 Marianne van Hage, Olivier Vandenplas, TuulaVasankari, DafinaVassileva, Cesar Velasco Munoz, Maria Teresa  
522 Ventura, Cécilia Vera-Munoz, DilyanaVicheva, PakitVichyanond, Petra Vidgren, Giovanni Viegi, Claus Vogelmeier,  
523 Leena Von Herten, Theodoros Vontetsianos, Dimitris Vourdas, Vu Tran Thien Quan, MartinWagenmann, Samantha  
524 Walker, Dana Wallace, De Yun Wang, Susan Wasserman, Magnus Wickman, Sian Williams, Dennis Williams, Nicola  
525 Wilson, Gary Wong, Kent Woo, John Wright, Piotr Wroczynski, ParaskeviXepapadaki, PlamenYakovliev, Masao  
526 Yamaguchi, Kwok Yan, Yoke Yeow Yap, Barbara Yawn, Panayiotis Yiallourous, ArzuYorgancioglu, Shigemi Yoshihara,  
527 Ian Young, Osman B Yusuf, Asghar Zaidi, Fares Zaitoun, Heather Zar, M.T. Zedda, Mario E Zernotti, Luo Zhang,  
528 Nanshan Zhong, Mihaela Zidarn, TorstenZuberbier, Celia Zubrinich.

529

531 **Conflict of interest** :the authors have no conflict of interest to declare.

533 **References**

1. Stafford N. Covid-19: Why Germany's case fatality rate seems so low. *BMJ* 2020;369:m1395.
2. Bousquet J, Czarlewski W, Blain H, Zuberbier T, Anto J. Rapid Response: Why Germany's case fatality rate seems so low: Is nutrition another possibility. *bmj* 2020;https://www.bmj.com/content/369/bmj.m1395/rr-12.
3. Bousquet J, Anto JM, Iaccarino G, et al. Is diet partly responsible for differences in COVID-19 death rates between and within countries? *Clin Transl Allergy* 2020;10:16.
4. Iddir M, Brito A, Dingeo G, et al. Strengthening the Immune System and Reducing Inflammation and Oxidative Stress through Diet and Nutrition: Considerations during the COVID-19 Crisis. *Nutrients* 2020;12.
5. Cena H, Chieppa M. Coronavirus Disease (COVID-19-SARS-CoV-2) and Nutrition: Is Infection in Italy Suggesting a Connection? *Front Immunol* 2020;11:944.
6. Infusino F, Marazzato M, Mancone M, et al. Diet Supplementation, Probiotics, and Nutraceuticals in SARS-CoV-2 Infection: A Scoping Review. *Nutrients* 2020;12.
7. Adams KK, Baker WL, Sobieraj DM. Myth Busters: Dietary Supplements and COVID-19. *Ann Pharmacother* 2020;54:820-6.
8. Sunyer J, Jarvis D, Pekkanen J, et al. Geographic variations in the effect of atopy on asthma in the European Community Respiratory Health Study. *J Allergy Clin Immunol* 2004;114:1033-9.
9. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science* 2020.
10. Rosenthal PJ, Breman JG, Djimde AA, et al. COVID-19: Shining the Light on Africa. *Am J Trop Med Hyg* 2020.
11. Fonseca S, Rivas I, Romaguera D, et al. Association between consumption of fermented vegetables and COVID-19 mortality at a country level in Europe. *MEDRXIV/2020/147025* 2020.
12. Fonseca S, Rivas I, Romaguera D, et al. Association between consumption of vegetables and COVID-19 mortality at a country level in Europe. *MedRxiv* 2020.
13. Baker P, Friel S. Food systems transformations, ultra-processed food markets and the nutrition transition in Asia. *Global Health* 2016;12:80.

14. Santulli G, Pascale V, Finelli R, et al. We are What We Eat: Impact of Food from Short Supply Chain on Metabolic Syndrome. *J Clin Med* 2019;8.
15. Peters A, Krumbholz P, Jager E, et al. Metabolites of lactic acid bacteria present in fermented foods are highly potent agonists of human hydroxycarboxylic acid receptor 3. *PLoS Genet* 2019;15:e1008145.
16. Azam-Ali S. Fermented fruits and vegetables. A global perspective. 1998.
17. Marco ML, Heeney D, Binda S, et al. Health benefits of fermented foods: microbiota and beyond. *Curr Opin Biotechnol* 2017;44:94-102.
18. Rhee SJ, Lee JE, Lee CH. Importance of lactic acid bacteria in Asian fermented foods. *Microb Cell Fact* 2011;10 Suppl 1:S5.
19. Patra JK, Das G, Paramithiotis S, Shin HS. Kimchi and Other Widely Consumed Traditional Fermented Foods of Korea: A Review. *Front Microbiol* 2016;7:1493.
20. Jung JY, Lee SH, Jeon CO. Kimchi microflora: history, current status, and perspectives for industrial kimchi production. *Appl Microbiol Biotechnol* 2014;98:2385-93.
21. Chen YS, Otaguro M, Lin YH, et al. *Lactococcus formosensis* sp. nov., a lactic acid bacterium isolated from yan-tsai-shin (fermented broccoli stems). *Int J Syst Evol Microbiol* 2014;64:146-51.
22. Han X, Yi H, Zhang L, et al. Improvement of fermented Chinese cabbage characteristics by selected starter cultures. *J Food Sci* 2014;79:M1387-92.
23. Yoon KY, Woodams EE, Hang YD. Production of probiotic cabbage juice by lactic acid bacteria. *Bioresour Technol* 2006;97:1427-30.
24. Slattery C, Cotter PD, O'Toole PW. Analysis of Health Benefits Conferred by *Lactobacillus* Species from Kefir. *Nutrients* 2019;11.
25. Shiby VK, Mishra HN. Fermented milks and milk products as functional foods--a review. *Crit Rev Food Sci Nutr* 2013;53:482-96.
26. Sanders ME, Merenstein DJ, Reid G, Gibson GR, Rastall RA. Probiotics and prebiotics in intestinal health and disease: from biology to the clinic. *Nat Rev Gastroenterol Hepatol* 2019;16:605-16.
27. Sanlier N, Gokcen BB, Sezgin AC. Health benefits of fermented foods. *Crit Rev Food Sci Nutr* 2019;59:506-27.
28. Laveffe L, Marasini D, Carbonero F. Microbial Ecology of Fermented Vegetables and Non-Alcoholic Drinks and Current Knowledge on Their Impact on Human Health. *Adv Food Nutr Res* 2019;87:147-85.
29. Melini F, Melini V, Luziatelli F, Ficca AG, Ruzzi M. Health-Promoting Components in Fermented Foods: An Up-to-Date Systematic Review. *Nutrients* 2019;11.
30. Azam M, Mohsin M, Ijaz H, et al. Review - Lactic acid bacteria in traditional fermented Asian foods. *Pak J Pharm Sci* 2017;30:1803-14.

31. Dimidi E, Cox SR, Rossi M, Whelan K. Fermented Foods: Definitions and Characteristics, Impact on the Gut Microbiota and Effects on Gastrointestinal Health and Disease. *Nutrients* 2019;11.
32. Riggioni C, Comberiati P, Giovannini M, et al. A compendium answering 150 questions on COVID-19 and SARS-CoV-2. *Allergy* 2020.
33. Ruokolainen L, Lehtimäki J, Karkman A, Haahtela T. Holistic view on health : two protective layers of biodiversity. *Ann Zool Fennici* 2017;54:39-49.
34. Septembre-Malaterre A, Remize F, Poucheret P. Fruits and vegetables, as a source of nutritional compounds and phytochemicals: Changes in bioactive compounds during lactic fermentation. *Food Res Int* 2018;104:86-99.
35. Kok CR, Hutkins R. Yogurt and other fermented foods as sources of health-promoting bacteria. *Nutr Rev* 2018;76:4-15.
36. De Filippis F, Pasolli E, Ercolini D. The food-gut axis: lactic acid bacteria and their link to food, the gut microbiome and human health. *FEMS Microbiol Rev* 2020.
37. Mobeen F, Sharma V, Tulika P. Enterotype Variations of the Healthy Human Gut Microbiome in Different Geographical Regions. *Bioinformation* 2018;14:560-73.
38. Bibbo S, Ianiro G, Giorgio V, et al. The role of diet on gut microbiota composition. *Eur Rev Med Pharmacol Sci* 2016;20:4742-9.
39. Tian S, Liu X, Lei P, Zhang X, Shan Y. Microbiota: a mediator to transform glucosinolate precursors in cruciferous vegetables to the active isothiocyanates. *J Sci Food Agric* 2018;98:1255-60.
40. Segata N. Gut Microbiome: Westernization and the Disappearance of Intestinal Diversity. *Curr Biol* 2015;25:R611-3.
41. Vangay P, Johnson AJ, Ward TL, et al. US Immigration Westernizes the Human Gut Microbiome. *Cell* 2018;175:962-72 e10.
42. Zuo T, Kamm MA, Colombel JF, Ng SC. Urbanization and the gut microbiota in health and inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol* 2018;15:440-52.
43. Wilson AS, Koller KR, Ramaboli MC, et al. Diet and the Human Gut Microbiome: An International Review. *Dig Dis Sci* 2020;65:723-40.
44. Yamashita M, Okubo H, Kobuke K, et al. Alteration of gut microbiota by a Westernized lifestyle and its correlation with insulin resistance in non-diabetic Japanese men. *J Diabetes Investig* 2019;10:1463-70.
45. Angelakis E, Yasir M, Bachar D, et al. Gut microbiome and dietary patterns in different Saudi populations and monkeys. *Sci Rep* 2016;6:32191.
46. Mitsou EK, Kakali A, Antonopoulou S, et al. Adherence to the Mediterranean diet is associated with the gut microbiota pattern and gastrointestinal characteristics in an adult population. *Br J Nutr* 2017;117:1645-55.

47. Saad MJ, Santos A, Prada PO. Linking Gut Microbiota and Inflammation to Obesity and Insulin Resistance. *Physiology (Bethesda)* 2016;31:283-93.
48. Chen X, Devaraj S. Gut Microbiome in Obesity, Metabolic Syndrome, and Diabetes. *Curr Diab Rep* 2018;18:129.
49. Lee CJ, Sears CL, Maruthur N. Gut microbiome and its role in obesity and insulin resistance. *Ann N Y Acad Sci* 2020;1461:37-52.
50. Zuo T, Zhang F, Lui GCY, et al. Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization. *Gastroenterology* 2020.
51. Xu K, Cai H, Shen Y, et al. [Management of Corona Virus disease-19 (COVID-19): The Zhejiang Experience]. *Zhejiang Da Xue Xue Bao Yi Xue Ban*2020;49.
52. Finucane FM, Davenport C. Coronavirus and Obesity: Could Insulin Resistance Mediate the Severity of Covid-19 Infection? *Front Public Health* 2020;8:184.
53. Guzik TJ, Cosentino F. Epigenetics and Immunometabolism in Diabetes and Aging. *Antioxid Redox Signal* 2018;29:257-74.
54. Miedema MD, Maziarz M, Biggs ML, et al. Plasma-free fatty acids, fatty acid-binding protein 4, and mortality in older adults (from the Cardiovascular Health Study). *Am J Cardiol* 2014;114:843-8.
55. Hurre S, Hsu WH. The etiology of oxidative stress in insulin resistance. *Biomed J* 2017;40:257-62.
56. Wen H, Gwathmey JK, Xie LH. Oxidative stress-mediated effects of angiotensin II in the cardiovascular system. *World J Hypertens* 2012;2:34-44.
57. Bhatt SR, Lokhandwala MF, Banday AA. Vascular oxidative stress upregulates angiotensin II type I receptors via mechanisms involving nuclear factor kappa B. *Clin Exp Hypertens* 2014;36:367-73.
58. Dalan R, Bornstein SR, El-Armouche A, et al. The ACE-2 in COVID-19: Foe or Friend? *Horm Metab Res* 2020;52:257-63.
59. Sarzani R, Giulietti F, Di Pentima C, Giordano P, Spannella F. Disequilibrium between the Classic Renin-Angiotensin System and Its Opposing Arm in Sars-Cov-2 Related Lung Injury. *Am J Physiol Lung Cell Mol Physiol* 2020.
60. Bousquet J, Anto J, Czarlewski W, et al. Sulforaphane: from death rate heterogeneity in countries to candidate for prevention of severe COVID-19 Allergy 2020;submitted.
61. Ren H, Yang Y, Wang F, et al. Association of the insulin resistance marker TyG index with the severity and mortality of COVID-19. *Cardiovasc Diabetol* 2020;19:58.
62. Jain S, Buttar HS, Chintameneni M, Kaur G. Prevention of Cardiovascular Diseases with Anti-Inflammatory and Anti- Oxidant Nutraceuticals and Herbal Products: An Overview of Pre-Clinical and Clinical Studies. *Recent Pat Inflamm Allergy Drug Discov* 2018;12:145-57.
63. Razmpoosh E, Javadi M, Ejtahed HS, Mirmiran P. Probiotics as beneficial agents in the management of diabetes mellitus: a systematic review. *Diabetes Metab Res Rev* 2016;32:143-68.

64. Serino A, Salazar G. Protective Role of Polyphenols against Vascular Inflammation, Aging and Cardiovascular Disease. *Nutrients* 2018;11.
65. Zabetakis I, Lordan R, Norton C, Tsoupras A. COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation. *Nutrients* 2020;12.
66. Tonelli C, Chio IIC, Tuveson DA. Transcriptional Regulation by Nrf2. *Antioxid Redox Signal* 2018;29:1727-45.
67. Yamamoto M, Kensler TW, Motohashi H. The KEAP1-NRF2 System: a Thiol-Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis. *Physiol Rev* 2018;98:1169-203.
68. !!! INVALID CITATION !!! 45.
69. Wardyn JD, Ponsford AH, Sanderson CM. Dissecting molecular cross-talk between Nrf2 and NF-kappaB response pathways. *Biochem Soc Trans* 2015;43:621-6.
70. Jimenez-Osorio AS, Gonzalez-Reyes S, Pedraza-Chaverri J. Natural Nrf2 activators in diabetes. *Clin Chim Acta* 2015;448:182-92.
71. Pall ML, Levine S. Nrf2, a master regulator of detoxification and also antioxidant, anti-inflammatory and other cytoprotective mechanisms, is raised by health promoting factors. *Sheng Li Xue Bao* 2015;67:1-18.
72. Senger DR, Li D, Jaminet SC, Cao S. Activation of the Nrf2 Cell Defense Pathway by Ancient Foods: Disease Prevention by Important Molecules and Microbes Lost from the Modern Western Diet. *PLoS One* 2016;11:e0148042.
73. Uruno A, Yagishita Y, Yamamoto M. The Keap1-Nrf2 system and diabetes mellitus. *Arch Biochem Biophys* 2015;566:76-84.
74. Vasileva LV, Savova MS, Amirova KM, Dinkova-Kostova AT, Georgiev MI. Obesity and NRF2-mediated cytoprotection: Where is the missing link? *Pharmacol Res* 2020;156:104760.
75. Guo Z, Mo Z. Keap1-Nrf2 signaling pathway in angiogenesis and vascular diseases. *J Tissue Eng Regen Med* 2020;14:869-83.
76. Zhang H, Davies KJA, Forman HJ. Oxidative stress response and Nrf2 signaling in aging. *Free Radic Biol Med* 2015;88:314-36.
77. Rojo de la Vega M, Dodson M, Gross C, et al. Role of Nrf2 and Autophagy in Acute Lung Injury. *Curr Pharmacol Rep* 2016;2:91-101.
78. Chen B, Lu Y, Chen Y, Cheng J. The role of Nrf2 in oxidative stress-induced endothelial injuries. *J Endocrinol* 2015;225:R83-99.
79. McCord JM, Hybertson BM, Cota-Gomez A, Gao B. Nrf2 Activator PB125(R) as a Potential Therapeutic Agent Against COVID-19. *bioRxiv* 2020.



80. Fang Y, Gao F, Liu Z. Angiotensin-converting enzyme 2 attenuates inflammatory response and oxidative stress in hyperoxic lung injury by regulating NF-kappaB and Nrf2 pathways. *QJM* 2019;112:914-24.
81. Palliyaguru DL, Yuan JM, Kensler TW, Fahey JW. Isothiocyanates: Translating the Power of Plants to People. *Mol Nutr Food Res* 2018;62:e1700965.
82. Oliviero T, Verkerk R, Dekker M. Isothiocyanates from Brassica Vegetables-Effects of Processing, Cooking, Mastication, and Digestion. *Mol Nutr Food Res* 2018;62:e1701069.
83. Vanduchova A, Anzenbacher P, Anzenbacherova E. Isothiocyanate from Broccoli, Sulforaphane, and Its Properties. *J Med Food* 2019;22:121-6.
84. Quirante-Moya S, Garcia-Ibanez P, Quirante-Moya F, Villano D, Moreno DA. The Role of Brassica Bioactives on Human Health: Are We Studying It the Right Way? *Molecules* 2020;25.
85. Luang-In V, Deeseenthum S, Udomwong P, Saengha W, Gregori M. Formation of Sulforaphane and Iberin Products from Thai Cabbage Fermented by Myrosinase-Positive Bacteria. *Molecules* 2018;23.
86. Yagishita Y, Fahey JW, Dinkova-Kostova AT, Kensler TW. Broccoli or Sulforaphane: Is It the Source or Dose That Matters? *Molecules* 2019;24.
87. Hindson J. Brassica vegetable metabolism by gut microbiota. *Nat Rev Gastroenterol Hepatol* 2020;17:195.
88. Houghton CA. Sulforaphane: Its "Coming of Age" as a Clinically Relevant Nutraceutical in the Prevention and Treatment of Chronic Disease. *Oxid Med Cell Longev* 2019;2019:2716870.
89. Horowitz RI, Freeman PR. Three novel prevention, diagnostic, and treatment options for COVID-19 urgently necessitating controlled randomized trials. *Med Hypotheses* 2020;143:109851.
90. Yang L, Palliyaguru DL, Kensler TW. Frugal chemoprevention: targeting Nrf2 with foods rich in sulforaphane. *Semin Oncol* 2016;43:146-53.
91. Bai Y, Wang X, Zhao S, Ma C, Cui J, Zheng Y. Sulforaphane Protects against Cardiovascular Disease via Nrf2 Activation. *Oxid Med Cell Longev* 2015;2015:407580.
92. Zhou S, Wang J, Yin X, et al. Nrf2 expression and function, but not MT expression, is indispensable for sulforaphane-mediated protection against intermittent hypoxia-induced cardiomyopathy in mice. *Redox Biol* 2018;19:11-21.
93. Xu L, Nagata N, Ota T. Glucoraphanin: a broccoli sprout extract that ameliorates obesity-induced inflammation and insulin resistance. *Adipocyte* 2018;7:218-25.
94. Teng W, Li Y, Du M, Lei X, Xie S, Ren F. Sulforaphane Prevents Hepatic Insulin Resistance by Blocking Serine Palmitoyltransferase 3-Mediated Ceramide Biosynthesis. *Nutrients* 2019;11.
95. Sun Y, Zhou S, Guo H, et al. Protective effects of sulforaphane on type 2 diabetes-induced cardiomyopathy via AMPK-mediated activation of lipid metabolic pathways and NRF2 function. *Metabolism* 2020;102:154002.

96. Perez S, Talens-Visconti R, Rius-Perez S, Finamor I, Sastre J. Redox signaling in the gastrointestinal tract. *Free Radic Biol Med* 2017;104:75-103.
97. An H, Zhai Z, Yin S, Luo Y, Han B, Hao Y. Coexpression of the superoxide dismutase and the catalase provides remarkable oxidative stress resistance in *Lactobacillus rhamnosus*. *J Agric Food Chem* 2011;59:3851-6.
98. Serata M, Iino T, Yasuda E, Sako T. Roles of thioredoxin and thioredoxin reductase in the resistance to oxidative stress in *Lactobacillus casei*. *Microbiology* 2012;158:953-62.
99. Kong Y, Olejar KJ, On SLW, Chelikani V. The Potential of *Lactobacillus* spp. for Modulating Oxidative Stress in the Gastrointestinal Tract. *Antioxidants (Basel)* 2020;9.
100. Lee E, Jung SR, Lee SY, Lee NK, Paik HD, Lim SI. *Lactobacillus plantarum* Strain Ln4 Attenuates Diet-Induced Obesity, Insulin Resistance, and Changes in Hepatic mRNA Levels Associated with Glucose and Lipid Metabolism. *Nutrients* 2018;10.
101. Koutnikova H, Genser B, Monteiro-Sepulveda M, et al. Impact of bacterial probiotics on obesity, diabetes and non-alcoholic fatty liver disease related variables: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* 2019;9:e017995.
102. Suzumura EA, Bersch-Ferreira AC, Torreglosa CR, et al. Effects of oral supplementation with probiotics or synbiotics in overweight and obese adults: a systematic review and meta-analyses of randomized trials. *Nutr Rev* 2019;77:430-50.
103. Barengolts E, Smith ED, Reutrakul S, Tonucci L, Anothaisintawee T. The Effect of Probiotic Yogurt on Glycemic Control in Type 2 Diabetes or Obesity: A Meta-Analysis of Nine Randomized Controlled Trials. *Nutrients* 2019;11.
104. Li B, Evivie SE, Lu J, et al. *Lactobacillus helveticus* KLDS1.8701 alleviates d-galactose-induced aging by regulating Nrf-2 and gut microbiota in mice. *Food Funct* 2018;9:6586-98.
105. Xu H, Wang J, Cai J, et al. Protective Effect of *Lactobacillus rhamnosus* GG and its Supernatant against Myocardial Dysfunction in Obese Mice Exposed to Intermittent Hypoxia is Associated with the Activation of Nrf2 Pathway. *Int J Biol Sci* 2019;15:2471-83.
106. Zhao Z, Wang C, Zhang L, et al. *Lactobacillus plantarum* NA136 improves the non-alcoholic fatty liver disease by modulating the AMPK/Nrf2 pathway. *Appl Microbiol Biotechnol* 2019;103:5843-50.
107. Qian Y, Zhang J, Zhou X, et al. *Lactobacillus plantarum* CQPC11 Isolated from Sichuan Pickled Cabbages Antagonizes d-galactose-Induced Oxidation and Aging in Mice. *Molecules* 2018;23.
108. El-Baz AM, Khodir AE, Adel El-Sokkary MM, Shata A. The protective effect of *Lactobacillus* versus 5-aminosalicylic acid in ulcerative colitis model by modulation of gut microbiota and Nrf2/Ho-1 pathway. *Life Sci* 2020;256:117927.

109. Chen YT, Lin YC, Lin JS, Yang NS, Chen MJ. Sugary Kefir Strain *Lactobacillus mali* APS1 Ameliorated Hepatic Steatosis by Regulation of SIRT-1/Nrf-2 and Gut Microbiota in Rats. *Mol Nutr Food Res* 2018;62:e1700903.
110. Xu C, Qiao L, Ma L, et al. Biogenic selenium nanoparticles synthesized by *Lactobacillus casei* ATCC 393 alleviate intestinal epithelial barrier dysfunction caused by oxidative stress via Nrf2 signaling-mediated mitochondrial pathway. *Int J Nanomedicine* 2019;14:4491-502.
111. Mu G, Li H, Tuo Y, Gao Y, Zhang Y. Antioxidative effect of *Lactobacillus plantarum* Y44 on 2,2'-azobis(2-methylpropionamide) dihydrochloride (ABAP)-damaged Caco-2 cells. *J Dairy Sci* 2019;102:6863-75.
112. Kobatake E, Nakagawa H, Seki T, Miyazaki T. Protective effects and functional mechanisms of *Lactobacillus gasseri* SBT2055 against oxidative stress. *PLoS One* 2017;12:e0177106.
113. Pistol GC, Marin DE, Dragomir C, Taranu I. Synbiotic combination of prebiotic grape pomace extract and probiotic *Lactobacillus* sp. reduced important intestinal inflammatory markers and in-depth signalling mediators in lipopolysaccharide-treated Caco-2 cells. *Br J Nutr* 2018;1-15.
114. Xia L, Yang Y, Wang J, Jing Y, Yang Q. Impact of TGEV infection on the pig small intestine. *Virol J* 2018;15:102.
115. Kumar R, Seo BJ, Mun MR, et al. Putative probiotic *Lactobacillus* spp. from porcine gastrointestinal tract inhibit transmissible gastroenteritis coronavirus and enteric bacterial pathogens. *Trop Anim Health Prod* 2010;42:1855-60.
116. Zhang X, Li P, Zheng Q, Hou J. *Lactobacillus acidophilus* S-layer protein-mediated inhibition of PEDV-induced apoptosis of Vero cells. *Vet Microbiol* 2019;229:159-67.
117. Hassan SM, Jawad MJ, Ahjel SW, et al. The Nrf2 Activator (DMF) and Covid-19: Is there a Possible Role? *Med Arch* 2020;74:134-8.
118. Romero A, San Hipolito-Luengo A, Villalobos LA, et al. The angiotensin-(1-7)/Mas receptor axis protects from endothelial cell senescence via klotho and Nrf2 activation. *Aging Cell* 2019;18:e12913.
119. Cai SM, Yang RQ, Li Y, et al. Angiotensin-(1-7) Improves Liver Fibrosis by Regulating the NLRP3 Inflammasome via Redox Balance Modulation. *Antioxid Redox Signal* 2016;24:795-812.
120. Liu Q, Gao Y, Ci X. Role of Nrf2 and Its Activators in Respiratory Diseases. *Oxid Med Cell Longev* 2019;2019:7090534.
121. Zhao H, Eguchi S, Alam A, Ma D. The role of nuclear factor-erythroid 2 related factor 2 (Nrf-2) in the protection against lung injury. *Am J Physiol Lung Cell Mol Physiol* 2017;312:L155-L62.
122. Keleku-Lukwete N, Suzuki M, Yamamoto M. An Overview of the Advantages of KEAP1-NRF2 System Activation During Inflammatory Disease Treatment. *Antioxid Redox Signal* 2018;29:1746-55.
123. Mitchell F. Vitamin-D and COVID-19: do deficient risk a poorer outcome? *Lancet Diabetes Endocrinol* 2020;8:570.

124. Hati S, Bhattacharyya S. Impact of Thiol-Disulfide Balance on the Binding of Covid-19 Spike Protein with Angiotensin-Converting Enzyme 2 Receptor. *ACS Omega* 2020;5:16292-8.
125. Tarvainen M, Fabritius M, Yang B. Determination of vitamin K composition of fermented food. *Food Chem* 2019;275:515-22.
126. An SY, Lee MS, Jeon JY, et al. Beneficial effects of fresh and fermented kimchi in prediabetic individuals. *Ann Nutr Metab* 2013;63:111-9.
127. Kim EK, An SY, Lee MS, et al. Fermented kimchi reduces body weight and improves metabolic parameters in overweight and obese patients. *Nutr Res* 2011;31:436-43.
128. Kim SA, Joung H, Shin S. Dietary pattern, dietary total antioxidant capacity, and dyslipidemia in Korean adults. *Nutr J* 2019;18:37.
129. Das G, Paramithiotis S, Sundaram Sivamaruthi B, et al. Traditional fermented foods with anti-aging effect: A concentric review. *Food Res Int* 2020;134:109269.
130. Bousquet J, Anto J, Czarlewski W, et al. Sulforaphane: from death rate heterogeneity in countries to candidate for prevention of severe COVID-19 Allergy 2020;in press.
131. Hong E, Kim GH. GC-MS Analysis of the Extracts from Korean Cabbage (*Brassica campestris* L. ssp. *pekinensis*) and Its Seed. *Prev Nutr Food Sci* 2013;18:218-21.
132. Park CH, Yeo HJ, Park SY, Kim JK, Park SU. Comparative Phytochemical Analyses and Metabolic Profiling of Different Phenotypes of Chinese Cabbage (*Brassica Rapa* ssp. *Pekinensis*). *Foods* 2019;8.
133. Raghuvanshi R, Grayson AG, Schena I, Amanze O, Suwintono K, Quinn RA. Microbial Transformations of Organically Fermented Foods. *Metabolites* 2019;9.
134. O'Dea K. Westernization and non-insulin-dependent diabetes in Australian Aborigines. *Ethn Dis* 1991;1:171-87.
135. Kopp W. How Western Diet And Lifestyle Drive The Pandemic Of Obesity And Civilization Diseases. *Diabetes Metab Syndr Obes* 2019;12:2221-36.
136. Mirabelli M, Chiefari E, Arcidiacono B, et al. Mediterranean Diet Nutrients to Turn the Tide against Insulin Resistance and Related Diseases. *Nutrients* 2020;12.
137. Martucci M, Ostan R, Biondi F, et al. Mediterranean diet and inflammaging within the hormesis paradigm. *Nutr Rev* 2017;75:442-55.
138. Darwiche G, Høglund P, Roth B, et al. An Okinawan-based Nordic diet improves anthropometry, metabolic control, and health-related quality of life in Scandinavian patients with type 2 diabetes: a pilot trial. *Food Nutr Res* 2016;60:32594.
139. Van Belle S, Affun-Adegbulu C, Soors W, et al. COVID-19 and informal settlements: an urgent call to rethink urban governance. *Int J Equity Health* 2020;19:81.

140. Correa-Agudelo E, Mersha T, Hernandez A, Branscum AJ, MacKinnon NJ, Cuadros DF. Identification of Vulnerable Populations and Areas at Higher Risk of COVID-19 Related Mortality in the U.S. medRxiv 2020.
141. Abedi V, Olulana O, Avula V, et al. Racial, Economic and Health Inequality and COVID-19 Infection in the United States. medRxiv 2020.
142. Mode NA, Evans MK, Zonderman AB. Race, Neighborhood Economic Status, Income Inequality and Mortality. PLoS One 2016;11:e0154535.
143. Abuelgasim E, Saw LJ, Shirke M, Zeinah M, Harky A. COVID-19: Unique public health issues facing Black, Asian and minority ethnic communities. Curr Probl Cardiol 2020;45:100621.
144. Lassale C, Gaye B, Hamer M, Gale CR, Batty GD. Ethnic disparities in hospitalisation for COVID-19 in England: The role of socioeconomic factors, mental health, and inflammatory and pro-inflammatory factors in a community-based cohort study. Brain Behav Immun 2020.
145. Raisi-Estabragh Z, McCracken C, Bethell MS, et al. Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank. J Public Health (Oxf) 2020.
146. Rubin D, Huang J, Fisher BT, et al. Association of Social Distancing, Population Density, and Temperature With the Instantaneous Reproduction Number of SARS-CoV-2 in Counties Across the United States. JAMA Netw Open 2020;3:e2016099.
147. SARS-CoV2 seroprevalence study in Mumbai: NTI Asayog-BMC-TIFR study-First round report. Municipal Corporation of greater Mumbai, Public Realtion Department, 28-07-2020 2020;https://www.livemint.com/news/india/mumbai-sero-prevalence-of-57-found-in-slums-and-16-in-residential-societies-11595952896909.html.
148. Pereira RJ, Nascimento G, Gratao LHA, Pimenta RS. The risk of COVID-19 transmission in favelas and slums in Brazil. Public Health 2020;183:42-3.
149. Smith RE. The Effects of Dietary Supplements that Overactivate the Nrf2/ARE System. Curr Med Chem 2020;27:2077-94.
150. Textor J, van der Zander B, Gilthorpe MS, Liskiewicz M, Ellison GT. Robust causal inference using directed acyclic graphs: the R package 'dagitty'. Int J Epidemiol 2016;45:1887-94.
151. Haahtela T, von Hertzen L, Anto JM, et al. Helsinki by nature: The Nature Step to Respiratory Health. Clin Transl Allergy 2019;9:57.
152. O'Callaghan C, Anto J. COVID-19: The Disease of the Anthropocene. Env Res 2020;187:109683.doi: 10.1016/j.envres.2020.. Epub 2020 May 15.
153. Vandana UK, Barlaskar NH, Gulzar ABM, et al. Linking gut microbiota with the human diseases. Bioinformatics 2020;16:196-208.

154. McCall LI, Callewaert C, Zhu Q, et al. Home chemical and microbial transitions across urbanization. Nat Microbiol 2020;5:108-15.
155. Haahtela T, Anto J, Bousquet J. Slow Health Catastrophe of Homo urbanicus – Loss of Resilience. Porto Med J 2020.
156. Haahtela T, Valovirta E, Bousquet J, Makela M, Allergy Programme Steering G. The Finnish Allergy Programme 2008-2018 works. Eur Respir J 2017;49.

**Table 1: Possible risk factors for COVID-19 infection explaining geographical differences**

		Individual level	Country/region level
A	Contact with a SARS-CoV-2 infected individual	++++	Case zero identified ++++ e.g. Lombardy
A	Intensity of social contacts	++	+++
A	Intensity of occupational contacts	+++	++
A	Confinement (level)	+++	+++ e.g. US versus EU Sweden vs Nordic countries
A	Confinement (early measures)	+++	+++ e.g. UK versus EU
A	Climatic conditions (temperature, humidity)	?	++ Hot and humid temperature may reduce infection but epidemic bursts in Brazil, Peru and Ecuador
A	GDP of a country/region	?	+
A	Vitamin D	?	+
B	Diet	?	+ The map of COVID-19 deaths in Europe and the low prevalence in Asia and Africa suggest a role for diet
B	Food	++?	+ Bibliographic analysis suggests a role for some fermented foods. Raw cabbage can be fermented in the intestine. Kefir is largely used in many low-prevalence countries.
B	Long food chain supply	++?	+ In Italy and Spain, there may be an association with long chain supply. This may be relevant since food quality differs.
B	Traditional fermented food (example of food)	++?	++ This may be a relevant issue. In former Eastern European countries, in the Balkans, in Africa and in many Asian countries with low-COVID-19 prevalence, traditional fermented foods are common (in line with short food chain supply)
B	Air pollution	+?	+?
B	Underserved area	++	++
C	Age	+++	
C	Comorbidities (severity of	+++	++

	COVID-19)		
C	Sex	++	
C	Institutionalized person	++	

A: Risk factors at a country level, B: Environment, nutrition, C: individual level

+ to ++++: Proposed relative importance



**Figure 1: COVID-19 deaths per million inhabitants**(from Johns Hopkins Coronavirus Center)

**Figure 2: Regional differences of death rates in Italy** (from Worldometer)

**Figure 3: Regional differences of death rates (May 20)** (from *Office fédéral de la santé publique*, Switzerland, Gouvernement français, Lander Bade Wurtemberg))

**Figure 4: Consumption of head cabbage and COVID-19 death rate at a country level**(from Fonseca et al, 12)


**Figure 5: Putative mechanisms of fermented or Brassica vegetables against COVID-19**

A: Oxidative stress induced by SARS-CoV-2 after its binding to ACE2


B: Preventive effects of cabbage and fermented vegetables through Nrf2

Figure 6 : Putative role of diet in COVID-19

May 20



July 18




Death rate/million






- <50
- 50-100
- 100-200
- 200-400
- 400-800
- 800-1600
- >1600




Switzerland






May 20, 2020	Deaths	Population	Deaths/million
Basel city (Sw)	49	198,000	25
Basel-Lands (Sw)	30	298,000	10
Freiburg (DE)	15	213,000	7
Haut-Rhin (F)	715	765,000	935

Deaths per million



- 1: French speaking canton
- 2: Partly French speaking canton
- 3: Italian speaking canton



**A****B**

# Social distancing, age (population, individual), lockdown, sex, other factors

