



Threatened fish spawning area revealed by specific metabarcoding identification of eggs and larvae in the Beni River, upper Amazon

Guido Miranda-Chumacero, Cédric Mariac, Fabrice Duponchelle, Lilian Painter, Robert Wallace, Gérard Cochonneau, Jorge Molina-Rodriguez, Carmen Garcia-Davila, Jean-François Renno

► To cite this version:

Guido Miranda-Chumacero, Cédric Mariac, Fabrice Duponchelle, Lilian Painter, Robert Wallace, et al.. Threatened fish spawning area revealed by specific metabarcoding identification of eggs and larvae in the Beni River, upper Amazon. *Global Ecology and Conservation*, 2020, 24, pp.e01309. 10.1016/j.gecco.2020.e01309 . hal-03411049

HAL Id: hal-03411049

<https://hal.umontpellier.fr/hal-03411049>

Submitted on 8 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Original Research Article

Threatened fish spawning area revealed by specific metabarcoding identification of eggs and larvae in the Beni River, upper Amazon



Guido Miranda-Chumacero ^{a, b, e, *}, Cédric Mariac ^{c, e}, Fabrice Duponchelle ^{d, e},
Lilian Painter ^a, Robert Wallace ^a, Gérard Cochonneau ^f, Jorge Molina-
Rodriguez ^b, Carmen Garcia-Davila ^{e, g}, Jean-François Renno ^{c, e}

^a Wildlife Conservation Society, Bolivia Program, La Paz, Bolivia

^b Unidad de Limnología, Instituto de Ecología, Universidad Mayor de San Andrés, La Paz, Bolivia

^c DIADE, Univ Montpellier, IRD, Montpellier, France

^d MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France

^e Laboratoire Mixte International - Evolution et Domestication de l'Ictyofaune Amazonienne (LMI - EDIA), IIAP - UAGRM - IRD, UMR DIADE, 911 avenue d'Agropolis, BP 64501, 34394 Montpellier cedex 5, France

^f GET, CNRS, IRD, Université de Toulouse, Toulouse, France

^g IIAP, Instituto de Investigaciones de La Amazonía Peruana, Laboratorio de Biología y Genética Molecular (LBGM), Carretera Iquitos-Nauta km 4.5, Iquitos, Peru

ARTICLE INFO

Article history:

Received 5 June 2020

Received in revised form 3 September 2020

Accepted 2 October 2020

Keywords:

Ichthyoplankton
Prochilodus
Brachyplatystoma
Zungaro
Dams
Arid extraction

ABSTRACT

Identifying fish spawning areas is of great ecological and conservation importance as fishes are suffering increasing threat levels from anthropogenic activities. However, to date very few studies have done so in the Amazon basin. In the Beni River located in the upper Madeira basin, fishers reported that a particular ecotone near the town of Rurrenabaque was a fish reproduction area. To test the importance of this zone as a spawning site, we conducted an ichthyoplankton survey during the month when reproduction is most likely to occur. The specific identification of larvae and eggs was made with a metabarcoding analysis. With this approach 13 different fish species of high importance for regional and local fisheries were identified, including the long-distance migratory gilded catfish (*Brachyplatystoma rousseauxii*), considered as endangered in the upper Madeira. Combining the development time of morula-gastrula egg stages with the integrated current velocity of the river, we identified a spawning area that ranges between the last gravel beaches close to Altamarani community and the San Miguel del Bala community on top of Suse strait, the last foothill of the Andes. This spawning area further extends upstream in the Madidi and Pilon Lajas protected areas and Tacana and Tsimane indigenous communities, when considering eggs in final embryo stage. This portion of the Beni River, at the Andean foothills, is heavily impacted by anthropogenic activities, ranging from unmanaged fisheries to contamination coming from the extraction of fluvial aggregates, upstream gold mining, deforestation associated with a sugar cane mill, an existing downstream dam, and projected upstream dam projects. Some urgent solutions for the conservation of this already impacted area are proposed. Confirming fish spawning zones, although difficult, is crucial to inform the definition of priority areas for conservation and management measures, in particular when these sites host endangered species.

* Corresponding author. Wildlife Conservation Society, Bolivia Program, La Paz, Bolivia.

E-mail address: gmiranda@wcs.org (G. Miranda-Chumacero).

1. Introduction

The Amazon basin, which hosts the largest freshwater fish diversity on earth, is facing increasing threats from anthropogenic activities such as overexploitation, habitat degradation, deforestation, hydropower dam construction, invasive species and pollution (Castello et al., 2013; Castello and Macedo, 2016; Hurd et al., 2016). Climate-induced modifications of Amazonian ecosystems are expected to further exacerbate these threats (Xenopoulos et al., 2005; Freitas et al., 2013; Frederico et al., 2016). Beside their exceptional diversity, fish play a major role in the life of Amazonian people (Barthem and Goulding, 2007) with some of the world's highest fish consumption rates per capita: 30–40 kg year⁻¹ for urban populations and 70–120 kg year⁻¹ for rural populations (Isaac and Almeida, 2011), with maximum of ~200 kg year⁻¹ (Batista et al., 1998).

In the Amazon basin, the life cycles of many fishes, including most commercial species, involve a broad range of migrations for feeding and/or reproductive purposes, from a few kilometers to several thousand kilometers (Barthem and Goulding, 1997; Lucas and Baras, 2001; Carolsfeld et al., 2003). As these larger migrations can span over several international boundaries, managing fisheries and conserving migratory species require a good knowledge of their movements and life history strategies. One of the most important issues is to understand precisely where and when they spawn, so as to take appropriate measures to avoid habitat degradation and overfishing. As reproduction is key to species survival, conserving spawning sites is a priority. Identifying reproductive zones and determining which species use them allows the definition of precise conservation actions based on the specific characteristics of the zone (Castello et al., 2013). Unfortunately, such knowledge is seldom available for Amazonian species.

Many studies have investigated fish reproduction in the Amazon basin using different methodologies, such as the monitoring of reproductive adults (Goulding, 1980, 1981; Vazzoler and Menezes, 1992; Ruffino and Isaac, 1995; Duponchelle et al. 2007; García-Dávila et al., 2015, Córdoba et al., 2013; Bonilla-Castillo et al., 2018), the morphological (Cañas and Pine, 2011; Barthem et al., 2014, 2017; Cella-Ribeiro et al., 2015) or molecular identification of larvae in ichthyoplankton samples (García-Dávila et al., 2015; Maggia et al., 2017), and, more recently, approaches based on citizen science (www.ictio.org). Yet, there is very little information on spawning areas and existing data does not go further than the family level (Oliveira and Ferreira, 2008; Cañas and Pine, 2011).

Recent conservation scenarios have evaluated the vulnerability of the main river basins of the Amazon and their evolution until 2050 under current and planned threats (Jézéquel et al., 2020a). The Beni River basin is currently ranked within the second most vulnerable watershed category in the Amazon, and is predicted to negatively evolve into the most vulnerable category by 2050. Another recent study highlighted the knowledge gap in fish diversity for the upper Beni watershed (Jézéquel et al., 2020b).

In spite of this lack of information about ichthyofauna, two of the most important national protected areas of the tropical Andes are located in this part of the Beni watershed, just above Rurrenabaque: the Madidi National Park and Natural Area of Integrated Management and the Pilon Lajas Biosphere Reserve and Indigenous Territory (Fig. 1). This area also hosts Tacana, T'simane-Mosetene and Uchupiamonas indigenous territories. These protected areas and indigenous territories play a critical role in the conservation of ichthyofauna and Bolivian biodiversity in general (SERNAP & CRTM, 2009; SERNAP & Uchupiamonas, 2009; CIPTA & CIMTA, 2014; Identidad Madidi & SERNAP, 2018). The natural boundary between the Madidi and Pilon Lajas protected areas is the Beni River, which crosses the core of the conservation area that these national parks represent. This portion of the Beni River is the main aquatic corridor linking both protected areas to the rest of the watershed, and hence the main migration route for the fish, but also for the indigenous communities that inhabit them, such as Asuncion del Quiquibey (Pilon Lajas) and San Jose de Uchupiamonas (Madidi). This route also crosses a series of marked ecological transitions including the Chepete, Beu, Bala and Suse mountain ranges (200–300 m a.s.l.) and the valleys in between (Fig. 1). Another important transition is downstream of Rurrenabaque at Altamarani, a Tacana indigenous community, where a marked ecotone between the Andean foothills and the Amazonian alluvial plain is characterized by a change from a stone and gravel substrate upstream towards fine sand downstream (Figs. 1 and 2). Local fishermen have identified this transition zone as a breeding site for many species of fish. Currently, this zone is highly impacted by several anthropogenic activities, including arid extraction, overfishing, and contamination by human and industrial waste, and could be impacted by dam projects in the near future.

The aim of this research was to confirm if this threatened ecotone is important for fish reproduction. A metabarcoding approach was implemented for taxonomic identification of fish species and resulting frequency estimation in plankton samples (Mariac et al., 2018).




Fig. 1. Upstream delimitation of the spawning area from the sampling point (yellow point), estimated from eggs at the morula-gastrula stages (red line), and from eggs at the final embryo stage (orange line). The map also indicates existing and potential threats in the area, as well as the Madidi and Pilon Lajas protected area limits and the location of indigenous territories. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

2. Material and methods

2.1. Study area

The specific study site is located in the Beni basin, near the Altamarani village (67.555° W, 14.333° S, altitude 185 m a.s.l.), between the last Andean foothills and the beginning of the Beni floodplain, and 22.8 km downstream from the nearest town: Rurrenabaque (Fig. 1). The Beni River is one of the largest rivers in Bolivia, at nearly 1000 km in length with a mean discharge of $2050 \text{ m}^3 \text{s}^{-1}$ at the Bala strait (Gautier et al., 2006, 2010). At the sampling site, the Beni River has a mean width of 269 m. The average depth is 7 m, reaching 45 m at the Suse strait (Fig. 1). Sampling was conducted at the lowest point of this sand-gravel ecotone, thereby ensuring that egg and larvae drift from the majority of the watershed upstream was included in the sample (Lechner et al., 2016). The hydrological regime registered at this location shows high amplitude variations related to rainfall (Molina-Carpio et al., 2017). Owing to ongoing human activities that threaten its integrity, this site was prioritized for the collection of ichthyoplankton (eggs and larvae). It was then sampled in February 2017, during the breeding season of most fish species in the upper Madeira (Cañas and Pine, 2011; Van Damme et al., 2011; Barthem et al., 2014; Cella-Ribeiro et al., 2015).

2.2. Sampling

Ichthyoplankton samples were collected (eggs and larvae) using ichthyoplankton nets, with a length of $2.5 \text{ m} \times 0.5 \text{ m}$ in diameter and $250 \mu\text{m}$ mesh. Three nets were used simultaneously, tied to a string to sample at 1, 2 and 3 m depths. The nets were horizontally towed seven times for 15 min each during daylight. The bottom of the three nets were weighted down and were held against the current from a boat running approximately at the same speed as the water flow (Barthem et al., 2014). The larvae and eggs collected were macroscopically separated from the organic matter retained in the nets (by adding alcohol in the water to turn them an easier to identify whitish color) directly in the field, preserved in ethanol 95% and stored at 4°C for later determination of developmental stages and metabarcoding analysis.



Fig. 2. Ecological transition close to the sampling point (67.5555 W, 14.3329 S) between sand beaches (a), and stone beaches (b). Extraction activity 10 km upstream from the sampling point, at the spawning area, showing the machinery (c) used for extracting gravel for construction materials. Photos by Pascal Ardit.

2.3. Estimation of the upstream limit of the spawning area

The upstream limit for spawning was estimated using the relation between the current velocity and the timing of specific embryonic stages. Water flow was estimated using 2015 and 2016 hydrological data from the HYBAM project station at Rurrenabaque. Owing to the construction of the Rurrenabaque – San Buenaventura bridge (Fig. 1), the historical HYBAM sampling station had to be moved to another location, which explains the unavailability of data for 2017 (Timouk, 2017). Hence, the average current velocity was calculated from the 11 days preceding the collection date (February 25, 2017) taken over the years 2015 and 2016. From the average of these 22 velocity measurements (1.74 m/s), the range of values at 95% CI (i.e. average speed $\pm 1.96 \text{ SE}$) was calculated: 1.21 to 2.28 ms^{-1} . These two range values were then used to define the lower and upper distance of a first part of the spawning area using the age, in hours, of the morula-gastrula developmental times. The second part of the spawning area corresponds to the ranges calculated from the development time of the final embryo stage (15h).

For that, it was assumed that eggs can be assimilated to inactive particles, that they drift downstream as soon as they are spawned, and that they do so at the same speed as the water flow in the river (Barros and Rosman, 2018). Eggs, but not larval development stages, were used for this calculation because the length of egg developmental stages is more constant between species (Nakatani et al., 2001; Andrade et al., 2016). As several species were involved, we used the approximate mean development times provided in the literature for some of these taxa (at the genus level at least) and some close relatives (Table 1). Water temperature is generally warmer in controlled conditions (Table 1) than in the river (in our case $\sim 24^\circ\text{C}$), which might also delay the development time, albeit more for developed embryo and larvae than for cleavage stages (morula

and gastrula). For this reason, the development time selected has been rounded up to 2 h for the morula stage, 4 h for the gastrula stage and 15 h for the final embryonic stage (means provided in Table 1).

In order to delimit a spawning area as conservatively (extensively) as possible, the lower limit of the spawning area was approximated using the shortest development time for the morula-gastrula stages (2h) and the slowest current velocity (1.21 m/s). We then defined two upper limits, the first using the longest development time (4h) of the morula-gastrula stages and the second using the development time of the final embryo stage (15h), both with the fastest current velocity (2.28 m/s).

2.4. Sequencing and taxonomic assignation

2.4.1. DNA extractions and NGS libraries preparation

Two DNA bulk extractions were performed, one each for the larvae and egg samples, following the rapid isolation of mammalian DNA procedures (Sambrook et al., 1989). The single COI (cytochrome c oxidase subunit I) probe production, as well as all steps and conditions for enriched libraries by capture preparations (DNA shearing, DNA end repair, adapters ligation, Bst Polymerase treatment, hybridizations and real time PCR) followed published protocols (Mariac et al., 2018). One negative (blank) and two positive controls (mock community samples with known species composition) were included during NGS (Next Generation Sequencing) libraries preparation. Indexed libraries were pooled at an equimolar ratio for the capture enrichment step, followed by paired-end sequencing using MiSeq v2 reagents and 2 × 150 bp. Sequencing was carried out at the CIRAD facilities (Montpellier, France).

2.4.2. Data cleaning and taxonomic assignation

Demultiplexing based on the 6-bp internal index was performed using the PYTHON script DEMULADAPT (<https://github.com/Maillo/demultadapt>). Adapters were removed using CUTADAPT 1.2.1 (Martin, 2011). The NGS sequences have been aligned to a COI database (command lines in S1) with the MALT version 0.3.8 program (Herbig et al., 2016) to generate a file in blast format. The reference database used contains 160,387 Actinopterygii COI sequences extracted from Genebank and Bold (Ratnasingham and Hebert, 2007) on February 21, 2017 (reference database S2). Among the 16,270 species or subspecies it contains, at least 445 are species described in the Amazon region.

Taxonomic assignation of the blast results was performed with MEGAN software version 6.12.3 (Huson et al., 2016) using the paired reads option and the weighted LCA method. Reads mapping COI were assigned to a species if a minimum alignment score value of 150 and 98% identity was reached with a reference sequence in the database. If a read mapped with the same score to more than one species it was assigned to their lowest common ancestor. Additional files containing synonymous and disabled taxa (S3) were included for assignment under MEGAN software. These files were used in order to control mis-assignments related to typographical errors and taxonomic revisions. The percentage of reads assigned, not assigned (reads with a hit but failing to meet the score and identity requirements) and reads with no hits to the COI database are reported (S4). A minimum frequency threshold was defined using the MaxSS method (Manel et al., 2001; Liu et al., 2013) with the two mock control samples. The value of this threshold, set at the maximum sum of specificity and sensitivity, was determined by comparing the frequencies estimated by NGS in mock samples to their known frequencies (established by an individual Sanger sequencing). As previously described in Mariac et al. (2018), species identified in egg and larvae samples whose frequencies were below this threshold were discarded.

3. Results

The number of larvae and eggs collected at the sampling point in February 2017 was 404 and 49, respectively. Among the eggs, 8.7% were in the morula stage (~2 h post-fertilization), 87% in the gastrula stage (~4 h post-fertilization), and 4.3% in the final embryo stage (~15 h post fertilization) (Table 2), which implies that the corresponding species spawned very close to the sampling point. Among the larvae, the highest proportion of individuals were found in early stages of development with 60% in the *early flexion* stage, 37.5% in the *pre-flexion* stage and 2.4% were yolk-sac larva (newly hatched individuals); or equivalent to ~62 h, 26 h and 14 h of total life, respectively. The larvae of more advanced stages were found in a lower proportion: most of them were more than 48 h old, with a variation depending on the species.

Table 1

Development times of ontogenetic stages in some Amazonian taxa. Reference 1: Nogueira et al. (2012); 2: Souza et al. (2016); 3: Nakauth et al. (2016); 4: Andrade et al. (2016).

Taxa	Temperature	Developmental time			Ref
		Morula (M)	Gastrula (G)	Final embryo (FE)	
Zungaro	26.7	2h10	4h40	13h30	1
Prochilodus	26 (FE)-28 (M-G)	1h	4h	22h	2
Brycon	30	1h10	1h50	9h30	3
Pseudoplatystoma	28.2	2h	4h	12h	4
Mean (h)		1.6	3.6	14.25	

3.1. Sequencing results and taxonomic assignation

The total number of raw reads was 1,230,896 from which 1,089,701 (88.5%) were successfully paired and then processed for the taxonomic identification of eggs and larvae. The number of reads per library ranged between 86,040 and 227,368 (see S3 for details) with a mean percentage of reads assigned against the COI reference database of 66.6% (SE = 4.7). Only 176 reads were retrieved in the negative control and none of them were assigned. Taxa with a frequency of less than 0.6% were discarded. This minimum frequency threshold was determined by the maxSSS method using mock control samples (supplementary data S5 and S6). Overall fifteen different taxa were identified, 13 at the species level and 2 at the genus level. These taxa belong to eight genera and two orders (Table 3). Eleven and five fish species were identified in eggs and larvae samples, respectively. The percentage of assigned reads was highest for the genus *Prochilodus*, with 33.1% in the egg sample and 62.35% in the larva sample. *Prochilodus nigricans* is the most represented species in both eggs and larvae with 18.15 and 26.1% respectively (Table 3). *Brachyplatystoma rousseauxii* and *Zungaro zungaro* were identified only in the egg samples and represent 17% and 8% of the assigned reads, respectively. The remaining reads were assigned to the genus *Pseudopimelodus* and to ten other species, including five species of *Prochilodus* in both eggs and larvae stages (Table 3). Of the 15 taxa recorded, 8 were observed only in egg samples (*Brachyplatystoma rousseauxii*, *Zungaro zungaro*, *Potamorhina altamazonica*, *Psectogaster rutiloides*, *Pimelodus blochii*, *Mylossoma albiscopum*, *Prochilodus aff. argenteus*, *Rhinodoras boehlkei*), 3 by both eggs and larvae (*P. nigricans*, *P. aff. lineatus*, *P. aff. rubrotaeniatus*) and 2 only in larvae samples (*P. aff. costatus*, *Pseudopimelodus* sp. LBP2397) (Table 3). It can therefore be assumed that at least the 11 species whose eggs were sampled reproduce in the spawning area closest to the sampling point (red stretch in Fig. 1).

3.2. Estimation of the spawning area

Linking the development time for the most abundant gastrula-morula stages (2h–4h) with the range of current velocity of the Beni river during the sampling period (1.21–2.28 m/s; Table 1), we estimate that the collected eggs were drifting from a zone located between 8.71 km (2h * 1.21 m/s * 3.6) and 32.83 km (4h * 2.28 m/s * 3.6) upstream from the sampling point, illustrated as a red stretch in Fig. 1. The final embryo stages (15h) would be drifting from a zone between 65.3 and 123.1 km above the sampling point through the Chepete and Beu Straits, illustrated as an orange stretch in Fig. 1. These two stretches delimit the whole spawning area (SPA) based on egg samples collected in the present study. This SPA integrates parts of the Madidi and Pilon Lajas protected areas and indigenous territories of the Tacana, T'simane Mosetene and Uchupiamonas people. Regarding the larvae, most were more than 48 h old, which also suggests spawning areas further upstream inside the Madidi and Pilon Lajas protected areas and indigenous territories.

4. Discussion

Our study is based on a single sampling date and locality, which constrains the extent of its conclusions. It is impossible, for instance, to state how many species exactly spawn in the identified spawning area, or to precisely determine the extent of the spawning area. We cannot assess either the relative importance of this spawning area for the different species, as they were sampled in very different proportions, probably because of the sampling scheme. Answering these questions would require a broader temporal and spatial sampling coverage. But in spite of these inherent limitations, our study nevertheless provides clear evidence that our sampling area is used by several fish species, including species of major commercial and cultural importance, as a spawning ground, and that it is under immediate threat by anthropogenic activities. Our results also allow, for the first time in the Bolivian Amazon, and one of the first in the Amazon basin, the delimitation of a spawning area for at least 15 taxa around Rurrenabaque in the Beni River, including 13 species, with 11 sampled at the egg stage. It is likely indeed, that further sampling at different times of the hydrological cycle would increase the number of species using the identified spawning area. Contrary to the few studies that have identified reproduction areas in the Amazon basin (e.g. Oliveira and Ferreira, 2008; Cañas and Pine, 2011), the delimitation of the spawning area in the present study was based on the specific molecular identification of eggs in early development stages and not of larvae, which allows for a more precise estimation of the spawning area. Approximations relative to current velocity and water temperature have to be made in both cases, as eggs and even more so larvae that can have swimming abilities, are not homogenously distributed in the water column or in the width of the rivers (Oliveira and Ferreira, 2008; Barthem et al., 2014; Cella-Ribeiro et al., 2015), and current velocity

Table 2

Stages and reference sizes of all eggs registered at the spawning area in February 2017.

Stage	Life Stage (h) ^a	N	(%)	Diameter (mm)
Morula	2	5	(8.7%)	1.24
Gastrula	4	41	(87.0%)	1.75
Final embryo	15	3	(4.3%)	2.16
Total		49	100	1.73

^a Adapted from Andrade et al. (2016)..

Table 3

Taxa identified in egg and larvae samples at the spawning area, and their percentage. Only the taxa with a percentage above the threshold value of 0.6% are reported.

Order	Family	Genus	Species	Eggs	Larvae	
Characiformes	Curimatidae	Prochilodus	<i>Potamorhina altamazonica</i>	6.39	0.00	
			<i>Psectrogaster rutiloides</i>	6.02	0.00	
	Prochilodontidae		<i>Prochilodus</i> aff. <i>argenteus</i>	33.15	62.35	
			<i>Prochilodus</i> aff. <i>costatus</i>	1.16	0.00	
			<i>Prochilodus</i> aff. <i>lineatus</i>	0.00	0.88	
			<i>Prochilodus</i> aff. <i>rubrotaeniatus</i>	0.92	2.95	
			<i>Prochilodus</i> aff. <i>nigricans</i>	0.62	0.76	
			<i>Prochilodus</i> <i>nigricans</i>	18.15	26.10	
	Serrasalmidae		<i>Mylossoma albiscopum</i>	1.39	0.00	
			<i>Rhinodoras boehlkei</i>	0.68	0.00	
			<i>Brachyplatystoma rousseauxii</i>	17.61	0.00	
Siluriformes	Doradidae	Pseudopimelodidae	<i>Pimelodus</i> <i>blochii</i>	2.94	0.00	
	Pimelodidae		<i>Zungaro zungaro</i>	8.81	0.00	
	<i>Pseudopimelodus</i> sp. LBP2397		0.00	4.41		
	<i>Pseudopimelodus</i> sp. LBP2397		0.00	0.70		
	Taxa below threshold			2.16	1.86	
Total (%)				100	100	

strongly varies in the river section. Current velocity can also vary among stretches of the river, especially when the slope or the width of the riverbed varies. However, early egg developments are counted in a few hours, which limits potential approximation errors to a few kilometers, as opposed to tens of km with larvae that develop over a few days (Nakatani et al., 2001; Nogueira et al., 2012; Andrade et al., 2016; Nakauth et al., 2016; Souza et al., 2016).

The identified spawning area extends between Altamarani below Rurrenabaque and well over 100 km upstream. Previous studies had suggested spawning area for migratory catfishes between 200 and 400 masl (Van Damme et al., 2011; Barthem et al., 2017). Although the spawning area identified in the present study is extensive, our results also indicate that at least 11 species reproduce in an area close to Rurrenabaque, between the Altamarani and San Miguel indigenous Tacana communities (~40 km) (the red stretch in Fig. 1). This relatively restricted portion of the spawning area is exposed to high anthropogenic threats. Indeed, the portion of the Beni River just downstream from Rurrenabaque is used intensively to extract gravel and stone for local infrastructure like roads and bridges (Fig. 1). This activity was already identified as a threat to the migration of the small and culturally important *Trichomycterus barbouri* catfishes, which migrate along the water edge (Miranda-Chumacero et al., 2015). It is also likely to alter the nature of the substrate, as well as physico-chemical conditions, in this particular ecotone, threatening its suitability as a spawning ground for fishes. Additionally, waste from a sugar mill 2 km from the sampling site, gold mining extraction and deforestation upstream of the spawning area, represent supplementary water pollution risks.

Another major threat is the accelerated construction of hydroelectric dams in the Amazon basin (Finer and Jenkins, 2012; Lees et al., 2016; Winemiller et al., 2016; Latrubblesse et al., 2017; Anderson et al., 2018). The dams planned between the protected areas of the Madidi and Pilon Lajas at the Bala, Chepete and Beu straits, would result in important alterations in the volume and flow of sediments, homogenizing downstream habitats (Forsberg et al., 2017), including the newly identified spawning area. There are also plans to build other dams in Bolivian territory, such as that on the border with Brazil in Nova Mamoré-Guayaramirim. Downstream, two large hydroelectric dams, Santo Antonio and Jirau, were recently (2011–2012) completed on the Brazilian portion of the river, between Porto Velho and the Bolivian border (Fearnside, 2014). Although located well downstream from the spawning area, these two major hydroelectric dams strongly disrupt the migration patterns of fish species, in particular upstream movements towards reproduction zones, including for several Pimelodidae like goliath catfishes (*Brachyplatystoma* spp: Duponchelle et al., 2016; Hauser, 2018; Hauser et al., 2019).

One particularly interesting fact about the newly identified spawning area is that it also hosts the reproduction of *Brachyplatystoma rousseauxii*, one of the most emblematic and commercially important fishes for the food security and economic sustainability of riverine human populations in the Amazon basin (Barthem and Goulding, 2007), and the Madeira in particular (Doria et al., 2018; Santos et al., 2018; Lima et al., 2020). As a migratory top predator (Barthem and Goulding, 1997), *B. rousseauxii* plays a crucial role for the stability of food webs (Angelini et al., 2006; Estes et al., 2011; Bauer and Hoyer, 2014). This species also performs the longest migration known in fresh water: a round trip of up to 12,000 km (7500 km for this specific spawning area) between its breeding areas in the Andean foothills and its nursery in the Amazon estuary (Barthem and Goulding, 1997, 2017; Duponchelle et al., 2016). Moreover, this exceptional migration involves a natal homing behavior (Batista and Alves-Gomes, 2006), in which most fish return to spawn in the specific geographic area where they were hatched in the upper Amazon (Hauser et al., 2020), including the upper Madeira (Duponchelle et al., 2016). Recent analyzes of otolith microchemistry have shown that the construction of Santo Antônio and Jirau dams on the Brazilian portion of the Madeira have profoundly altered its migration pattern, preventing pre-adults to home back to their breeding areas in the upper Madeira and reducing the downstream drift of larvae to the estuary (Hauser, 2018; Hauser et al. 2018). The populations of *B.*

rousseauxii are now totally isolated in the upper Madeira basin: all the individuals caught in the upper Madeira after dam construction were hatched and bred within the Upper Madeira, without ever reaching their nursery area in the lower Amazon-estuary, a behavior rarely observed before the dams (Duponchelle et al., 2016; Hauser, 2018). This situation has already led to a drastic post-dam reduction in the catches of *B. rousseauxii* in Puerto Villarroel, located in another major river basin of the Bolivian Amazon (Van Damme et al., 2019). This species is the most vulnerable *Brachyplatystoma* species, classified as Vulnerable (VU) (Van Damme et al., 2009), and with a proposal to categorize it as Critically Endangered (CR) in the Madeira basin (Van Damme, in prep).

Interestingly, we identified 6 different genetic entities of *Prochilodus* in the geographic area, with a genetic differentiation at a specific level according to the identification criterion in our methodology: *P. nigricans* already known in the Madeira basin (Castro and Vari, 2004), but also *P. aff. rubrotaeniatus*, *P. aff. argenteus*, *P. aff. costatus* and *P. aff. lineatus*, as well as a sixth entity *Prochilodus* (without valide assignation at the species level). Only *P. nigricans* and *P. rubrotaeniatus* are supposed to be present in the Amazon basin (Castro and Vari, 2004; Melo et al., 2016, 2018). These results suggest cryptic diversity in the Upper Madeira with more naturally occurring species. Although these cryptic taxa are most likely not *P. argenteus*, *P. lineatus* and *P. costatus*, which belong to other river basin of South America (Castro and Vari, 2004; Melo et al., 2016, 2018), these newly identified taxa are nevertheless genetically close to these species, hence our use of *affinis* in their names (Table 3). This genetic proximity is not surprising given the very low divergence in the COI barcode among lineages of the genus *Prochilodus* (Melo et al., 2018). The existence of several putative species of *Prochilodus* in Bolivia was reported by Loubens et al. (1991), in a review in which 6 species of *Prochilodus* were identified in Bolivia, including a new species: *P. labeo*. However, this diversity was neglected by a posterior revision of the Prochilodontidae in which only one fish from Bolivia was analyzed (Castro and Vari, 2004). Nevertheless, the existence of greater species diversity in the genus *Prochilodus*, including possible endemism in the Beni watershed, is also supported by the local knowledge of the T'simane indigenous people (Daillant, 1999, Sarmiento J., pers. comm.). These results emphasize the need for molecular studies on *Prochilodus* in the upper Madeira to shed light on these taxonomic issues.

Among the identified species, many are of major importance for local and regional fisheries (Van Damme et al., 2011; Doria et al., 2018), e.g. *Brachyplatystoma rousseauxii*, *Prochilodus nigricans* and *Zungaro zungaro*. These species are also critical for the livelihoods of the indigenous Tacana communities, whose catches are dominated (85%) by migratory catfishes (CIPTA & WCS, 2010). These species also constitute a large part of the annual income of Tacana communities. In these communities ~65% of the fish are sold whereas the other 35% are for communal consumption (Miranda-Chumacero et al., 2011). Alterations in the reproductive cycles of these species could affect the food security and sovereignty of hundreds of indigenous families living in this spawning area. Furthermore, a large part of the fish traded in the principal Bolivian cities come from the Beni watershed, and therefore underpin the livelihoods of hundreds of commercial fisher people in Rurrenabaque and San Buenaventura (Carvajal-Vallejos et al., 2011).

Preserving the integrity and function of the endangered spawning area around Rurrenabaque is paramount for several reasons: it is used by many important species for local and regional fisheries; it is used as a corridor by migratory species, including the culturally important *T. barburi* (Miranda-Chumacero et al., 2015); it is located within two regional protected areas and indigenous territories; and it is the first precisely identified spawning area of *B. rousseauxii* in the Amazon basin, and likely one of the last remaining spawning areas for the already strongly diminished and isolated population of *B. rousseauxii* in the upper Madeira (Van Damme et al., 2019). Indeed, the breeding areas above Puerto Maldonado in the Peruvian portion of the Madre de Dios (Cañas and Pine, 2011) are now severely compromised by the illegal mining activities that have drastically altered the river's geomorphological and chemical properties (Diringer et al., 2015; Martinez et al., 2018).

Fish life cycles face several natural and anthropogenic threats (dams, pollution, fishing) and the destruction of their spawning sites magnifies their vulnerability. To guarantee the permanence of the species that use this spawning area, it will be important to: 1) maintain the connectivity within and between the still free-flowing Beni, Madre de Dios and Mamoré sub-basins, avoiding the construction of dams with large reservoirs in the lowlands, such as the binational dam planned at Nova Mamore-Guayaramerim, or the Bala, Chepete, Beu dams projected upstream of Rurrenabaque (Anderson et al., 2018), 2) declare a general fishing ban in the upper Madeira for the most threatened fish species, e.g. the goliath catfishes (Ruffino et al., 2018), to allow the remaining individuals to recruit (Hauser et al., 2020), and accompany this with a fishing monitoring program in the Rurrenabaque area, 3) regulate and zone the extraction of gravel and stone excluding it from the breeding area to avoid alterations to the hydro-geomorphological characteristics that migratory species seek for their reproduction, 4) conduct studies to determine the impact of agro-industrial waste and gold extraction on the Beni River, 5) initiate a system for monitoring the density of larvae and their identification using NGS metabarcoding, 6) establish information campaigns with a citizen science approach, like the Ciencia Ciudadana para la Amazonia project (<https://www.amazoniacienciaciudadana.org>) for local residents and fishermen to raise awareness of the importance of this spawning area for local fisheries and, finally, 7) strengthen territorial management in both protected areas and indigenous lands by increasing their role in fish and aquatic biodiversity protection, highlighting the importance of rivers and their connectivity for conservation.

Data accessibility

DNA sequences in.fastq file format and MEGAN project file (rma) used in this study are available from NCBI's Sequence Read Archive (SRA). Bioproject number PRJNA507878.

Biosample accession numbers SAMN10511183-SAMN10511185.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Gabriel Tarifa, Camila Ramallo, Debora Alvestegui and Eddy Ocampo for their help with field and laboratory work. This study was financed by IRD, the Gordon and Betty Moore Foundation and the Wildlife Conservation Society. We are extremely grateful to CIPTA and the Altamarani community for allowing the research to be carried out within their ancestral lands. The authors acknowledge the IRD itrop HPC (South Green Platform) at IRD Montpellier for providing HPC resources that contributed to the research results reported within this paper (URL: <https://bioinfo.ird.fr/>- <http://www.southgreen.fr>).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.gecco.2020.e01309>.

References

- Anderson, E.P., Jenkins, C.N., Heilpern, S., Maldonado-Ocampo, J.A., Carvajal-Vallejos, F.M., Encalada, A.C., Rivadeneira, J.F., Hidalgo, M., Cañas, C.M., Ortega, H., Salcedo, N., Maldonado, M., Tedesco, P.A., 2018. Fragmentation of Andes-to-Amazon connectivity by hydropower dams, 2018 Sci. Adv. 4, eaao1642.
- Andrade, F.F., Lima, A.F., Assumpção, L., Makrakis, S., Kasai, R.I.D., Makrakis, M.C., 2016. Characterization of the early development of *Pseudoplatystoma reticulatum* eigenmann & eigenmann, 1889 (siluriformes: Pimelodidae) from the Paraguay river basin. Neotrop. Ichthyol. 14, e150032.
- Angelini, R., Fabré, N.N., Lopes da Silva-Jr, U., 2006. Trophic analysis and fishing simulation of the biggest Amazonian catfish. Afr. J. Agric. Res. 1, 151–158.
- Barthem, R., Goulding, M., 1997. *The Catfish Connection: Ecology, Migration, and Conservation of Amazon Predators*. Columbia University Press, 155 pp.
- Barros, M.d.C., Rosman, P.C.C., 2018. A study on fish eggs and larvae drifting in the Jirau reservoir, Brazilian Amazon. J. Braz. Soc. Mech. Sci. Eng. 40, 62. <https://doi.org/10.1007/s40430-017-0951-1>.
- Barthem, R., da Costa, M.C., Cassemiro, F., Leite, R.G., da Silva Jr., 2014. Diversity and abundance of fish larvae drifting in the Madeira river, Amazon basin: sampling methods comparison. In: Grillo, O. (Ed.), *Biodiversity - the Dynamic Balance of the Planet*, pp. 137–158. InTech, Available from: <http://www.intechopen.com/books/biodiversity-the-dynamic-balance-of-the-planet/diversity-and-abundance-of-fish-larvae-drifting-in-the-madeira-river-amazon-basin-sampling-methods-c>.
- Barthem, R., Goulding, M., 2007. Un ecosistema inesperado: la Amazonía revelada por la pesca. Lima, Peru: Museu Paraense Emilio Goeldi, Amazon Conservation Association (ACA).
- Barthem, R.B., Goulding, M., Leite, R.G., Cañas, C., Forsberg, B., Venticinque, E., Petry, P., Ribeiro, M., Chuctaya, J., Mercado, A., 2017. Goliath catfish spawning in the far western Amazon confirmed by the distribution of mature adults, drifting larvae and migrating juveniles. Sci. Rep. 7, 41784.
- Batista, J.S., Alves-Gomes, J.A., 2006. Phylogeography of *Brachyplatystoma rousseauxii* (Siluriformes - Pimelodidae) in the Amazon Basin offers preliminary evidence for the first case of "homing" for an Amazonian migratory catfish. Genet. Mol. Res. 5, 723–740.
- Batista, V.S., Inhamuns, A.J., Freitas, C.E.C., Freire-Brasil, D., 1998. Characterization of the fishery in river communities in the low-Solimões/high-Amazon region. Fish. Manag. Ecol. 5, 419–435.
- Bauer, S., Hoye, B.J., 2014. Migratory animals couple biodiversity and ecosystem functioning worldwide. Science 344, 1242552.
- Bonilla-Castillo, C.A., Agudelo, E.C., Gómez, G., Duponchelle, F., 2018. Population dynamics of *Prochilodus nigricans* (characiformes: Prochilodontidae) in the putumayo river. Neotrop. Ichthyol. 16, e170139.
- Cañas, C.M., Pine, W.E., 2011. Documentation of the temporal and spatial patterns of Pimelodidae catfish spawning and larvae dispersion in the Madre de Dios River (Peru): insights for conservation in the Andean-Amazon headwaters. River Res. Appl. 27, 602–611.
- Carolsfeld, J., Harvey, B., Ross, C., Baer, A., 2003. *Migratory Fishes of South America: Biology, Fisheries and Conservation Status*. World Bank, Washington, USA: IDRC.
- Carvajal-Vallejos, F., Van Damme, P.A., Muñoz, H., 2011. Composición de las capturas comerciales y de subsistencia en la amazonía boliviana. In: Van Damme, P., Carvajal-Vallejos, F. (Eds.), *Los peces y delfines de la amazonía boliviana*, pp. 209–234 (Cochabamba, Bolivia: Editorial INIA).
- Castello, L., Macedo, M.N., 2016. Large-scale degradation of Amazonian freshwater ecosystems. Global Change Biol. 22, 990–1007.
- Castello, L., McGrath, D.G., Hess, L.L., Coe, M.T., Lefebvre, P.A., Petry, P., Macedo, M.N., Renó, V.F., Arantes, C.C., 2013. The vulnerability of Amazon freshwater ecosystems. Conserv. Lett. 6, 217–229.
- Castro, R.M.C., Vari, R.P., 2004. Detritivores of the South American Fish Family Prochilodontidae (Teleostei: Ostariophysi: Characiformes): a Phylogenetic and Revisionary Study. Washington DC. Smithsonian Contributions to Zoology No. 622, Smithsonian Institute.
- Cella-Ribeiro, A., Assakawa, L.F., Torrente-Vilara, G., Zuanon, J., Leite, R.G., Doria, C., Duponchelle, F., 2015. Temporal and spatial distribution of young *Brachyplatystoma* spp. (Siluriformes: Pimelodidae) along the rapids stretch of the Madeira River (Brazil) before the construction of two hydroelectric dams. J. Fish. Biol. 86, 1429–1437.
- CIPTA & CIMTA, 2014. Plan de Gestión Territorial Indígena del Pueblo Tacana. Kema Ejudhes'a Jakuastas'ati S'aiddha Enime 2015–2025. Consejo Indígena del Pueblo Tacana (CIPTA), Consejo Indígena de Mujeres Tacanas (CIMTA) y Wildlife Conservation Society (WCS). Bolivia, La Paz.
- CIPTA & WCS, 2010. La pesca en el territorio Takana. WCS & CIPTA, La Paz, Bolivia, p. 69.
- Córdoba, E.A., León, Á.V.J., Bonilla-Castillo, C.A., Petreche Junior, M., Peláez, M., Duponchelle, F., 2013. Breeding, growth and exploitation of *Brachyplatystoma rousseauxii* Castelnau, 1855 in the Caquetá River, Colombia. Neotrop. Ichthyol. 11 (3), 637–647. <https://doi.org/10.1590/S1679-62252013000300017>.
- Daillant, I., 1999. Bajar y arribar: etno-ictiología de las migraciones de peces. Rev. Bol. Ecol y Cons. Amb. 6, 87–94.
- Diringer, S.E., Feingold, B.J., Ortiz, E.J., Gallis, J.A., Araujo-Flores, J.M., Berký, A., Pan, W.K.Y., Hsu-Kim, H., 2015. River transport of mercury from artisanal and small-scale gold mining and risks for dietary mercury exposure in Madre de Dios, Peru. Environ. Sci. Process Impacts 17, 478–487.
- Doria, C.R.C., Duponchelle, F., Lima, M.A.L., Garcia, A., Carvajal-Vallejos, F.M., Méndez, C.C., Catarino, M.F., de Carvalho Freitas, C.E., Vega, B., Miranda-Chumacero, G., Van Damme, P.A., 2018. Review of fisheries resource use and status in the Madeira River basin (Brazil, Bolivia, and Peru) before hydroelectric dam completion. Rev. Fish. Sci. Aquac. 26, 494–514.
- Duponchelle, F., Lino, F., Hubert, N., Panfili, J., Renno, J.-F., Baras, E., Torrico, J., Dugue, R., Nuñez, J., 2007. Environment-related life history trait variations of the red-bellied piranha, *Pygocentrus nattereri*, in two river basins of the Bolivian Amazon. J. Fish. Biol. 71, 1113–1134.
- Duponchelle, F., Pouilly, M., Péchéyran, C., Hauser, M., Renno, J.-F., Panfili, J., Darnaude, A.M., García-Vásquez, A., Carvajal-Vallejos, F., García-Dávila, C., Doria, C., Bérial, S., Donard, A., Sondag, F., Santos, R.V., Núñez, J., Point, D., Labonne, M., Baras, E., 2016. Trans-Amazonian natal homing in giant catfish. J. Appl. Ecol. 53, 1511–1520.
- Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Wardle, D.A., 2011. Trophic downgrading of planet Earth. Science 333, 301–306.
- Fearnside, P.M., 2014. Impacts of Brazil's Madeira River dams: unlearned lessons for hydroelectric development in Amazonia. Environ. Sci. Pol. 38, 164–172.

- Finer, M., Jenkins, C.N., 2012. Proliferation of hydroelectric dams in the Andean Amazon and implications for Andes-Amazon connectivity. *PLoS One* 7, e35126.
- Forsberg, B.R., Melack, J.M., Dunne, T., Barthem, R.B., Goulding, M., Paiva, R.C.D., Weisser, S., 2017. The potential impact of new Andean dams on Amazon fluvial ecosystems. *PLoS One* 12 e0182254.
- Frederico, R.G., Olden, J.D., Zuanon, J., 2016. Climate change sensitivity of threatened, and largely unprotected, Amazonian fishes. *Aquat. Conserv.* 26, 91–102.
- Freitas, C.E.C., Siqueira-Souza, F.K., Humston, R., Hurd, L.E., 2013. An initial assessment of drought sensitivity in Amazonian fish communities. *Hydrobiologia* 705, 159–171.
- García-Dávila, C., Castro-Ruiz, D., Renno, J.F., Chota-Macuyama, W., Carvajal-Vallejos, F.M., Sanchez, H., Angulo, C., Nolorbe, C., Alvarado, J., Estivals, G., Núñez-Rodríguez, J., Duponchelle, F., 2015. Using barcoding of larvae for investigating the breeding seasons of pimelodid catfishes from the Marañon, Napo and Ucayali rivers in the Peruvian Amazon. *J. Appl. Ichthyol.* 31, 40–51.
- Gautier, E., Brunstein, D., Vauchel, P., Roulet, M., Fuertes, O., Guyot, J., Darozzes, J., Bourrel, L., 2006. Temporal relations between meander deformation, water discharge and sediment fluxes in the floodplain of the Rio Beni (Bolivian Amazonia). *Earth Surf. Process. Landforms* 32, 230–248.
- Gautier, E., Brunstein, D., Vauchel, P., Jouanneau, J.-M., Roulet, M., Garcia, C., Guyot, J.L., Castro, M., 2010. Channel and floodplain sediment dynamics in a reach of the tropical meandering Rio Beni (Bolivian Amazonia). *Earth Surf. Process. Landforms* 35, 1838–1853.
- Goulding, M., 1980. The Fishes and the Forest, Explorations in Amazonian Natural History. University of California Press, Berkeley.
- Goulding, M., 1981. Man and Fisheries on an Amazon Frontier. Boston – London: Junk Publishers.
- Hauser, M., 2018. Migração dos grandes bagres Amazônicos pela perspectiva dos isotópos de Estrônio em otólitos. PhD Thesis. Programa de Pós Graduação da rede de Biodiversidade e Biotecnologia da Amazônia legal Porto Velho. Universidade Federal do Rondônia (UNIR), Brazil, 159 pp.
- Hauser, M., Doria, C.R.C., Melo, L.R.C., Santos, A.R., Ayala, D.M., Nogueira, L.D., Amadio, S., Fabré, N., Torrente-Vilara, G., García Vasquez, A., Renno, J.F., Carvajal-Vallejos, F.M., Alonso, J.C., Núñez, J., Duponchelle, F., 2018. Age and growth of the Amazonian migratory catfish *Brachyplatystoma rousseauxii* in the Madeira River basin before the construction of dams. *Neotrop. Ichthyol.* 16, e170130.
- Hauser, M., Doria, C.R.C., Santos, R.V., García Vasquez, A., Pouilly, M., Péchéyran, C., Ponzevera, E., Torrente-Vilara, G., Bérai, S., Panfili, J., Darnaude, A., Renno, J.-F., García-Dávila, C., Núñez, J., Ferraton, F., Vargas, G., Duponchelle, F., 2019. Shedding light on the migratory patterns of the Amazonian goliath catfish, *Brachyplatystoma platynemum*, using otolith 87Sr:86Sr analyses. *Aquat. Conserv.* 29, 397–408.
- Hauser, M., Duponchelle, F., Hermann, T.W., Limburg, K.E., Castello, L., Stewart, D.J., Torrente-Vilara, G., García Vasquez, A., García-Dávila, C., Pouilly, M., Péchéyran, C., Ponzevera, E., Renno, J.-F., Moret, A.S., Doria, C.R.C., 2020. Unmasking continental natal homing in goliath catfish from the upper Amazon. *Freshw. Biol.* 65, 325–336.
- Herbig, A., Maixner, F., Bos, K.I., Zink, A., Krause, J., Huson, D.H., 2016. MALT: Fast alignment and analysis of metagenomic DNA sequence data applied to the Tyrolean Iceman. *2089*, 73.
- Hurd, L.E., Sousa, R.G.C., Siqueira-Souza, F.K., Cooper, G.J., Kahn, J.R., Freitas, C.E.C., 2016. Amazon floodplain fish communities: habitat connectivity and conservation in a rapidly deteriorating environment. *Biol. Conserv.* 195, 118–127.
- Huson, D.H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.-J., Tappu, R., 2016. MEGAN community edition – interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS Comput. Biol.* 12, e1004957.
- Identidad Madidi & Sernap, 2018. Informe de la expedición científica Identidad Madidi 2017, relevamientos de biodiversidad en el Parque Nacional y Área Natural de Manejo Integrado Madidi, Bolivia, La Paz, 227 pp.
- Isaac, V.J., Almeida, M.C., 2011. El consumo de pescado en la Amazonía brasileña. COPESCAALC Documento Ocasional. No. 13. FAO, Kingdom, WWF, IUCN and the Earth Institute at Columbia University. Retrieved from Rome.
- Jézéquel, C., Tedesco, P.A., Bigorne, R., Maldonado-Ocampo, J.A., Ortega, H., Hidalgo, M., Martens, K., Torrente-Vilara, G., Zuanon, J., Acosta, A., Agudelo, E., Barrera Maure, S., Bastos, D.A., Bogotá Gregory, J., Cabeceira, F.G., Canto, A.L.C., Carvajal-Vallejos, F.M., Carvalho, L.N., Cella-Ribeiro, A., Covain, R., Donascimiento, C., Doria, C.R.C., Duarte, C., Ferreira, E.J.G., Galuch, A.V., Giarrizzo, T., Leitão, R.P., Lundberg, J.G., Maldonado, M., Mojica, J.I., Montag, L.F.A., Ohara, W.M., Pires, T.H.S., Pouilly, M., Prada-Pedreros, S., de Queiroz, L.J., Rapp Py-Daniel, L., Ribeiro, F.R.V., Ríos Herrera, R., Sarmiento, J., Sousa, L.M., Stegmann, L.F., Valdiviezo-Rivera, J., Villa, F., Yunoki, T., Oberdorff, T., 2020 a. A database of freshwater fish species of the Amazon Basin. *Sci. Data* 7, 96. <https://doi.org/10.1038/s41597-020-0436-4>.
- Jézéquel, C., Tedesco, P.A., Darwall, W., Dias, M.S., Frederico, R.G., Hidalgo, M., Hugueny, B., Maldonado-Ocampo, J., Martens, K., Ortega, H., Torrente-Vilara, G., Zuanon, J., Oberdorff, T., 2020. Freshwater fish diversity hotspots for conservation priorities in the Amazon Basin. *Conserv. Biol.* 34 (4), 956–965. <https://doi.org/10.1111/cobi.13466>.
- Latrubesse, E.M., Arima, E.Y., Dunne, T., Park, E., Baker, V.R., d'Horta, F.M., Wight, C., Wittmann, F., Zuanon, J., Baker, P.A., Ribas, C.C., Norgaard, R.B., Filizola, N., Ansar, A., Flyvbjerg, B., Stevax, J.C., 2017. Damming the rivers of the Amazon basin. *Nature* 546, 363–369.
- Lechner, A., Keckes, H., Humphries, P., 2016. Patterns and processes in the drift of early developmental stages of fish in rivers: a review. *Rev. Fish Biol. Fish.* 26, 471–489.
- Lees, A.C., Peres, C.A., Fearnside, P.M., Schneider, M., Zuanon, J.A.S., 2016. Hydropower and the future of Amazonian biodiversity. *Biodivers. Conserv.* 25, 451–466.
- Lima, M.A.L., Carvalho, A.R., Nunes, M.A., Angelini, R., Doria, C.R.C., 2020. Declining fisheries and increasing prices: the economic cost of tropical rivers impoundment. *Fish. Res.* 221, 105399–9.
- Liu, C., White, M., Newell, G., 2013. Selecting thresholds for the prediction of species occurrence with presence-only data. *J. Biogeogr.* 40, 778–789.
- Loubens, G., Lauzanne, L., Géry, J., 1991. Contribution à la systématique des Prochilodus boliviens (Pisces, Characiformes, Prochilodontidae). *Rev. Hydrobiol. Trop.* 24, 217–239.
- Lucas, M.C., Baras, E., 2001. Migration of Freshwater Fishes. Blackwell Science Ltd., Oxford.
- Maggia, M.E., Vigouroux, Y., Renno, J.F., Duponchelle, F., Desmarais, E., Núñez, J., García-Dávila, C., Carvajal-Vallejos, F.M., Paradis, E., Martin, J.F., Mariac, C., 2017. DNA metabarcoding of Amazonian ichthyoplankton swarms. *PLoS One* 12, e0170009–e0170014.
- Manel, S., Williams, H.C., Ormerod, S.J., 2001. Evaluating presence-absence models in ecology: the need to account for prevalence. *J. Appl. Ecol.* 38, 921–931.
- Mariac, C., Vigouroux, Y., Duponchelle, F., García-Dávila, C., Núñez, J., Desmarais, E., Renno, J.F., 2018. Metabarcoding by capture using a single COI probe (MCSP) to identify and quantify fish species in ichthyoplankton swarms. *PLoS One* 13 e0202976.
- Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17, 10–12.
- Martinez, G., McCord, S., Driscoll, C., Todorova, S., Wu, S., Araújo, J., Vega, C., Fernandez, L., 2018. Mercury contamination in riverine sediments and fish associated with artisanal and small-scale gold mining in Madre de Dios, Peru. *Int. J. Environ. Res. Publ. Health* 15, 1584.
- Melo, B.F., Dorini, B.F., Foresti, F., Oliveira, C., 2018. Little divergence among mitochondrial lineages of Prochilodus (teleostei, characiformes). *Front. Genet.* 9, 107.
- Melo, B.F., Sidlauskas, B.L., Hoekzema, K., Frable, B.W., Vari, R.P., Oliveira, C., 2016. Molecular phylogenetics of the neotropical fish family Prochilodontidae (teleostei: characiformes). *Mol. Phylogenet. Evol.* 102, 189–201.
- Miranda-Chumacero, G., Álvarez, G., Luna, V., Wallace, R.B., Painter, L., 2015. First observations on annual massive upstream migration of juvenile catfish *Trichomycterus* in an Amazonian River. *Environ. Biol. Fish.* 98, 1913–1926.
- Miranda-Chumacero, Guido, Terrazas, Aizar, Wallace, Robert, 2011. Importancia económica de la ictiofauna para comunidades indígenas takanas del río Beni. In: Van Damme, Paul, Cavajala-Vallejos, Fernando (Eds.), Los peces y delfines de la amazonía boliviana, first ed., pp. 235–245 (Cochabamba, Bolivia: Editorial INIA).
- Molina-Carpio, J., Espinoza, J.C., Vauchel, P., Ronchail, J., Caloir, BG, Guyot, JL, Noriega, L., Caloir, BG, 2017. Hydroclimatology of the Upper Madeira River basin: spatio-temporal variability and trends. *Hydrol. Sci. J. – J. Sci. Hydrol.* 62 (6), 911–927. <https://doi.org/10.1080/02626667.2016.1267861>.

- Nakatani, K., Agostinho, A.A., Baumgartner, G., Bialestzki, A., Sanches, P.V., Makrakis, M.C., Pavanello, C.S., 2001. Ovos e Larvas De Água Doce: Desenvolvimento e Manual de Identificação. [Freshwater Eggs and Larvae: Development and Identification Manual]. Maringá: EDUEM, 378pp.
- Nakauth, A.C.S.S., Villacorta-Correa, M.A., Figueiredo, M.R., Bernardino, G., França, J.M., 2016. Embryonic and larval development of *Brycon amazonicus* (spix & agassiz, 1829). *Braz. J. Biol.* 76, 109–116.
- Nogueira, L.B., Azevedo, P.G., Canelhas, M.R., Bedore, A.G., Lopes, J.M., Godinho, H.P., 2012. Induced spawning and early ontogeny in hatchery-reared catfish *Zungaro jahu* (Siluriformes: Pimelodidae). *Neotrop. Ichthyol.* 10, 89–98.
- Oliveira, E.C.d., Ferreira, E.J.G., 2008. Spawning areas, dispersion and microhabitats of fish larvae in the anavilhanas ecological station, rio negro, amazonas state. Brazil. *Neotrop. Ichthyol.* 6, 559–566.
- Ratnasingham, S., Hebert, P.D.N., 2007. BOLD: the barcode of life data system. *Mol. Ecol. Notes* 7, 355–364. <http://www.barcodinglife.org>.
- Ruffino, M., Isaac, V., 1995. Life cycle and biological parameters of several Brazilian Amazon fish species. NAGA. The ICLARM Quarterly 18, 41–45.
- Ruffino, L.M., Baigún, C., Vitule, J., Cañas, C., Miranda, G., MacNaughton, A., Doria, C., Hauser, M., Cordova, L., Echeverría, A., Hahn, L., Hallwas, G., Ortúñoz, O., Van Damme, P., 2018. Acciones urgentes para la conservación del dorado (*Brachyplatystoma rousseauxii*) en la cuenca del río Madera. Policy Brief, 7 pp. Cochabamba.
- Sambrook, J., Fritsch, E.F., Maniatis, T., 1989. Molecular cloning: a laboratory manual. <https://www.cabdirect.org/cabdirect/abstract/19901616061>.
- Santos, R.E., Pinto-Coelho, R.M., Fonseca, R., Simões, N.R., Zanchi, F.B., 2018. The decline of fisheries on the Madeira River, Brazil: the high cost of the hydroelectric dams in the Amazon Basin. *Fish. Manag. Ecol.* 25, 380–391.
- Sernap, Crtm, 2009. Plan de manejo y plan de vida de la Reserva de la Biosfera y Tierra Comunitaria de Origen Pilón Lajas 2007 – 2017.
- Sernap, Uchupiamonas, 2009. Plan de Desarrollo y Gestión Territorial Indígena de San José de Uchupiamonas. Bolivia, La Paz.
- Souza, G., Melo, E.J.T., Caramaschi, E.P., Andrade, D.R., Monteiro, L.R., 2016. Early development and allometric growth patterns of the grumatá (*Prochilodus vimboides* Kner, 1859). *Zygote* 24, 428–441.
- Timouk, F., 2017. Compte rendu des tournées hydrométriques du SNO HYBAM en Bolivie. De Mars 2013 à Avril 2017. HYBAM, IRD, SENAMHI, p. 22.
- Van Damme, P., Carvajal-Vallejos, F., Sarmiento, J., Barrera, S., Osinaga, K., Miranda-Chumacero, G., 2009. Peces, in: MMAYA. MMAyA, La Paz, pp. 29–42. Libro Rojo de Vertebrados de Bolivia.
- Van Damme, P.A., Carvajal-Vallejos, F., Camacho, J., Muñoz, H., Coronel, J.S., 2011. Peces migratorios de la Amazonía boliviana. In: Van Damme, P.A., Carvajal-Vallejos, F., Molina Carpio, J. (Eds.), Los peces y delfines de la Amazonía Boliviana - Hábitats, potencialidades y amenazas, pp. 149–200 (Cochabamba, Bolivia: Editorial INIA).
- Van Damme, P.A., Córdoba-Clavijo, L., Baigún, C., Hauser, M., Doria, C.R.d.C., Duponchelle, F., 2019. Upstream dam impacts on gilded catfish *Brachyplatystoma rousseauxii* (Siluriformes: Pimelodidae) in the Bolivian Amazon. *Neotrop. Ichthyol.* 17, 1119–9.
- Vazzoler, A.E.d.M., Menezes, N.A., 1992. Síntese de conhecimentos sobre o comportamento reprodutivo dos Characiformes da América do Sul (Teleostei, Ostariophysi). *Braz. J. Biol.* 52, 627–640.
- Winemiller, K.O., McIntyre, P.B., Castello, L., Fluet-Chouinard, E., Giarrizzo, T., Nam, S., Sáenz, L., 2016. Balancing hydropower and biodiversity in the Amazon, Congo, and mekong. *Science* 351, 128–129.
- Xenopoulos, M.A., Lodge, D.M., Alcamo, J., Märker, M., Schulze, K., Van Vuuren, D.P., 2005. Scenarios of freshwater fish extinctions from climate change and water withdrawal. *Global Change Biol.* 11, 1557–1564.