

Unirationality of Certain Universal Families of Cubic Fourfolds

Michele Bolognesi, Hanine Awada

▶ To cite this version:

Michele Bolognesi, Hanine Awada. Unirationality of Certain Universal Families of Cubic Fourfolds. Rationality of Varieties, 342, Springer International Publishing, pp.97-112, 2021, Progress in Mathematics, $10.1007/978-3-030-75421-1_4$. hal-03413180

HAL Id: hal-03413180 https://hal.umontpellier.fr/hal-03413180

Submitted on 15 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNIRATIONALITY OF CERTAIN UNIVERSAL FAMILIES OF CUBIC FOURFOLDS.

HANINE AWADA AND MICHELE BOLOGNESI

ABSTRACT. The aim of this short note is to define the universal cubic fourfold over certain loci of their moduli space. Then, we propose two methods to prove that it is unirational over the Hassett divisors C_d , in the range $8 \le d \le 42$. By applying inductively this argument, we are able to show that, in the same range of values, $C_{d,n}$ is unirational for all integer values of n. Finally, we observe that for explicit infinitely many values of d, the universal cubic fourfold over C_d can not be unirational.

1. INTRODUCTION

Since the celebrated result of Clemens and Griffiths [CG72] on the irrationality of smooth cubic threefolds, at least three generations of algebraic geometers have been working hard on the problem of rationality of cubic fourfolds. The search for the good invariant that would detect (ir)rationality has involved Hodge theoretical [Has16], homological [Kuz10], K-theoretic [GS14] and motivic [BP20] methods. While the general suspicion is that the generic cubic fourfold should be non rational as well, no explicit example of irrational cubic fourfold is known at the moment. A little more is known, and the expectation are slightly more precise. about rationality. Let $\mathcal C$ denote the moduli space of smooth cubic fourfolds and $\mathcal{C}_d \subset \mathcal{C}$ the Hassett divisor of discriminant d, then a well known conjecture due to Harris-Hassett-Kuznetsov (see [Kuz10, Has16, AT14]) states that there should be a countable infinity of Hassett divisors that parametrize rational cubic fourfolds. In particular these divisors should correspond to cubic fourfolds that have an associated K3 surface, in a Hodge theoretical or derived categorical sense (see [Kuz10, Has00]). In the second decade of this century, the search for rational examples of cubic fourfolds has concentrated more on cubics containing some special surfaces [ABBVA14, Nue17, AHTVA19, Awa19]. This has been developed further in the works of Russo-Staglianó, and at the moment the HHK conjecture is known to be true for four low values of d [BD85, BRS19, RS19b, RS19a]. In all these cases, the special cubic fourfolds involved contain rational surfaces.

In this note we are also interested in the birational geometry of cubic fourfolds, but under a slightly different point of view. Universal families are well known and studied objects in the realm of moduli spaces of curves, where forgetful maps are the algebraic geometer's everyday tools. Also in the context of K3 surfaces several results about these objects have appeared in the literature, also very recently [FV18, FV19, Ma19]. Probably because the birational geometry of a single cubic fourfold is already so difficult to understand, the problem of studying the same problems for universal families of cubics seems to have been overtaken. Thanks to the recent advances in the study of rationality of a single cubic fourfold, it seems

H. AWADA AND M. BOLOGNESI

however natural to ask what can be said in families, in particular over certain relevant divisors C_d in the moduli space. In this paper we prove the following fact.

Theorem 1.1. The universal cubic fourfolds $C_{d,1}$ are unirational in the range $8 \le d \le 42$.

More precisely, we propose two different methods, since each one seems to have its own interest, concerning different aspects of the geometry of cubic fourfolds. On one hand, we use some recent results of Farkas-Verra about universal K3 surfaces and use the relation with cubic fourfolds given by associated K3s. On the other hand, we use the presentation of Hassett divisors as cubics containing certain rational surfaces, as done by Nuer and Russo-Staglianó, and Kollar's theorem on unirationality of cubics. The second argument, applied inductively, also allows us to prove the following fact.

Theorem 1.2. The universal families $C_{d,n}$ are universal for all values of n if $8 \le d \le 42$.

Of course, though the intuitive definition is clear, this requires a few checks that actually the universal family exists, which we make in Sect. 2. In the same Section, we also collect some important properties of cubic fourfolds, the rational scrolls they contain, their associated K3 surfaces and their Fano varieties of lines, that are then necessary in Section 3. Section 3 is devoted to the proof of unirationality of $C_{26,1}$ and $C_{42,1}$ via universal K3s, which is based on the construction of a couple of rational/unirational moduli quotients involving the families of scrolls inside our cubic fourfolds.

In Sect. 4 we develop the second argument, which is slightly more general, and allows us to complete the missing cases of Thm. 1.1. We use thoroughly the recent theory of rational surfaces contained in special cubic fourfolds of low discriminants, and combine this with Kollar's Theorem on unirationality of cubic hypersurfaces.

In Section 5 we underline the special nature of the cases we are considering, by comparing them with preceding results of Gritsenko-Hulek-Sankaran [GHS13] and Várilly-Alvarado-Tanimoto [TVA19].

In fact, in [GHS13] it is proven that the moduli space of polarized K3 surfaces of degree 2m has positive Kodaira dimension or is even of general type for an infinity of values (see Prop. 5.2 for details).

Furthermore one expects that the gaps in Prop. 5.2 can be filled in by using automorphic form techniques. Várilly-Alvarado and Tanimoto [TVA19] have started pursuing this project obtaining another relevant range of values where the Kodaira dimension of C_d is positive (see Prop. 5.3).

These results give us *negative* information about the divisors C_d where the universal cubic $C_{d,1}$ can not be unirational.

Proposition 1.3. The universal cubic fourfold $C_{d,1}$ is not unirational if:

- (1) d > 80, $d \equiv 2 \pmod{6}$, 4 //d and such that for any odd prime p, p/d implies $p \equiv 1 \pmod{3}$;
- (2) d = 6n + 2 for n > 18 and $n \neq 20, 21, 25$;
- (3) d = 6n + 2, n > 13 and $n \neq 15$;
- (4) d = 6n, and n > 18, $n \neq 20, 22, 23, 25, 30, 32$;
- (5) d = 6n for n > 16 and $n \neq 18, 20, 22, 30$.

We wish to thank Francesco Russo, Giovanni Staglianó and Sandro Verra for lots of discussions on related topics in the last few months. Thanks to Shouhei Ma and Zhiwei Zheng for giving us precise references and information.

2. The existence of the universal cubic fourfold, and some PROPERTIES OF SCROLLS AND ASSOCIATED K3 SURFACES.

As it is customary when discussing moduli spaces, one of the first questions one considers is the actual existence of a universal family. The invariant theory of cubic fourfolds, with respect to the natural PGL(6)-action, is known by the work of Laza [Laz09, Laz10].

Theorem 2.1. [Laz09, Thm. 1.1] A cubic fourfold $X \subset \mathbb{P}^5$ is not GIT stable if and only if one of the following conditions holds:

- (1) X is singular along a curve C spanning a linear subspace of \mathbb{P}^5 of dimension at most 3:
- (2) X contains a singularity that deforms to a singularity of class P_8, X_9 or J_{10} .

In particular, if X is a cubic fourfold with isolated singularities, then X is stable if and only if X has at worst simple singularities.

The GIT quotient that we are considering is slightly different than the one constructed in [Laz09], since we only consider smooth cubic fourfolds. That is, we consider the complement $U \subset |\mathcal{O}_{\mathbb{P}^5}(3)|$ of the discriminant and take the quotient $\mathcal{C} := U//PGL(6)$. The resulting quotient is a quasi-projective variety of dimension 20 and Thm. 2.1 assures that all point are stable in the GIT sense. We are in particular concerned by the divisors \mathcal{C}_d inside \mathcal{C} , with $8 \leq d \leq 42$. By the seminal work of Hassett [Has00], and further developments [Nue17, RS19b, RS19a, FV18, FV19], they can be characterized in terms of *special* surfaces - not homologous to linear sections - contained in the generic cubic fourfold in each divisor. Recall in fact that for the general cubic fourfold $H^{2,2}(X,\mathbb{Z}) = \mathbb{Z}$, generated by a linear section, and the locus where the rank of this group is bigger than one is the union of a countable infinity of divisors in \mathcal{C} . For example, we will need the following descriptions.

Definition 2.2. We have:

- \$\mathcal{C}_{26} := {Cubic fourfolds containing a 3 nodal septic scroll}
 \$\mathcal{C}_{42} := {Cubic fourfolds containing a 8 nodal degree 9 scroll}.

Moreover, cubic fourfolds X inside these two divisors have an associated K3 surface¹. As it is well known, this means that the *Kuznetsov component* of the bounded derived category of X is equivalent to the derived category of a K3 surface, or equivalently that the Hodge structure of the nonspecial cohomology of the cubic fourfold is essentially the Hodge structure of the primitive cohomology of a K3 surface. We refrain to give more details about this, since these notions are now well known and the references in the literature are very good [Kuz10, Has00, Has16]. The associated K3 surfaces for cubics from C_{26} have genus 14 and degree 26, whereas for cubics from C_{42} they have genus 22 and degree 42.

More generally, thanks in particular to the work of Nuer [Nue17] (see also [RS19b, Sect. 4]), it is known that if $d \leq 42$, then the cubics in each C_d can be defined as

¹Actually cubics in \mathcal{C}_{42} have two associated K3s, we will see more details about this later

the cubics containing certain rational surfaces. These rational surfaces, that are introduced in [Nue17, Sect. 3], are obtained as images of \mathbb{P}^2 via linear systems with prescribed fixed locus. For each value of d we have a different linear system. We will see more about this in Sect. 4.

Let \mathcal{F}_g denote the moduli space of genus g K3 surfaces. For $X \in \mathcal{C}_d$ with discriminant $d = 2(n^2 + n + 1)$, with $n \geq 2$ (remark that 26 and 42 all verify this equality), Hasset shows [Has00, Sect. 6] that there is an isomorphism

between the Fano variety of lines F(X) and the Hilbert scheme $S^{[2]}$ of couple of points of (S, H) a polarized K3 surface with $H^2 = d$, the K3 associated to X. The moduli spaces of the corrisponding K3 surfaces have dimension 19, the same dimension as divisors in C, and this assignment induces a rational map

(2.4)
$$\mathcal{F}_{\frac{d+2}{2}} \dashrightarrow \mathcal{C}_d$$

that is birational for $d \equiv 2 \pmod{6}$ and degree two for $d \equiv 0 \pmod{6}$. In particular, for C_{26} and C_{42} , we have rational maps

$$\begin{array}{ccc} \mathcal{F}_{14} & \stackrel{\sim}{\dashrightarrow} & \mathcal{C}_{26}; \\ \mathcal{F}_{22} & \stackrel{2:1}{\dashrightarrow} & \mathcal{C}_{42}. \end{array}$$

Another upshot of the isomorphism (2.3) is that, once we fix a smooth cubic fourfold in C_{26} or C_{42} , the family of scrolls contained inside X is precisely parametrized by the associated K3. The construction, roughly speaking, goes as follows. For each $p \in S$, one defines a rational curve

$$\Delta_p := \{ y \in S^{[2]} : \{ p \} = Supp(y) \}.$$

The image of Δ_p inside F(X) then defines a (possibly singular) scroll $R_p \subset X$.

Proposition 2.5. [FV18, FV19]

- (1) The family of septimic 3-nodal scrolls inside a generic cubic fourfold X in C_{26} is the genus 14 K3 surface associated to X.
- (2) The family of 8-nodal degree 9 scrolls inside a generic cubic fourfold X in C_{42} is the genus 22 K3 surface associated to X.

Recall that the generic K3 surface in \mathcal{F}_{14} and \mathcal{F}_{22} has no automorphism, hence universal families exist:

$$\begin{array}{rccc} f_{14}:\mathcal{F}_{14,1} & \to & \mathcal{F}_{14}; \\ f_{22}:\mathcal{F}_{22,1} & \to & \mathcal{F}_{22}. \end{array}$$

at least over an open subset of each moduli space. This has allowed the study of $\mathcal{F}_{14,1}$ and $\mathcal{F}_{22,1}$ by Faraks-Verra. They considered universal K3s as moduli spaces for couples (X, R), where X is a cubic fourfold, and R is a scroll - belonging to a given class of surfaces - contained in X.

Proposition 2.6. [FV18, FV19] The universal K3s $\mathcal{F}_{8,1}$ and $\mathcal{F}_{22,1}$ are unirational.

5

Let us now come to the definition of the object that we will study in Sect. 3. Similarly to the case of curves and K3 surfaces we give the following definition.

Definition 2.7. By *universal cubic fourfold* over a divisor C_d we mean the moduli space $C_{d,1}$ of 1-pointed cubic fourfolds.

As it is customary over moduli spaces, the mere existence of a universal cubic fourfold needs a little bit of justification.

Proposition 2.8. The generic cubic fourfold in any divisor C_d does not have projective automorphisms, hence a universal family of cubic fourfolds exists over an open subset of each divisor.

Proof. This appears to be an easy corollary of [GAL11, Thm 3.8] where it is proven that the families of cubic fourfolds with a non-trivial projective automorphism have dimension at most 14. Hence clearly also the generic element of any divisor has no automorphism

Remark 2.9. Given the result of [GAL11], Prop. 2.8 is straightforward and very general. For d = 26, and 42, since the isomorphism (2.3) holds, one could also argue in a more "geometric" way as follows. Any automorphism $\phi \in \operatorname{Aut}(X)$ of a smooth cubic fourfold X induces an automorphism of the Fano variety of lines F(X). Then, via the isomorphism $F(X) \cong S^{[2]}$ with the symmetric square of the associated K3, one can use the fact that the generic K3 has no non-trivial automorphism to conclude that ϕ is the identity. Remark also that, by Prop. 2.5, the fact that the associated K3 has no non-trivial automorphism implies that any $\phi \in \operatorname{Aut}(X)$ should send each scroll onto itself.

3. Unirationality for $C_{26,1}$ and $C_{42,1}$ via universal K3 surfaces.

By the results of the preceding section, at least over a dense subset of C_{26} and C_{42} , there exists a universal family of cubic fourfolds. For simplicity, and since we are however working in the birational category, we will still denote the two universal families by $C_{26,1} \rightarrow C_{26}$ and $C_{42,1} \rightarrow C_{42}$, without stressing the fact that they may not be defined everywhere.

In this section we will show that:

Theorem 3.1. The universal cubic fourfolds $C_{26,1}$ and $C_{42,1}$ are universal.

The strategy will be the same for the two cases, hence we will resume here below shortly the properties, that hold for both divisors, that we will need. In order to keep the notation not too tedious, when we will say that a certain property holds for C_n we will assume n = 26, 42.

Let X be a generic smooth cubic fourfold in \mathcal{C}_n . We will denote by K(X) the associated K3 surface and by S(X) the family of scrolls (as defined in Def. 2.2) contained in $X \subset \mathbb{P}^5$. For n = 26 septimic 3-nodal scrolls, for n = 42 8-nodal degree 9 scrolls. One can rephrase Prop. 2.5 by saying that K(X) is isomorphic to S(X). More generally, we will denote by \mathcal{S}_n the Hilbert scheme of scrolls contained in cubics in \mathcal{C}_n (those appearing in Def. 2.2), and by S_n the PGL(6)-quotient of \mathcal{S}_n . These moduli quotients have been also considered in [FV18, FV19, Lai17]. Taking example from these papers let us give the following definition.

Definition 3.2. Let us denote by

 $\mathcal{H}_n = \{ (X, R) : R \subset X, \ [X] \in \mathcal{C}_n, \ R \in S(X) \}$

the "nested" Hilbert scheme given by the couples (X cubic fourfold whose class lives in C_n) + (R rational normal scroll in S(X)); and by $H_n := \mathcal{H}_n//PGL(6)$ the corresponding moduli quotient.

From [FV18, FV19] we have the following result.

Proposition 3.3. The universal K3 surfaces $\mathcal{F}_{n,1}$ are birational to H_n .

In particular, in [FV18, FV19] the authors show that S_n is unirational and \mathcal{H}_n is birational to a projective bundle over S_n , and hence unirational as well. We add just one more character to the plot, that is:

Definition 3.4. We denote by $\mathcal{H}_{n,1}$ the Hilbert scheme of the triples (X, R, p) as follows:

$$\mathcal{H}_{n,1} = \{ (X, R, p) \} : R \subset X, \ [X] \in \mathcal{C}_n, \ R \in S(X), \ p \in X \}.$$

We will denote by $H_{n,1}$ the corresponding moduli quotient by $PGL(6)$.

Of course, as a consequence of Prop. 2.8, this is generically a fibration in cubic hypersurfaces over \mathcal{H}_n . Finally we will denote by \mathcal{H}_n^3 the open subset complementary to the discriminant inside $|\mathcal{O}_{\mathbb{P}^5}(3)|^2$, and by $\mathcal{H}_{n,1}^3$ the universal family over \mathcal{H}_n^3 .

Theorem 3.5. The Hilbert scheme $\mathcal{H}_{n,1}$ is unirational.

Proof. Let us try to give a global picture of the situation.



 $^{^2\}mathrm{the}$ was previously denoted by U - but it seems more coherent to use this notation here

Here above all vertical arrows shall be intended as quotients by the PGL(6)action. On the *upper* level of the diagram all spaces are Hilbert schemes, whereas on the *lower* they are moduli quotients (H.S and M.S. for short). By the unirationality of S_n we have the unirationality of $\mathbb{P}^5 \times S_n$. Then we observe that there is a natural forgetful map

(3.7)
$$\phi_n: \mathcal{H}_{n,1} \to \mathbb{P}^5 \times \mathcal{S}_n;$$

$$(3.8) (X, R, p) \mapsto (p, R).$$

Namely, the fiber over a given couple (R, p) is exactly the linear system $\mathbb{P}H^0(\mathbb{P}^5, \mathcal{I}_{R\cup p}(3))$ of cubics in \mathbb{P}^5 containing R and p. Over an open subset of $\mathbb{P}^5 \times S_n$ (i.e. the locus where $p \notin R$) the rank does not change and it is 11, and 5 respectively for d = 26 and 42. This in turn implies that $\mathcal{H}_{n,1}$ is unirational. \Box

Proposition 3.9. The universal cubic fourfold $C_{n,1} \to C_n$ is unirational.

Proof. Recall from Prop. 2.8 that the generic cubic fourfold in the divisors C_n has no projective automorphism. Call $V \subset C_n$ the dense locus where cubics have no automorphism. It is straightforward to see - and we have already implicitely used this in Diag. 3.6 - that, over V, the universal cubic has a natural quotient structure

(3.10)
$$C_{n,1} = \mathcal{H}_{n,1}^3 / / PGL(6).$$

Hence $C_{n,1}$ is the natural moduli space for couples (X, p), up to the action of PGL(6). The upshot is that there exist a natural rational surjective forgetful map (up to PGL(6)-action)

(3.11)
$$\varphi_R: H_{n,1} \to \mathcal{C}_{n,1};$$

$$(3.12) (X,R,p) \mapsto (X,p);$$

that forgets the scrolls contained in X. By Thm. 3.5, the variety $H_{n,1}$ is unirational, and it dominates $C_{n,1}$, thus also $C_{n,1}$ is unirational.

4. UNIRATIONALITY THROUGH RATIONAL SPECIAL SURFACES.

4.1. Special cubics in C_d in the range $8 \le d \le 38$. The goal of this Section is to prove the unirationality of the non-empty families of universal cubics $C_{d,1}$ for $8 \le d \le 38$ by generalizing some results of Nuer, and using a celebrated result of Kollar [Kol02]. Then, by applying inductively the same argument, we will show that $C_{d,n}$ - the universal cubic with n marked points - is also unirational in the same range. Let us recall shortly the results we need.

In his paper [Kol02], extending a result of Segre that held only over \mathbb{Q} , Kollar shows the following

Theorem 4.1. Let k be a field and $X \subset \mathbb{P}^{n+1}$ a smooth cubic hypersurface of dimension $n \geq 2$ over k. Then the following are equivalent:

(1) X is unirational (over k);

(2) X has a k-point.

We are going to use this in a relative setting. In order to do this we recall some results due to Nuer.

In [Nue17], the author studies the birational geometry of the divisors C_d , for $12 \leq d \leq 38$, by considering an open subset $U_d \subset (\mathbb{P}^2)^p$ parametrizing generic *p*-tuples of distinct points that give certain cohomological invariants that appear in [Nue17, Table 2]. Let *S* be the rational surface obtained as the blow-up of \mathbb{P}^2 along the *p* points x_1, \ldots, x_p , and let us embed it into \mathbb{P}^5 via the linear series $|aL - (E_1 + \cdots + E_i) - 2(E_{i+1} + \cdots + E_{i+j}) - 3(E_{i+j+1} + \cdots + E_p)|$, where the parameters (a, i, j, p) of the linear system are those displayed in [Nue17, Table 1]. From [Nue17, Thm. 3.1] we know that there exists a vector bundle $V_d \to U_d$ such that the fiber over (x_1, \ldots, x_p) is the space of global sections $H^0(\mathcal{I}_{S|P5}(3))$. The natural classifying morphism $\mathbb{P}(V_d) \to C_d$, obtained by the universal property of C_d is dominant and this implies the unirationality of C_d .

Let us now denote by \mathcal{X}_d the universal cubic over $\mathbb{P}(V_d)$, and by $\pi_{p+1} : (\mathbb{P}^2)^{p+1} \to (\mathbb{P}^2)^p$ the forgetful map that forgets the last point. Now we can claim :

Theorem 4.2. The universal cubic fourfolds $C_{d,1}$ are unirational if $12 \le d \le 38$.

Proof. By pulling back through π , we have the following commutative diagram.



Actually the LHS column is only defined on $\pi^{-1}(U_d)$ but for simplicity we will not write this. Working in the birational category, this does not affect our results.

We observe that the pulled-back family $\pi^* \mathcal{X}_d$, seen as universal family over $\mathbb{P}(\pi^* V_d)$ has a tautological rational section *s*, that is the image in *S* of the $(p+1)^{th}$ point from $(\mathbb{P}^2)^{p+1}$. This means that $\pi^* \mathcal{X}_d$ has a rational point over the function field of $\mathbb{P}(\pi^* V_d)$. By Kollar's Theorem $\pi^* \mathcal{X}_d$ is then unirational over this field. This in turn is equivalent to the existence of a \mathbb{P}^4 -bundle on $\mathbb{P}(\pi^* V_d)$ that dominates $\pi^* \mathcal{X}_d$. The \mathbb{P}^4 -bundle is rational, $\pi^* \mathcal{X}_d$ dominates \mathcal{X}_d and this in turn dominates $\mathcal{C}_{d,1}$, which is then unirational.

Remark 4.4. An easy adaptation of this argument allows us to show the unirationality of $\mathcal{C}_{8,1}$ as well. Of course, all the planes in \mathbb{P}^5 are projectively equivalent, and the linear system of cubics through a given 2-plane P is $|\mathcal{I}_{P/\mathbb{P}^5}(3)| \cong \mathbb{P}^{45}$. The plane P has rational points over \mathbb{C} and the universal cubic $\mathcal{X}_8 \subset \mathbb{P}^{45} \times \mathbb{P}^5$ contains the "constant" plane $P \times \mathbb{P}^{45}$. The variety \mathcal{X}_8 also dominates $\mathcal{C}_{8,1}$, by definition. Hence \mathcal{X}_8 has rational points over the function field of \mathbb{P}^{45} and hence is unirational over this field. By the same argument as in Thm. 4.2, this implies the unirationality of $\mathcal{C}_{8,1}$.

4.2. Special cubics in C_{42} . The unirationality of universal cubics $C_{42,1}$ goes along the same lines as in the previous section, but we need to extract a couple of quite subtle results from [RS19b] about rational surfaces contained in cubics belonging to these divisor.

In [RS19b], the authors construct a 48-dimensional unirational family S_{42} of 5nodal surfaces of degree 9 and sectional genus 2, such that the divisor C_{42} can be described as the locus of cubics containing surfaces from this family. We reconstruct briefly their argument.

More precisely (see [RS19b, Rmk. 4.3]) they construct a family \mathcal{W} of del Pezzo quintics with some special intersection theoretical properties. The construction of such surfaces depend on certain choices (5 points on \mathbb{P}^2 , a line secant to a particular rational surface, etc.) which all depend on free, rational parameters. This implies that the family \mathcal{W} is unirational. By construction, each del Pezzo quintic $S \in \mathcal{W}$ is contained in one smooth del Pezzo fivefold $F \subset \mathbb{P}^8$, that is an hyperplane section of the Grassmannian Gr(2,5). The fivefold F contains a rational, 3-dimensional, family of planes, whose members all have class (2, 2) inside the Chow ring $CH^{\bullet}(Gr(2,5))$ of the Grassmannian. The projection with center any one of these planes defines a birational morphism between any del Pezzo quintic from \mathcal{W} and some 5-nodal $S \in S_{42}$. As observed in [RS19b], this in turn implies that S_{42} is unirational. Let us denote by \mathbb{S} the rational parameter space that dominates S_{42} . Other relevant work about C_{42} has been done in [Lai17] and [HS20].

Remark 4.5. We observe that the construction above implies that any $S \in S_{42}$ is birational to a del Pezzo quintic over $\mathbb{C}(\mathbb{S})$, the function field of \mathbb{S} . This means that on the same field there exist a birational map $\mathbb{P}^2 \dashrightarrow S$, as we explain in the following Lemma.

Lemma 4.6. Let S be any surface in S_{42} , then S is rational over the function field of \mathbb{S} .

Proof. As we have observed in Rmk. 4.5, S is birational to a del Pezzo quintic, over $\mathbb{C}(\mathbb{S})$. By a well-known result of Enriques [Enr97], a del Pezzo quintic is rational over any field.

Proposition 4.7. The universal cubic $C_{42,1}$ is unirational.

Proof. Let us denote by $\mathbb{P}(V_{42})$ the relative linear system over S_{42} of cubics through $S \in S_{42}$, and by \mathcal{X}_{42} the universal family over $\mathbb{P}(V_{42})$, that dominates $\mathcal{C}_{42,1}$. Then the situation is the following.



Here $\pi : \mathbb{S} \to S_{42}$ is the unirational parametrization described above, ρ the second projection and φ is the natural classifying map. Now, all cubics that are

the fibers of the fibration $\mathcal{X}_{42} \to \mathbb{P}(V_{42})$ contain by construction a surface $S \in \mathcal{S}_{42}$, and since these surfaces are rational over $\mathbb{C}(\mathbb{S})$ (they are birational to del Pezzo quintics), they have rational points over the same field. Summing up, there exists a section $s : \mathbb{P}(\rho^* \pi^* V_{42}) \to \rho^* \pi^* \mathcal{X}_{42}$, that - thanks to Kollar's Theorem - makes $\rho^* \pi^* \mathcal{X}_{42}$ unirational over $\mathbb{P}^2 \times \mathbb{S}$ and hence implies the unirationality of $\mathcal{C}_{42,1}$. \Box

Remark 4.9. As the reader may observe, the universal cubic $C_{44,1}$ is missing from our description. Cubics in this divisor contain a Fano model of an Enriques surface [Nue17, Thm. 3.2]. The divisor C_{44} is unirational, so in order to apply Kollar's Theorem and show that $C_{44,1}$ is unirational, one should show that the generic Enriques surface has a rational point.

4.3. Unirationality of $C_{d,n}$. In this section we are going to use the inductivity of the construction of $C_{d,n}$ - the moduli space of cubic fourfolds in C_d with n marked points - and Kollar's theorem in order to show the unirationality of $C_{d,n}$ for all n.

Recall in fact that $C_{d,n}$ is just the universal cubic fourfold over $C_{d,n-1}$. This allows us to use our machinery to prove inductively the following.

Theorem 4.10. The moduli spaces $C_{d,n}$ are unirational for all n, if $8 \le d \le 42$.

Proof. We will work inductively on n. For n = 0 (or respectively 1) the claim is true thanks to [Nue17] and [RS19b] (or resp. Sections 4.1 and 4.2). We will denote by S_d the unirational family of rational surfaces contained in cubics in C_d , $\mathbb{P}(V_d) \to S_d$ the relative ideal of of cubic hypersurfaces through each surface and $\mathcal{X}_{d,1} \to \mathbb{P}(V_d)$ the universal cubic over the relative linear system. As seen in Thm. 4.2 and Prop. 4.7, the variety $\mathcal{X}_{d,1}$ is unirational and dominates $C_{d,1}$ through the natural classifying map of the coarse moduli space $C_{d,1}$. The situation is described by the following diagram:



where $\mathbb{P}^2 \times \hat{\mathcal{S}}_d$ is the appropriate rational space that assures the existence of the section s. As seen in Thms. 4.2 and 4.7, this boils down to taking $\mathbb{P}^2 \times (\mathbb{P}^2)^p$ for $8 \leq d < 42$, and the rational parameter space $\mathbb{P}^2 \times \mathbb{S}$ for d = 42.

We remark that s is the section that makes $\pi^* \mathcal{X}_{d,1}$ unirational following Kollar's Theorem. Call σ the classifying map to $\mathcal{C}_{d,1}$. Then, we can add one point and consider the universal cubic $\mathcal{C}_{d,2} \to \mathcal{C}_{d,1}$, that fits in the following diagram



where γ is the second projection and σ the dominant map from Diagram 4.11. Recall that $\pi^* \mathcal{X}_{d,1}$ is unirational, and dominates $\hat{\mathcal{S}}_d$.

We observe that, exactly as we have done in Diag 4.11 for $\mathcal{C}_{d,1}$, since $\pi^* \mathcal{X}_{d,1}$ dominates $\hat{\mathcal{S}}_{d,1}$, we are able to define a natural rational section $s : \mathbb{P}^2 \times \pi^* \mathcal{X}_d \to \gamma^* \sigma^* \mathcal{C}_{d,2}$. Hence the universal cubic has rational points over $\mathbb{P}^2 \times \pi^* \mathcal{X}_{d,1}$ and this space is unirational. By Kollar's theorem and since $\gamma^* \sigma^* \mathcal{C}_{d,2}$ dominates $\mathcal{C}_{d,2}$, we get that $\mathcal{C}_{d,2}$ is unirational.

Now it is straightforward to see how to continue the argument by induction; we just draw the diagram for the following step for clarity. Recall that $\gamma^* \sigma^* C_{d,2} = \tau^* C_{d,2}$.



5. Some results of non-unirationality.

In this last section we collect some recent results about the Kodaira dimension of moduli spaces of K3 surfaces and of Hassett divisors C_d . These allow us to show that for an infinite range of values of d, the universal cubic fourfold over C_d can not be unirational. Let us first claim the following straightforward Lemma, in which we need to assume that d is even.

Lemma 5.1. Suppose that the Kodaira dimension of $\mathcal{F}_{\frac{d+2}{2}}$ or of \mathcal{C}_d is positive, then the universal cubic fourfold $\mathcal{C}_{d,1}$ is not unirational.

We will apply Lemma 5.1 to the following two Propositions, due to Gritsenko-Hulek-Sankaran and Várilly-Alvarado-Tanimoto. Prop. 5.2 was initially conceived for moduli of K3 surfaces; following [Nue17] we write its "translation" in terms of cubic fourfolds via the rational map of (2.4).

Proposition 5.2. Let d > 80, $d \equiv 2 \pmod{6}$, $4 \not| d$ be such that for any odd prime $p, p \mid d$ implies $p \equiv 1 \pmod{3}$. Then the Kodaira dimension of C_d is non-negative. If moreover d > 122, then C_d is of general type.

Proposition 5.3. The divisor C_{6n+2} is of general type for n > 18 and $n \neq 20, 21, 25$ and has nonnegative Kodaira dimension for n > 13 and $n \neq 15$. Moreover, C_{6n} is of general type for n > 18, $n \neq 20, 22, 23, 25, 30, 32$ and has nonnegative Kodaira dimension for n > 16 and $n \neq 18, 20, 22, 30$.

By combining Prop. 5.3 and 5.2 with Lemma 5.1 we obtain the following Proposition.

Proposition 5.4. Under the hypotheses on d of Prop. 5.3 and 5.2, the universal cubic fourfold $C_{d,1}$ is not unirational.

The full, lenghty, statement is in Prop. 1.3 in the Introduction.

5.1. **Open questions.** It would be interesting to start a systematic study of the birational geometry of universal cubic fourfolds. Some very natural questions rise, and they are of course related to the classical birational geometry of cubic fourfolds.

Questions:

- (1) What is the Kodaira dimension of the universal cubic fourfold over C_d , for different values of d.
- (2) In particular, is $C_{44,1}$ unirational?
- (3) Is there a relation between the Kodaira dimension of the universal cubic fourfold over C_d and that of the universal associated K3?
- (4) What is the Kodaira dimension of the universal cubic fourfold over the full moduli space C?
- (5) Are there divisors C_d , where the geometry of special rational surfaces inside cubic fourfolds can help to describe the birational geometry of universal associated K3 surfaces?

References

- [ABBVA14] A. Auel, M. Bernardara, M. Bolognesi, and A. Várilly-Alvarado, Cubic fourfolds containing a plane and a quintic del Pezzo surface, Algebr. Geom. 1 (2014), no. 2, 181–193.
- [AHTVA19] Nick Addington, Brendan Hassett, Yuri Tschinkel, and Anthony Várilly-Alvarado, Cubic fourfolds fibered in sextic del pezzo surfaces, Amer. J. Math. 141 (2019), no. 6, 1479–1500.
- [AT14] Nicolas Addington and Richard Thomas, Hodge theory and derived categories of cubic fourfolds, Duke Math. J. 163 (2014), no. 10, 1885–1927.
- [Awa19] Hanine Awada, Rational fibered cubic fourfolds with nontrivial brauer classes, arXiv preprint (2019), 1–20.
- [BD85] A. Beauville and R. Donagi, La variété des droites d'une hypersurface cubique de dimension 4, C. R. Acad. Sci. Paris Sér. I Math. 301 (1985), no. 14, 703–706.
- [BP20] Michele Bolognesi and Claudio Pedrini, The transcendental motive of a cubic fourfold, J. Pure Appl. Algebra 224 (2020), no. 8, 106333.
- [BRS19] Michele Bolognesi, Francesco Russo, and Giovanni Staglianò, Some loci of rational cubic fourfolds, Math. Ann. 373 (2019), no. 1-2, 165–190.
- [CG72] C. Herbert Clemens and Phillip A. Griffiths, The intermediate Jacobian of the cubic threefold, Ann. of Math. (2) 95 (1972), 281–356. MR 302652

- [Enr97] Federigo Enriques, Sulle irrazionalità da cui può farsi dipendere la risoluzione d'un' equazione algebrica f(xyz) = 0 con funzioni razionali di due parametri, Math. Ann. **49** (1897), no. 1, 1–23.
- [FV18] Gavril Farkas and Alessandro Verra, The universal K3 surface of genus 14 via cubic fourfolds, J. Math. Pures Appl. (9) 111 (2018), 1–20.
- [FV19] _____, The unirationality of the moduli space of k3 surfaces of genus 22, arXiv preprint (2019), 1–17.
- [GAL11] Víctor González-Aguilera and Alvaro Liendo, Automorphisms of prime order of smooth cubic n-folds, Arch. Math. (Basel) 97 (2011), no. 1, 25–37.
- [GHS13] V. Gritsenko, K. Hulek, and G. K. Sankaran, Moduli of K3 surfaces and irreducible symplectic manifolds, Handbook of moduli. Vol. I, Adv. Lect. Math. (ALM), vol. 24, Int. Press, Somerville, MA, 2013, pp. 459–526.
- [GS14] S. Galkin and E. Shinder, The fano variety of lines and rationality problem for a cubic hypersurface, arxiv preprint https://arxiv.org/abs/1405.5154 (2014).
- [Has00] B. Hassett, Special cubic fourfolds, Comp. Math. 120 (2000), no. 1, 1–23.
- [Has16] Brendan Hassett, Cubic fourfolds, K3 surfaces, and rationality *questions, Rationality problems in algebraic geometry, Lecture Notes in Math., vol. 2172, 2016, pp. 29– 66.
- [HS20] M. Hoff and G. Staglianó, New examples of rational gushel-mukai fourfolds, Math. Zeit. (2020).
- [Kol02] János Kollár, Unirationality of cubic hypersurfaces, J. Inst. Math. Jussieu 1 (2002), no. 3, 467–476.
- [Kuz10] A. Kuznetsov, Derived categories of cubic fourfolds, Cohomological and Geometric Approaches to Rationality Problems, Progress in Mathematics, vol. 282, Birkhäuser Boston, 2010, pp. 219–243.
- [Lai17] Kuan-Wen Lai, New cubic fourfolds with odd-degree unirational parametrizations, Algebra Number Theory 11 (2017), no. 7, 1597–1626.
- [Laz09] Radu Laza, The moduli space of cubic fourfolds, J. Algebraic Geom. 18 (2009), no. 3, 511–545.
- $[Laz10] \qquad \qquad , The moduli space of cubic fourfolds via the period map, Ann. of Math. (2) 172 (2010), no. 1, 673-711.$
- [Ma19] Shouhei Ma, Mukai models and borcherds products, arXiv preprint (2019), 1–37.
- [Nue17] Howard Nuer, Unirationality of moduli spaces of special cubic fourfolds and K3 surfaces, Algebr. Geom. 4 (2017), no. 3, 281–289.
- [RS19a] Francesco Russo and Giovanni Staglianò, Congruences of 5-secant conics and the rationality of some admissible cubic fourfolds, Duke Math. J. 168 (2019), no. 5, 849–865.
- [RS19b] _____, Trisecant flops, their associated k3 surfaces and the rationality of some fano fourfolds, arXiv preprint (2019), 1–30.
- [TVA19] Sho Tanimoto and Anthony Várilly-Alvarado, Kodaira dimension of moduli of special cubic fourfolds, J. Reine Angew. Math. 752 (2019), 265–300.

INSTITUT MONTPELLIERAIN ALEXANDER GROTHENDIECK, UNIVERSITÉ DE MONTPELLIER, CNRS CASE COURRIER 051 - PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 5, FRANCE *E-mail address:* hanine.awada@umontpellier.fr

INSTITUT MONTPELLIERAIN ALEXANDER GROTHENDIECK, UNIVERSITÉ DE MONTPELLIER, CNRS CASE COURRIER 051 - PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 5, FRANCE *E-mail address*: michele.bolognesi@umontpellier.fr