



**HAL**  
open science

# Spatial management can significantly reduce dFAD beachings in Indian and Atlantic Ocean tropical tuna purse seine fisheries

Taha Imzilen, Christophe Lett, Emmanuel Chassot, David M. Kaplan

► **To cite this version:**

Taha Imzilen, Christophe Lett, Emmanuel Chassot, David M. Kaplan. Spatial management can significantly reduce dFAD beachings in Indian and Atlantic Ocean tropical tuna purse seine fisheries. *Biological Conservation*, 2021, 254, pp.108939. 10.1016/j.biocon.2020.108939 . hal-03413528

**HAL Id: hal-03413528**

<https://hal.umontpellier.fr/hal-03413528v1>

Submitted on 3 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Spatial management can significantly reduce dFAD beachings in Indian and Atlantic Ocean tropical tuna purse seine fisheries

Taha Imzilen<sup>1,2,3a\*</sup>, Christophe Lett<sup>1,2b</sup>, Emmanuel Chassot<sup>4,5c</sup>, David M. Kaplan<sup>1,2d</sup>

1. Institut de Recherche pour le Développement (IRD), Avenue Jean Monnet,  
CS30171, 34203 Sète cedex, France

2. MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France

3. Sorbonne Université, Collège Doctoral, 75005 Paris, France

4. IRD, PO BOX 570, Victoria, Seychelles

5. Seychelles Fishing Authority, PO BOX 449, Victoria, Seychelles

**Author email addresses:** (a) [taha.imzilen@ird.fr](mailto:taha.imzilen@ird.fr); (b) [christophe.lett@ird.fr](mailto:christophe.lett@ird.fr); (c) [emmanuel.chassot@ird.fr](mailto:emmanuel.chassot@ird.fr); (d) [david.kaplan@ird.fr](mailto:david.kaplan@ird.fr)

**\*Corresponding author:** Taha Imzilen; Telephone: +33 (0)6 51 75 87 54; Email: [taha.imzilen@ird.fr](mailto:taha.imzilen@ird.fr)

**Total word count:** 6148

**Abstract:** 254

**Figures:** 6

## Acknowledgments

This work was funded by the Research Project INNOV-FAD (European Maritime and Fisheries Fund, measure n°39, OSIRIS #PFEA390017FA1000004, and France Filière Pêche), the European Research Project CECOFA2 (Specific Contract No 9 of

28 EASME/EMFF/2016/008) and the Ob7 Exploited Tropical Pelagic Ecosystems  
29 Observatory of the IRD. We express our sincere thanks to ORTHONGEL for making  
30 their dFAD tracking data available and to the Ob7 for data management and  
31 preparation. We are particularly grateful to L. Dagorn, D. Gaertner, A. Maufroy, L.  
32 Floch and M. Goujon for their assistance. We also thank two anonymous reviewers  
33 for their useful comments.

# 1 **Abstract**

2 Debris from fisheries pose significant threats to coastal marine ecosystems  
3 worldwide. Tropical tuna purse seine fisheries contribute to this problem via the  
4 construction and deployment of thousands of human-made drifting fish aggregating  
5 devices (dFADs) annually, many of which end up beaching in coastal areas. Here, we  
6 analyzed approximately 40 000 dFAD trajectories in the Indian Ocean and 12 000  
7 dFAD trajectories in the Atlantic Ocean deployed over the decade 2008-2017 to  
8 identify where and when beachings occur. We find that there is tremendous promise  
9 for reducing beaching events by prohibiting deployments in areas most likely to lead  
10 to a beaching. For example, our results indicate that 21% to 40% (depending on effort  
11 redistribution after closure) of beachings can be prevented if deployments are  
12 prohibited in areas in the south of 8°S latitude, the Somali zone in winter, and the  
13 western Maldives in summer for the Indian Ocean, and in an elongated strip of areas  
14 adjacent to the western African coast for the Atlantic Ocean. In both oceans, the  
15 riskiest areas for beaching are not coincident with areas of high dFAD deployment  
16 activity, suggesting that these closures could be implemented with relatively minimal  
17 impact to fisheries. Furthermore, the existence of clear hotspots for beaching  
18 likelihood and the high rates of putative recovery of dFAD buoys by small-scale  
19 fishers in some areas suggests that early warning systems and dFAD recovery  
20 programs may be effective in areas that cannot be protected via closures if appropriate  
21 incentives can be provided to local partners for participating in these programs.

22

23 **Keywords:** Marine pollution, Fishing debris, Coral reefs, Fish aggregating device  
24 (FAD), Ocean currents

# 25 1 Introduction

26 Debris from fisheries pose significant threats to coastal marine ecosystems worldwide  
27 (Tavares et al. 2017; Parton et al. 2019). Tropical tuna purse seine fisheries contribute  
28 to this problem via their extensive use of drifting fish aggregating devices (dFADs)  
29 (Consoli et al. 2020). Whereas historically purse seine vessels divided their fishing  
30 effort between free-swimming fish schools and schools associated with naturally-  
31 occurring floating objects (FOBs), they increasingly focus principally on FOB fishing  
32 (Galland et al. 2016; Taconet et al. 2018). The attachment to FOBs of, first, radio  
33 beacons in the mid-1980's and 1990's and then satellite-tracked, GPS-equipped buoys  
34 from the early 2000's, and most recently the integration of echo-sounders in satellite-  
35 tracked buoys have made this approach to catching tunas increasingly attractive to  
36 fishers (Chassot et al. 2014; Lopez et al. 2014). These technological developments  
37 have led purse seiners to manufacture and deploy large numbers of their own, human-  
38 made dFADs (Maufroy et al. 2017), and today it is believed that over 100 000 of these  
39 devices are deployed annually worldwide (Scott & Lopez 2014). dFADs typically  
40 consist of a floating structure and of a submerged substructure stretching up to 80 m  
41 below the surface (Imzilen et al. 2019). Some of the materials regularly used in dFAD  
42 construction include non-biodegradables such as PVC and metal tubes for the raft  
43 frames, ethylene vinyl acetate floats and plastic containers for buoyancy, and old  
44 nylon nets and pieces of salt bags for the subsurface structure. The massive increase  
45 in dFAD use poses a number of major concerns regarding ecological disturbance,  
46 overfishing, increased bycatch and creation of marine debris (Amandè et al. 2010;  
47 Dagorn et al. 2013; Filmalter et al. 2013; Maufroy et al. 2015). Most importantly for  
48 the context of this paper, a significant fraction of these dFADs end up beaching (i.e.,

49 stranding in coastal environments) (Maufroy et al. 2015), potentially damaging  
50 sensitive habitats such as coral reefs, and contributing to coastal marine debris and  
51 ghost fishing (Balderson & Martin 2015; Stelfox et al. 2016; Zudaire et al. 2018). This  
52 is of particular concern in a context of growing awareness of the extent of marine  
53 plastic pollution, with abandoned and lost fishing gears having been shown to be a  
54 major component of marine litter worldwide (Haward 2018; Lebreton et al. 2018;  
55 Richardson et al. 2019).

56

57 Given these concerns, dFAD beachings are a major area of interest for science,  
58 management and conservation. An initial examination of French dFAD spatio-  
59 temporal use in the tropical Indian Ocean and Atlantic Ocean over the period 2007-  
60 2011 indicated that ~10% of deployed dFADs ended up beached (Maufroy et al.  
61 2015), highlighting the potential for considerable impacts on fragile coastal habitats  
62 due to these events. A similar examination in the Western and Central Pacific Ocean  
63 found that ~6% of all trajectories were likely to have beached over a two year period  
64 (2016-2017; Escalle et al. 2019). However, given the significant differences in  
65 bathymetry and circulation between the western and central Pacific Ocean , Indian  
66 Ocean and Atlantic Ocean, and the more than four-fold increase in the number of  
67 dFADs deployed by purse seiners in the Indian and Atlantic Oceans since 2011  
68 (Katara et al. 2018; Floch et al. 2019), the extent to which existing literature applies to  
69 current patterns of dFAD use is an important open question. Moreover, the French  
70 fleet switched to almost exclusively using echo-sounder equipped dFAD tracking  
71 buoys around 2012 (Chassot et al. 2014; Floch et al. 2019) and other major purse  
72 seine fleets also started using this new technology on or before this date, potentially

73 altering the spatio-temporal distribution of dFAD deployments, fishing activity and  
74 associated beaching events. In parallel, management measures have been taken by the  
75 tuna regional fisheries management organizations to limit the total number of GPS  
76 buoys used by each purse seine vessel in both the Indian and Atlantic Oceans, but  
77 these measures have not directly addressed the spatial and temporal dynamics of  
78 beachings and, therefore, their efficacy for reducing this problem is unknown. A new  
79 analysis of dFAD beachings focused on spatio-temporal patterns that might be useful  
80 for identifying appropriate mitigation measures to avoid beachings is therefore  
81 urgently needed.

82

83 The goal of this paper is to quantify the impacts of dFAD beachings and identify  
84 strategies for mitigating these impacts in the tropical Indian and Atlantic Oceans.  
85 Using a large dataset of over 50 000 dFAD buoy trajectories, we first extend and  
86 improve upon the analysis of Maufroy et al. (2015, 2018), estimating beachings for  
87 the decade 2008-2017. We then identify deployment locations likely to lead to  
88 beaching events, and, using this information, we are able to estimate the impact of  
89 closing high beaching risk areas to dFAD deployments on the overall beaching rate  
90 under a pair of reasonable fishing effort redeployment strategies. Results indicate that  
91 there is indeed much promise in the Indian and Atlantic Oceans for reducing dFAD  
92 beachings by implementing sensible spatial limitations on deployment locations.

93

## 94 **2 Materials and methods**

### 95 **2.1 Data collection**

96

97 Through a collaboration with the French frozen tuna producers' organization  
98 (ORTHONGEL), the French Institute of Research on Development (IRD) has access  
99 to data on the locations of thousands of distinct GPS buoys attached to FOBs  
100 deployed by the French and associated flags (Mauritius, Italy, Seychelles) purse seine  
101 fleets operating in the tropical Indian and Atlantic Oceans from ~2007 onward  
102 (coverage ~75-86% before 2010 and ~100% after that date; Maufroy et al. 2015).

103 Though GPS buoys can be attached to both natural FOBs and human-made dFADs,  
104 the vast majority of FOBs in both oceans are now human-made dFADs (>90% of  
105 buoy deployments in both oceans based on observer data for 2013-2017), and,  
106 therefore, we will refer to these buoy trajectories as dFAD trajectories even though a  
107 small fraction of them are for other types of objects. GPS buoys are attached to  
108 dFADs deployed at sea by purse seine fishing vessels and their associated support  
109 vessels. Buoys can also be exchanged on FOBs encountered at sea and the buoys  
110 retrieved from the water are generally brought back to port where they can be  
111 recovered by the owner vessel for reuse. A single GPS buoy may therefore be  
112 redeployed several times, potentially on different dFADs. It is therefore important to  
113 note that, in this paper, we use the term 'dFAD' to refer to the entire device consisting  
114 of the floating object itself and the attached GPS buoy, whereas, the term 'buoy'  
115 designates solely the GPS buoy.

116



117 Buoy location data are transmitted with a periodicity that varies along the buoy  
118 trajectory, generally ranging from 15 minutes to 2 days. Buoy positions were filtered  
119 to remove those that were emitted while the buoy was onboard using a Random Forest  
120 classification algorithm that is an improvement over that developed in Maufroy et al.  
121 (2015) (Appendix A). This improved classification algorithm is estimated to have an  
122 error rate of ~ 2% when predicting onboard positions and ~ 0.2% for at sea positions  
123 (Supplementary Table A4).

124

125 In this study, we used data of dFAD positions covering the decade 2008-2017. This  
126 data set consists of ~15 million Indian Ocean positions representing a total of 38 845  
127 distinct buoys and ~6 million Atlantic Ocean positions representing a total of 12 147  
128 distinct buoys. Separately, locations and times for dFAD deployments are available in  
129 French logbook data from 2013 onward.

## 130 **2.2 Identification of dFAD beaching events**

131

132 dFAD beachings were identified in two steps. The first step was to find dFADs that  
133 had an abnormally small rate of movement for an extended period of time, whereas  
134 the second step removed false positives (e.g., buoys onboard or at port) from this list  
135 of potential beachings. A given dFAD position was considered to be a potential  
136 beaching if: (1) at least 2 other later positions were within 200 m, and (2) all these  
137 close positions span a time period exceeding 1 day. The 200 m threshold is based on a  
138 dFAD snagged on the very bottom of its <100 m length nets hanging below the dFAD  
139 swinging at most 100 m in each direction. The time span of at least 1 day is required  
140 to avoid identifying as beachings multiple position emissions from a single buoy over

141 a short time period, such as occurs when the emission periodicity of dFAD positions  
142 is modified to 15 min to facilitate detection by vessels before a fishing set.

143

144 In the second step, the putative beachings identified in this first step were filtered to  
145 remove non-beaching events based on 4 tests: (1) the beaching is more than 10 km  
146 from a major fishing port to avoid cases where dFAD buoys are at a port; (2) the  
147 beaching event is <5 km from land or the water column depth is <100 m; (3) all  
148 positions are classified at sea and there are no gaps in location emission exceeding 2  
149 days over the 5 days preceding the beaching; (4) greater than 90% of all positions of a  
150 given buoy within the time span of the potential beaching event are associated with  
151 the beaching event (i.e., meet the distance criteria described above; this condition  
152 avoids cases where a buoy happens to pass multiple times through the same area,  
153 because of an eddy for example). Only beaching events meeting these 4 conditions  
154 were considered for further analyses.

155

156 About half of the beachings identified by the conditions described above occurred in  
157 the water. The other half were generally located on land close to small fishing ports or  
158 coastal villages (Supplementary Fig. B2, Fig. B3 and Fig. B4). This suggests that  
159 these buoys were retrieved by small-scale boats, likely fishers. As these boats  
160 generally intercept dFADs in coastal areas and only collect the buoy for its valuable  
161 electronics, leaving the raft and netting to drift, it is entirely possible that these dFADs  
162 (without the buoy) later ended up beaching. Nevertheless, given the uncertainty  
163 regarding the fate of these dFADs, calculations in this paper have been carried out  
164 both including all beachings and including only beachings in the water. Unless

165 otherwise stated, statistics reported in the paper are for all beachings including those  
166 on land. In the rest of this paper, beachings located in water and on land are  
167 respectively referred to as “beachings along shore” and “recoveries displaced to  
168 shore”.

169

170 The number of beaching events per km of the continental shelf was calculated by  
171 counting all beachings occurring in each 5°x5° grid cell and then dividing that number  
172 by the kilometers of continental shelf edge, defined by the 200 m isobath, within the  
173 cell. The continental shelf edge was used instead of the coastline to avoid  
174 anomalously high beaching rates for some very small islands surrounded by large  
175 continental shelf areas.

176

177 For identifying beachings, classifying beachings as on land or at sea and determining  
178 the continental shelf edge, coastline data were obtained from OpenStreetMap land  
179 polygons (available at <https://osmdata.openstreetmap.de/data/land-polygons.html>;  
180 accessed 2020-02-19) and bathymetry was obtained from the 30-arcsecond-resolution  
181 General Bathymetric Chart of the Oceans (GEBCO v.2014; available at  
182 [https://www.gebco.net/data\\_and\\_products/gridded\\_bathymetry\\_data](https://www.gebco.net/data_and_products/gridded_bathymetry_data); accessed 2020-  
183 02-19).

## 184 **2.3 Drift locations leading to beachings**

185

186 In order to identify dFAD drift locations that had a high risk to lead to a beaching  
187 event, we calculated the fraction of buoys that beach within 3 months of a passage  
188 through a given 1°x1° grid cell. This analysis was carried out over the entire study

189 period, but also by season to estimate seasonal variability in beaching risk. We  
190 selected 3 months as the time limit as it is intermediate between the mean timespan of  
191 at sea trajectories and that of the lifespan of a buoy in the dataset (i.e. 25 and 196  
192 days, respectively), and because 3 months was considered a reasonable timespan over  
193 which fishers and managers could reasonably be expected to predict and mitigate for  
194 beaching likelihood. To ensure that results are not strongly sensitive to this choice,  
195 additional analyses were carried out to calculate the fraction of buoys that beach  
196 within 12 months. Note that individual buoy trajectories were separated into multiple  
197 in water trajectories using breaks defined by gaps of more than 2 days or positions  
198 classified as onboard representing more than 1 minute of trajectory time. The 1  
199 minute limit was imposed to remove very short trajectory segments that were  
200 problematic for the classification algorithm (Appendix A).

201

202 Since beachings threaten fragile marine habitats, especially coral habitats, we carried  
203 out the same analyses focusing exclusively on beachings in coral reef areas. Data on  
204 the global distribution of coral reefs were obtained from UNEP-WCMC, WorldFish  
205 Centre, WRI, TNC (2018 ,version 4.0; available at [https://data.unep-](https://data.unep-wcmc.org/datasets/1)  
206 [wcmc.org/datasets/1](https://data.unep-wcmc.org/datasets/1); accessed April 30, 2019).

## 207 **2.4 Deployment risk**

208

209 To assess potential for reducing the dFAD beaching rate, we investigated closing areas  
210 of high beaching risk to dFAD deployments. Deployment locations were obtained  
211 from logbook data, whereas probability of beaching for a given deployment location  
212 was estimated as described above. Logbook deployment data was used instead of

213 putative deployments from reconstructed dFAD trajectories because, though the  
214 random forest position classification model has a very high accuracy rate and  
215 predicted deployment locations do approximately follow the spatial distribution of  
216 logbook deployment locations (Maufroy et al. 2015), accurately predicting  
217 deployment locations is quite difficult and error prone given that a single error  
218 anywhere in the trajectory will split the trajectory, generating a new false deployment  
219 (Maufroy et al. 2015). Given the high quality of logbook data, it was considered that  
220 this was the most accurate estimate of recent dFAD deployment locations.

221

222 Multiplying dFAD deployments by the proportion of devices beaching allowed us to  
223 predict the reduction in beachings that would result from closing a given area.  
224 Different sized areas corresponding to specific percentages of all pre-closure  
225 deployments were closed in order of beaching risk going from highest to lowest. Two  
226 hypotheses were considered regarding the number and spatial distribution of  
227 deployments after closing an area to deployments: (1) closures eliminate deployments  
228 that would have occurred in closed areas (i.e., fishing effort reduction occurs), and (2)  
229 closures displace deployments formerly in closed areas to remaining unclosed areas in  
230 proportion to the relative density of deployments prior to implementation of closures  
231 (i.e., “fishery squeeze” occurs; Halpern et al. 2004).

### 232 **3 Results**

233 The number of French buoys deployed per year has increased dramatically and  
234 continuously over the decade 2008-2017, especially in the Indian Ocean (Fig. 1a).  
235 Over that period, more GPS buoys were deployed in the Indian Ocean (~ 40 000) than

236 in the Atlantic Ocean (~ 12 000). The percentage of all deployed dFADs that ended up  
237 beaching has also dramatically increased from ~3.5% in 2008 to ~20% in 2013 (Fig.  
238 1b; these numbers are roughly halved if we count only beachings along shore). After  
239 2013, the percentage of dFADs that beached stabilizes at ~15-20% in the Indian  
240 Ocean and ~19-22% in the Atlantic Ocean. In total, we obtained 7187 beaching events  
241 for the Indian Ocean and 2283 for the Atlantic Ocean.

242

243 Maps of these 9470 beaching locations clearly identify coastal beaching hotspots (Fig.  
244 2a and Supplementary Fig. C1a). Beachings occur in several zones in the Indian  
245 Ocean, including southern Somalia, Kenya, Tanzania, Seychelles and the Maldives. In  
246 the Atlantic Ocean, they occur mainly along the West African coast and the Gulf of  
247 Guinea between 20°N and 20°S. In both oceans, beachings also sporadically occur in  
248 more remote areas outside typical purse-seine fishing grounds (Maufroy et al. 2017),  
249 such as Indonesia, South Africa, Brazil and the Caribbean. Including only beachings  
250 that occur along the shore, the number of beaching decreases mostly along the  
251 western and north-eastern African coasts and in the Maldives (Fig. 2b and  
252 Supplementary Fig. C1b), indicating that significant rates of putative recovery of  
253 dFAD buoys occur in those areas.

254

255 In both oceans, the proportion of dFADs beaching within 3 months of passing through  
256 a 1°x1° grid cell shows high spatial heterogeneity, with hotspots of beaching  
257 likelihood clearly visible (Fig. 3a). In the Indian Ocean, the Gulf of Aden, Oman,  
258 Mozambique Channel, eastern and northern Madagascar, northern Maldives, western  
259 India, Sri Lanka and western Indonesia are all high risk areas for beaching. In the

260 Atlantic Ocean, the Gulf of Guinea, southern West Africa, the northern coast of South  
261 America and Caribbean have high proportion of beaching. Including only beachings  
262 that occur along shore reduces beaching proportions in all areas and reduces the  
263 importance of some coastal areas characterized by a high density of small-scale  
264 fishers, such as in the vicinity of the Arabian Peninsula, the northern Gulf of Guinea  
265 and West Africa (Fig. 3b). Increasing the temporal window from 3 months to 12  
266 months increases somewhat the spatial area over which proportion of beaching is non-  
267 negligible, but overall spatial patterns remain the same (Supplementary Fig. C2).  
268 Seasonal variability in dominant currents impacts beaching risk in predictable ways.  
269 For example, in the Indian Ocean, during the winter monsoon, onshore currents create  
270 an area of high proportion of beaching east of Somalia, but this high risk area  
271 disappears during the upwelling favorable period of the summer monsoon  
272 (Supplementary Fig. C4). However, seasonal variability in the Atlantic Ocean was  
273 weak (Supplementary Fig. C5). Finally, focusing exclusively on dFAD beachings on  
274 coral reefs narrowed the areas of high beaching risk to the north-west of the Maldives,  
275 Seychelles, northern Madagascar, the Mozambique Channel and the Caribbean  
276 (Supplementary Fig. C6).

277

278 Major areas of dFAD deployments during 2013-2017 spanned the whole fishing  
279 grounds of the French and associated flags purse seine fishery (Fig. 4a-b). In the  
280 Atlantic Ocean, dFADs were deployed all along the coast of West Africa, from  
281 Mauritania down to Angola with the most intense activity being observed along the  
282 equator and off the coasts of Mauritania, Gabon and Angola. In the Indian Ocean,  
283 dFADs were deployed in the Western Indian Ocean, including the Exclusive

284 Economic Zones of the Seychelles, Comoros, Kenya, French overseas territories and  
285 northwest of Madagascar in the northern Mozambique Channel. dFADs deployments  
286 were particularly frequent North-West of the Seychelles.

287

288 Combining spatial proportions of dFADs that beached (Fig. 3a-b) with observed  
289 dFAD deployment positions (Fig. 4a-b), we estimated the expected change in  
290 beachings and dFAD deployments due to prohibiting dFAD deployments in the  
291 highest risk areas for both oceans. Under all scenarios of dFAD deployment  
292 redistribution, spatial prohibitions are predicted to significantly reduce beaching rates.  
293 For example, if we prohibit dFAD deployments in areas corresponding to the 20% of  
294 deployments with highest beaching risk, we can prevent 37% of beachings in the  
295 Indian Ocean and 40% in the Atlantic Ocean in the absence of dFAD deployment  
296 effort redistribution, and 21% and 25% of beachings in the Indian Ocean and Atlantic  
297 Ocean, respectively, even if we allow for dFAD deployment redistribution to areas  
298 with less beaching risk (Fig. 5a). These percentages are even higher when we focus on  
299 the proportion of beaching including only beachings that happen along shore, with up  
300 to a 52% reduction in beachings in the Atlantic Ocean even if the total number of  
301 deployments is conserved via effort redistribution (grey dashed line in Fig. 5b).

302

303 Spatial prohibitions can be optimized to account for seasonal variability in beaching  
304 risk. For example, if areas corresponding to the 20% of deployments in areas with the  
305 highest beaching risk for each quarter are closed to dFAD deployments  
306 (Supplementary Fig. C7), we predict a 27% and 28% reduction in the Indian Ocean



307 and Atlantic Ocean, respectively, even if dFAD deployment redistribution is allowed  
308 (Fig. 5c).

309 Focusing exclusively on beachings in coral reefs, prohibiting the 20% of deployments  
310 in the Indian Ocean with the highest beaching risk to corals reduces coral reef  
311 beachings by 27% assuming dFAD deployment redistribution (Supplementary Fig.  
312 C8b), but the zones prohibited differ significantly from those that would be prohibited  
313 to reduce all beaching events (compare Fig. 4a and Supplementary Fig. C8a).

314

315 Closing the highest beaching risk areas to dFAD deployments is particularly effective  
316 at reducing beaching events in the south-western Indian Ocean and in the eastern Gulf  
317 of Guinea in the Atlantic Ocean (Fig. 6). If one focuses exclusively on coral reef  
318 beaching, then significant beaching reductions in the Indian Ocean are also seen in the  
319 Maldives and off Indonesia (Supplementary Fig. C9). These results apply both with  
320 and without dFAD deployment redistribution post closure implementation.

## 321 **4 Discussion**

322 The overriding conclusion to be drawn from our results is that there is potentially a lot  
323 to be gained in terms of reduction in the rate of dFAD beachings from spatio-temporal  
324 closures for dFAD deployments by purse seine fishing vessels in the Indian and  
325 Atlantic Oceans. We examined a wide range of scenarios for closure objectives,  
326 implementation, and post-closure effects: considering all beachings versus just  
327 strandings along shore; considering all coastal zones versus just coral reefs;  
328 implementing static versus quarterly varying closures; and post-closure effort  
329 reduction versus effort redistribution to remaining open areas. In all cases, closing the

330 riskiest areas for beaching is predicted to produce a tremendous reduction in  
331 beachings. Analyses of recent dFAD deployments in the Indian Ocean by the Spanish  
332 fleet (the dominant other fleet in both oceans) indicate that Spanish and French  
333 deployments have quite similar spatial distributions (Katara et al. 2018). This suggests  
334 that our results may be applicable to all fleets, though access to dFAD trajectory data  
335 should be enhanced to confirm this. Perhaps most encouraging, high risk areas  
336 generally are relatively coherent in space so that it should be feasible from a  
337 management perspective to implement closures (e.g., south of 8°S in the Indian Ocean  
338 and coastal zones in the Gulf of Guinea in the Atlantic Ocean). In both oceans, the  
339 riskiest areas for beaching are not coincident with areas of high dFAD deployment  
340 activity or fishing activities (Maufroy et al. 2015), suggesting that these closures  
341 could be implemented with relatively minimal impact to fisheries. The beaching  
342 reduction across coastal areas spared by the closures for dFAD deployment is highest  
343 in the south-western Indian Ocean and in the eastern Gulf of Guinea in the Atlantic  
344 Ocean, suggesting that our proposed deployment closure strategy is particularly  
345 efficient to protect these areas. The north-western Indian Ocean and the northern Gulf  
346 of Guinea, which both represent hotspots of beaching, are less protected by the  
347 closures for dFAD deployments. However, high rates of putative recovery of dFAD  
348 buoys by coastal boats in these areas indicate that beaching early warning systems and  
349 dFAD recovery programs may be effective in areas that cannot be protected via  
350 closures if appropriate incentives can be provided to local partners for participating in  
351 these programs.  
352

353 As reported elsewhere (Maufroy et al. 2015; Floch et al. 2017, 2019), the number of  
354 dFADs deployed in both oceans has dramatically increased over the last decade. More  
355 surprising, the fraction of dFADs that end up beaching increased significantly over the  
356 period 2008-2013, after which time the fraction stabilizes. As this 2008-2013 period is  
357 coincident with a number of changes in the fishery, such as the switch to echosounder  
358 buoys (2010-2012), an increase in the prevalence of dFAD fishing as opposed to  
359 fishing on free-swimming schools (Assan et al. 2019; Floch et al. 2019) and the  
360 fallout from Somali piracy (~2007-2011), it is hard to assign a specific cause to this  
361 pattern. One hypothesis is that as the number of dFADs has increased, the fraction of  
362 dFADs that are never fished upon has become more and more important to the point  
363 that after 2013 the fraction beaching simply reflects the balance one would expect in  
364 the absence of fishing between dFADs that beach versus dFADs that sink at sea. The  
365 stabilization of the beaching rate after 2013 may also be partially due to the  
366 implementation after 2014 of industry and/or regional fisheries management  
367 organizations' limits on the number of buoys monitored by purse seine vessels  
368 (ICCAT 2019; IOTC 2019a) as fishers may remotely deactivate non-productive  
369 dFADs to remain under industry limits, resulting in the loss of location information  
370 for these dFADs that continue to drift at sea and may later beach.

371

372 The risk of beaching depends strongly on upper ocean circulation and its seasonal  
373 variability. In the Indian Ocean, the African coast south of the equator represents a  
374 high beaching risk area throughout the year due to the westward flowing Northern  
375 Equatorial Madagascar Current (Schott et al. 2009) that drives dFADs to the coasts of  
376 Mozambique and Tanzania. In the northern Indian Ocean, high beaching risk areas

377 change with monsoon regimes. The Somali coast represents a high beaching risk area  
378 in the winter when the Somali Current flows westwards (Schott & McCreary 2001),  
379 but not during the summer, when the western Maldives become a high risk area due to  
380 monsoon driven eastward circulation. There is less effect of seasonality on beaching  
381 risk in the Atlantic Ocean, where areas of high beaching risk are driven by more-  
382 stable dominant circulation patterns. Along the western coast of Africa, beachings are  
383 related to the North Equatorial Countercurrent and the Guinea Current flowing  
384 eastwards, whereas high risk areas along the northern coast of South America and the  
385 Caribbean are linked to the South Equatorial, North Equatorial, North Brazil and  
386 Caribbean Currents flowing westwards (Bourles et al. 1999).

387

388 Our estimates of dFAD beaching rates after 2013 are higher than those estimated in  
389 the western central Pacific (Escalle et al. 2019) and in previous examinations in the  
390 Indian and Atlantic Oceans (Maufroy et al. 2015; Zudaire et al. 2018). Escalle et al.  
391 (2019) examined an area of the Pacific characterized principally by many small island  
392 chains, perhaps explaining lower beaching rates with respect to the continental land  
393 masses of the Indian and Atlantic Oceans. In the Indian and Atlantic Oceans, Maufroy  
394 et al. (2015) examined the period prior to 2013 for which we also find lower beaching  
395 rates. Zudaire et al. (2018) were principally concerned with the more-limited area of  
396 the Seychelles Archipelago, which is composed of a large set of small islands similar  
397 to the area examined by Escalle et al. (2019) in the western central Pacific. They also  
398 considered a somewhat more restrictive definition of beaching.

399

400 There have been several recent management changes regarding the use of dFADs that  
401 may alter future dFAD beaching patterns, highlighting the importance of continuous  
402 monitoring of dFAD trajectories. The Indian Ocean Tuna Commission (IOTC) and the  
403 International Commission for the Conservation of Atlantic Tunas (ICCAT) currently  
404 limit the number of buoys monitored by an individual purse seine vessel at any given  
405 time to 300 (ICCAT 2019) and 350 (IOTC 2019a) buoys in the Atlantic Ocean and  
406 Indian Ocean, respectively, and these limits are may decrease over time. The IOTC  
407 has also implemented a resolution to progressively reduce and phase out the number  
408 of support vessels that assist purse seiners with the management of dFADs (IOTC  
409 2019b). These changes may lead purse seine vessels to optimize their use of dFADs in  
410 a number of ways. One potential outcome would be that fishers remotely deactivate  
411 dFADs that are likely to beach or drift outside of areas of interest so as to remain  
412 under industry limits. This practice is of much concern as it would result in the loss of  
413 information on the extent and location of dFAD beachings currently made available  
414 via fishing companies on a voluntary basis. Tuna regional fisheries management  
415 organizations should put in place appropriate incentives or other measures to assure  
416 that this information loss does not occur.

417

418 This study would not have been possible without access to a long and extensive time  
419 series of data on French dFAD trajectories. Though access to these extensive datasets  
420 is still quite limited for most fishing fleets worldwide, there are a number of  
421 encouraging signs of increased reporting of dFAD deployments and other dFAD  
422 activities to tuna regional fisheries management organizations (IOTC 2019a; Escalle  
423 et al. 2020). We are hopeful that comprehensive datasets from all purse seine fishing

424 fleets will be available in the near future, permitting better estimates of the impacts of  
425 management options and the development of real-time tools for the management of  
426 dFAD impacts on marine ecosystems.

## 427 **5 Supporting information**

428

429 Supporting information available online comprises details of the new classification  
430 model for onboard and at sea states of dFAD trajectory data (Appendix A), and  
431 quantification of beachings occurred in water (beachings along shore) and on land  
432 (recoveries displaced) (Appendix B), as well as additional figures presenting the  
433 number of French dFADs beached in each 5°x5° cell, proportions of beaching using a  
434 12 month time window, seasonal variability in beaching risks and beaching risks for  
435 coral reefs (Appendix C).

## 436 **6 References**

- 437 Amandè MJ, Ariz J, Chassot E, Delgado de Molina A, Gaertner D, Murua H, Pianet  
R, Ruiz J, Chavance P. 2010. Bycatch of the European purse seine tuna fishery  
in the Atlantic Ocean for the 2003–2007 period. *Aquatic Living Resources*  
**23**:353–362.
- Assan C, Lucas J, Chassot E. 2019. Statistics of the Seychelles purse seine fleet  
targeting tropical tunas in the Indian Ocean (2000-2018). Page 18p. IOTC, San  
Sebastian, Spain, 21-26 October 2019.
- Balderson SD, Martin LEC. 2015. Environmental impacts and causation of “beached”  
drifting fish aggregating devices around Seychelles Islands: A preliminary  
report on data collected by Island Conservation Society. Page 15. IOTC,  
Olhao, Portugal, 7-11 September 2015.
- Bourles B, Molinari RL, Johns E, Wilson WD, Leaman KD. 1999. Upper layer  
currents in the western tropical North Atlantic (1989–1991). *Journal of  
Geophysical Research: Oceans* **104**:1361–1375.
- Chassot E, Goujon M, Maufroy A, Cauquil P, Fonteneau A, Gaertner D. 2014. The  
use of artificial fish aggregating devices by the French tropical tuna purse  
seine fleet: historical perspective and current practice in the Indian Ocean.  
Page 17p Sixteenth Session of the Working Party on Tropical Tunas. IOTC,  
Victoria. Available from <http://www.documentation.ird.fr/hor/fdi:010063284>  
(accessed January 13, 2015).

- Consoli P, Sinopoli M, Deidun A, Canese S, Berti C, Andaloro F, Romeo T. 2020. The impact of marine litter from fish aggregation devices on vulnerable marine benthic habitats of the central Mediterranean Sea. *Marine Pollution Bulletin* **152**:110928.
- Dagorn L, Holland KN, Restrepo V, Moreno G. 2013. Is it good or bad to fish with FADs? What are the real impacts of the use of drifting FADs on pelagic marine ecosystems? *Fish and Fisheries* **14**:391–415.
- Escalle L, Phillips JS, Brownjohn M, Brouwer S, Gupta AS, Sebille EV, Hampton J, Pilling G. 2019. Environmental versus operational drivers of drifting FAD beaching in the Western and Central Pacific Ocean. *Scientific Reports* **9**:1–12.
- Escalle L, Muller B, Hare S, Hamer P, Pilling G, PNAO, 2020. Report on analyses of the 2016/2020 PNA FAD tracking programme. WCPFC Scientific Committee WCPFC-SC16-2020/MI-IP-14.
- Filmalter JD, Capello M, Deneubourg J-L, Cowley PD, Dagorn L. 2013. Looking behind the curtain: quantifying massive shark mortality in fish aggregating devices. *Frontiers in Ecology and the Environment*:130627131409009.
- Floch L, Billet N, Dewals P, Irié D, Cauquil P, Gaertner D, Chassot E. 2017. Statistics of the French purse seine fishing fleet targeting tropical tunas in the Atlantic Ocean (1962-2015). *ICCAT Col. Vol. Sci. Pap.* **73**:755–778.
- Floch L, Depetris M, Dewals P, Duparc A, Kaplan DM, Lebranchu J, Marsac F, Pernak M, Bach P. 2019. Statistics of the French purse seine fishing fleet targeting tropical tunas in the Indian Ocean (1981-2018). Page 27p. San Sebastian, Spain, 21-26 October 2019. Available from [https://www.iotc.org/sites/default/files/documents/2019/10/IOTC-2019-WPTT21-11\\_Rev1.pdf](https://www.iotc.org/sites/default/files/documents/2019/10/IOTC-2019-WPTT21-11_Rev1.pdf).
- Galland G, Rogers A, Nickson A. 2016. Netting billions: A global valuation of tuna. Page 22. The PEW Charitable Trusts, Washington D.C., U.S.A.
- Halpern BS, Gaines SD, Warner RR. 2004. Confounding Effects of the Export of Production and the Displacement of Fishing Effort from Marine Reserves. *Ecological Applications* **14**:1248–1256.
- Haward M. 2018. Plastic pollution of the world's seas and oceans as a contemporary challenge in ocean governance. *Nature Communications* **9**:1–3. Nature Publishing Group.
- ICCAT. 2019. Compendium management recommendations and resolutions adopted by iccat for the conservation of atlantic tunas and tuna-like species. Available from [https://iccat.int/Documents/Recs/COMPENDIUM\\_ACTIVE\\_ENG.pdf](https://iccat.int/Documents/Recs/COMPENDIUM_ACTIVE_ENG.pdf) (accessed March 12, 2020).
- Imzilen T, Chassot E, Barde J, Demarcq H, Maufroy A, Roa-Pascuali L, Ternon J-F, Lett C. 2019. Fish aggregating devices drift like oceanographic drifters in the near-surface currents of the Atlantic and Indian Oceans. *Progress in Oceanography* **171**:108–127.
- IOTC. 2019a. Procedures on a Fish Aggregating Devices (FADs) management plan. Page 19/02. Available from <https://iotc.org/cmm/resolution-1902-procedures-fish-aggregating-devices-fads-management-plan> (accessed March 12, 2020).
- IOTC. 2019b. On an interim plan for rebuilding the indian ocean yellowfin tuna stock in the IOTC area of competence. Page 19/01. Available from [https://www.iotc.org/sites/default/files/documents/compliance/cmm/iotc\\_cmm\\_1901.pdf](https://www.iotc.org/sites/default/files/documents/compliance/cmm/iotc_cmm_1901.pdf).

- Katara I, Gaertner D, Marsac F, Grande M, Kaplan DM, Urtizberea A, Guery L, Depetris M, Duparc A, Floch L, Lopez J, Abascal F. 2018. Standardisation of yellowfin tuna CPUE for the EU purse seine fleet operating in the Indian Ocean. Page 19. IOTC–2018–WPTT20–36\_Rev1. Indian Ocean Tuna Commission 20th Working Party on Tropical Tunas (WPTT), Mahé, Seychelles. Available from [https://www.iotc.org/sites/default/files/documents/2018/10/IOTC-2018-WPTT20-36\\_Rev1.pdf](https://www.iotc.org/sites/default/files/documents/2018/10/IOTC-2018-WPTT20-36_Rev1.pdf).
- Lebreton L, Slat B, Ferrari F, Sainte-Rose B, Aitken J, Marthouse R, Hajbane S, Cunsolo S, Schwarz A, Levivier A, Noble K, Debeljak P, Maral H, Schoeneich-Argent R, Brambini R, Reisser J. 2018. Evidence that the Great Pacific Garbage Patch is rapidly accumulating plastic. *Scientific Reports* **8**:1–15. Nature Publishing Group.
- Lopez J, Moreno G, Sancristobal I, Murua J. 2014. Evolution and current state of the technology of echo-sounder buoys used by Spanish tropical tuna purse seiners in the Atlantic, Indian and Pacific Oceans. *Fisheries Research* **155**:127–137.
- Maufroy A, Chassot E, Joo R, Kaplan DM. 2015. Large-scale examination of spatio-temporal patterns of drifting Fish Aggregating Devices (dFADs) from tropical tuna fisheries of the Indian and Atlantic Oceans. *PLOS ONE* **10**:e0128023.
- Maufroy A, Kaplan DM, Bez N, Molina D, Delgado A, Murua H, Floch L, Chassot E. 2017. Massive increase in the use of drifting Fish Aggregating Devices (dFADs) by tropical tuna purse seine fisheries in the Atlantic and Indian oceans. *ICES Journal of Marine Science* **74**:215–225.
- Maufroy A, Kaplan D, Chassot E, Goujon M. 2018. Drifting fish aggregating devices (dFADs) beaching in the Atlantic Ocean: an estimate for the French purse seine fleet (2007-2015). *ICCAT Collective Volume of Scientific Papers* **74**:2219–2229.
- Parton K, Galloway T, Godley B. 2019. Global review of shark and ray entanglement in anthropogenic marine debris. *Endangered Species Research* **39**.
- Richardson K, Asmutis-Silvia R, Drinkwin J, Gilardi KVK, Giskes I, Jones G, O'Brien K, Pragnell-Raasch H, Ludwig L, Antonelis K, Barco S, Henry A, Knowlton A, Landry S, Mattila D, MacDonald K, Moore M, Morgan J, Robbins J, van der Hoop J, Hogan E. 2019. Building evidence around ghost gear: Global trends and analysis for sustainable solutions at scale. *Marine Pollution Bulletin* **138**:222–229.
- Schott FA, McCreary JP. 2001. The monsoon circulation of the Indian Ocean. *Progress in Oceanography* **51**:1–123.
- Schott FA, Xie S-P, McCreary Jr. JP. 2009. Indian Ocean circulation and climate variability. *Reviews of Geophysics* **47**. Available from <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2007RG000245> (accessed April 12, 2019).
- Scott GP, Lopez J. 2014. *The Use of Fads in Tuna Fisheries*. Brussels: European parliament. Available from [https://www.europarl.europa.eu/thinktank/en/document.html?reference=IPOL-PECH\\_NT\(2014\)514002](https://www.europarl.europa.eu/thinktank/en/document.html?reference=IPOL-PECH_NT(2014)514002) (accessed August 9, 2020).
- Stelfox M, Hudgins J, Sweet M. 2016. A review of ghost gear entanglement amongst marine mammals, reptiles and elasmobranchs. *Marine Pollution Bulletin* **111**:6–17.



- Taconet P, Chassot E, Barde J. 2018, February 1. Global monthly catch of tuna, tuna-like and shark species (1950-2015) aggregated by 1° or 5° squares (IRD level 2). Zenodo. Available from <https://zenodo.org/record/1164128#.XmoV3nVKhuQ> (accessed March 12, 2020).
- Tavares D, de Moura J, Merico A, Siciliano S. 2017. Incidence of marine debris in seabirds feeding at different water depths. *Marine Pollution Bulletin* **119**:68–73.
- Zudaire I, Santiago J, Grande M, Murua H, Adam PA, Nogués P, Collier T, Morgan M, Kahn N, Baguette F, Moron J, Moniz I, Herrera M. 2018. FAD Watch: a collaborative initiative to minimize the impact of FADs in coastal ecosystems. *IOTC Proceedings* **IOTC-2018-WPEB14-12**:21.

## 438 **Figure Captions**

439 **Fig. 1** (a) Annual number of new buoys deployed by the French and associated flags  
440 purse seine fleet in the Atlantic (grey) and Indian (black) oceans over the period 2008-  
441 2017 and (b) percentage of these buoys that beached. The lines in (b) with solid  
442 circles include all beachings, whereas the lines with solid triangles include only  
443 beachings identified along shore. Beachings along shore and recoveries displaced to  
444 shore were separated via intersection with OpenStreetMap land polygons.


445 **Fig. 2** The number of French dFAD beachings recorded in our data per km of  
446 continental shelf edge in each  $5^\circ \times 5^\circ$  grid cell for the period 2008-2017. Darker areas  
447 indicate higher rates of beaching. In (a), all beachings are considered, whereas in (b)  
448 only beachings along shore are included. Beachings along shore and recoveries  
449 displaced to shore were separated via intersection with OpenStreetMap land polygons.  
450 Note that our dFAD trajectory data is incomplete before  $\sim 2010$ , so the absolute  
451 number of beachings per kilometer is likely somewhat higher than values shown in  
452 the figure, though differences are likely to be small as the number of dFADs was far  
453 lower before 2010 than after 2010.

454 **Fig. 3** Maps of the proportion of dFADs that beached within 3 months after passing  
455 through each  $1^\circ \times 1^\circ$  grid cell over the period 2008-2017. In (a), all beachings are  
456 considered, whereas in (b) only beachings along shore are included. Beachings along  
457 shore and recoveries displaced to shore were separated via intersection with  
458 OpenStreetMap land polygons. Note that the color intervals are unevenly distributed  
459 to highlight the low values.


460 **Fig. 4** Density maps representing the number of dFAD deployments in each  $1^{\circ} \times 1^{\circ}$  cell  
461 recorded in logbook data for the period 2013-2017. The thick, solid curves delimit  
462 areas representing the 20% of deployments most likely to produce a beaching within 3  
463 months of a dFAD passing through those areas. In (a), all beachings are considered,  
464 whereas in (b), only beachings along shore are included. Beachings along shore and  
465 recoveries displaced to shore were separated via intersection with OpenStreetMap  
466 land polygons.

467 **Fig. 5** Predicted reduction in beaching rate as a function of the amount of area put  
468 aside in annual (a-b) or quarterly (c-d) closures to dFAD deployments. Areas are  
469 closed from most likely to least likely to produce a beaching within 3 months of  
470 deployment, with area being quantified along the x-axis in terms of the fraction of  
471 deployments that occurred in closed areas prior to their closure. Black and grey dotted  
472 lines correspond to the null expectation of what the corresponding black and grey  
473 curves would look like if all areas had the same beaching risk, and are the same in the  
474 Indian and Atlantic Oceans. In (a) and (c), all beachings are considered, whereas in  
475 (b) and (d), only beachings occurring along shore are included. Beachings along shore  
476 and recoveries displaced to shore were separated via intersection with OpenStreetMap  
477 land polygons.

478 **Fig. 6** Map representing the predicted reduction in beaching when the 20% of dFAD  
479 deployments most likely to produce a beaching within 3 months are prohibited (see  
480 areas in Fig 4a), without (values on the left of the colorbar) and with (values on the  
481 right of the colorbar) dFAD deployment effort redistribution to non-prohibited areas.




**Fig. 1** (a) Annual number of new buoys deployed by the French and associated flags purse seine fleet in the Atlantic (grey) and Indian (black) oceans over the period 2008-2017 and (b) percentage of these buoys that beached. The lines in (b) with solid circles include all beachings, whereas the lines with solid triangles include only beachings identified along shore. Beachings along shore and recoveries displaced to shore were separated via intersection with OpenStreetMap land polygons.




**Fig. 2** The number of French dFAD beachings recorded in our data per km of continental shelf edge in each 5°x5° grid cell for the period 2008-2017. Darker areas indicate higher rates of beaching. In (a), all beachings are considered, whereas in (b) only beachings along shore are included.


Beachings along shore and recoveries displaced to shore were separated via intersection with OpenStreetMap land polygons. Note that our dFAD trajectory data is incomplete before ~2010, so the absolute number of beachings per kilometer is likely somewhat higher than values shown in the figure, though differences are likely to be small as the number of dFADs was far lower before 2010 than after 2010.



**Fig. 3** Maps of the proportion of dFADs that beached within 3 months after passing through each  $1^\circ \times 1^\circ$  grid cell over the period 2008-2017. In (a), all beachings are considered, whereas in (b) only beachings along shore are included. Beachings along shore and recoveries displaced to shore were separated via intersection with OpenStreetMap land polygons. Note that the color intervals are unevenly distributed to highlight the low values.




**Fig. 4** Density maps representing the number of dFAD deployments in each  $1^\circ \times 1^\circ$  cell recorded in logbook data for the period 2013-2017. The thick, solid curves delimit areas representing the 20% of deployments most likely to produce a beaching within 3 months of a dFAD passing through those areas. In (a), all beachings are considered, whereas in (b), only beachings along shore are included. Beachings along shore and recoveries displaced to shore were separated via intersection with OpenStreetMap land polygons.



**Fig. 5** Predicted reduction in beaching rate as a function of the amount of area put aside in annual (a-b) or quarterly (c-d) closures to dFAD deployments. Areas are closed from most likely to least likely to produce a beaching within 3 months of deployment, with area being quantified along the x-axis in terms of the fraction of deployments that occurred in closed areas prior to their closure. Black and grey dotted lines correspond to the null expectation of what the corresponding black and grey curves would look like if all areas had the same beaching risk, and are the same in the Indian and Atlantic Oceans. In (a) and (c), all beachings are considered, whereas in (b) and (d), only beachings occurring along shore are included. Beachings along shore and recoveries displaced to shore were separated via intersection with OpenStreetMap land polygons.





**Fig. 6** Map representing the predicted reduction in beaching when the 20% of dFAD deployments most likely to produce a beaching within 3 months are prohibited (see areas in Fig 4a), without (values on the left of the colorbar) and with (values on the right of the colorbar) dFAD deployment effort redistribution to non-prohibited areas.