Jellyfish degradation in a shallow coastal Mediterranean lagoon

Raquel Marques, Marta Rufino, Audrey M. Darnaude, Frédérique Carcaillet,
Marie Meffre, Delphine Bonnet

► To cite this version:

Raquel Marques, Marta Rufino, Audrey M. Darnaude, Frédérique Carcaillet, Marie Meffre, et al..
Jellyfish degradation in a shallow coastal Mediterranean lagoon. Estuarine, Coastal and Shelf Science,
2021, 261, pp.107527. 10.1016/j.ecss.2021.107527 . hal-03415608

HAL Id: hal-03415608

<https://hal.umontpellier.fr/hal-03415608>

Submitted on 10 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Jellyfish degradation in a shallow coastal Mediterranean lagoon

Raquel Marques^{1,2}, Marta Rufino^{3,4}, Audrey M. Darnaude¹, Frédérique Carcaillet¹, Marie Meffre¹, Delphine Bonnet¹*

¹ MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD, Montpellier, France

²German Center for Marine Biodiversity Research (DZMB), Senckenberg am Meer, Martin-Luther-King Platz 3, D-20146 Hamburg, Germany

³ Instituto Português do Mar e da Atmosfera (IPMA, I.P.), Divisão de Modelação e Gestão de Recursos Pesqueiros, Av. Dr. Alfredo Magalhães Ramalho, 6, 1495-165 Lisboa

⁴ Centro de Ciências do Mar (CCMAR), Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

***Corresponding author:** raquel.marques@umontpellier.fr / raquel.marques@senckenberg.de / marques.rfs@gmail.com

Present address : ²German Center for Marine Biodiversity Research (DZMB), Senckenberg am Meer, Martin-Luther-King Platz 3, D-20146 Hamburg, Germany

Key words: *Aurelia coerulea*, Macrobenthic community, Thau lagoon, Sediment, Seagrass

Present address : ²German Center for Marine Biodiversity Research (DZMB), Senckenberg am Meer, Martin-Luther-King Platz 3, D-20146 Hamburg, Germany

Abstract

Recurrent jellyfish blooms in the coastal zone call for understanding the impacts of jelly-falls on the functioning of benthic communities, especially in shallow enclosed ecosystems where their biomass can affect local carbon cycling and productivity. Each year, blooms of the jellyfish *Aurelia coerulea* appear and collapse in a semi-enclosed coastal Mediterranean lagoon (the Thau lagoon, south of France). Although the lagoon is shallow, large accumulations of dead jellyfish are never observed on its bottom, so it was hypothesized that decaying jellyfish were rapidly consumed by local macrobenthic organisms. The current work aimed to test this hypothesis, by estimating the impact of the presence of dead *A. aurelia* medusae on local macrobenthic community composition and assessing their biomass loss rates under different scenarios of accessibility by the macrobenthos. Unexpectedly, our results revealed a limited role of macrobenthic scavengers in the disappearance of dead medusae, although this later was particularly fast (19 to 78h). Only one taxon (*Tritia* sp., Nassariidae family) showed a significant response to the presence of dead *A. coerulea* medusae on the seabed. Thus, our results suggest that the fast disappearance of dead jellyfish biomass in Thau results from its rapid degradation and consumption by local microorganisms, likely due to the combined effects of high local temperatures and the small size of *A. coerulea* medusae. Thus, the important biomass produced during *A. aurelia* blooms in Thau might essentially boost its microbial food web. The potential role of jellyfish blooms in controlling biogeochemical cycles and food web functioning in shallow lagoons is discussed, underlying the need to include this process in ecosystem-based models.

1. Introduction

Jellyfish (in particular scyphozoans) are famous for their conspicuous blooms, which may locally generate biomasses exceeding 10 t wet weight 100 m^{-3} (Lilley et al. 2011). The population dynamics of jellyfish at the pelagic stage is frequently described as ‘bloom and bust’, since jellyfish blooms collapse rapidly, usually within a few weeks or months (Pitt et al. 2014). This might cause large accumulations of sinking dead jellyfish (referred to as *jelly-falls*) on the seafloor (Lebrato et al. 2012), which can be particularly impressive, especially in deep-sea habitats (Billett et al. 2006; Yamamoto et al. 2008; Lebrato and Jones 2009; Sweetman and Chapman 2011, 2015) where they can form localized layers of up to 70 cm in thickness (78 g C m^{-2}) on the seabed (Billett et al. 2006). These massive accumulations of jelly-falls are likely sporadic and considered as anomalies when compared with the mean annual jellyfish fluxes in those regions (Luo et al. 2020). However, jellyfish are common worldwide and represent a global biomass of 290 Tg C (Luo et al. 2020) with very high sinking speeds (Lebrato et al. 2013) which calls for a better understanding of the impacts of jelly-falls on the functioning and productivity of benthic communities.

When jellyfish blooms collapse, this accumulated organic matter has several possible fates. First, it can be consumed or fragmented by pelagic predators and scavengers (Cardona et al. 2012; Bos et al. 2017; Hays et al. 2018; Marques et al. 2019). Otherwise, carcasses sink through the water column (Lebrato et al. 2012) where they can be degraded by pelagic microbial communities (Titelman et al. 2006; Blanchet et al. 2015; Tinta et al. 2016, 2020). The amount of jellyfish biomass that reaches the seafloor depends, thus, on their decay rate, the sinking speed of the carcasses, the depth at which the jellyfish die, and the depth of the water column itself (Lebrato et al. 2011, 2019). Decay rates for jellyfish depend on the temperature (Lebrato et al. 2011), while their sinking speed is a function of their size, diameter, bio-volume, geometry, density, and drag coefficients (Yamamoto et al. 2008; Lebrato et al. 2011, 2012, 2013). If not degraded in the water column (Titelman et al. 2006; Tinta et al. 2016, 2020), jelly-falls accumulate on the seabed, with potentially important impacts on both the biogeochemical cycling and the functioning of benthic ecosystems (Sweetman et al. 2016). Jellyfish biomass tend to be rapidly degraded because it is

characterized by protein-rich organic matter, low C:N ratio, and no hard exoskeleton, being described as highly bioavailable for some particular microbial organisms with very high growth efficiency (Tinta et al. 2012, 2020). This may lead to high consumption of dissolved oxygen and a drastic decrease of its concentrations in the vicinity of jellyfish carcasses (West et al. 2009; Sweetman et al. 2016; Chelsky et al. 2016; Guy-Haim et al. 2020), which might induce inhospitable conditions for the benthic macrofauna, decreasing its activity, causing local emigrations or even massive mortalities (Sweetman et al. 2016; Chelsky et al. 2016). Nevertheless, dead jellyfish can also potentially provide suitable food for many benthic species, including fishes, echinoderms, anthozoans, polychaetes, gastropods, and crustaceans (Lebrato et al. 2012; Sweetman et al. 2014; Chelsky et al. 2016; Ates 2017). These later can consume considerable amounts of jelly-falls biomass within few hours, which considerably boosts benthic productivity (Sweetman et al. 2014). This scavenging behaviour plays a key role in benthic ecosystem functioning as it determines the fate of the organic matter that reaches the seafloor, *i.e.* whether the organic material from jelly-falls contributes to the microbial loop or enters the macrofaunal food web (Sweetman et al. 2014, 2016).

So far, the microbial degradation of jelly-falls has been relatively well described (Titelman et al. 2006; West et al. 2009; Tinta et al. 2010, 2012, 2020; Condon et al. 2011; Frost et al. 2012; Blanchet et al. 2015; Sweetman et al. 2016). However, studies investigating the impact of jelly-falls on benthic macrofaunal communities are still scarce (Sweetman et al. 2014; Chelsky et al. 2016; Dunlop et al. 2017). Such studies are imperative for coastal habitats, where the highest jellyfish biomass values were reported (Luo et al. 2020) and the anthropogenic impacts are intense and pointed out as likely promoters of jellyfish blooms (Purcell 2012). This is particularly evident in coastal lagoons, where jellyfish blooms occur regularly (e.g. Fuentes et al., 2011; Marques et al., 2015a), with abundances that can overcome 530 tonnes km⁻² (Pitt and Kingsford 2003). These ecosystems are very productive enclosed systems, supporting important ecological processes and providing numerous ecosystem services (Newton et al. 2014; De Wit et al. 2017), but very little information is available regarding the fate of jelly-falls within these shallow environments (Chelsky et al. 2016). Despite the frequent occurrence of jellyfish blooms, to our knowledge,

massive accumulations of jelly-falls on coastal lagoons' floor were never reported. Although jellyfish can be rapidly degraded by pelagic microbial communities (Tinta et al. 2020), jelly-falls should still be observed on lagoons' seabed because, in such shallow habitats, sinking jellyfish are likely to reach the bottom even before they die (Lebrato et al. 2012). Therefore, the absence of dead jellyfish on the seabed in these environments suggest that jelly-falls are rapidly eaten by local benthic scavengers and/or decomposed by local microorganisms.

The current work aims to investigate this issue and evaluate the contribution of macrobenthic scavengers in the disappearance of jellyfish biomass in coastal lagoons. To this aim, we studied the fate of dead medusae of the jellyfish *Aurelia coreulea* when ending on the seafloor in a shallow lagoon located in the south of France: the Thau lagoon. *In situ* experiments were carried out by adding dead medusae on the seabed of the lagoon under different scenarios of accessibility by macrobenthic scavengers. The experiments were performed in two contrasting types of habitat, typical of this shallow ecosystem, assuming that the different macrobenthic community composition associated with each habitat would have a different impact on the biomass loss rate of the jelly-falls. In particular, we tested whether if the addition of dead jellyfish on the seabed altered the composition of macrobenthic communities by, for instance, attracting benthic scavengers (Chelsky et al. 2016).

2. Material and Methods

2.1. Study site and jellyfish collection

The Thau lagoon (43°25'31.1"N; 03°42'0.9"E) is a semi-enclosed coastal lagoon of 75 km², connected to the Mediterranean Sea by three narrow channels. It is shallow, with a mean of 4 m depth, and is highly influenced by strong wind events (Fiandrino et al. 2012). The local tidal range is weak (< 1 m), which promotes a high water residence time (1 - 4 months, Fiandrino et al., 2012). With regards to jellyfish, the Thau lagoon has the particularity to harbour its own population of *Aurelia coerulea*, isolated from those in the Mediterranean Sea (Bonnet et al. 2012; Marques et al. 2015b) which offers a rare occasion to study the fate of the blooms of this species. In the lagoon, *A. coerulea* ephyrae first appear in the early winter (November – December), to

give rise to medusae at the beginning of spring (April – May), when temperature increases (Marques et al. 2015a). High abundances of medusae, associated with high growth rates generate the annual jellyfish bloom, which usually collapses in the early summer (June-July). Although sparse decaying medusae are regularly seen on the lagoon floor, either on bare sediment or entangled in seagrass leaves (R. Marques, personal observation), large accumulations of *A. coerulea* carcasses have never been observed so far.

To try to elucidate the local fate of *A. coerulea* jelly-falls, two different *in situ* experiments were performed: one (1) to assess medusae biomass loss rates under different scenarios of medusae accessibility for benthic scavengers, and one (2) to study the impact of jellyfish presence on the seafloor on the composition of local macro-benthic communities (Fig. 1). Both experiments were performed in a shallow area (< 1m depth), where dead jellyfish had already been observed on the seabed and repeated in the two most common habitats found in the lagoon (Plus et al. 2003): on bare sediments and in seagrass (*Zostera noltii*) meadows.

This study was conducted in 2018, during the collapse of the annual bloom of *A. coerulea*. Due to logistic constraints, the two experiments were performed on different days (on May 30th and June 07th, 2018). However, all the medusae used were collected alive on the same day (May 28th, 2018). This was done using hand nets to avoid damaging the medusae and they were immediately transported to the laboratory in ambient seawater. All medusae were then kept alive for 2 to 10 days, in 1 m³ tanks (ca. 100 ind.m⁻³) with open seawater circulation system (i.e. seawater from the lagoon) to ensure similar rearing conditions as *in situ*. They were all fed once per day with newly hatched *Artemia* to ensure their survival. A few hours before each experiment, live, healthy and active medusae were selected from the husbandry tanks and equally distributed in 30 L cold boxes, filled with ambient seawater. Medusae were killed by sparging the water with nitrogen gas for ca. 3h following Chelsky et al. (2016). The medusae were then immediately transported (20 min) to the experimental site and placed in the experimental bags.

Fig. 1: Schematic representation of the two *in situ* experiments performed in this study: a) assessing jellyfish biomass loss rates under different scenarios of medusae accessibility for benthic scavengers; b) assessing the impact of jellyfish degradation on the macrobenthic community composition.

2.2. Jellyfish biomass loss rates

2.2.1. Experimental set-up

Jellyfish biomass loss rates were assessed both on the bare sediment and seagrass meadows habitats under two different scenarios of accessibility to dead medusae for the macrobenthic organisms of the lagoon (Fig.1a).

The first scenario (Micro) involved placing individual medusae in 20×15 cm net bags with a mesh size of $200 \mu\text{m}$ so they were accessible only to microorganisms (*e.g.* bacteria, microzooplankton, and small mesozooplankton species). In the second scenario (Macro), medusae were placed in 20×15 cm net bags with a mesh size of 1 cm, which allowed both microorganisms and macroorganisms (*e.g.* gastropods, amphipods, crustaceans) to access them. In both scenarios, the net bags containing the dead medusae were protected with a $1.50 \times 2.00 \times 0.15$ m net cage with a coarse mesh size of 2.5 cm, to prevent medusae consumption by large organisms (*e.g.* large fish, echinoderms, crustaceans). However, the Macro scenario was also replicated without the protection net cage to assess if medusae consumption by large scavengers was significant. Since no significant effect of the cage could be evidenced (generalised nonlinear least square model, $p\text{-value} = 0.17$ and 0.62 for bare sediment and seagrass habitats, respectively), data from Macro scenarios with and without cage were pooled.

Before each experiment, dead medusae were partially dried on a paper towel to absorb the excess water and mucus from their surface, weighted (wet weight in g, to the nearest 0.1 g), measured (bell diameter in cm), and placed individually in a bag. In each scenario, 24 bags were fixed on the sediment using tent pegs, with a minimum distance of 1.5 m between them. For each of the two scenarios (Micro and Macro) and irrespective of the habitat (bare sediment or seagrass meadow), three replicates of individual medusa were collected at regular time intervals: every 1h at the beginning of the experiment (when medusae degradation is usually faster, Titelman et al., 2006), and every 2 to 5h afterwards, adjusted in each habitat type to ensure all the medusae biomass had disappeared from the net bags by the end of the experiment. The total experimental time was, therefore, different between habitats (24h and 30h for the seagrass meadows and the bare sediment habitats, respectively). At each sampling time, collected bags were immediately placed inside a hermetic plastic bag to avoid the loss of biological material. The remaining medusa biomass within each bag was partially dried on a paper towel to absorb the excess water (when possible) and weighed to the nearest 0.1 g (wet weight). *In situ* temperature was measured (EC 300 VWR international/ WTW model 350i) right before the start and end of each experiment to detect differences in temperature between habitats.

2.2.2. Data analysis

For each habitat (bare sediment and seagrass meadow) and accessibility scenarios (Micro and Macro), jellyfish biomass loss rates were determined by fitting exponential decay models to the non-averaged wet weights of medusae (percentage of the initial biomass) as a function of time:

$$M_t = M_0 e^{-\lambda t}$$

where M_t is the percentage of medusa biomass at time t (in hours), M_0 and λ are the model coefficients representing the initial ($t = 0$) medusa biomass (in percentage) and the loss rate, respectively. The biomass loss rates were then used to calculate the degradation time (D_t in hours), *i.e.* the time required to achieve a loss of 50% ($t = 0.5$) and 99% ($t = 0.01$) of the initial biomass of medusa, according to the following equation (Lebrato et al. 2011):

$$D_t = \frac{-\ln(t)}{\lambda}$$

Differences, according to the accessibility scenarios and the habitats, were tested by fitting generalised nonlinear least square models (GNLS) using “nlme” package (Pinheiro et al. 2019), which allows fitting the model to zero values, using 100 and 0.01 as starting parameters, for M_0 and λ , respectively.

Significant differences in the initial medusae biomass and environmental conditions (temperature) between habitats and treatments were assessed by Kruskal-Wallis and T-tests, after verifying the normality assumptions.

2.3. Impact on macrobenthic community composition

2.3.1. Experimental set-up

To test the impact of dead jellyfish presence on benthic community composition, a second experiment was carried out on both habitats.

The experiments started (t_0) at 16h30 and 15h40 on the seagrass meadows and the bare sediment, respectively, and samples were collected at five sampling times (every 3h) for 15h (Fig. 1b), based on the preliminary results of the jellyfish biomass loss rates experiments. In each habitat, three different treatments were performed. The *medusa* (M) treatment was a replication of the Macro

scenario from the previous experiment: one dead medusa was placed within a 1 cm mesh net bag, thereby being accessible to both micro- and macroorganisms. The *procedure control* (PC) treatment aimed to test the effect of the experimental setup and therefore the M treatment was reproduced without any medusa in the net bag. The last treatment was for *control* (C). In this case, the sampling was performed on undisturbed areas of each habitat. To assess differences in macrobenthic community composition between treatments, the substrate (sediment and seagrass) below each bag was collected, as well as the organisms present on its surface and top of the bags. This sampling was performed immediately upon medusae (in M) or empty bag (in PC) collection. Three replicates were collected per combination of habitat, treatment, and sampling time. In each case, the sediment was sampled using a shovel (0.03 m², 4 cm deep) and placed inside a hermetic plastic bag, ensuring a minimum sample loss. Samples were stored in cold boxes and frozen within 6h, until later laboratory analysis. *In situ* temperature was measured (EC 300 VWR international/ WTW model 350i) before the start of the experiment and at each sampling time to detect differences in temperature between habitats. Once in the laboratory, the volume of sediment in each sample was measured using a graduated beaker to standardize the abundance of organisms by sampling area (m²). The sediment was sieved (1 mm mesh size) and its macrofauna was sorted, counted, and identified under a dissecting microscope according to D'Angelo and Gargiullo (1978), Fauvel (1927), and Fauvel (1923). The organisms were identified to the lowest taxonomic level. However, since identification at the species level was not possible for all organisms, species of the same genus were grouped. Annelids and Decapods were identified down to the family level only.

2.3.2.Data analysis

Only taxa representing more than 1% of the total community biomass in each habitat were considered for data analysis, to reduce the influence of rare organisms. Diversity indices (Shannon and Pielou's evenness indices) were calculated using the "BiodiversityR" package (Kindt and Coe 2005), based on "vegan" package in R (Oksanen et al. 2019). The changes in total abundance (after logarithmic transformation) and diversity indices, among habitats, scenarios, and sampling

times were tested using linear models. For each variable (*i.e.* total abundance, Shannon, and Pielou's evenness indices) a full model was produced, with all main terms and respective interactions (index ~ habitat * treatment * sampling time). Model selection was then carried out using the Akaike information criterion (AIC), following Zuur et al. (2009). Visual inspection of residual plots did not reveal any obvious deviations from homoscedasticity or normality. Differences between each combination of treatment and sampling time within each habitat were tested using post hoc Tukey HSD tests for multiple comparisons.

Changes in community composition among habitats, treatment, and sampling time were analysed using three different complementary approaches. First, the community composition was represented through a principal component analysis (PCA) of the abundances ($\log(x+1)$). Second, differences between community composition among habitats, treatments, and sampling times were assessed through a permutational multivariate analysis of variance (PERMANOVA, with 9999 permutations), using Bray-Curtis distance. Since homogeneity of dispersion between factors is an assumption of the PERMANOVA analysis, multivariate dispersion was first tested using BETADISPER. When significant differences were observed, a pairwise comparison was performed (PERMUTEST, with 9999 permutations). Nonmetric Multidimensional Scaling (NMDS) plots were used to visualize the results of BETADISPER and PERMANOVA, as recommended (Anderson 2017). These analyses were performed using the package "vegan" (Oksanen et al. 2019).

Third, to cope with the limitations reported for PERMANOVA, which does not take into consideration the influence of the mean-variance structure of each species (Warton et al. 2012), we also ran a model-based approach using a multi-taxa generalized linear model (ManyGLM, "manyglm" function, from package "mvabund"; Wang et al., 2012). A two fixed factor model structure (sampling time and treatment) was used, separately for each habitat, with a negative binomial distribution and a log-link function. The examination of residual plots of the model showed the absence of a clear pattern, validating the model. This analysis was also used to determine which species contributed most to the differences observed.

3. Results

3.1. Jellyfish biomass loss rates

The initial wet weights of the *A. coerulea* medusae used in the experiments (Table 1) were similar among scenarios within each habitat (Kruskal-Wallis, $\chi^2 = 0.32$, $df = 1$, p -value = 0.57 and $\chi^2 = 0.61$, $df = 1$, p -value = 0.44 for bare sediment and seagrass, respectively), but differed between habitats (Kruskal-Wallis, $\chi^2 = 81.33$, $df = 1$, p -value < 0.001). Indeed, as the experiments started on different days for the two habitats and medusae were kept alive between the two experiments, differences in the time spent in captivity likely induced a bias on the initial weight of the medusae. The initial and final experimental temperatures were similar for the two habitats (23.5 ± 0.7 °C and 23.4 ± 0.5 °C on the seagrass meadows and bare sediment, respectively; T-test, p -value = 0.8), suggesting a limited effect of temperature on the results.

The disappearance of *A. coerulea* biomass was fast irrespective of the scenario or habitat (Fig. 2), with λ coefficient ranging from -0.24 to -0.06 per hour (*i.e.* -1.42 to -5.8 per day, Table 1). All model fits and correspondent coefficients were statistically significant (GNLS, p -value < 0.001, Table 2). On the bare sediment habitat, the jellyfish biomass loss rate was not affected by the accessibility scenarios (GNLS, p -value = 0.798), while on the seagrass meadow the biomass loss rate was significantly higher (GNLS, p -value < 0.001) in the Micro scenario than in the Macro one. Irrespective of the scenario, medusae degradation was significantly faster on the bare sediments (GNLS, p -value < 0.001), where 99% of the initial medusae biomass was lost in about 19 hours in both accessibility scenarios (Table 1). Medusae biomass loss was slower on seagrass meadows: under the Macro scenario, R_t was estimated at 78 h, while microorganisms alone degraded 99% of the biomass in 32 h. However, medusae biomass loss was consistently faster during the first few hours of the experiments, with a 50% loss of the initial biomass in 3h on bare sediment (both scenarios), against 5 and 12h on seagrass meadows, for Micro and Macro scenarios, respectively.

Table 1: Mean initial wet weights (W) and bell diameters (BD) of the dead *A. coerulea* medusae used in each experiment and resulting estimates of biomass loss rate (λ in hours) and degradation time (Dt in hours).

Scenario	Initial W (g $\pm$ SD)	Initial BD (cm $\pm$ SD)	λ (h)	Dt (h)
----------	------------------------	--------------------------	---------------	----------

	Bare sediment			
Macro	34.8 ± 12.4	8.0 ± 1.3	-0.24	19.51
Micro	32.8 ± 14.9	8.1 ± 1.5	-0.24	19.05
	Seagrass meadows			
Macro	107.5 ± 20.0	12.8 ± 0.9	-0.06	78.04
Micro	111.8 ± 19.0	13.2 ± 1.0	-0.15	31.56

Fig. 2: Dynamic of *A. coerulea* medusae biomass loss in both habitats (bare sediment and seagrass meadows) under both accessibility scenarios (Macro and Micro). Exponential decay models (lines) were fitted to the non-averaged data of biomass (in % of the initial medusae biomass), with 95% confidence intervals (shadow areas).

Table 2: Estimation of the parameters (M_0 and λ) by the GNLS models used to assess differences between scenarios within each habitat. Significant differences (p -value < 0.05) are indicated in bold.

	Value	Std.Error	t-value	p-value
Bare sediment				
M_0				
Macro (Intercept)	99.514	2.944	33.805	< 0.01
Micro	-1.228	5.025	-0.244	0.807
λ				
Macro (Intercept)	0.236	0.013	18.509	< 0.01
Micro	0.006	0.023	0.257	0.798
Seagrass meadows				
M_0				
Macro (Intercept)	92.834	2.511	36.974	< 0.01
Micro	8.045	6.095	1.320	0.192
λ				

Macro (Intercept)	0.059	0.004	14.525	< 0.01
Micro	0.087	0.017	5.142	< 0.01

3.2. Benthic community changes

3.2.1. General composition of macrobenthic communities

A total of 9478 macrobenthic organisms, belonging to 34 different taxa, were identified during the study. The two types of habitats investigated differed in terms of species richness with a higher average number of taxa on seagrass meadows (29) than on bare sediments (20). However, macrobenthic communities on seagrass meadows were clearly dominated by nine taxa only (Fig. 3). On this type of habitat, the gastropods *Bittium* sp. and the bivalves *Ruditapes* sp. represented together more than 80% of the total abundance recorded in all treatments: Medusae (M, 67.8 and 17.4%, respectively), Procedure Control (PC, 58.5 and 25.4%, respectively) and Control (C, 41.6 and 41.3%, respectively). On bare sediments, macrobenthic communities were more balanced: in the C treatment, 81.7% of the total abundance was represented by the annelid Glyceridae (28.2%), and the gastropods *Bittium* sp. (23.6%), *Tricolia* sp. (17.9%), and *Rissoa* sp. (12.1%); in the M treatment, the contribution of Glyceridae dropped to 12.5%, while taxa like *Ruditapes* sp. and *Tritia* sp. increased their importance representing 15.8% and 8.9% of total abundance, respectively; and in the PC treatment the most abundant taxa were *Bittium* sp. (24.4%), *Tricolia* sp. (16.5%), *Rissoa* sp. (15.4%) and Glyceridae (12.2%).

Fig. 3: Contribution of the most important taxa to the total abundance of the community on the seagrass meadows and bare sediment, in each treatment (M: Medusae, C: Control, PC: Procedure control). Only species that contributed to more than 1% of the total abundance are presented.

3.2.2. Abundance and diversity indices

The abundance of macrobenthic organisms was, on average, 10 times higher (p-value < 0.001, Fig.4A and B, Table 3) on the seagrass meadows than on the bare sediments ($6\,800 \pm 8830$ and $637 \pm 717 \text{ ind.m}^{-2}$, respectively). It was significantly affected by all factors considered (habitat, sampling time, and treatment, Table 3), but the overall interaction among these factors was not significant, indicating that treatment and time similarly affected the abundance of organisms within each habitat. Differences in total macrobenthic abundance among treatments were only observed at 3h on the seagrass meadows, and 3 and 9h on the bare sediments (Tukey HSD, p-value < 0.05; Fig. 4A and B). This was particularly evident at 3h, when a peak of macrobenthos abundance was detected for both M and PC (both over 1500 ind.m^{-2} and 190 ind.m^{-2} , on seagrass meadows and bare sediment, respectively), with significantly higher values than in C (Tukey HSD, p-value < 0.05). However, no differences in abundance were observed between M and PC (Tukey HSD, p-value > 0.05) over the whole experiment time. In both habitats, the abundance of macrobenthic organisms in the controls did not vary significantly over time (Tukey HSD, p-value

> 0.05), except at 12h in the seagrass habitat, when it was significantly lower than at t_0 (Tukey HSD, p-value = 0.02) (Fig. 4A and B).

The diversity of macrobenthic organisms (Shannon diversity index) appeared to vary differently depending on the habitat (Fig. 4C and D, Table 3). In seagrass meadows, it increased significantly at the end of the study period (9, 12, and 15h; Tukey HSD, p-value < 0.05; Fig. 4C), but did not vary among treatments while, on bare sediments, differences were only observed among treatments, with higher diversities in M and PC than in C at 3h (Tukey HSD, p-value = 0.01; Fig. 4D).

For Pielou's evenness index, only the sampling time and habitat factors were retained in the linear model (Table 3), showing that the treatment did not affect community evenness. Differences in the evenness were only observed on the seagrass meadows, where it was lower at 3h, suggesting a possible disturbance of the community (Tukey HSD, p-value < 0.05; Fig. 4E and F).

Table 3: Results of the linear models and the effect of each factor (Habitat, Time, and Treatment), on each variable (Abundance, Shannon, and Evenness diversity indices). Bold values indicate significant differences between at least two groups, at $\alpha = 0.05$.

Abundance	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Time	5	18.294	3.659	8.687	<0.001
Treatment	2	24.407	12.204	28.974	<0.001
Habitat	1	156.490	156.490	371.541	<0.001
Time:Treatment	8	19.163	2.395	5.687	<0.001
Time:Habitat	5	8.792	1.758	4.175	0.002
Treatment:Habitat	2	4.745	2.372	5.633	0.006
Time:Treatment:Habitat	8	6.233	0.779	1.850	0.084
Shannon					
Time	5	1.304	0.261	1.853	0.115
Treatment	2	2.986	1.493	10.610	<0.001
Habitat	1	2.528	2.528	17.962	<0.001
Time:Treatment	8	2.195	0.274	1.949	0.068
Time:Habitat	5	3.180	0.636	4.519	0.001
Treatment:Habitat	2	2.612	1.306	9.281	<0.001
Time:Treatment:Habitat	8	2.176	0.272	1.933	0.070
Evenness					
Time	5	0.201	0.040	6.956	<0.001
Habitat	1	2.332	2.332	403.462	<0.001
Time:Habitat	5	0.085	0.017	2.936	0.017

Fig. 4: Abundance of macrobenthic organisms (A and B, note the different scales for the two habitats), Shannon diversity index (C and D), and Pielou's equitability index (Evenness) on the seagrass meadow (A, C, and E) and bare sediment (B, D, and F) habitats. For each variable, dots represent average values and vertical bars standard deviations for each treatment (M: medusae, PC: procedure control, C: control). Asterisks indicate significant differences between treatments at $\alpha = 0.05$.

3.2.3. Differences in the community composition

In the principal component analysis (PCA), only the first two axes (PC1 and PC2, Fig. 5) were retained since they represent the majority of the variability of the data (69.7%). These axes discriminated the samples from the two habitats (Fig. 5A), showing that habitat type is the main

383 driver of the variation in macrobenthic community composition observed between samples. In
 384 seagrass meadows, the community was characterized by high abundances of *Bittium* sp. (B),
 385 *Ruditapes* sp. (Ru), and *Loripes* sp. (L), whereas on the bare sediment Glyceridae (Gl), *Rissoa* sp.
 386 (R), and *Tricolia* sp. (Tri) highly contributed to differentiate these groups. The effect of the
 387 treatment or time on the community composition was not evident in the PCA (Fig. 5B and C).
 388 However, 17 samples presented high Euclidean distance from the centre (outside the grey lines,
 389 Fig. 5A, B, and C), suggesting that they had a different community composition. Most of these
 390 samples (10) were collected at 3h (Fig. 5C) and only 3 were C samples (Fig. 5B).

391
 392 Fig. 5: Biplots of Principal Component Analysis (PCA). Samples (points) and taxa (arrows) are presented. In A, B,
 393 and C, samples are identified according to habitat, treatment (M: medusae, PC: procedure control, C: control), and
 394 sampling time (in h), respectively. Grey dotted lines are indicative thresholds to identify the samples with high
 395 Euclidean distance from the centre. For the sake of simplicity, taxa names are abbreviated (see Fig. 3 with taxa codes).
 396

397 The results from BETADISPER did not detect significant changes in community dispersion over
 398 time ($F = 0.7$, $p = 0.6$) or among treatments ($F = 2.5$, $p = 0.08$). However, the community dispersion

was different between the two habitats ($F= 13.2$, $p\text{-value} < 0.001$), and, therefore, BETADISPER and PERMANOVA were performed for each habitat separately.

On the seagrass meadows, community dispersion was homogeneous among treatments ($F = 0.5$, $p\text{-value} = 0.6$, Fig. 6A), but not across sampling time ($F = 2.5$, $p\text{-value} = 0.045$, Fig. 6B). Pairwise comparisons identified the samples collected at 3h as being different from those collected from 6 to 15h ($p\text{-value} < 0.05$). The analysis was, therefore, repeated without the samples collected at 3h, to ensure homogeneity of dispersion among samples ($F = 0.3$, $p\text{-value} = 0.9$). Accordingly, a PERMANOVA was performed to determine changes in the community for the seagrass meadow, between treatments, sampling times, and the respective interaction (full model), omitting the samples collected at 3h. The results indicate that the community composition did not vary between treatments ($F = 0.6$, $p\text{-value} = 0.8$, Fig. 6A), but showed significant differences over time ($F= 9.3$, $p\text{-value} < 0.01$, Fig. 6B).

In the bare sediment habitat, community dispersion was homogeneous across sampling times and among treatments ($F = 0.6$, $p\text{-value} = 0.7$ and $F = 0.4$, $p\text{-value} = 0.7$, respectively, Fig. 6C and D). Therefore, a full factorial model of PERMANOVA (*i.e.* treatment, sampling time, and interaction) was performed for this habitat. The results show that both factors significantly affected macrobenthic community composition ($F = 2.8$, $p\text{-value} < 0.01$ and $F = 4.8$ and $p\text{-value} < 0.01$, respectively), with PC and M presenting similar values, whereas treatment C was significantly different (Fig. 6C). However, the interaction between both factors was not significant ($F = 1.4$ and $p\text{-value} = 0.09$) indicating that changes in community composition among treatments were not affected by sampling time.

Fig. 6: Results from the Nonmetric Multidimensional Scaling (NMDS) analysis on the seagrass meadows (A and B) and the bare sediment habitats (C and D), showing the dispersion of the samples (lines) around the centroid of each group, by treatment (A and C): M: medusae, PC: procedure control, C: control; and by sampling time in hours (B and D).

The results of the two-factor multivariate linear model (mvabund), for each habitat, were consistent with the results of the PERMANOVA (reported above). In the seagrass meadows, the treatment alone did not affect the community composition, but it was significant when combined with sampling time (significant interaction, Dev = 175.6, p-value = 0.007). On the bare sediment, both the treatment and sampling time affected the community composition (p-value < 0.05).

This analysis allowed the identification of the taxa that contributed most to the observed differences between treatments (Table 4): *Bittium* sp. in the seagrass meadows and *Tricolia* sp., *Hexaplex* sp., *Tritia* sp. and *Ruditapes* sp. on the bare sediment habitats (Fig. 7). These organisms exhibited higher abundances in M and PC at 3h but no difference of abundances between M and PC treatments were observed ($P_{adj} > 0.05$). Indeed, differences between M and PC treatments were only observed for *Tritia* sp. ($P_{adj} = 0.04$), which showed higher abundances for M during the first 9h of the study period (up to $14.9 \pm 3.6 \times 10 \text{ ind.m}^{-2}$, Fig. 7D). Therefore, although *Bittium*

sp. (in seagrass meadows), *Hexaplex* sp., *Tricolia* sp., and *Ruditapes* sp. (on the bare sediment) appeared to have positively responded to the presence of jelly-falls, especially at 3h, only the *Tritia* sp. (on the bare sediment) revealed statistical evidence of a positive response to the presence of dead *A. coerulea* medusae on the bottom.

Table 4: Results of the ‘species-by-species’ two-factor multivariate linear model (ManyGLM), with the terms and the significance of each term (adjusted p-values) in the model (treatment, sampling time, and interaction). Bold values indicate significant differences at $\alpha = 0.05$.

	Treatment		Sampling time		Treatment: Sampling time	
	Dev	P _{adj}	Dev	P _{adj}	Dev	P _{adj}
Seagrass meadows						
<i>Cerastoderma</i> sp.	0.433	0.982	26.982	0.008	5.557	0.945
<i>Gastrana</i> sp.	0.676	0.982	18.624	0.055	5.012	0.945
<i>Loripes</i> sp.	1.876	0.958	60.183	0.001	19.451	0.359
<i>Ruditapes</i> sp.	0.249	0.982	41.143	0.001	9.649	0.918
<i>Bittium</i> sp.	7.987	0.449	41.265	0.001	40.549	0.001
<i>Cerithium</i> sp.	2.013	0.958	7.663	0.768	4.902	0.945
<i>Gibbula</i> sp.	6.677	0.548	20.578	0.030	25.757	0.100
<i>Hexaplex</i> sp.	3.726	0.936	7.681	0.768	7.366	0.935
<i>Jujubinus</i> sp.	0.000	1.000	0.000	1.000	0.000	1.000
<i>Rissoa</i> sp.	3.409	0.941	9.682	0.555	8.813	0.918
<i>Tricolia</i> sp.	7.972	0.449	6.472	0.768	7.810	0.935
Gastropoda NI	0.000	1.000	0.000	1.000	0.000	1.000
<i>Sphaeroma</i> sp.	2.843	0.941	11.816	0.381	15.583	0.582
<i>Microdeutopus</i> sp.	3.262	0.941	4.412	0.850	8.919	0.918
Cirratulidae	2.402	0.941	3.750	0.850	0.001	0.945
Glyceridae	0.000	1.000	0.000	1.000	0.000	1.000
<i>Tritia</i> sp.	1.452	0.958	14.332	0.214	16.203	0.582
Bare sediment						
<i>Cerastoderma</i> sp.	0.391	0.843	7.686	0.661	9.641	0.794
<i>Gastrana</i> sp.	0.000	1.000	0.000	1.000	0.000	1.000
<i>Loripes</i> sp.	1.764	0.713	3.137	0.948	6.730	0.794
<i>Ruditapes</i> sp.	24.699	0.001	18.268	0.036	8.739	0.794
<i>Bittium</i> sp.	9.271	0.109	32.619	0.001	20.626	0.140
<i>Cerithium</i> sp.	6.039	0.395	2.509	0.948	4.872	0.794
<i>Gibbula</i> sp.	7.441	0.251	15.723	0.067	8.634	0.794
<i>Hexaplex</i> sp.	12.639	0.024	3.401	0.948	1.955	0.794
<i>Jujubinus</i> sp.	6.794	0.313	17.327	0.048	10.152	0.794
<i>Rissoa</i> sp.	8.154	0.177	22.319	0.014	19.967	0.145
<i>Tricolia</i> sp.	4.182	0.576	16.865	0.051	25.929	0.046
Gastropoda NI	4.280	0.576	15.542	0.067	2.883	0.794
<i>Sphaeroma</i> sp.	0.000	1.000	0.000	1.000	0.000	1.000
<i>Microdeutopus</i> sp.	4.852	0.567	5.147	0.892	0.000	0.830
Cirratulidae	4.720	0.567	24.968	0.003	8.259	0.794

Glyceridae	3.377	0.596	5.275	0.892	10.271	0.794
<i>Tritia</i> sp.	21.770	0.001	12.924	0.136	12.333	0.638

Fig. 7: Abundance of the macrobenthic organisms which showed significant contributions to the observed differences among treatments and sampling time (ManyGLM) on bare sediment (A to D) and seagrass meadows (E) habitats. Note the differences in scale between graphs.

4. Discussion

4.1. The contribution of macrobenthic consumption to the fast degradation of jellyfish in *Thau*

Local scavengers might play a significant role in jelly-falls disappearance from the benthic environments by consuming jellyfish biomass that sinks or accumulates on the seafloor (Sweetman et al. 2014). In our study, though, irrespective of the habitat, macrobenthic

communities appear to have a limited impact on the biomass loss of dead *A. coerulea* medusae in the Thau lagoon. This is supported by the lack of significant differences in jellyfish biomass loss rates between the Micro and Macro scenarios and by the limited response of the benthic communities to the addition of jelly-falls to the seabed (except for one particular taxon, the *Tritia* sp.). Therefore, our results suggest that, in the Thau lagoon, jelly-falls are likely mainly degraded by microorganisms (*e.g.* bacteria, microzooplankton, and small mesozooplankton species) with a limited contribution of macrobenthic consumption in their disappearance.

The *in situ* degradation of *A. coerulea* medusae in the Thau lagoon was very fast, with biomass loss rates ranging from 0.06 to 0.24 h⁻¹ (1.42 to 5.8 d⁻¹) and a 99% degradation of dead jellyfish biomass in about 19 hours on bare sediments and less than 3.5 days on seagrass meadows. High jellyfish biomass loss rates have been previously reported, with complete degradation of fresh dead jellyfish occurring within 5 to 14 days (Titelman et al. 2006; West et al. 2009; Qu et al. 2015). These rapid degradations of dead jellyfish (mainly by microbial decomposition) were attributed to the biochemical composition of their tissues, with a high proportion of proteins, low C:N ratio, and lack of hard structures, providing high-quality substrate for specific bacteria (Titelman et al. 2006; Tinta et al. 2012, 2020). Indeed, simulated scenarios performed with *Aurelia aurita* showed that about half of its dead organic matter is instantly available as dissolved organic matter and rapidly consumed by microbes (within 1.5 days, Tinta et al., 2020). Decay rates are also known to vary with seawater temperature and jellyfish size (Titelman et al. 2006; Lebrato et al. 2011, 2012). Temperature is probably one of the most important factors driving differences in jellyfish decay rates (Lebrato et al. 2011). In the Thau lagoon, the collapse of the *A. coerulea* bloom coincides with the peak of summer temperatures (>20 °C; Marques et al. 2015a). This might promote fast medusae degradation by the microbenthos, but also in the water column (Tinta et al. 2020), thereby reducing the amount of jellyfish biomass that reaches the seabed. The biomass loss rates observed in Thau are in the range of those decay rates estimated for tropical shallow environments, where less than one day is required to decompose 99% of jellyfish organic matter (Lebrato et al. 2011). However, degradation rates also depend on the initial medusae biomass, with smaller individuals decaying faster than larger ones (Titelman et al. 2006).

The medusae of *A. coerulea* in Thau are usually smaller than those of other *Aurelia* species (Marques et al. 2015a) or scyphozoans (Pitt 2000; Fuentes et al. 2011; Prieto et al. 2013). Therefore, it is a combination of small-sized medusae and high local temperature which likely promotes fast biomass loss rates of jelly-falls in this shallow ecosystem, through the action of microorganisms.

We initially hypothesised that differences in macrobenthic community composition associated with different habitats would have a distinct impact on the biomass loss rate of the jelly-falls. The consumption of dead jellyfish by macroorganisms can be expected to be more important in habitats with lower food availability (Sweetman et al. 2014) since local organisms depend on less frequent inputs of new sources of organic matter (Holmer et al. 2004). Indeed, in the Thau lagoon and as expected (Thouzeau et al. 2007; Rueda et al. 2009), macrobenthic communities differed between the two tested habitats, and higher biomass loss rates were observed on bare sediments, where the amount of available organic matter is lower (Plus et al. 2003). Furthermore, it was on bare sediments that the only macrobenthic organism was significantly attracted by jelly-falls (gastropods from the *Tritia* genus). These results suggest a possible contribution of the macroorganisms consumption to the disappearance of *A. coerulea* jelly-falls in the lagoon. However, these results must be considered with caution since the effect of the habitat in our study was not completely independent from the initial biomass of *A. coerulea* used in each experiment (higher on the seagrass meadows than on the bare sediment), which has been shown to affect jellyfish decay rates (Titelman et al. 2006). The experiments were performed on different days and, although individuals were randomly selected, the captivity time between experiments might have induced a bias on the initial weight of the medusae. Furthermore, although this was not tested in the present work, this captivity time might have also affected the biochemical composition of jellyfish tissues, potentially affecting our results.

Within each habitat, our results suggest a limited consumption of jelly-falls by macrobenthic organisms. On the bare sediments, the results from the degradation experiment showed that jellyfish decay rates were not affected by the accessibility scenarios, while on the seagrass meadows, the biomass loss of dead medusae was even faster for the Micro than for the Macro

scenario, which was unexpected. This might partially be caused by our experimental setup. Indeed, the small mesh of the net bags (200 μm) used in this scenario, might have promoted the physical retention of microorganisms, protecting them against local currents, thereby avoiding their advection and dilution in the surrounding water. Moreover, by eliminating large organisms, the retention of microorganisms within the bag might have modified the trophic interactions, by simultaneously providing high concentrations of organic matter and decreasing the grazing pressure on microorganisms, boosting their proliferation. If this is true, it is possible that the decay rates obtained for the Micro scenario are greater than those actually occurring in the lagoon.

The results from our second experiment corroborate the limited contribution of macrobenthic organisms in the disappearance of jelly-falls in Thau. The lack of significant difference in community composition among the M and PC treatments indicates that the addition of jelly-falls to the seabed did not significantly disturb the community composition, either by attracting or repelling organisms. However, the species-specific analysis (i.e. ManyGLM), revealed potential positive responses of some particular taxa. Among those are the *Bittium* sp. in the seagrass meadows and the *Tritia* sp., *Hexaplex* sp., and *Ruditapes* sp. on the bare sediment. Indeed, peaks of their abundance were recorded, especially, for the M treatment at 3h, which co-occurred with the maximum in jellyfish biomass loss rates in our study. This might suggest a limited but still possible contribution of these macroorganisms consumption to the disappearance of dead medusae in the lagoon. Nevertheless, only the *Tritia* sp. (Nassariidae family; Galindo et al., 2016) revealed significant differences between the M and the PC treatments, indicating that, jelly-falls only significantly attracted this particular species, as also reported by Chelsky et al. (2016). Nassariidae species are common on soft sediment habitats and reported as herbivorous, carnivorous, but mainly as scavengers, feeding opportunistically on the available dead organic matter (Morton 2011). These organisms rapidly detect carrions from long distances and move fast towards the carcasses, but they leave it once they are satiated to avoid potential predators (Morton 2011). They appear to eat large amounts of organic matter (20 to 60% of their weight) in as fast as 8 minutes (Morton 2011; Lucena et al. 2012) and the amount of time they spend on feeding appears to be a function of their hunger, with individuals living in habitats with lower food supply,

545 eating a larger amount of food and spending more time feeding (Morton and Chan 1999). This
546 might explain the peak of *Tritia* sp. abundance observed during the first hours of the experiment
547 on the bare sediment habitat. Therefore, although our results suggest a limited consumption of *A.*
548 *coerulea* carcasses by the macrobenthos after the annual blooms of the jellyfish in Thau, the
549 scavenging activity of the gastropods from the *Tritia* genus might still contribute to the fast
550 disappearance of its jelly-falls on the lagoon's bare sediment habitats.

552 4.2. Potential ecological impacts of jellyfish degradation in Thau

553 Our results suggest that the rapid biomass loss of *A. coerulea* jelly-falls in the Thau lagoon is
554 mostly caused by a fast degradation of its dead medusae by local microorganisms, with a possible
555 contribution of some particular species of scavenger on the bare sediment habitat. This might
556 have several ecological implications in Thau, but also in other shallow coastal habitats.

557 The increase of dissolved inorganic nutrients in the surroundings of decaying jellyfish might
558 enhance the local phytoplankton and algal production through direct assimilation of dissolved
559 inorganic compounds (Pitt et al. 2009; Blanchet et al. 2015). Likewise, bacterial production might
560 be enhanced during the jellyfish degradation process (Tinta et al. 2010, 2012, 2020), which
561 represents an important food source for microzooplankton (Rassoulzadegan and Sheldon 1986).
562 This supports the hypothesis that the available energy of jelly-falls can be directly (if consumed
563 by some scavengers) or indirectly (through microorganisms) transferred to higher trophic levels.
564 However, the ecological consequences of the rapid degradation of *A. coerulea* blooms by the
565 bacterial community in Thau might also be negative. During the summer, anoxic crisis episodes,
566 known as ‘malaïgues’, occasionally occur in the lagoon. They are caused by the bacterial
567 degradation of high concentrations of organic matter as the combination of high water
568 temperatures, weak winds, and important water residence times which promotes stratification of
569 the water column and decreases oxygen exchanges at the surface or with the sea (Harzallah and
570 Chapelle 2002). During jellyfish degradation, large amounts of highly bioavailable dissolved
571 organic matter are released and quickly metabolized by the microbial community, decreasing the
572 dissolved oxygen concentrations in the vicinity of jellyfish carcasses (West et al. 2009; Pitt et al.

2009; Sweetman et al. 2016; Chelsky et al. 2016; Guy-Haim et al. 2020). Therefore, the collapse of the *A. coerulea* bloom in the early summer might amplify the magnitude of summer anoxic crises, potentially leading to massive benthic community mortalities. Lastly, the summer collapse of the jellyfish bloom and its degradation by local bacteria might partially contribute to the summer mortalities of the cultivated oyster *Crassostrea gigas* that sporadically occur in Thau (Pernet et al. 2012). Indeed, the bacterial degradation of *A. coerulea* medusae has been shown to enhance abundances of *Vibrio* spp. in the surrounding water (Tinta et al. 2012; Blanchet et al. 2015), and peaks of these microorganisms (from *Vibrionaceae* family) have been associated with *C. gigas* mortalities (Pernet et al. 2012; Cantet et al. 2013). Although this scenario is very speculative, it needs to be investigated, because shellfish farming is the most important local economic activity in the lagoon, and shellfish mortality events have dramatic consequences on the local economy (Pernet et al. 2012).

5. Conclusion

Evaluating the ecological impacts of jellyfish blooms requires identifying the fate of their organic matter, *i.e.* whether they are scavenged by demersal or benthic predators, decomposed by microorganisms, or both. In Thau, the absence of large accumulations of dead medusae of *A. coreulea* on the seafloor probably partially results from their ingestion by several local fish species (Marques et al. 2019) and rapid degradation in the water column (Tinta et al. 2020). However, we show that, upon their arrival on the sea bed, their fast biomass loss is mainly caused by their rapid degradation by local microorganisms, favoured by the high local summer temperatures and the small size of the medusae. Ingestion by benthic scavengers is possible but limited. Therefore the collapse of the jellyfish blooms in the lagoon has a limited impact on its macrobenthic communities. Instead, they have the potential to significantly modify local biogeochemical cycles, reshape ecosystem functioning and, ultimately, affect ecosystem services with important implications on several economic activities. This supports the need for further

600 investigations on jellyfish degradation in coastal lagoons and calls for incorporating this process
601 in ecosystem-based models.

6. Acknowledgments

We thank Sebastien Colantoni, Perline Bastide, and Sadjia Belkacemi for their collaboration during field and laboratory work. We also thank Prof. Stephan Baghdiguian for his precious help during molluscs' identification. We also thank the comments of the reviewers which largely contributed to the improvement of the manuscript.

This work was funded by MARBEC laboratory (internal resources).

The authors declare that they have no competing interests.

References:

- Anderson, M. J. 2017. Permutational Multivariate Analysis of Variance (PERMANOVA), p. 1–15. In Wiley StatsRef: Statistics Reference Online. John Wiley & Sons, Ltd.
- Ates, R. M. L. 2017. Benthic scavengers and predators of jellyfish, material for a review. *Plankton and Benthos Research* **12**: 71–77. doi:10.3800/pbr.12.71
- Billett, D. S. M., B. J. Bett, C. L. Jacobs, I. P. Rouse, and B. D. Wigham. 2006. Mass deposition of jellyfish in the deep Arabian Sea. *Limnology and Oceanography* **51**: 2077–2083. doi:10.4319/lo.2006.51.5.2077
- Blanchet, M., O. Pringault, M. Bouvy, and others. 2015. Changes in bacterial community metabolism and composition during the degradation of dissolved organic matter from the jellyfish *Aurelia aurita* in a Mediterranean coastal lagoon. *Environ Sci Pollut Res* **22**: 13638–13653. doi:10.1007/s11356-014-3848-x
- Bonnet, D., J.-C. Molinero, T. Schohn, and M. N. Daly-Yahia. 2012. Seasonal changes in the population dynamics of *Aurelia aurita* in Thau lagoon. *Cahier de Biologie Marine* **53**: 343–347.
- Bos, A. R., E. Cruz-Rivera, and A. M. Sanad. 2017. Herbivorous fishes *Siganus rivulatus* (Siganidae) and *Zebbrasoma desjardinii* (Acanthuridae) feed on Ctenophora and Scyphozoa in the Red Sea. *Mar Biodiv* **47**: 243–246. doi:10.1007/s12526-016-0454-9
- Cantet, F., D. Hervio-Heath, A. Caro, and others. 2013. Quantification of *Vibrio parahaemolyticus*, *Vibrio vulnificus* and *Vibrio cholerae* in French Mediterranean coastal lagoons. *Research in Microbiology* **164**: 867–874. doi:10.1016/j.resmic.2013.06.005
- Cardona, L., I. Á. de Quevedo, A. Borrell, and A. Aguilar. 2012. Massive Consumption of Gelatinous Plankton by Mediterranean Apex Predators. *PLOS ONE* **7**: e31329. doi:10.1371/journal.pone.0031329
- Chelsky, A., K. A. Pitt, A. J. P. Ferguson, W. W. Bennett, P. R. Teasdale, and D. T. Welsh. 2016. Decomposition of jellyfish carrion *in situ*: Short-term impacts on infauna, benthic nutrient fluxes and sediment redox conditions. *Science of The Total Environment* **566–567**: 929–937. doi:10.1016/j.scitotenv.2016.05.011

- Condon, R. H., D. K. Steinberg, P. A. del Giorgio, and others. 2011. Jellyfish blooms result in a major microbial respiratory sink of carbon in marine systems. *Proceedings of the National Academy of Sciences* **108**: 10225–10230. doi:10.1073/pnas.1015782108
- D'Angelo, G., and S. Gargiullo. 1978. Guida alle conchiglie mediterranee: Conoscerle, cercarle, collezionarle, I. S. Gruppo Editoriale Fabbri, Bompiani [ed.]. Fabbri.
- Dunlop, K. M., D. O. B. Jones, and A. K. Sweetman. 2017. Direct evidence of an efficient energy transfer pathway from jellyfish carcasses to a commercially important deep-water species. *Scientific Reports* **7**: 17455. doi:10.1038/s41598-017-17557-x
- Fauvel, P. 1923. Faune de France. Polychetes Errantes, P. Lechevalier [ed.].
- Fauvel, P. 1927. Faune de France. Polychetes Errantes, P. Lechevalier [ed.].
- Fiandrino, A., A. Giraud, S. Robin, and C. Pinatel. 2012. Validation d'une méthode d'estimation des volumes d'eau échangés entre la mer et les lagunes et définition d'indicateurs hydrodynamiques associés.
- Frost, J. R., C. A. Jacoby, T. K. Frazer, and A. R. Zimmerman. 2012. Pulse perturbations from bacterial decomposition of *Chrysaora quinquecirrha* (Scyphozoa: Pelagiidae). *Hydrobiologia* **690**: 247–256. doi:10.1007/s10750-012-1042-z
- Fuentes, V., I. Straehler-Pohl, D. Atienza, and others. 2011. Life cycle of the jellyfish *Rhizostoma pulmo* (Scyphozoa: Rhizostomeae) and its distribution, seasonality and inter-annual variability along the Catalan coast and the Mar Menor (Spain, NW Mediterranean). *Marine Biology* **158**: 2247–2266. doi:10.1007/s00227-011-1730-7
- Galindo, L. A., N. Puillandre, J. Utge, P. Lozouet, and P. Bouchet. 2016. The phylogeny and systematics of the Nassariidae revisited (Gastropoda, Buccinoidea). *Molecular Phylogenetics and Evolution* **99**: 337–353. doi:10.1016/j.ympev.2016.03.019
- Guy-Haim, T., M. Rubin-Blum, E. Rahav, N. Belkin, J. Silverman, and G. Sisma-Ventura. 2020. The effects of decomposing invasive jellyfish on biogeochemical fluxes and microbial dynamics in an ultra-oligotrophic sea. *Biogeosciences* **17**: 5489–5511. doi:10.5194/bg-17-5489-2020

- Harzallah, A., and A. Chapelle. 2002. Contribution of climate variability to occurrences of anoxic crises ‘malaïgues’ in the Thau lagoon (southern France). *Oceanologica Acta* **25**: 79–86.
- Hays, G. C., T. K. Doyle, and J. D. R. Houghton. 2018. A Paradigm Shift in the Trophic Importance of Jellyfish? *Trends in Ecology & Evolution* **33**: 874–884. doi:10.1016/j.tree.2018.09.001
- Holmer, M., C. M. Duarte, H. T. S. Boschker, and C. Barrón. 2004. Carbon cycling and bacterial carbon sources in pristine and impacted Mediterranean seagrass sediments. *Aquatic Microbial Ecology* **36**: 227–237. doi:10.3354/ame036227
- Kindt, R., and R. Coe. 2005. Tree diversity analysis. A manual and software for common statistical methods for ecological and biodiversity studies., World Agroforestry Centre (ICRAF).
- Lebrato, M., and D. O. B. Jones. 2009. Mass deposition event of *Pyrosoma atlanticum* carcasses off Ivory Coast (West Africa). *Limnology and Oceanography* **54**: 1197–1209. doi:10.4319/lo.2009.54.4.1197
- Lebrato, M., P. de J. Mendes, D. K. Steinberg, J. E. Cartes, B. M. Jones, L. M. Birsa, R. Benavides, and A. Oschlies. 2013. Jelly biomass sinking speed reveals a fast carbon export mechanism. *Limnology and Oceanography* **58**: 1113–1122. doi:https://doi.org/10.4319/lo.2013.58.3.1113
- Lebrato, M., M. Pahlow, J. R. Frost, M. Küter, P. de J. Mendes, J.-C. Molinero, and A. Oschlies. 2019. Sinking of Gelatinous Zooplankton Biomass Increases Deep Carbon Transfer Efficiency Globally. *Global Biogeochemical Cycles* **33**: 1764–1783. doi:https://doi.org/10.1029/2019GB006265
- Lebrato, M., M. Pahlow, A. Oschlies, K. A. Pitt, D. O. B. Jones, J. C. Molinero, and R. H. Condon. 2011. Depth attenuation of organic matter export associated with jelly falls. *Limnology and Oceanography* **56**: 1917–1928. doi:10.4319/lo.2011.56.5.1917
- Lebrato, M., K. A. Pitt, A. K. Sweetman, and others. 2012. Jelly-falls historic and recent observations: A review to drive future research directions. *Hydrobiologia* **690**: 227–245. doi:10.1007/s10750-012-1046-8

- Lilley, M. K. S., S. E. Beggs, T. K. Doyle, V. J. Hobson, K. H. P. Stromberg, and G. C. Hays. 2011. Global patterns of epipelagic gelatinous zooplankton biomass. *Marine Biology* **158**: 2429–2436. doi:10.1007/s00227-011-1744-1
- Lucena, J., C. Meirelles, and H. Matthews-Cascon. 2012. Feeding behavior of *Nassarius vibex* (Gasteropoda: Nassariidae). *Arquivos de Ciências do Mar* **45**: 60–67. doi:10.32360/acmar.v45i2.134
- Luo, J. Y., R. H. Condon, C. A. Stock, C. M. Duarte, C. H. Lucas, K. A. Pitt, and R. K. Cowen. 2020. Gelatinous Zooplankton-Mediated Carbon Flows in the Global Oceans: A Data-Driven Modeling Study. *Global Biogeochemical Cycles* **34**: e2020GB006704. doi:https://doi.org/10.1029/2020GB006704
- Marques, R., S. Albouy-Boyer, F. Delpy, C. Carré, É. Le Floch, C. Roques, J.-C. Molinero, and D. Bonnet. 2015a. Pelagic population dynamics of *Aurelia* sp. in French Mediterranean lagoons. *Journal of Plankton Research* **37**: 1019–1035. doi:10.1093/plankt/fbv059
- Marques, R., M. Cantou, S. Soriano, J.-C. Molinero, and D. Bonnet. 2015b. Mapping distribution and habitats of *Aurelia* sp. polyps in Thau lagoon, north-western Mediterranean Sea (France). *Mar Biol* **162**: 1441–1449. doi:10.1007/s00227-015-2680-2
- Marques, R., A. M. Darnaude, S. Crochemore, C. Bouvier, and D. Bonnet. 2019. Molecular approach indicates consumption of jellyfish by commercially important fish species in a coastal Mediterranean lagoon. *Marine Environmental Research* **152**: 104787. doi:10.1016/j.marenvres.2019.104787
- Morton, B. 2011. Behaviour of *Nassarius bicallosus* (Caenogastropoda) on a northwestern Western Australian surf beach with a review of feeding in the Nassariidae. *Molluscan Research* **31**: 89–94.
- Morton, B., and K. Chan. 1999. Hunger rapidly overrides the risk of predation in the subtidal scavenger *Nassarius siquijorensis* (Gastropoda: Nassariidae): an energy budget and a comparison with the intertidal *Nassarius festivus* in Hong Kong. *Journal of Experimental Marine Biology and Ecology* **240**: 213–228. doi:10.1016/S0022-0981(99)00060-X

- Newton, A., J. Icely, S. Cristina, and others. 2014. An overview of ecological status, vulnerability and future perspectives of European large shallow, semi-enclosed coastal systems, lagoons and transitional waters. *Estuarine, Coastal and Shelf Science* **140**: 95–122. doi:10.1016/j.ecss.2013.05.023
- Oksanen, A. J., F. G. Blanchet, M. Friendly, and others. 2019. *vegan*: Community Ecology Package.
- Pernet, F., J. Barret, P. Le Gall, C. Corporeau, L. Dégremont, F. Lagarde, J. Pépin, and N. Keck. 2012. Mass mortalities of Pacific oysters *Crassostrea gigas* reflect infectious diseases and vary with farming practices in the Mediterranean Thau lagoon, France. *Aquaculture Environment Interactions* **2**: 215–237. doi:10.3354/aei00041
- Pinheiro, J., Bates, S. DebRoy, D. Sarkar, and R Core Team. 2019. *nlme*: Linear and Nonlinear Mixed Effects Models.
- Pitt, K. A. 2000. Life history and settlement preferences of the edible jellyfish *Catostylus mosaicus* (Scyphozoa: Rhizostomeae). *Marine Biology* **136**: 269–279. doi:10.1007/s002270050685
- Pitt, K. A., A. C. Budarf, J. G. Browne, and R. H. Condon. 2014. Bloom and Bust: Why Do Blooms of Jellyfish Collapse?, p. 79–103. In K.A. Pitt and C.H. Lucas [eds.], *Jellyfish Blooms*. Springer Netherlands.
- Pitt, K. A., and M. J. Kingsford. 2003. Temporal and spatial variation in recruitment and growth of medusae of the jellyfish, *Catostylus mosaicus* (Scyphozoa : Rhizostomeae). *Marine and Freshwater Research* **54**: 117. doi:10.1071/MF02110
- Pitt, K. A., D. T. Welsh, and R. H. Condon. 2009. Influence of jellyfish blooms on carbon, nitrogen and phosphorus cycling and plankton production. *Hydrobiologia* **616**: 133–149. doi:10.1007/s10750-008-9584-9
- Plus, M., A. Chapelle, P. Lazure, and others. 2003. Modelling of oxygen and nitrogen cycling as a function of macrophyte community in the Thau lagoon. *Continental Shelf Research* **23**: 1877–1898. doi:10.1016/j.csr.2003.03.001

- Prieto, L., A. Armani, and D. Macías. 2013. Recent strandings of the giant jellyfish *Rhizostoma*
luteum Quoy and Gaimard, 1827 (Cnidaria: Scyphozoa: Rhizostomeae) on the Atlantic
and Mediterranean coasts. *Marine Biology* **160**: 3241–3247. doi:10.1007/s00227-013-
2293-6
- Purcell, J. E. 2012. Jellyfish and Ctenophore Blooms Coincide with Human Proliferations and
Environmental Perturbations. *Annual Review of Marine Science* **4**: 209–235.
doi:10.1146/annurev-marine-120709-142751
- Qu, C.-F., J.-M. Song, N. Li, X.-G. Li, H.-M. Yuan, L.-Q. Duan, and Q.-X. Ma. 2015. Jellyfish
(*Cyanea nozakii*) decomposition and its potential influence on marine environments
studied via simulation experiments. *Marine Pollution Bulletin* **97**: 199–208.
doi:10.1016/j.marpolbul.2015.06.016
- Rassoulzadegan, F., and R. W. Sheldon. 1986. Predator-prey interactions of nanozooplankton and
bacteria in an oligotrophic marine environment¹. *Limnology and Oceanography* **31**:
1010–1029. doi:10.4319/lo.1986.31.5.1010
- Rueda, J. L., S. Gofas, J. Urrea, and C. Salas. 2009. A highly diverse molluscan assemblage
associated with eelgrass beds (*Zostera marina* L.) in the Alboran Sea: Micro-habitat
preference, feeding guilds and biogeographical distribution. *Scientia Marina* **73**: 679–
700. doi:10.3989/scimar.2009.73n4679
- Sweetman, A. K., and A. Chapman. 2011. First observations of jelly-falls at the seafloor in a
deep-sea fjord. *Deep Sea Research Part I: Oceanographic Research Papers* **58**: 1206–
1211. doi:10.1016/j.dsr.2011.08.006
- Sweetman, A. K., and A. Chapman. 2015. First assessment of flux rates of jellyfish carcasses
(jelly-falls) to the benthos reveals the importance of gelatinous material for biological C-
cycling in jellyfish-dominated ecosystems. *Frontiers in Marine Science* **2**: 1–7.
doi:10.3389/fmars.2015.00047
- Sweetman, A. K., A. Chelsky, K. A. Pitt, and others. 2016. Jellyfish decomposition at the seafloor
rapidly alters biogeochemical cycling and carbon flow through benthic food-webs.
Limnology and Oceanography **61**: 1449–1461. doi:10.1002/lno.10310

776 Sweetman, A. K., C. R. Smith, T. Dale, and D. O. B. Jones. 2014. Rapid scavenging of jellyfish
 777 carcasses reveals the importance of gelatinous material to deep-sea food webs.
 778 Proceedings of the Royal Society B: Biological Sciences **281**: 20142210–20142210.
 779 doi:10.1098/rspb.2014.2210

780 Thouzeau, G., J. Grall, J. Clavier, and others. 2007. Spatial and temporal variability of benthic
 781 biogeochemical fluxes associated with macrophytic and macrofaunal distributions in the
 782 Thau lagoon (France). Estuarine, Coastal and Shelf Science **72**: 432–446.
 783 doi:10.1016/j.ecss.2006.11.028

784 Tinta, T., T. Kogovšek, A. Malej, and V. Turk. 2012. Jellyfish Modulate Bacterial Dynamic and
 785 Community Structure. PLoS ONE **7**: e39274. doi:10.1371/journal.pone.0039274

786 Tinta, T., T. Kogovšek, V. Turk, T. A. Shiganova, A. S. Mikaelyan, and A. Malej. 2016. Microbial
 787 transformation of jellyfish organic matter affects the nitrogen cycle in the marine water
 788 column — A Black Sea case study. Journal of Experimental Marine Biology and Ecology
 789 **475**: 19–30. doi:10.1016/j.jembe.2015.10.018

790 Tinta, T., A. Malej, M. Kos, and V. Turk. 2010. Degradation of the Adriatic medusa *Aurelia* sp.
 791 by ambient bacteria. Hydrobiologia **645**: 179–191. doi:10.1007/s10750-010-0223-x

792 Tinta, T., Z. Zhao, A. Escobar, K. Klun, B. Bayer, C. Amano, L. Bamonti, and G. J. Herndl. 2020.
 793 Microbial Processing of Jellyfish Detritus in the Ocean. Front. Microbiol. **11**.
 794 doi:10.3389/fmicb.2020.590995

795 Titelman, J., L. Riemann, T. Sørnes, T. Nilsen, P. Griekspoor, and U. Båmstedt. 2006. Turnover
 796 of dead jellyfish: stimulation and retardation of microbial activity. Marine Ecology
 797 Progress Series **325**: 43–58. doi:10.3354/meps325043

798 Wang, Y., U. Naumann, S. T. Wright, and D. I. Warton. 2012. mvabund - an R package for model-
 799 based analysis of multivariate abundance data. Methods in Ecology and Evolution **3**: 471–
 800 474. doi:10.1111/j.2041-210X.2012.00190.x

801 Warton, D. I., S. T. Wright, and Y. Wang. 2012. Distance-based multivariate analyses confound
 802 location and dispersion effects. Methods in Ecology and Evolution **3**: 89–101.
 803 doi:10.1111/j.2041-210X.2011.00127.x

- West, E. J., D. T. Welsh, and K. A. Pitt. 2009. Influence of decomposing jellyfish on the sediment oxygen demand and nutrient dynamics, p. 151–160. *In* K.A. Pitt and J.E. Purcell [eds.], Jellyfish Blooms: Causes, Consequences, and Recent Advances: Proceedings of the Second International Jellyfish Blooms Symposium, held at the Gold Coast, Queensland, Australia, 24–27 June, 2007. Springer Netherlands.
- De Wit, R., H. Rey-Valette, J. Balavoine, V. Ouisse, and R. Lifran. 2017. Restoration ecology of coastal lagoons: new methods for the prediction of ecological trajectories and economic valuation. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**: 137–157. doi:10.1002/aqc.2601
- Yamamoto, J., M. Hirose, T. Ohtani, and others. 2008. Transportation of organic matter to the sea floor by carrion falls of the giant jellyfish *Nemopilema nomurai* in the Sea of Japan. *Marine Biology* **153**: 311–317. doi:10.1007/s00227-007-0807-9
- Zuur, A., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. *Mixed Effects Models and Extensions in Ecology with R*, Springer New York.