Parasitism and host dispersal plasticity in an aquatic model system

Giacomo Zilio, Louise S Nørgaard, Giovanni Petrucci, Nathalie Zeballos, Claire Gougat-barbera, Emanuel A Fronhofer, Oliver Kaltz

► To cite this version:

Giacomo Zilio, Louise S Nørgaard, Giovanni Petrucci, Nathalie Zeballos, Claire Gougat-barbera, et al.. Parasitism and host dispersal plasticity in an aquatic model system. *Journal of Evolutionary Biology*, 2021, 34 (8), pp.1316-1325. 10.1111/jeb.13893 . hal-03426187

HAL Id: hal-03426187

<https://hal.umontpellier.fr/hal-03426187>

Submitted on 12 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Parasitism and host dispersal plasticity in an aquatic model system

Running title: Parasitism and dispersal plasticity

Authors details:

Giacomo Zilio^{1*}, gcm.zilio@gmail.com, ORCID ID: <https://orcid.org/0000-0002-4448-3118>

Louise S. Nørgaard^{1,2}, lnorga10@hotmail.com, ORCID ID: <https://orcid.org/0000-0002-0938-0017>

Giovanni Petrucci¹, giovannipetrucci82@gmail.com, ORCID ID: unavailable

Nathalie Zeballos^{1,3}, nathalie.zeballos@cefe.cnrs.fr, ORCID ID: unavailable

Claire Gougat-Barbera¹, claire.gougat-barbera@umontpellier.fr, ORCID ID: unavailable

Emanuel A. Fronhofer¹, emanuel.fronhofer@umontpellier.fr, ORCID ID: <https://orcid.org/0000-0002-2219-784X>

Oliver Kaltz^{1*}, oliver.kaltz@umontpellier.fr, ORCID ID: <https://orcid.org/0000-0002-7154-0456>

¹ ISEM, University of Montpellier, CNRS, EPHE, IRD, Montpellier, France.

² School of Biological Sciences, Monash University, Melbourne 3800, Australia.

³ CEFE, University of Montpellier, CNRS, EPHE, IRD, Montpellier, France.

*Corresponding authors: gcm.zilio@gmail.com, oliver.kaltz@umontpellier.fr

Acknowledgements section

This work was supported by the Swiss National Science Foundation (grant no. P2NEP3_184489) to GZ, and by the 2019 Godfrey Hewitt mobility award granted to LN by ESEB, and by the. This is publication ISEM-XXXX-XXX of the Institut des Sciences de l'Evolution.

Author Contributions section

OK, GZ, LN, NZ and EAF conceived the study. OK, GZ, LN and NZ designed the experiments. GZ, LN, NZ, CGB and OK performed the experimental work. GZ, EAF, OK, GP and NZ performed the statistical analysis. All authors interpreted the results and contributed to the writing of the manuscript.

Conflict of Interest statement

The authors have no conflict of interest to declare.

Abstract

Dispersal is a central determinant of spatial dynamics in communities and ecosystems, and various ecological factors can shape the evolution of constitutive and plastic dispersal behaviours. One important driver of dispersal plasticity is the biotic environment. Parasites, for example, influence the internal condition of infected hosts and define external patch quality. Thus state-dependent dispersal may be determined by infection status and context-dependent dispersal by the abundance of infected hosts in the population. A prerequisite for such dispersal plasticity to evolve is a genetic basis on which natural selection can act. Using interconnected microcosms, we investigated dispersal in experimental populations of the freshwater protist *Paramecium caudatum* in response to the bacterial parasite *Holospora undulata*. For a collection of 20 natural host strains, we found substantial variation in constitutive dispersal, and to a lesser degree in dispersal plasticity. First, infection tended to increase or decrease dispersal relative to uninfected controls, depending on strain identity, potentially indicative of state-dependent dispersal plasticity. Infection additionally decreased host swimming speed compared to the uninfected counterparts. Second, for certain strains, there was a weak negative association between dispersal and infection prevalence, such that uninfected hosts tended to disperse less when infection was more frequent in the population, indicating context-dependent dispersal plasticity. Future experiments may test whether the observed differences in dispersal plasticity are sufficiently strong to react to natural selection. The evolution of dispersal plasticity as a strategy to mitigate parasite effects spatially may have important implications for epidemiological dynamics.

Keywords:

Condition-dependent dispersal, dispersal plasticity, eco-evolution, epidemiology, *Holospora undulata*, host-parasite interactions, *Paramecium caudatum*, reaction norms, spatial dynamics

54 Introduction

55 Dispersal, broadly defined as the movement of individuals with consequences for gene flow, is a key life-
 56 history trait (Bonte & Doherty, 2017) driving metapopulation and metacommunity dynamics as well as the
 57 geographic distribution of species (Hanski, 1999). In recent years, the study of dispersal and dispersal
 58 syndromes have received increasing interest (Clobert *et al.*, 2012; Stevens *et al.*, 2014), as landscapes are
 59 seeing large-scale environmental alterations and fragmentation, rendering dispersal crucial to potentially
 60 mitigate these changes (Parmesan & Yohe, 2003; Cote *et al.*, 2017). Although dispersal is often considered a
 61 constitutive trait, plastic dispersal behaviour represents a flexible alternative, responding to changes in the
 62 internal condition of an individual (state-dependent dispersal) and to external environmental factors
 63 (context-dependent dispersal) (Clobert *et al.*, 2009). State-dependent dispersal has been associated with
 64 variation in factors such as body size, the developmental stage or sex of individuals (Bowler & Benton, 2005).
 65 In contrast, context-dependent dispersal decisions may be based on cues that provide information on biotic
 66 and abiotic patch properties, such as food availability, population density, or kin competition (see Ronce,
 67 2007 and references therein).

68
 69 In communities, dispersal plasticity may be advantageous in mitigating adverse interactions with other
 70 species (Fronhofer *et al.*, 2015a). Parasites are particularly interesting in this respect: they are ubiquitous and
 71 impose strong selection pressures, and potentially drive the evolution of both state-dependent and context-
 72 dependent dispersal of their hosts (Iritani & Iwasa, 2014; Iritani, 2015; Narayanan *et al.*, 2020; Deshpande *et al.*,
 73 2021). Empirical studies have investigated aspects of parasite-related dispersal (see below), but still little
 74 is known about the genetic basis of this kind of dispersal plasticity and its adaptive significance.

75
 76 State-dependent dispersal may relate to morphological or physiological changes induced by parasites. The
 77 exploitation of host resources might decrease general activity levels, and thereby reduce movement and
 78 dispersal. Such negative effects have been documented for various organisms (Binning *et al.*, 2017; Nørgaard
 79 *et al.*, 2019; Baines *et al.*, 2020), even though it is not necessarily a general rule (Nelson *et al.*, 2015; Csata *et al.*,
 80 2017). While in many examples the observed effects may represent side effects, theory has identified
 81 conditions under which increased (but also decreased) dispersal when infected is adaptive, namely under kin
 82 selection (Iritani & Iwasa, 2014; Iritani, 2015) or when infection can be lost during dispersal (Shaw & Binning,
 83 2016; Daversa *et al.*, 2017). Indeed, increased dispersal of infected hosts is not uncommon (Suhonen *et al.*,
 84 2010; Brown *et al.*, 2016), although it may also be the result of parasite manipulation (Lion *et al.*, 2006;
 85 Martini *et al.*, 2015).

86
 87 Natural enemies may also produce context-dependent dispersal, as a means to reduce immediate predation
 88 or infection risk. For example, herbivores or predators can induce the production of specific dispersal morphs

(Weisser *et al.*, 1999; de la Pena *et al.*, 2011). A recent multi-species study further showed that chemical predator-related cues increase dispersal probability (Fronhofer *et al.*, 2018). Such cues may also exist in host-parasite systems, where infection-avoidance behaviour is well known (Behringer *et al.*, 2006; Curtis, 2014). Recent theory shows that hosts may indeed evolve reaction norms, with dispersal being a function of the parasite infection prevalence (Deshpande *et al.*, 2021). To date, few if any empirical studies have tested for the existence of such plastic population-level responses (French & Travis, 2001).

Adaptive phenotypic plasticity is a powerful solution in many situations (Chevin *et al.*, 2013; Stamp & Hadfield, 2020), and just like constitutive traits, it has a genetic basis on which selection can act (Pigliucci, 2005; Garland & Kelly, 2006; Laitinen & Nikoloski, 2019). Dispersal-related traits have such a genetic basis (Saastamoinen *et al.*, 2018) and constitutive dispersal can evolve rapidly in a parasite context (Koskella *et al.*, 2011; Zilio *et al.*, 2020). However, the genetics and evolution of dispersal plasticity is less well studied. In fact, how plastic dispersal varies between different genotypes under parasite challenge is rarely evaluated in empirical studies (Suhonen *et al.*, 2010; Fellous *et al.*, 2011), or the genetic diversity is treated as a random effect (Csata *et al.*, 2017). Moreover, the number of genotypes evaluated is usually small, making it difficult to draw general conclusions (Leggett *et al.*, 2013).

Here, using interconnected microcosms, we tested a collection of 20 natural strains of *Paramecium caudatum* for dispersal in the presence and absence of the bacterial parasite *Holospira undulata*. Previous work in this system had shown that infection reduces dispersal for a small number of strains (Fellous *et al.*, 2011; Nørgaard *et al.*, 2021). The first objective of the present study was to test whether this negative effect was general, or whether strains varied in infection-state dependent dispersal. Second, we tested for genetic variation in context-dependent dispersal by comparing the dispersal of uninfected hosts over a range of infection prevalences that had naturally established in the experimental populations. We found that parasite reduced or increased dispersal levels depending on strain identity, indicating a state-dependent plastic response of the infected hosts, but no general negative effect of infection. Furthermore, increasing infection prevalence tended to reduce host dispersal for certain strains, suggesting context-dependent dispersal plasticity of uninfected hosts. Such genetic variation in dispersal plasticity may provide the raw material for parasite-mediated selection, in natural settings or for the purpose of experimental evolution.

Materials and methods

Study system

Paramecium caudatum is a freshwater filter-feeding protist from stagnant waters of the Northern hemisphere (Wichterman, 2012). Like all ciliates, paramecia have a macronucleus for somatic gene expression and a germ-line micronucleus, used for sexual reproduction. The micronucleus can be infected by

Holospira undulata, a gram-negative alpha-proteobacterium (Fokin, 2004). Infectious spores are released for horizontal transmission after host cell division or upon host death. Infectious spores are immobile and therefore rely on host movement or water current for their own dispersal. Vertical transmission occurs when hosts divide mitotically. Infection reduces *P. caudatum* division and survival (Restif & Kaltz, 2006) and also host dispersal (Fellous *et al.*, 2011; Nørgaard *et al.*, 2021).

Experimental setup

Preparation of replicates. We established mass cultures for a collection of 20 genetically distinct strains of *P. caudatum* from different geographical regions (provided by S. Krenek, TU Dresden, Germany; Table S1, Supplementary Information). Distributed over two experimental blocks, 6 infected replicate cultures were established for each strain (20 strains x 2 blocks x 3 replicates = 120 replicates). Inocula were prepared from a mix of infected stock cultures in the lab, all originating from a single isolate of *H. undulata* established in 2001 (Dohra *et al.*, 2013). Following standard protocols for the extraction of infectious spores (e.g., Nørgaard *et al.*, 2021) we used c. 10^4 spores to inoculate samples of c. $3\text{--}5 \times 10^3$ host cells in 1.5 mL per assay replicate. Four days after inoculation, when infections have established, we expanded the cultures by regular addition of lettuce medium (supplemented with the food bacterium *Serratia marcescens*), until a volume of 50 mL was reached. In the same way, we set up three uninfected control populations per strain, giving a total of 180 experimental cultures. After three weeks, prior to the dispersal assay, population size (mean: $190 \text{ mL}^{-1} \pm 9 \text{ SE}$; 95% range [172; 208]) and infection prevalence (mean: $26.8 \% \pm 2.1$; 95% range [3.1; 90.7]) had settled naturally in each experimental replicate.

Dispersal assay. We assayed the dispersal of infected and uninfected replicates in dispersal arenas, as described in Nørgaard *et al.* (2021). A dispersal arena consisted of three 50-mL Falcon tubes, linearly connected by 5-cm long silicon tubing (inner diameter: 0.8 cm). The 3-patch system was filled with 75 mL of medium to establish connections. Then the connections were blocked with clamps and 20 mL of a given replicate culture added into the middle tube. The lateral tubes received 20 mL of *Paramecium*-free medium. Connections were then opened, and the *Paramecium* allowed to disperse to the lateral tubes for 3h. After blocking the connections, we counted the individuals in samples from the middle tube (500- μ l) and from the combined lateral tubes (3 mL) to estimate the number of non-dispersing and dispersing individuals (dissecting microscope, 40x). From the same samples, we also made lacto-aceto-orcin fixations (Görtz & Wiemann, 1989) and determined the infection status (infected / uninfected) of up to 30 dispersing and non-dispersing individuals, respectively (light microscope, phase contrast, 1000x). From the cell counts and the infection status data, we estimated the population density and infection prevalence in the middle tube at the beginning of the assay. From the same data, we also estimated the proportion of infected and uninfected dispersers for each replicate, referred to as per-3h “dispersal rate” or dispersal, hereafter.

In addition, to investigate a potential link between dispersal and movement (Banerji *et al.*, 2015; Pennekamp *et al.*, 2019), we assayed swimming behaviour. For each strain, 1 infected and 1 uninfected individual were isolated from arbitrarily selected assay replicates, and allowed to replicate in a 2-mL plastic tubes for 8 days. For the resulting 40 monoclonal cultures (20 strains x 2 infection status) we placed 200- μ L samples (10-20 individuals) on a microscope slide and recorded individual movement trajectories under a Perflex Pro 10 stereomicroscope, using a Perflex SC38800 camera (15 frames per second; duration: 10 s; total magnification: 10x). For each sample, average swimming speed (μ m/s) and swimming tortuosity (standard deviation of the turning angle distribution, describing the extent of swimming trajectory change) were determined using video analysis ("BEMOVI" package; Pennekamp *et al.*, 2015).

Statistical analysis

Statistical analyses were conducted in R, v. 3.6.3 (R Core Team, 2020) using Bayesian models with the 'rstan' (version 2.19.3) and 'rethinking' (version 2.0.1) packages (McElreath 2020). For state-dependent dispersal, we compared the dispersal of the infected individuals (in infected replicates) with the dispersal in the uninfected control replicates. We fitted four models, from the intercept to the full interaction model, using a binomial regression with logit link function (chain length: warmup = 20,000 iterations, chain = 40,000 iterations). In the full model, the explanatory factors were infection status (infected or uninfected control), *Paramecium* strain identity, and the strain x status interaction. Experimental block only explained a negligible fraction of the dispersal variation (preliminary analysis, not shown) and was omitted from all further analyses. We fitted the models choosing vaguely informative priors; the intercepts and slope parameters followed a normal distribution with mean -2 and standard deviation 3 for the first, and mean 0 and standard deviation 1.75 for the latter. To account for overdispersion we included an observation-level random effect. The mean and standard deviation of the observation-level hyperprior followed a normal and half-normal distribution respectively, with mean 0 and standard deviation 1. The four state-dependent models were compared and ranked using the Watanabe-Akaike information criterion, WAIC (Watanabe, 2010), a generalized version of the Akaike information criterion (Gelman *et al.*, 2014). The posterior predictions of the models were then averaged based on WAIC weights, and the relative importance (RI) of the explanatory variables was calculated as the sum of the respective WAIC model weights in which that variable was included. Due to loss of replicates, low population density, and/or very low levels of infection, 159 replicates (from 20 strains) of the 180 initial replicates were available for this analysis.

For context-dependent dispersal, we analysed the dispersal of uninfected *Paramecium* in infected assay replicates. We fitted 6 models, from the intercept to the full interaction model, using the same binomial regression with logit link function, chain lengths and prior specifications as above. The explanatory factors of the full model (varying intercept and slope) were infection prevalence, strain identity and the strain x

infection prevalence interaction. The posterior predictions were averaged and ranked, and the RI calculated based on WAIC model weights as described above. For this analysis, 99 assay replicates (from 19 strains) of the initially 120 inoculated replicates were available.

We used similar analyses to test whether swimming speed and tortuosity varied as a function of strain identity and infection status. We standardized the response variable and fitted four models (from the intercept to the additive model, see Table S2 and S3, Supplementary Information) using a linear regression (chain length: warmup = 20,000 iterations, chain = 40,000 iterations) with an exponentially distributed prior (rate = 1) for standard deviation. As for the dispersal analysis, the parameter priors were vaguely informative; the intercept and slope parameters followed a normal distribution with mean 0 and standard deviation 2. We averaged and ranked the posterior predictions, and we obtained RI based on WAIC model weights. We further tested for correlations between these two swimming traits and mean strain dispersal, for infected and uninfected *Paramecium* (chain length: warmup = 2,000 iterations, chain = 10,000 iterations). Due to missing data, only 17 of the 20 strains were used for these analyses.

Results

State-dependent dispersal

Our analysis revealed substantial variation in constitutive dispersal among the 20 *P. caudatum* strains (relative importance, RI, of strain identity = 0.85; Table 1), ranging from 1% (95% compatibility Interval [0.001; 0.135]) to 41% ([0.02; 0.80]) of the individuals moving from the central to the lateral tubes (Fig. 1A).

Our models provided limited evidence for state-dependent dispersal plasticity. Infection status (RI = 0.57) was retained in the best model fit (lowest WAIC; Model 3 in Table 1), indicating a general trend of infection to increase host dispersal. Even though the signal of the strain x infection status interaction (RI = 0.22) was only weak, patterns in Fig. 1B indicate that effects of infection varied with strain identity: several strains indeed dispersed more when infected (Fig. 1B right side of panel), but in at least half of the strains, infection had little effect or decreased host dispersal.

Context-dependent dispersal

As in the above analysis, we found substantial genotypic variation in overall constitutive levels of dispersal for uninfected *Paramecium* (RI of strain identity = 0.80; Table 2). The best model (model 4 in Table 2) included an effect of infection prevalence (RI = 0.67), and thus context-dependent dispersal. Namely, uninfected individuals tended to disperse less at higher parasite infection prevalence in the population (Fig. 2): such negative dispersal-prevalence relationships were predicted for all but one strain (negative median slope values; Fig. 2B). To some degree, however, the strength of this relationship varied between strains (RI of infection prevalence x strain interaction = 0.21). As shown in Fig. 2B, distributions of predicted slopes show

considerable variation and for the majority of strains there is considerable overlap with 0. Only a small number of strains (e.g., C139, C116, C083) show clearly negative slopes (Fig. 2B).

Swimming behaviour

The analysis of standardized swimming speed revealed strong effects of strain identity (RI = 0.9; Table S2) and infection status (RI = 1; Table S2). Namely, standardized swimming speed of uninfected *Paramecium* (median = 0.57, 95% CI [-0.64; 2.34]) was generally higher than that of infected ones (median = -1.20, 95% CI [-1.63; -0.77]), corresponding to a difference of almost 40% (median = 0.39, 95% CI [0.10; 0.68]; Fig. S1A-B). Swimming tortuosity was not affected by strain and weakly affected by infection status (RI strain = 0; RI status = 0.28; Table S3). Neither swimming speed (uninfected: $r = 0.08$, 95% CI [-0.39; 0.52]; infected: $r = 0.07$, 95% CI [-0.41; 0.54]) nor swimming tortuosity (uninfected: $r = 0.15$, 95% CI [-0.29; 0.56]; infected: $r = -0.10$, 95% CI [-0.55; 0.39]) were strongly correlated with dispersal.

Discussion

Dispersal affects epidemiology and host-parasite (co)evolution in metapopulations (Lion & Gandon, 2015; Parratt *et al.*, 2016), but how dispersal itself evolves due to antagonistic species interactions is less well known (Poethke *et al.*, 2010; Drown *et al.*, 2013; Deshpande *et al.*, 2021). Here we focused on dispersal plasticity in response to parasitism, which may evolve as a means to reduce infection risk of the dispersing individuals and/or their relatives (Iritani & Iwasa, 2014; Iritani, 2015; Deshpande *et al.*, 2021). Our study takes a first step towards an understanding of population-level processes, by measuring dispersal of infected and uninfected hosts in experimental microcosms and by exploring the genetic variation in plasticity for a collection of host strains. Overall, signals of dispersal plasticity were weak. Both infection status and infection prevalence modified dispersal to some degree, with at least some strains showing indications of state-dependent dispersal (i.e., when infected) and/or context-dependent dispersal (i.e., in response to infection prevalence).

State-dependent plasticity: the dispersal of infected hosts

In previous studies, infection by *H. undulata* reduced dispersal in *P. caudatum* for a small set of strains (Fellous *et al.*, 2011; Nørgaard *et al.*, 2021). Here we used strains from a worldwide collection (Table S1) and find the entire range of trends, from negative or no impact of infection to even positive effects on host dispersal (Fig. 1). Reduced host dispersal may be explained by general negative effects of infection, through the energetic demand of an immune response, the diversion of host resources by the parasite or direct physical damage (Mideo, 2009). Indeed, *H. undulata* consumes nuclear proteins and nucleotides (Garushyants *et al.*, 2018) and also causes massive interior swelling of the infected micronucleus, which would explain the clear and pervasive reduction in swimming speed observed in the complementary

experiment (Fig. S1A-B). However, dispersal reductions were far from being universal, suggesting that the amount of host damage differs between genotypes. Differential fitness effects (virulence) and variation in resistance are known for this system (Restif & Kaltz, 2006), indicating the strong potential for genotypic-specificity in the responses to this parasite.

Moreover, it should be noted that the absence of a difference between infected and uninfected dispersal does not necessarily mean the absence of plasticity. Infected hosts may compensate parasite damage by re-allocating resources to maintain vital functions, such as foraging and feeding activity, and this may lead to a net-zero effect of infection on dispersal. Interestingly, some of our strains even seemed to 'overcompensate' and dispersed more when infected. Such a positive state-dependent dispersal may be selectively favoured in a metapopulation because it can reduce kin competition and kin infection (Iritani & Iwasa, 2014; Iritani, 2015; Deshpande *et al.*, 2021). However, increased host dispersal may equally well reflect parasite manipulation, enhancing its dispersal to novel infection sites (Kamo & Boots, 2006; Lion *et al.*, 2006; Martini *et al.*, 2015).

The main purpose here was to quantify the (variation in) population-level effects of infection on dispersal. More work is needed to better understand the links between parasite action, host movement and dispersal. This concern, for example the relationship between parasite load, virulence and dispersal. Furthermore, unlike in other protists (Pennekamp *et al.*, 2019), swimming speed was not a good predictor of dispersal. Other aspects of swimming behaviour (Ricci, 1989) may be more relevant in our system. Namely, *Paramecium* show a characteristic vertical distribution (Fels *et al.*, 2008) relating to food and oxygen availability (Wichterman, 2012). Parasites are known to affect the position of hosts in the water column (Cezilly *et al.*, 2000; Fels *et al.*, 2004), and this may directly influence the probability of infected individuals finding the dispersal corridors in our microcosms.

Context-dependent plasticity: the dispersal of uninfected hosts

Predator chemical signals induce dispersal in various organisms, including *P. caudatum* (Fronhofer *et al.* 2018). We tested for a similar parasite effect in our microcosm populations, by measuring the dispersal of uninfected hosts at different infection prevalences, with the assumption that higher prevalence equals a stronger signal of 'parasite presence'. Unlike in the predator-cues study, we found little evidence for a positive dispersal-inducing effect. Dispersal decreased at higher infection prevalence, at least for certain strains. Interestingly, Deshpande *et al.*'s model (2021) predicts the evolution of such negative prevalence-dependent dispersal, as the result of complex spatio-temporal variations in eco-evolutionary processes. We do not know the evolutionary history of the strains, but our results suggest a possible genetic basis of context-dependent dispersal in this system and hence genetic variation that might be seen by natural selection.

Our experimental approach of using naturally established infection prevalence may not have produced strong enough signal variation for all strains. This could be remedied via more artificial designs, by mixing of infected and uninfected individuals to establish well-defined gradients. Infected cultures or inocula may also be filtered to specifically test for chemical cues (see Fronhofer *et al.*, 2018). Finally, we made the simplifying assumption of linear dispersal reaction norms. However, dispersal responses may well follow non-linear rules, e.g., if there are signal thresholds (Fronhofer *et al.*, 2015b), as observed for other traits (Morel-Journel *et al.*, 2020) and predicted by Deshpande *et al.* (2021). Tests for non-linear relationships would require a much finer resolution (i.e., more replication) on the signal axis.

Conditions for plasticity selection: outlook

The heritability of phenotypic plasticity of morphological or behavioural traits is generally lower than their constitutive heritability (Scheiner, 1993; Stirling *et al.*, 2002). In line with this, we find much less among-strain differentiation for parasite-related dispersal plasticity than for constitutive dispersal, suggesting a weaker potential for responding to selection. However, the available genetic variation alone does not determine the relative importance of phenotypic plasticity in shaping evolutionary trajectories (Stamp & Hadfield, 2020). Phenotypic plasticity is generally favoured in variable, but nonetheless predictable environments (Leung *et al.*, 2020). In a parasite context, dispersal plasticity evolution may thus depend on the spatio-temporal predictability of parasite encounter rates across a metapopulation (Deshpande *et al.*, 2021). Additional factors are parasite virulence, the cost of dispersal (or its advantage if parasite release is possible during dispersal), or correlations with other traits (Iritani & Iwasa, 2014). For example, a recent experiment with the protist *Tetrahymena* revealed few genetic constraints on the concurrent evolution of plasticity across various traits (Morel-Journel *et al.*, 2020). Indeed, state- and context-dependent dispersal might also evolve simultaneously in the presence of parasites, even though not necessarily in a correlated fashion (Deshpande *et al.*, 2021). Our data indicate no genetic correlation between state- and context-dependent plasticity ($r = -0.11$, 95% CI [-0.55; 0.36]; based on strain averages), suggesting that independent responses to selection are possible, as shown in the model.

Our study represents one of the first accounts of the naturally existing genetic variation for state-dependent and context-dependent dispersal plasticity in relation to parasites. The signals of plasticity are weak and there are many open questions regarding the mechanistic and physiological basis of trait expression or information use. Nonetheless, in microbial systems such as ours, the observed variation opens promising avenues for future experiments. In microcosm landscapes, allowing the free interplay between dispersal and epidemiological processes, we can assess how dispersal plasticity affects parasite spread at the metapopulation level. Over longer time spans, we can also explore dispersal evolution and test evolutionary predictions on dispersal plasticity and its adaptive role in host-parasite interactions.

333 **Data availability statement**

334 The experimental data will be made available upon potential acceptance (via Dryad/Figshare repository).

335

336 **References**

337 Baines, C.B., Diab, S. & McCauley, S.J. 2020. Parasitism Risk and Infection Alter Host Dispersal. *Am. Nat.* 000–
338 000. The University of Chicago Press.

339 Banerji, A., Duncan, A.B., Griffin, J.S., Humphries, S., Petchey, O.L. & Kaltz, O. 2015. Density- and trait-
340 mediated effects of a parasite and a predator in a tri-trophic food web. *J. Anim. Ecol.* **84**: 723–733.

341 Behringer, D.C., Butler, M.J. & Shields, J.D. 2006. Avoidance of disease by social lobsters. *Nature* **441**: 421–
342 421. Nature Publishing Group.

343 Binning, S.A., Shaw, A.K. & Roche, D.G. 2017. Parasites and Host Performance: Incorporating Infection into
344 Our Understanding of Animal Movement. *Integr. Comp. Biol.* **57**: 267–280. Oxford Academic.

345 Bonte, D. & Doherty, M. 2017. Dispersal: a central and independent trait in life history. *Oikos* **126**: 472–479.

346 Bowler, D.E. & Benton, T.G. 2005. Causes and consequences of animal dispersal strategies: relating individual
347 behaviour to spatial dynamics. *Biol. Rev.* **80**: 205–225.

348 Brown, G.P., Kelehear, C., Pizzatto, L. & Shine, R. 2016. The impact of lungworm parasites on rates of dispersal
349 of their anuran host, the invasive cane toad. *Biol. Invasions* **18**: 103–114.

350 Cezilly, F., Gregoire, A. & Bertin, A. 2000. Conflict between co-occurring manipulative parasites? An
351 experimental study of the joint influence of two acanthocephalan parasites on the behaviour of
352 *Gammarus pulex*. *Parasitology* **120**: 625–630. Cambridge University Press.

353 Chevin, L.-M., Gallet, R., Gomulkiewicz, R., Holt, R.D. & Fellous, S. 2013. Phenotypic plasticity in evolutionary
354 rescue experiments. *Philos. Trans. R. Soc. B Biol. Sci.* **368**.

355 Clobert, J., Baguette, M., Benton, T.G. & Bullock, J.M. 2012. *Dispersal Ecology and Evolution*. Oxford
356 University Press.

357 Clobert, J., Galliard, J.-F.L., Cote, J., Meylan, S. & Massot, M. 2009. Informed dispersal, heterogeneity in
358 animal dispersal syndromes and the dynamics of spatially structured populations. *Ecol. Lett.* **12**: 197–
359 209.

- 360 Cote, J., Bestion, E., Jacob, S., Travis, J., Legrand, D. & Baguette, M. 2017. Evolution of dispersal strategies
361 and dispersal syndromes in fragmented landscapes. *Ecography* **40**: 56–73.
- 362 Csata, E., Bernadou, A., Rakosy-Tican, E., Heinze, J. & Marko, B. 2017. The effects of fungal infection and
363 physiological condition on the locomotory behaviour of the ant *Myrmica scabrinodis*. *J. Insect Physiol.*
364 **98**: 167–172.
- 365 Curtis, V.A. 2014. Infection-avoidance behaviour in humans and other animals. *Trends Immunol.* **35**: 457–
366 464.
- 367 Daversa, D.R., Fenton, A., Dell, A.I., Garner, T.W.J. & Manica, A. 2017. Infections on the move: how transient
368 phases of host movement influence disease spread. *Proc. R. Soc. B Biol. Sci.* **284**: 20171807.
- 369 de la Pena, E., D’hondt, B. & Bonte, D. 2011. Landscape structure, dispersal and the evolution of antagonistic
370 plant-herbivore interactions. *Ecography* **34**: 480–487.
- 371 Deshpande, J.N., Kaltz, O. & Fronhofer, E.A. 2021. Host–parasite dynamics set the ecological theatre for the
372 evolution of state- and context-dependent dispersal in hosts. *Oikos* **130**: 121–132.
- 373 Dohra, H., Suzuki, H., Suzuki, T., Tanaka, K. & Fujishima, M. 2013. Draft Genome Sequence of *Holosporea*
374 *undulata* Strain HU1, a Micronucleus-Specific Symbiont of the Ciliate *Paramecium caudatum*.
375 *Genome Announc.* **1**.
- 376 Drown, D.M., Dybdahl, M.F. & Gomulkiewicz, R. 2013. Consumer–Resource Interactions and the Evolution of
377 Migration. *Evolution* **67**: 3290–3304.
- 378 Fellous, S., Quillery, E., Duncan, A.B. & Kaltz, O. 2011. Parasitic infection reduces dispersal of ciliate host. *Biol.*
379 *Lett.* **7**: 327–329.
- 380 Fels, D., Lee, V.A. & Ebert, D. 2004. The impact of microparasites on the vertical distribution of *Daphnia*
381 *magna*. *Arch. Für Hydrobiol.* **161**: 65–80.
- 382 Fels, D., Vignon, M. & Kaltz, O. 2008. Ecological and genetic determinants of multiple infection and
383 aggregation in a microbial host-parasite system. *Parasitology* **135**: 1373–1383. Cambridge University
384 Press.
- 385 Fokin, S.I. 2004. Bacterial endocytobionts of ciliophora and their interactions with the host cell. *Int. Rev. Cytol.*
386 **236**: 181–250. New York, NY: Academic Press, 1952-.
- 387 French, D. & Travis, J. 2001. Density-dependent dispersal in host-parasitoid assemblages. *Oikos* **95**: 125–135.

388 Fronhofer, E.A., Klecka, J., Melián, C.J. & Altermatt, F. 2015a. Condition-dependent movement and dispersal
389 in experimental metacommunities. *Ecol. Lett.* **18**: 954–963.

390 Fronhofer, E.A., Kropf, T. & Altermatt, F. 2015b. Density-dependent movement and the consequences of the
391 Allee effect in the model organism *Tetrahymena*. *J. Anim. Ecol.* **84**: 712–722.

392 Fronhofer, E.A., Legrand, D., Altermatt, F., Ansart, A., Blanchet, S., Bonte, D., *et al.* 2018. Bottom-up and top-
393 down control of dispersal across major organismal groups. *Nat. Ecol. Evol.* **2**: 1859–1863. Nature
394 Publishing Group.

395 Garland, T. & Kelly, S.A. 2006. Phenotypic plasticity and experimental evolution. *J. Exp. Biol.* **209**: 2344–2361.

396 Garushyants, S.K., Beliavskaia, A.Y., Malko, D.B., Logacheva, M.D., Rautian, M.S. & Gelfand, M.S. 2018.
397 Comparative Genomic Analysis of *Holospira spp.*, Intranuclear Symbionts of Paramecia. *Front.*
398 *Microbiol.* **9**. Frontiers.

399 Gelman, A., Hwang, J. & Vehtari, A. 2014. Understanding predictive information criteria for Bayesian models.
400 *Stat. Comput.* **24**: 997–1016.

401 Görtz, H.-D. & Wiemann, M. 1989. Route of infection of the bacteria *Holospira elegans* and *Holospira obtusa*
402 into the nuclei of *Paramecium caudatum*. *Eur. J. Protistol.* **24**: 101–109.

403 Hanski, I. 1999. *Metapopulation Ecology*. Oxford University Press, Oxford, New York.

404 Iritani, R. 2015. How parasite-mediated costs drive the evolution of disease state-dependent dispersal. *Ecol.*
405 *Complex.* **21**: 1–13.

406 Iritani, R. & Iwasa, Y. 2014. Parasite infection drives the evolution of state-dependent dispersal of the host.
407 *Theor. Popul. Biol.* **92**: 1–13.

408 Kamo, M. & Boots, M. 2006. The evolution of parasite dispersal, transmission, and virulence in spatial host
409 populations. *Evol. Ecol. Res.* **8**: 1333–1347.

410 Koskella, B., Taylor, T.B., Bates, J. & Buckling, A. 2011. Using experimental evolution to explore natural
411 patterns between bacterial motility and resistance to bacteriophages. *ISME J.* **5**: 1809–1817.

412 Laitinen, R.A.E. & Nikoloski, Z. 2019. Genetic basis of plasticity in plants. *J. Exp. Bot.* **70**: 739–745.

413 Leggett, H.C., Benmayor, R., Hodgson, D.J. & Buckling, A. 2013. Experimental Evolution of Adaptive
414 Phenotypic Plasticity in a Parasite. *Curr. Biol.* **23**: 139–142.

- 415 Leung, C., Rescan, M., Grulois, D. & Chevin, L.-M. n.d. Reduced phenotypic plasticity evolves in less
416 predictable environments. *Ecol. Lett.* **23**: 1664-1672.
- 417 Lion, S. & Gandon, S. 2015. Evolution of spatially structured host-parasite interactions. *J. Evol. Biol.* **28**: 10–
418 28.
- 419 Lion, S., van Baalen, M. & Wilson, W.G. 2006. The evolution of parasite manipulation of host dispersal. *Proc.*
420 *R. Soc. B Biol. Sci.* **273**: 1063–1071. Royal Society.
- 421 Martini, X., Hoffmann, M., Coy, M.R., Stelinski, L.L. & Pelz-Stelinski, K.S. 2015. Infection of an Insect Vector
422 with a Bacterial Plant Pathogen Increases Its Propensity for Dispersal. *PLoS One* **10**.
- 423 McElreath, R. 2020. Statistical rethinking: a Bayesian course with examples in R and Stan. 2nd ed. Chapman
424 and Hall/CRC, Boca Raton, FL.
- 425 Mideo, N. 2009. Parasite adaptations to within-host competition. *Trends Parasitol.* **25**: 261–268.
- 426 Morel-Journel, T., Thuillier, V., Pennekamp, F., Laurent, E., Legrand, D., Chaine, A.S., *et al.* 2020. A
427 multidimensional approach to the expression of phenotypic plasticity. *Funct. Ecol.* **34**: 2338–2349.
- 428 Narayanan, N., Binning, S.A. & Shaw, A.K. 2020. Infection state can affect host migratory decisions. *Oikos* **129**:
429 1493–1503.
- 430 Nelson, F.B.L., Brown, G.P., Dubey, S. & Shine, R. 2015. The Effects of a Nematode Lungworm (*Rhabdias hylae*)
431 on its Natural and Invasive Anuran Hosts. *J. Parasitol.* **101**: 290–296. American Society of
432 Parasitologists.
- 433 Nørgaard, L.S., Phillips, B.L. & Hall, M.D. 2019. Infection in patchy populations: Contrasting pathogen invasion
434 success and dispersal at varying times since host colonization. *Evol. Lett.* **3**: 555–566.
- 435 Nørgaard, L.S., Zilio, G., Saade, C., Gougat-Barbera, C., Hall, M.D., Fronhofer, E.A., *et al.* 2021. An evolutionary
436 trade-off between parasite virulence and dispersal at experimental invasion fronts. *Ecol. Lett.*
437 ele.13692.
- 438 Parmesan, C. & Yohe, G. 2003. A globally coherent fingerprint of climate change impacts across natural
439 systems. *Nature* **421**: 37–42. Nature Publishing Group.
- 440 Parratt, S.R., Numminen, E. & Laine, A.-L. 2016. Infectious Disease Dynamics in Heterogeneous Landscapes.
441 *Annu. Rev. Ecol. Evol. Syst.* **47**: 283–306.

442 Pennekamp, F., Clobert, J. & Schtickzelle, N. 2019. The interplay between movement, morphology and
443 dispersal in Tetrahymena ciliates. *PeerJ* **7**: e8197. PeerJ Inc.

444 Pennekamp, F., Schtickzelle, N. & Petchey, O.L. 2015. BEMOVI, software for extracting behavior and
445 morphology from videos, illustrated with analyses of microbes. *Ecol. Evol.* **5**: 2584–2595.

446 Pigliucci, M. 2005. Evolution of phenotypic plasticity: where are we going now? *Trends Ecol. Evol.* **20**: 481–
447 486.

448 Poethke, H.J., Weisser, W.W. & Hovestadt, T. 2010. Predator-Induced Dispersal and the Evolution of
449 Conditional Dispersal in Correlated Environments. *Am. Nat.* **175**: 577–586. The University of Chicago
450 Press.

451 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical
452 Computing, Vienna, Austria.

453 Restif, O. & Kaltz, O. 2006. Condition-dependent virulence in a horizontally and vertically transmitted
454 bacterial parasite. *Oikos* **114**: 148–158.

455 Ricci, N. 1989. Locomotion as a criterion to read the adaptive biology of Protozoa and their evolution toward
456 Metazoa. *Boll. Zool.* **56**: 245–263. Taylor & Francis.

457 Ronce, O. 2007. How Does It Feel to Be Like a Rolling Stone? Ten Questions About Dispersal Evolution. *Annu.*
458 *Rev. Ecol. Evol. Syst.* **38**: 231–253.

459 Saastamoinen, M., Bocedi, G., Cote, J., Legrand, D., Guillaume, F., Wheat, C.W., *et al.* 2018. Genetics of
460 dispersal. *Biol. Rev.* **93**: 574–599.

461 Scheiner, S.M. 1993. Genetics and Evolution of Phenotypic Plasticity. *Annu. Rev. Ecol. Syst.* **24**: 35–68.

462 Shaw, A.K. & Binning, S.A. 2016. Migratory Recovery from Infection as a Selective Pressure for the Evolution
463 of Migration. *Am. Nat.* **187**: 491–501. The University of Chicago Press.

464 Stamp, M.A. & Hadfield, J.D. 2020. The relative importance of plasticity versus genetic differentiation in
465 explaining between population differences; a meta-analysis. *Ecol. Lett.* **23**: 1432–1441.

466 Stevens, V.M., Whitmee, S., Galliard, J.-F.L., Clobert, J., Böhning-Gaese, K., Bonte, D., *et al.* 2014. A
467 comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial animals. *Ecol. Lett.* **17**:
468 1039–1052.

469 Stirling, D.G., Réale, D. & Roff, D.A. 2002. Selection, structure and the heritability of behaviour. *J. Evol. Biol.*
470 **15**: 277–289.

471 Suhonen, J., Honkavaara, J. & Rantala, M.J. 2010. Activation of the immune system promotes insect dispersal
472 in the wild. *Oecologia* **162**: 541–547.

473 Watanabe, S. 2010. Asymptotic Equivalence of Bayes Cross Validation and Widely Applicable Information
474 Criterion in Singular Learning Theory. *J. Mach. Learn. Res.* **11**: 3571–3594.

475 Weisser, W.W., Braendle, C. & Minoretti, N. 1999. Predator-induced morphological shift in the pea aphid.
476 *Proc. R. Soc. Lond. B Biol. Sci.* **266**: 1175–1181. Royal Society.

477 Wichterman, R. 2012. *The Biology of Paramecium*. Springer Science & Business Media.

478 Zilio, G., Nørgaard, L.S., Gougat-Barbera, C., Hall, M.D., Fronhofer, E.A. & Kaltz, O. 2020. Travelling with a
479 parasite: the evolution of resistance and dispersal syndrome during experimental range expansion.
480 *bioRxiv* 2020.01.29.924498. Cold Spring Harbor Laboratory.

481

482

483

484

485

486

487

488

489

490

491

492

493

494

Tables

Table 1. Different statistical models and parameters included for the analysis and model averaging of the state-dependent dispersal. The rows represent the different models (the best model is highlighted in bold) and the columns the factors included in each model with the corresponding WAIC, standard error of the WAIC and WAIC weights. The RI row shows the relative importance of the explanatory variables.

	Strain	Status	Strain * Status	WAIC	SE	WAIC weight
Model 1				1165	7.36	0.15
Model 2	X			1163.8	27.76	0.27
Model 3	X	X		1163.3	27.67	0.35
Model 4	X	X	X	1164.3	7.68	0.22
RI	0.85	0.57	0.22			

Table 2. Statistical models and parameters for the analysis and model averaging of the context-dependent dispersal. Each row represents a different model, the best model is highlighted in bold and the last row indicates the relative importance (RI) of the explanatory variables. The columns are the variables included in the six models with the corresponding WAIC, standard error of the WAIC and WAIC weights.

	Strain	Prevalence	Strain * Prevalence	WAIC	SE	WAIC weight
Model 1				746.4	17.97	0.10
Model 2	X			744.6	18.62	0.23
Model 3		X		746.4	17.93	0.10
Model 4	X	X		744.2	18.77	0.28
Model 5	X	X		746.7	18.16	0.08
Model 6	X	X	X	744.8	18.61	0.21
RI	0.80	0.67	0.21			

Figures

Figure 1. State-dependent dispersal of 20 *Paramecium caudatum* strains, as a function of infection status (uninfected / infected with *Holospira undulata*). (A) Shaded bars and thick lines represent the 95% compatibility interval and the median of the averaged model predictions of the posterior distributions. Strains are ordered according to the difference between uninfected (grey) and infected (red) dispersal. Each circle represents an experimental replicate. (B) Difference between uninfected and infected averaged model posterior predictions for each strain (expressed in logits), the thick black line represents the median of the difference distribution. Distributions shifted below zero (dashed grey line) indicates higher dispersal in the infected (pointing-down arrow) compared to the uninfected (pointing-up arrow) treatment.

Figure 2. Context-dependent dispersal of 19 uninfected *Paramecium caudatum* strains, as a function of parasite (*Holospira undulata*) infection prevalence in the microcosm population. (A) Each panel represents a strain, and each circle an experimental replicate; the red shaded area and thick red lines are the 95% compatibility interval and median of the averaged model of the posterior distributions. (B) Averaging of the posterior distributions of the slope parameter calculated in logit (model 3-6, Table 2) with the thick black lines showing the median. Positive or negative slopes distributions (above or below zero, dashed grey line), indicate a higher or lower dispersal in response to increasing frequency of infected hosts.

Supplementary Information

Figure S1. Standardized swimming speed of 17 *Paramecium caudatum*, as a function of infection status (uninfected / infected with *Holospira undulata*). (A) Strains are ordered according to the difference between uninfected (grey) and infected (red) dispersal. Each circle represents an experimental replicate. Shaded bars and thick lines are the 95% compatibility interval and median of the averaged model predictions of the posterior distributions, and each circle represents the measured data of swimming speed per strain. (B) The difference in swimming speed between uninfected and infected averaged model posterior predictions for each strain. The thick black lines are the median of the difference distribution. Distributions shifted above zero (dashed grey line) indicates higher swimming speed in the uninfected treatment (pointing-up arrow) compared to the infected treatment (pointing-down arrow).

Table S1. Host strain identity, location of origin and GPS coordinates, (*) indicates approximate location.

Strain	Location code	Origin	GPS coordinates
C002	Hainberger See	Germany	51.035688, 12.285028
C021	Erzgebirge E1	Germany	50.647032, 13.256963
C023	Plön K 2	Germany	54.15021, 10.440307
C026	Fokin 1 UBR 42	USA, Louisiana	53.131996, 13.107747
C030	Österreich_Lahnalp	Austria	47.695948, 12.218064 (*)
C033	Peking 1_C3	China	39.128165, 117.185083 (*)
C034	Plön K 2	Germany	54.15021, 10.440307
C065	SWE 17.1	Sweden	60.105747, 15.966911
C083	USBL-511	USA, Indiana	39.069861, -86.414361
C105	Sp 10C	Spain	39.548852, -1.502887
C116	Frankreich 10-2.1	France	43.430297, 6.126616
C119	Peru	Peru	-12.046184, -77.040842 (*)
C139	My43c3d	Japan	38.480973, 141.372414 (*)
C147	KNZ5414	Japan	36.519469, 136.709415
C173	Greece 10.1	Greece	40.805947, 21.983306
GS1CI	Globsowsee	Germany	53.128590, 13.118740
GS6BI	Globsowsee	Germany	53.128590, 13.118740
KS10CI	Kochsee	Germany	53.131996, 13.107747
KS3AII	Kochsee	Germany	53.131996, 13.107747
KS7CII	Kochsee	Germany	53.131996, 13.107747

585 **Table S2.** Statistical models and parameters for the analysis and model averaging of the swimming speed. Each row
586 shows a different model, from the intercept to the additive model. The best model is highlighted in bold and the last
587 row indicates the relative importance (RI) of the strain and status effect. The columns are the explanatory variables
588 included in the four models with the corresponding WAIC, standard error of the WAIC and WAIC weights.
589

	Strain	Status	WAIC	SE	WAIC weight
Model 1			100.0	10.45	0.0
Model 2	X		114.1	5.84	0.0
Model 3		X	83.7	10.29	0.1
Model 4	X	X	79.4	7.56	0.9
RI	0.9	1.0			

590
591 **Table S3.** Models and parameters for the analysis and model averaging of the swimming tortuosity. Each row
592 corresponds to a different model used for the analysis, with the best model is highlighted in bold. The last row shows
593 the relative importance (RI) of the explanatory variables. The columns are the variables of the models with the
594 corresponding WAIC, standard error of the WAIC and WAIC weights.
595

	Strain	Status	WAIC	SE	WAIC weight
Model 1			99.1	6.52	0.72
Model 2	X		119.4	5.89	0.00
Model 3		X	101.0	6.39	0.28
Model 4	X	X	121.4	5.86	0.00
RI	0.00	0.28			

596
597
598
599
600