



**HAL**  
open science

## Niche breadth affects bacterial transcription patterns along a salinity gradient

Angel Rain-Franco, Nicolas Mouquet, Claire Gougat-Barbera, Thierry  
Bouvier, Sara Beier

► **To cite this version:**

Angel Rain-Franco, Nicolas Mouquet, Claire Gougat-Barbera, Thierry Bouvier, Sara Beier. Niche breadth affects bacterial transcription patterns along a salinity gradient. *Molecular ecology*, 2021, 10.1111/mec.16316 . hal-03589732

**HAL Id: hal-03589732**

**<https://hal.umontpellier.fr/hal-03589732>**

Submitted on 28 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Niche breadth affects bacterial transcription patterns along a salinity gradient

Angel Rain-Franco<sup>1</sup>  | Nicolas Mouquet<sup>2</sup>  | Claire Gougat-Barbera<sup>3</sup> |  
Thierry Bouvier<sup>2</sup>  | Sara Beier<sup>1,4</sup> 

<sup>1</sup>CNRS, Laboratoire d'Océanographie Microbienne, LOMIC, Sorbonne Université, Banyuls/mer, France

<sup>2</sup>MARBEC, CNRS, Ifremer, IRD, Université de Montpellier, Montpellier, France

<sup>3</sup>ISEM, CNRS, EPHE, IRD, University of Montpellier, Montpellier, France

<sup>4</sup>Leibniz Institute for Baltic Sea Research, Warnemünde, Germany

## Correspondence

Sara Beier, Leibniz Institute for Baltic Sea Research, Warnemünde, Germany.  
Email: sara.beier@io-warnemuende.de

## Funding information

Deutsche Forschungsgemeinschaft, Grant/Award Number: BE 5937/2-1; Chilean National Agency for Research and Development (ANID), Grant/Award Number: 2017 - 72180448

## Abstract

Understanding the molecular mechanisms that determine a species' life history is important for predicting their susceptibility to environmental change. While specialist species with a narrow niche breadth (NB) maximize their fitness in their optimum habitat, generalists with broad NB adapt to multiple environments. The main objective of this study was to identify general transcriptional patterns that would distinguish bacterial strains characterized by contrasted NBs along a salinity gradient. More specifically, we hypothesized that genes encoding fitness-related traits, such as biomass production, have a higher degree of transcriptional regulation in specialists than in generalists, because the fitness of specialists is more variable under environmental change. By contrast, we expected that generalists would exhibit enhanced transcriptional regulation of genes encoding traits that protect them against cellular damage. To test these hypotheses, we assessed the transcriptional regulation of fitness-related and adaptation-related genes of 11 bacterial strains in relation to their NB and stress exposure under changing salinity conditions. The results suggested that transcriptional regulation levels of fitness- and adaptation-related genes correlated with the NB and/or the stress exposure of the inspected strains. We further identified a short-list of candidate stress marker genes that could be used in future studies to monitor the susceptibility of bacterial populations or communities to environmental changes.

## KEYWORDS

generalists, niche width, salinity, specialists, stress marker genes, stress response, transcriptome

## 1 | INTRODUCTION

Environmental changes and disturbances caused by anthropogenic activities are an increasing threat to natural ecosystems (Grimm et al., 2013). To understand the effect of environmental disturbances on the functioning and composition of species assemblages

it is essential to examine the response and adaptations of individual species in a changing environment. In this context, the assessment of niche dimensions has been one of the most important questions in ecology for understanding the biological adaptation of individual species to environmental change (Slatyer et al., 2013). The ecological niche is a fundamental concept in ecology and is defined as the set of

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd.


environmental conditions, under which species can live and reproduce (Hutchinson, 1957). The niche breadth (NB) is defined as the width of an organism's fitness curve over an environmental gradient and can be used to discriminate between generalist and specialist species (Devictor et al., 2010; Lynch & Gabriel, 1987). Specialists with a narrow niche possess traits that optimize fitness in their optimum habitat at the expense of performance under suboptimal conditions, while generalists with a broad niche are considered to be less competitive under optimal growth conditions but feature higher resistance against changing conditions (Huey & Slatkin, 1976; Jasmin & Kassen, 2007; MacArthur, 1972). Accordingly, specialists are locally adapted organisms and are considered to be particularly endangered by environmental disturbances that are predicted to occur more frequently under climate change scenarios (Colles et al., 2009; Planton et al., 2008; Thuiller et al., 2005).

Different approaches for NB estimations have been used in previous research that were based for example on the number of different resources an organism can use, the number of abiotic associations and biological interactions an organism is involved in, or its tolerance against changing environmental conditions (Sexton et al., 2017). Among these aspects, environmental tolerance has been demonstrated to be one of the most important factors determining the geographical distribution of species (Slatyer et al., 2013).

Stress refers directly to the decrease of an organism's fitness caused by external constraints, such as nutrient limitation, but also physical factors, such as temperature or salinity changes (P. Grime J. & Pierce, 2012). Stress is accordingly linked to NB, where species with a narrow NB exhibit a larger variability in fitness under the same environmental change compared with species with a broad NB (e.g., Matias et al., 2013). However, while NB is a constant parameter for a given species and environmental parameter, the stress exposure of this species depends on the environmental gradient under inspection (Figure 1).

Microbes have been used widely over the past few decades to test general ecological theory (Bell et al., 2009; Ladau & Eloë-Fadrosch, 2019), and including the integration of functional perspectives has helped to elucidate the link between traits and niche-related processes (Krause et al., 2014). The short generation times of microbes and their small size allow for controlled and replicable experiments at different spatial and temporal scales (Jessup et al., 2004). In combination with powerful sequencing techniques, microbes can be used as model systems for studying molecular mechanisms that are associated with the adaptation of species to environmental change concerning their NB. Moreover, microorganisms are the main drivers of carbon and nutrient cycling in all environments and therefore are relevant to the function of ecosystems (Konopka et al., 2015).

The insurance hypothesis states that high species diversity provides insurance against disturbances, because diversity increases the probability that the performance of maladapted species is compensated for by others due to functional redundancy (Yachi & Loreau, 1999). However, it has been shown that different ecological strategies can shape biodiversity–insurance relationships (Matias et al., 2013). The impact of the NB distribution in communities



**FIGURE 1** Schematic illustration for NB estimation and stress exposure. The NB is a constant parameter for each species defined as the length of the environmental gradient covered by the fitness curve of the species at a given fitness value. Stress is quantified as the difference in fitness between any pair of environments E1 and E2. Depending on the position of Env.1 and Env.2 along the environmental gradient, stress can take all values between 0 and the maximum fitness value maxF. We hypothesized that the NB and accordingly the stress an organism is exposed to after environmental change is linked to general patterns of its transcriptional response, which eventually is manifested in the organism's physiological response to changing environmental conditions

on biodiversity–insurance and biodiversity–ecosystem–function relationships (Gravel et al., 2011; Matias et al., 2013) or dispersal and community assembly mechanisms (Shen et al., 2018; Szekely et al., 2013) underlines the importance of the NB concept in understanding community functional and compositional dynamics. Transcriptome data of microbial organisms represent a blueprint of their response and tolerance to environmental change and could be used as a tool to understand the molecular mechanisms behind resistance but also to identify NB distributions in more complex species assemblages. Our main objective was therefore to investigate whether there are commonly valid transcriptional regulation mechanisms that are related to the tolerance-based NB of microorganisms. We are aware of one earlier study addressing general differences in the transcriptional regulation patterns of oligotrophic vs. copiotrophic aquatic bacterial strains (Cottrell & Kirchman, 2016). However, to our knowledge, this is the first study that aims to identify transcriptional patterns that allow us to discriminate between generalist and specialist life histories or to rank stress exposure responses. This will make it possible in future studies to learn more about the susceptibility of bacterial organisms to environmental change.

By definition, the fitness of specialists is more variable along an environmental gradient than that of generalists (Lynch & Gabriel, 1987). We therefore hypothesized that genes that are directly involved in fitness-related traits, such as growth or biomass production, have a higher level of transcriptional regulation in specialists than in generalists under changing environmental conditions. Due to a negative relationship between NB and stress exposure and the causal link between stress and fitness (Figure 1), we expect to detect the opposite relationship between gene regulation and stress

exposure levels (H1). We furthermore hypothesized that generalist species would exhibit higher transcriptional regulation of genes encoding traits involved in the physiological adaptation to changing environmental conditions than specialists (H2). This hypothesis refers to all genes that prevent damage from cells under suboptimal environmental conditions.

To test these hypotheses we incubated 11 bacterial strains with varying tolerance against salinity changes at different salinity levels and tested correlations of transcriptional regulation patterns of fitness- and adaptation-related genes against NB and stress. We furthermore present a list of candidate stress marker genes whose transcriptional regulation correlated to stress exposure and which may be applied in future studies to monitor stress in microbial populations or communities.

We have chosen changing salinity as a stressor because it has been described as one of the major abiotic drivers of microbial community composition across several environments (Lozupone & Knight, 2007). Changing salinity conditions are furthermore environmentally relevant because climate change scenarios predict an increasing occurrence of salinity changes caused by droughts, storms, floods and river run-off (Seneviratne et al., 2012).

## 2 | METHODS

### 2.1 | Bacterial fitness curves

To assess the effect of the NB on bacterial transcriptional activity, we included bacterial model strains in our study that belong to a larger collection of 148 isolates originating from several aquatic environments with different salinities (Matias et al., 2013). The NB of all isolates had been estimated by Matias et al. (2013) via discrete optical density (OD) measurements after 48 h of growth and using a polynomial model. For our study, based on these earlier NB estimations, we selected 11 strains with contrasting halotolerance that exhibited a maximum at a salt concentration of  $\sim 30 \text{ g L}^{-1}$  NaCl (hereafter used as salinity) and that affiliated with different phylogenetic lineages (Table 1; Table S1).

To ensure the purity of all isolates, cryopreserved cells from all 11 strains were re-isolated after plating on standard LB agar to which  $20 \text{ g L}^{-1}$  NaCl was added (final salinity of  $30 \text{ g L}^{-1}$  NaCl). A single colony was picked and transferred into a tube containing 2 ml of liquid LB medium (final salinity of  $30 \text{ g L}^{-1}$  NaCl), incubated at room temperature, and then cryopreserved using glycerol as cryoprotectant for downstream applications.

The number of only 11 strains included in our study allowed us to apply a more labour-intensive but also more accurate protocol compared to an earlier approach (Matias et al., 2013) to assess their tolerance against salinity changes: cells from the strains were defrosted for 15 min at room temperature and used to inoculate sterile standard LB medium (Carl Roth; final salinity of  $30 \text{ g L}^{-1}$  NaCl), in which they were incubated for 2 days at  $25^\circ\text{C}$ . Next, 300  $\mu\text{l}$  sterile

standard LB medium (Carl Roth) with different NaCl concentrations (10, 20, 30, 40, 50, 60, 70, 80, 90 and  $100 \text{ g L}^{-1}$ ) was distributed in 96-well microplates and inoculated with the same number of cells from the individual strains, respectively (six replicates per strain and salt concentration). Growth of the strains at  $25^\circ\text{C}$  was monitored in each well by hourly measuring their cell densities via the OD at 600 nm (after shaking at medium intensity, with orbital shaking set to 20) in a Paradigm Microplate reader (Molecular Devices) until the plateau phase was reached (Figure S1).

The fitness of strains under the different salt concentrations was assessed based on their growth curves in each medium by using their maximum cell densities and their growth rates. Growth rates were estimated by fitting a logistic growth equation as detailed elsewhere (García et al., 2018) until maximal cell densities were reached. We further excluded the lag-phases defined by the period until initial OD values had doubled from growth rate estimations because the lag-phases were probably impacted by an acclimation phase only in those media that differed from the salinity in the preculture medium. The growth curves of some strains differed over the salinity gradient mainly in their maximal cell densities, while others differed more strongly in the incubation time after which the maximal growth density was reached and accordingly in their growth rates (Figure S1). To integrate these different aspects of fitness we created a combined fitness index from the product of maximum cell densities and growth rates for each strain in the given medium. This combined fitness index was used for all downstream analyses. Fitness curves were fitted by applying either a lognormal or a gaussian model to the fitness values along the salinity gradient, while the best model was selected by the Akaike information criterion (Figure 2; Table S1).

The NB of the strains was calculated after normalizing the fitness curves by dividing by the maximal modelled fitness value as the salinity range in  $\text{g L}^{-1}$  NaCl where the individual strains reached at least 50% of the extrapolated normalized fitness. The resulting fitness curves featured in most cases asymmetric shapes (Skewness  $>0$ ; Table 1), implying that the strains were characterized by differential tolerances against hypoosmotic and hyperosmotic stress (Figure 2). For this, we divided the NBs between the right (hyperosmotic) and left (hypoosmotic) sides relative to the modelled optimal salt concentration to obtain a side-dependent hyperosmotic and hypoosmotic NB for each strain (Figure 3; Table 1). Beyond NB estimations, fitness curves allow us also to delineate stress exposure levels in response to changing salinity, by subtracting the fitness values at the start and end points of a salinity gradient (Figure 3). NBs are negatively related to stress levels if the stress values are assessed for salinity gradients that are located symmetrically around the optimum fitness value (Figure 4). However, stress values can be determined for every arbitrary salinity gradient, within which the growth of a strain is detected. Accordingly, for any given strain and independently of its NB, all stress values between zero and one can be obtained in dependence of the position and strength of the salinity gradient of interest.

TABLE 1 Summary of strains used in this study and their niche breadth (NB) characteristics

Strain ID	Phylogeny (Class   Order   Family   Genus)	NB (NaCl g L <sup>-1</sup> )	Optimum modelled salinity (NaCl g L <sup>-1</sup> )	Skewness	Hypo-osmotic NB	Hyper-osmotic NB
S331	Gammaproteobacteria   Pseudomonadales   Halomonadaceae   <i>Halomonas</i>	143	-5	0.93	45	98
S337	Gammaproteobacteria   Pseudomonadales   Marinomonadaceae   <i>Marinomonas</i>	48	42	1.14	17	31
S338	Gammaproteobacteria   Pseudomonadales   Marinomonadaceae   <i>Marinomonas</i>	53	33	1.44	17	36
S366	Gammaproteobacteria   Pseudomonadales   Marinomonadaceae   <i>Marinomonas</i>	56	35	1.32	18	38
S374	Gammaproteobacteria   Pseudomonadales   Marinomonadaceae   <i>Marinomonas</i>	58	35	1.21	18	40
S432	Gammaproteobacteria   Enterobacteriales   Alteromonadaceae   <i>Pseudoalteromonas</i>	50	44	1.02	18	32
S479	Gammaproteobacteria   Enterobacteriales   Alteromonadaceae   <i>Pseudoalteromonas</i>	58	53	0.15	29	29
S490	Gammaproteobacteria   Enterobacteriales   Alteromonadaceae   <i>Pseudoalteromonas</i>	51	42	1.22	18	33
S599	Alphaproteobacteria   Rhodobacterales   Rhodobacteraceae   <i>Celeribacter</i>	29	49	0.49	12	17
S618	Gammaproteobacteria   Enterobacteriales   Vibrionaceae   <i>Vibrio</i>	67	21	1.23	14	53
S630	Actinobacteria   Actinomycetales   Micrococcaceae   <i>Nesterenkonia</i>	140	41	0.61	29	111


## 2.2 | Transcriptional response to changing salinity levels

For downstream RNA extractions, each strain was grown in duplicate at three salinity levels and 25°C (100 r.p.m.) in sterile glass tubes containing 15 or 50 ml LB medium (Table S2). We originally had planned to grow the strains for this experiment at their optimal salinity levels (S<sub>2</sub>), as well as two salinity levels, each located symmetrically 15 g L<sup>-1</sup>

NaCl at the hyper- and hypoosmotic side from the optimum (S<sub>1</sub>, S<sub>3</sub>) (Figure 3). However, in several cases, the incubation salinity levels were shifted relative to the optimum salinity, because we had used polynomial models for NB estimations to design the experiment as detailed earlier (Matias et al., 2013). We later decided to optimize the fitness models as described above (Figure 5).

To monitor the growth of the strains during the experiments, 300- $\mu$ l aliquots from the inoculated medium were pipetted in


96-well microplates. The microplates were incubated at the same temperature as in the glass tubes either in a Paradigm Microplate reader (Molecular Devices) or in a VICTOR Multilabel Plate Reader




**FIGURE 2** Fitness estimations and modelled curves. Black points indicate the fitness indices from six replicates per strain at each salinity level, respectively. Grey lines indicate the fitness curves fitted according to the Akaike information criterion either via a lognormal (L) or gaussian (G) model

(PerkinElmer), where hourly measurements for OD (600 nm) were performed after 5 s of shaking. During the exponential phase that was monitored in real-time for each bacterial strain via OD measurements (Figure S2), samples for RNA extraction and cell enumeration were harvested from the glass tubes. In order to stop the incubations, 13.5 or 48.5 ml of the cultured bacteria was fixed by adding 1.5 or 5.4 ml of a 5% solution of phenol/chloroform 5:1 (Sigma-Aldrich) diluted in absolute ethanol (Sigma) (Feike et al., 2012). The fixed cell cultures were centrifuged (8500 g for 5 min) and the supernatant was carefully discarded. Next, 300  $\mu$ l of RNA later (Sigma-Aldrich) was added to the pelleted cells that were resuspended by pipetting up and down. The cell suspension was transferred into a 2-ml tube, flash-frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  for later RNA extraction (Table S2 for incubation details).

From the remaining volume in the incubation tubes, 1200  $\mu$ l was fixed with glutaraldehyde (0.1% final concentration) and kept at  $-80^{\circ}\text{C}$  until later analysis for cell enumeration. The number of cells was estimated by flow cytometry in a Cytoflex Flow Cytometer (Beckman Coulter) using a standard procedure described elsewhere (Marie et al., 2000).



**FIGURE 3** Schematic illustration for side-dependent NBs and stress exposure. Most of the 11 model strains were characterized by asymmetric fitness curves. The hyper- and hypoosmotic NBs are indicated with red and blue arrows, respectively. Stress exposure levels along salinity gradients (here S1:S2, S2:3 and S1:S3) can be estimated by subtracting fitness values at the start and end point of the salinity gradients under consideration (here  $\Delta F_1$ ,  $\Delta F_2$  or  $\Delta F_3$ ). In contrast to the NBs, stress exposure is not a constant parameter for a given species but depends on the position and length of the salinity gradient under inspection (i.e., the position of S1, S2 or S3)



**FIGURE 4** Regression between side-dependent NB and stress estimations for the 11 model strains. The black cross-marks (and solid black trend line:  $R^2 = .73$ ,  $p < .001$ ) represent stress levels assessed for salinity gradients that were located  $\pm 15 \text{ g L}^{-1}$  NaCl symmetrically around the optimum fitness value and the corresponding side-dependent NBs that were used for mixed model regressions against the NB. The grey circles represent stress levels between salinity concentrations where the optimum was crossed and/or delineated from a salinity gradient of  $30 \text{ g L}^{-1}$  NaCl. The corresponding NB values were in this case estimated from the proportional contribution of the hypo- or hyperosmotic NB that was covered by the respectively considered salinity gradient. The dotted trend line was fitted by including all data points ( $R^2 = .23$ ,  $p < .021$ )

### 2.3 | DNA extractions for genome sequencing

The strains were incubated in 2 ml LB (salinity of  $30 \text{ g L}^{-1}$  NaCl) for 5 days at room temperature, pelleted and stored at  $-30^\circ\text{C}$  until DNA extraction. DNA was extracted following the standard protocol of the Genra Puregene Cell Kit (Qiagen). The concentration and quality of the eluted DNA were tested by electrophoresis (1% agarose gel) and using a Quantus fluorometer using the PicoGreen assay (Promega). The extracted DNA was sent for library preparation and genome sequencing (DNA shot gun library, insert size: 300 bp,  $2 \times 150$ -bp reads, Illumina NextSeq 500 V2). Library demultiplexing of libraries via the Illumina BCL2FASTQ 2.17.1.14 software (two mismatches allowed, minimum length  $<20$ ) was carried out by the sequencing company.

### 2.4 | RNA extractions for transcriptome sequencing

Cells harvested during their exponential growth phase from the incubation experiment at different salt concentrations were defrosted at room temperature and pelleted by centrifugation, and the supernatant containing RNA later solution was discarded. The RNA extractions were performed for all strains using the Direct-zol


RNA-Miniprep Plus (Zymo Research) following the manufacturer's instructions, except for strain S630. For downstream absolute transcript quantification (Satinsky et al., 2013), 5 ng of a spiked-in RNA standard ( $=9 \times 10^6$  standard molecules) was added to the cell lysis solution of each individual extraction (TRIzol reagent, the first step of the extraction protocol). This standard was obtained after linearizing the plasmid pFN18A HaloTag T7 Flexi (Promega) and transcribing a 970-bp part of it into RNA using the MEGAscript T7 Transcription Kit (ThermoFisher). The sequence of the used standard is given in File S1.

For strain S630, it was not possible to obtain sufficient RNA yield when using the Direct-zol RNA-Miniprep Plus kit for RNA extraction. For this reason, we implemented the SNAP protocol for gram-positive bacteria published elsewhere (Stead et al., 2012) with some modifications. In short, cell pellets were resuspended in  $300 \mu\text{l}$  SNAP RNA extraction solution and 5 ng of the internal RNA standard was added for downstream absolute transcript quantification. After the addition of low binding zirconium beads (OPS diagnostics) the suspension was treated at  $6 \text{ m s}^{-1}$  for  $2 \times 45 \text{ s}$  in a FastPrep-24 5G MP (Biomedicals) for mechanical cell disruption and subsequently heated for 7 min at  $95^\circ\text{C}$ . To enhance the efficiency of the SNAP method, we further placed tubes with the extraction solution in a beaker filled with 500 ml water and microwaved them for 45 s (Brandt Microwave, model SM2602B) at the defrosting mode ( $\sim 300 \text{ W}$ ). While a similar microwaving protocol was originally developed to permeabilize cell walls for downstream fluorescence *in situ* hybridization (FISH) applications (Tischer et al., 2012), this additional step also increased the RNA yield for strain S630. The extracted RNA in the SNAP RNA extraction solution was purified with the RNA Clean & Concentrator-5 kit including the optional on-column DNA digestion step (Zymo Research).

An additional DNA removal step was performed for all RNA extracts using the TURBO DNA-free Kit (Invitrogen). Then, the eluted RNA was concentrated by applying the RNA Clean & Concentrator-5 kit (Zymo Research). The concentration of the eluted RNA was estimated in a NanoVue Plus Spectrophotometer (Biochrom), while RNA quality was visually evaluated by inspecting RNA molecule length profiles using an Agilent 2100 Bioanalyzer (Agilent Technologies; RNA6000 NanoKit).

To reduce sequencing costs, RNA extracts from three of four strains that according to taxonomic annotations of genome sequences (see Bioinformatic processing) affiliated with different genera were pooled equimolarly (Table S2). For each consecutive RNA purification step, we noted the volume that was transferred from the preceding step. This allowed us to follow back the fraction of the initial lysis solution and, accordingly, the number of standard molecules from each individual extraction that were added to the respective RNA pool. Based on this information, we later assigned the corresponding fractions of sequenced reads encoding the added standard to the individual RNA extracts in order to obtain information about total transcripts per cell. The extracted RNA was sent for library preparation and genome sequencing (mRNA nonstranded library, rRNA depletion via the Ribo-Zero rRNA removal Kit for Bacteria, insert size: 300–500 bp,  $2 \times 150$ -bp reads, Illumina NextSeq 500 V2).





**FIGURE 5** Modelled fitness for the 11 strains used in this study. (a–k) Fitted fitness curves and the sampled salinity levels (dashed black lines, from left to right  $S_1$ ,  $S_2$ ,  $S_3$  in each panel) to estimate the transcriptional response to changing salinity relative to the fitness curves. The vertical black line indicates the optimal fitness of each strain

## 2.5 | Bioinformatic processing

Library demultiplexing of all obtained raw reads via the Illumina BCL2FASTQ 2.17.1.14 software (two mismatches allowed, minimum length <20) was carried out by the sequencing company. The resulting adapter-free paired-end reads were trimmed using the SICKLE software version 1.33 (quality-type sanger, quality-threshold 20, length-threshold 75) (Joshi & Fass, 2001).

The trimmed reads from the genome sequences were then assembled using the SPADES genome assembler version 3.13.0 (Bankevich et al., 2012) with the following settings: k-mers from 21 to 99, four nucleotide steps. Taxonomic assignments for the assembled genomes were performed using the GTDB-TK version 0.2.2 software (Chaumeil et al., 2020) against the GTDB database. Genes on the assembled contigs were predicted via the PRODIGAL version 2.6.3 software (Hyatt et al., 2010). The predicted genes were functionally annotated via DIAMOND BLAST version 0.8.22 against genes in the KEGG database (Kanehisa et al., 2007, downloaded May 2016) with an e-value cutoff of  $1e-5$ .

The SORTMERA version 1.9 software (Kopylova et al., 2012) was used to separate the trimmed nonprotein-coding RNA from protein-coding reads. Spike-in standard reads in each sample were detected using the LASTAL software version 393 (Kielbasa et al., 2011), and counted

for downstream absolute transcript quantification. Quality-trimmed protein-coding reads were mapped on the predicted genes from the assembled genomes using the BOWTIE2 version 2.3.4.3 software set to very-sensitive-local (Langmead & Salzberg, 2012) and summarized by the FEATURECOUNTS version 1.4.6-p2 software (Liao et al., 2014).

## 2.6 | Transcriptional regulation patterns

Per-cell transcriptional regulation was calculated as  $\log_2$  fold-changes of per-cell transcription levels between two salinity levels and was determined for the individual genes annotated from each strain and each pair of salinity conditions using the R package DESeq2 (Love et al., 2014). The regulation patterns were delineated from the absolute transcript per-cell transcription levels, as detailed elsewhere (Beier et al., 2019). In short, raw count data for individual genes were normalized using the count data of the added spike-in standard that were multiplied with the total number of cells subjected to RNA extraction and divided by the amount of added standard via the “controlGene” option of the DESeq2 “estimateSizeFactors” function. As expected, we found an almost perfect linear correlation between values for total transcripts per cell calculated with a previously published formula (Satinsky et al., 2013) and the total transcripts per cell



variable obtained by summing count data after the DESeq2 normalization step ( $r = 1.00$ ; Figure S3). We did not specifically select genes based on the DESeq2 significance levels for transcriptional regulation, because the downstream regression analyses were designed to also include genes that exhibited low or no transcriptional regulation.

We used the value for total transcripts per cell (DESeq2-estimated) to test the overall correlations between total per-cell transcription levels against NB and stress estimations, to screen for potential relationships that could bias our hypothesis testing.

## 2.7 | Definition of gene categories

To test the hypothesis H1 that genes with a direct link to an organism's fitness are more strongly regulated in strains with narrow NB, we built three categories of putatively fitness-related genes encoding the following proteins: (i) DNA polymerase enzymes, which are essential for DNA replication and cell proliferation. This category included all genes containing the text string "DNA polymerase" in the gene description. (ii) RNA polymerases, which transcribe genes into mRNA and are accordingly the first step in cellular biomass production via protein synthesis. RNA polymerase genes were defined as the genes classified as protein-coding genes in the KEGG pathway ko03020. (iii) Ribosomal proteins, which are essential units of the ribosomes that catalyse the translation of mRNA into proteins and are accordingly the second step in cellular biomass production via protein synthesis. We defined all genes annotated to protein-coding genes in the KEGG pathway ko03010 as ribosomal proteins.

These three categories comprise core genes that are highly conserved across organisms in all three domains of life (Kültz, 2003) and the individual encoded proteins in these categories (e.g., ribosomal proteins) exhibit their catalytic activity as subunits in a larger assembly of catalytic active units. We therefore selected the individual genes in each category from the pool of genes shared by all of the 11 model strains (253 shared genes; Table S3) and considered the transcriptional regulation of the individual genes in each category as random factors for downstream statistical analyses.

To test the hypothesis H2 that genes encoding cell adaptation against salt stress are more strongly regulated in strains with broader NB, we constructed another two categories with potential adaptation-related genes: (iv) Genes encoding the transport of osmoprotective compounds, mainly including compatible solutes, but also other osmolytes, such as urea. Compatible solutes are small organic molecules such as sugars or amino acids that do not have detrimental effects on cell functions (Welsh, 2000). However, the use of compatible solutes as osmoprotectants is highly species-specific (Bougouffa et al., 2014; Sévin et al., 2016). For instance, glycine betaine, which serves as an osmoprotectant in organisms of all domains (Empadinhas & Viète-Vallejo, 2008), was not observed to protect the archaeon *Sulfolobus solfataricus* against salt stress (Park & Lee, 2000). We therefore selected from all transporter genes that were expressed in at least one of the model strains a list of potential osmoprotectant transporters (Table S4). (v) Heat-shock proteins

(HSPs), which catalyse the folding and unfolding of macromolecules and are therefore involved in the repair of damaged macromolecules (Kültz, 2003; Lindquist & Craig, 1988). However, different from the osmoprotectant genes listed above, genes in this category do not initially prevent cells from damage but step into action only after damage has occurred. This category included all genes containing the text strings "heat-shock" or "chaperone" in the gene description.

Different from in the categories encoding fitness-related proteins, we assumed that the individual genes in the categories of the adaptation-related proteins act independently from each other and their protective effect is additive. For instance, the more different are osmoprotectants transported the stronger the protection against osmotic stress. Genes encoding the adaptive response against osmotic stress are furthermore and in contrast to fitness-related genes species-specific (Sévin et al., 2016), and only two among the selected potential osmoprotectant transporters (K15268, K02030) were shared among all model strains from this study (Table S3). We therefore considered the additive transcriptional response by summing the fold change of the individual genes in categories iv and v, respectively for downstream statistical analyses.

## 2.8 | Regression of gene regulation patterns against NB and stress levels

We applied mixed linear models (MLMs) to test if the transcriptional regulation levels of genes within each of the above-described categories were correlated in the predicted direction with either NB (log-transformed) or stress exposure along the respective salinity gradient of the individual strains. The MLMs were implemented in R using the package nlme (Pinheiro et al., 2020), and the R-package olsrr (Hebbali, 2020) was applied to verify the normal distribution of the residuals in all MLMs using Kolmogorov–Smirnov tests. In a few cases where a normal distribution of the residuals could not be verified, transcriptional regulation levels were subjected to a square root or inverse transformation.

In the case of regressions against NB, we considered regulation patterns along salinity gradients of  $15 \text{ g L}^{-1}$  NaCl, but only if the salinity optimum ( $\pm 4.5 \text{ g L}^{-1}$  NaCl, equivalent to 30% of the total change) of the respective strain was not passed. With this rule, a maximal of two values for gene regulation per strain and gene were considered (Table 1; Table S1). Because these regulation values referred to either the hyper- or the hypoosmotic side, an MLM was performed against the respective values for side-dependent NBs, with the side-dependent NB as a fixed factor. Replicate regulation values per strain (retrieved from different salinity gradients on either of the two sides and, where applicable, individual genes in a category that were shared by all 11 strains) were used as random factors. An analogous approach was used to test if the degree of gene regulation increased or decreased with increasing stress exposure. However, in this case, we considered stress levels between all sampled salinity gradients per strain as the replication level and accordingly as a random factor in the MLM (three regulation values per strain and gene:  $\Delta F_1$ ,  $\Delta F_2$ ,  $\Delta F_3$ ; Figure 3; Table 1). Stress levels showed a pronounced

negative correlation with NB, if only stress values from the salinity range  $15 \text{ g L}^{-1}$  NaCl and not crossing the optimum were considered (i.e., those data points considered for the regressions against NB; Figure 4). However, a reduced correlation strength between NB and stress levels was observed when all data points were considered (Figure 4).

While our hypotheses referred only to the extent of regulation activity regardless of whether genes were up- or downregulated, we also inspected regressions taking into account the regulation direction to see whether genes in the categories were either up- or downregulated in response to stress.

The slopes of the MLM, as well as  $p$ -values, were retrieved to describe the direction of the regression and test the hypotheses. For hypotheses H1 and H2, we specifically expected that gene regulation of fitness-related genes decreases with the increasing NB (slope  $< 0$ ), while we expected to find the opposite trend for adaptation-related genes (slope  $> 0$ ). Because the R-package nlme version 3.1-148 does not enable us to set up mixed models using one-tailed test designs we report here the more stringent  $p$ -value from the two-tailed test design. In the case of correlations with the hypothesized direction, however, we consider a significance level of  $p < .1$  to be valid. To evaluate the proportion of variance that was explained by the fitted MLM, pseudo  $R^2$  values for the overall model were estimated using the `r.squaredGLMM` function from the R-package `MunIn` (Bartoń, 2020).

## 2.9 | Detection of stress marker genes

To detect potential individual stress marker genes, we applied an MLM on each of the 253 genes shared by the 11 strains using stress as a fixed factor and the three replicate stress levels per strain as random factors. In this case, we performed the regression analyses considering the direction of the gene regulation under stress conditions (up- or downregulation). The resulting  $p$ -values were adjusted to account for false discovery rates after multiple comparisons (Benjamini & Hochberg, 1995).

The R codes used for the statistical evaluations that were performed for this study are available on GitHub (<https://github.com/sarabeier/Strains.NB>).

## 3 | RESULTS

### 3.1 | Strain characteristics and total transcript regulation

The bacterial model strains covered several phylogenetic lineages (Alphaproteobacteria, Gammaproteobacteria and Actinobacteria) with representatives affiliated to the orders Pseudomonadales, Enterobacterales, Rhodobacterales and Actinomycetales (Table 1). The strains exhibited side-dependent NBs that ranged from 12 to  $111 \text{ g L}^{-1}$  NaCl. Along the NB continuum, shorter NBs were dominated

by hypoosmotic estimations, while most broader NBs corresponded to hyperosmotic values (Figure 6).

Per-cell transcription levels were highly variable among strains, differing by up to a factor of 5 (i.e., S630 vs. S490, Figure S4). However, the overall per-cell transcriptional regulation did not correlate significantly with NB or with stress estimations and  $R^2$  values were consistently low (Table 2, Figure 7a–c).

### 3.2 | Regulation of fitness-related genes

We detected only two genes that were shared across strains in the DNA and RNA polymerase categories, respectively. In contrast, the ribosomal protein category contained 32 genes shared by all strains (Table 2). While no significant trend was noted for the regulation of genes encoding DNA polymerases (Table 2), the regulation of genes encoding ribosomal proteins and RNA polymerases decreased in agreement with H1 significantly with increasing NB ( $p = .003$  and  $p = .041$ , respectively; Figure 7g,j; Table 2). The direction of the relationship turned when considering stress exposure instead of NB as an explanatory variable for gene regulation. However, only genes encoding ribosomal proteins were significantly regulated with increasing stress levels ( $p < .001$ ; Figure 7i). An inspection of the direction of gene regulation revealed a significant upregulation of genes encoding ribosomal proteins and RNA polymerases under increasing stress exposure (Figure 7i; Table 2) and a nonsignificant trend for upregulation of DNA polymerase-encoding genes (Figure 7f; Table 2). However, in all three categories, individual genes were also downregulated under stress, particularly if stress levels were retrieved from salinity ranges crossing the salinity optimum (Figure 7c,f,i).

### 3.3 | Regulation of adaptation-related genes

The number of potential osmoprotectant transporter genes per strain varied between 20 and 93 (Table 2). The summed upregulation level of these genes along with increasing NB exhibited a positive slope, which was in agreement with the above defined significance level of  $p < .1$  ( $p = .076$ ; Table 2; Figure 7m). On the other hand, stress did not seem to affect the absolute regulation of these genes nor the direction of the regulation (Table 2; Figure 7n,o).

The regulation of genes encoding HSPs did not exhibit a significant correlation in the predicted direction with NB or with stress levels (Table 2; Figure 7p,q), and instead trends contrary to the prediction were observed. Similar to the osmoprotectant transporter category, no significant trend for an up- or downregulation of HSPs under stress was detected (Table 2; Figure 7r).

### 3.4 | Candidate marker genes for stress

Only one of the individual genes shared among all 11 model strains exhibited a significant relationship with stress levels after  $p$ -value



TABLE 2 Summary of mixed linear models (MLMs) for gene regulation against niche breadth (NB) and stress including genes grouped into categories

Classification	Functional categories	Number of considered genes per strain	Absolute gene regulation vs. NB			Absolute gene regulation vs. stress <sup>a</sup>			Gene regulation vs. stress <sup>a</sup>		
			Slope	R <sup>2</sup>	P	Slope	R <sup>2</sup>	P	Slope	R <sup>2</sup>	P
Fitness-related genes	Total transcripts	1 <sup>b</sup>	-0.06	.01	.768	0.38	.02	.458	0.28	.07	.749
	DNA polymerases	2 <sup>b</sup>	-0.19	.02	.399	0.42	.01	.399	1.19	.13	.151
	Ribosomal proteins	32 <sup>b</sup>	-0.08	.09	.003*	0.30	.02	.000***	1.19	.13	.000***
	RNA polymerases	2 <sup>b</sup>	-0.46	.12	.041*	0.87	.04	.113	1.84	.14	.030*
Adaptation-related genes	Transport of osmoprotectants	25–93 <sup>c</sup>	0.52	.19	.076●	-0.23	.00	.767	-0.44	.02	.940
	HSP	6–17 <sup>c</sup>	-0.24	.06	.332	1.26	.14	.030	2.06	.17	.571

Note: \*\*\*  $p < .001$ , \*  $p < .05$ , ●  $p < .1$ . All values considered in our analyses as significant ( $p < .1$  under the condition that the regression followed the predicted direction) are printed in bold.

<sup>a</sup>The displayed statistical output parameters refer to regressions including all stress exposure data points. The statistical output parameters for regressions including only data based on 15 NaCl L<sup>-1</sup> salinity gradients in which the performance optimum was not passed and which correspond to the data points included in the NB regressions are displayed in Table S5.

<sup>b</sup>MLMs were performed using the individual genes as a random factor.

<sup>c</sup>MLMs were performed using the additive transcriptional regulation on individual genes.

between transcriptional regulation levels and NB or stress, we have differentiated between genes encoding fitness-related traits and genes encoding traits related to the adaptation of organisms to changing environmental conditions.


Indeed, in agreement with our hypotheses, the results suggested that transcriptional regulation levels of either fitness- or adaptation-related genes if inspected separately correlated with NB, and therefore the life history of species as well as stress levels to which cells were exposed.

Although the regulation levels of DNA polymerases did not decrease significantly with increasing NB, our analyses confirmed as expected in H1 a significant decrease of RNA polymerases as well as ribosomal protein transcription levels along with increasing NB (Table 2). Both RNA polymerases and ribosomal proteins are directly involved in cellular translation activity and accordingly in biomass production. Unlike NB, which is a constant for a given strain, cellular fitness and stress levels are closely linked, even if environmental gradients of different lengths and exceeding the point of optimal cellular performance are taken into account (Figure 2), as was the case in our analyses. The direct involvement of transcriptional regulation of RNA polymerases and ribosomal proteins in cellular fitness was therefore also reflected in the positive relationship of gene regulation patterns with stress levels (Table 2). While a downregulation of genes encoding biomass is to be expected intuitively under increasing stress and correspondingly decreasing fitness levels, we observed the opposite trend. This was particularly pronounced if environmental ranges that did not pass the fitness optimum were considered (Figure 7f,i,l). This observation suggests that the detected transcriptional upregulation of genes involved in translation activity under elevated stress levels could be linked to mechanisms for replacing or repairing damaged proteins or other macromolecules, as has been outlined elsewhere (Evans & Hofmann, 2012).


Our results further give certain evidence in support of H2, where we expected an increased transcriptional regulation of genes encoding the adaptation to changing salinity along with increasing NB. The expected positive relationship was observed for a list of candidate genes encoding the transport of osmoprotectant substances (Figure 7m).

The nonsignificant relationship of the regulation of osmoprotectant transporters against cellular stress levels was not unexpected (Table 2), at least not if stress levels of salinity ranges passing the optimum were included in the regression because within the same organism different osmoregulation mechanisms may be relevant under hypo- or hyperosmotic stress (Deole & Hoff, 2020; Lin et al., 2017).

Also, as a consequence of the species-specific use of different osmoprotectants, the uptake of compounds such as certain amino acids serves in one species for osmoregulation (Park & Lee, 2000) while the uptake of the same amino acid could be instead associated with biomass production in another species. This blurred separation of transport mechanisms, which may either represent adaptation-related traits via their association with osmoregulation or be related to cellular fitness, probably introduced noise into the regression



**FIGURE 7** Regressions of gene regulation against side-dependent niche breadth and stress. MLM of the absolute gene regulation against NB (log-transformed, left panels) and stress (centre panels), and directional gene regulation against stress (right panels). (a–c) Total transcripts per cell, (d–f) DNA polymerases, (g–i) ribosomal proteins, (j–l) RNA polymerases, (m–o) transport of osmoprotectants and (p–r) HSPs. Black cross-marks (and the black solid trend lines) represent pairwise comparisons that did not cross the optimum fitness for NB and stress. Grey circles present all pairwise comparisons between salinity concentrations where the optimum was crossed and/or delineated from a salinity gradient of  $30 \text{ g L}^{-1} \text{ NaCl}$ . The dotted trend lines were fitted by including all data points



**FIGURE 8** Regression of candidate stress marker gene regulation against stress exposure levels. This graphic displays the regression of seven candidate stress marker genes that were selected because of their significant correlation ( $p < .05$ , no adjustments for multiple testing) against the exposure of the 11 model strains to osmotic stress

analyses performed. The positive regression found between regulation and NB (Figure 6m) shows, however, that the role of the listed candidate osmoprotectant transporters (Table S4) in our experimental setup actually laid largely in their participation in cellular osmoregulation.

We also inspected regulation patterns of HSPs that are recognized as classical stress response proteins that due to their folding and unfolding function catalyse the repair of damaged proteins (Roncarati & Scarlato, 2017; Sørensen et al., 2003). Their expression may increase the resistance of cells to stress and thereby impact an organism's life history (Sørensen et al., 2003). We therefore classified HSPs as genes encoding an adaptation to environmental change and which are assumed to detect a more pronounced regulation of HSPs in strains with broader NB (H2). This, however, was not the case. Even though not significant, an opposite than expected trend for the relationship between HSP regulation and NB was observed (Figure 7p). Although elevated HSP expression may, on the one hand, impact the resistance of cells against stress (Sørensen et al., 2003), expression levels of HSPs have also been shown to increase under stress (Evans & Hofmann, 2012): this was reflected in a trend for a positive correlation between stress levels and HSP regulation (Table 2). However, in our sample design, stress levels correlated negatively with NB (Figure 4), and obviously the stress-induced regulation of HSPs masked the possible effect on NB-dependent regulation. If applying a sample design in which we would have chosen

salinity ranges to keep the stress levels rather than the environmental distance constant, we might have been able to detect the predicted positive correlation between NB and HSP regulation.

Beyond testing hypotheses H1 and H2 we also screened absolute regulation patterns of the genes shared across all strains without any a priori hypothesis for their correlation with stress exposure and present a list of candidate stress marker genes. This was done to test if individual genes were better markers of stress compared to the pool of genes included in the categories detailed above. Except for the *phoH*-like ATPase *phoH2*, we did not observe any individual genes whose regulation correlated after adjustment for multiple comparisons significantly to stress levels ( $p_{\text{adj}} < .01$ ). Regardless, we reported here all genes with unadjusted  $p < .05$  to be considered as potential candidate stress marker genes. The gene *rpsI* encodes a ribosomal protein and is consequently involved in biomass production and was within the group of genes considered for hypothesis testing. The upregulation of *rpsI* under elevated stress levels is furthermore in agreement with the overall regulation patterns observed for other ribosomal proteins (Figure 7g) and might be explained by enhanced transcriptional activity under stress to induce the replacement of damaged proteins (Evans & Hofmann, 2012). Although not considered to be ribosomal, *ybeB* also encodes a ribosome-associated protein, which has been described as a ribosomal silencing factor that downregulates protein synthesis when cells enter the stationary phase (Häuser et al., 2012). This agrees with observations that mechanisms that induce growth arrest are typically upregulated under stress (Kültz, 2003). Both *rpsI* and *ybeB* can be considered as fitness-related genes to which hypothesis H2 applies, and meaningful mechanisms can explain the observed correlation of gene regulation patterns with stress levels. Accordingly, also the nonadjusted  $p$ -value may in these cases provide sufficient statistical evidence to support the classification of these genes as stress markers. Furthermore, both genes are highly conserved and encoded by almost all bacteria (Häuser et al., 2012; Lecompte et al., 2002), which allows the design of primers or probes to monitor transcriptional expression levels specifically of target genes, for example via quantitative polymerase chain reaction approaches (Beier, Gálvez, et al., 2015). On the other hand, it would also be possible to use metatranscriptome approaches to taxonomically assign different *rpsI* or *ybeB* transcript variants and thereby allow tracking of taxon-specific stress levels. Yet, in the case of *rpsI*, it must be taken into account that we detected transcriptional upregulation in response to stress exposure by comparing populations during their exponential phase. Stress due to nutrient restrictions and an associated switch from the exponential to the stationary growth phase can instead lead to a downregulation of ribosomal protein transcription



TABLE 3 Summary of significant mixed linear models of regulation of shared genes against stress levels

KEGG ID	Gene ID	Gene description	Slope	R <sup>2</sup>	<i>p</i>	<i>p</i> <sub>adj</sub>	Occurrence in prokaryotes (%) <sup>a</sup>
K07175	<i>phoH2</i>	PhoH-like ATPase	-5.14	.04	<b>.000***</b>	<b>.088</b>	34
K13953	<i>adhP</i>	Alcohol dehydrogenase, propanol-preferring	-3.57	.09	<b>.038*</b>	.982	49
K03522	<i>fixB, etfA</i>	Electron transfer flavoprotein	-2.10	.15	<b>.015*</b>	.982	53
K09710	<i>ybeB</i>	Ribosome-associated protein	2.06	.04	<b>.018*</b>	.982	86
K15268	<i>eamA</i>	O-acetylserine/cysteine efflux transporter	2.17	.06	<b>.025*</b>	.982	16
K02996	<i>rpsL</i>	Small subunit ribosomal protein S9	2.48	.08	<b>.045*</b>	.982	99
K06075	<i>slyA</i>	MarR family transcriptional regulator	2.50	.08	<b>.004*</b>	.450	20

Note: \*\*\**p* < .001, \**p* < .05. All values considered in our analyses as significant (*p* adj < .1) are printed in bold.

<sup>a</sup>The fraction of prokaryotic genomes in the KEGG database (version 2016) carrying the gene.

(Aseev et al., 2016), and regulation patterns of *rpsL* may not apply well as a stress marker in all situations. In contrast, analogously to results in our study, *ybeB* was also upregulated during the stationary growth phase in response to nutrient limitation (Häuser et al., 2012). The upregulation of *ybeB* may therefore be universally associated with exposure of bacterial cells to multiple stressors.

The presence of the remaining listed candidate stress marker genes was less conserved across prokaryote genomes (Table 3) and the exact mechanisms that could explain the detected correlations of regulation patterns with stress levels were less evident. However, several of them were observed earlier in connection with some kind of stress response (*syIA*: Alekshun & Levy, 1999; *eamA*: Ohtsu et al., 2015; *phoH2*: Andrews & Arcus, 2020). The genes *adhP* and *fixB* are both involved in anaerobic energy production processes (Rao & Stokes, 1953; Weidenhaupt et al., 1996) and their downregulation along with increasing stress levels is more difficult to interpret.

The 11 model strains considered in our study comprised members of the classes Alphaproteobacteria, Gammaproteobacteria and Actinobacteria (Table 1), which represent quantitatively important and ecologically relevant taxonomic groups in aquatic environments (Hoshino et al., 2020; Lambert et al., 2019; Rojas-Jimenez et al., 2021). However, we assume that these model strains represent rather copiotrophic strains, while marine habitats are particularly often dominated by oligotrophic strains (Giovannoni, 2017). This assumption was supported by the observation that the genome sizes of the 11 model strains ranged from 2.3 to 4.6 Mbp (Table S1), which does not reflect the genome size range of typically streamlined oligotroph aquatic prokaryotes. Previous research suggested that streamlined oligotroph marine bacterial strains may feature reduced transcriptional regulation compared to copiotrophic strains (Cottrell & Kirchman, 2016). Instead, post-transcriptional regulation mechanisms (e.g., mediated via riboswitches) seemed to be particularly common in the highly abundant and oligotroph SAR11 clade (Kazanov et al., 2007). On the other hand, at least the regulation of SAR11 ribosomal proteins has been shown to be under transcriptional control (Ottesen et al., 2013). Consequently, there is no particular reason to assume that the stress-related regulation of ribosomal

proteins in SAR11 should not follow the patterns that we detected in this study.

While we used changing NaCl concentrations to induce osmotic stress, organisms in natural habitats will be exposed to a large variety of different stressors. However, by definition all stressors interact with the fitness of organisms. It therefore seems plausible that the detected patterns concerning the fitness-related genes screened in this study are not specific for a certain stressor, but could be more universally linked to the stress tolerance (i.e., tolerance-related NB) of bacterial organisms. Also, the employment of compatible solutes has been shown to protect cells not only against osmotic stress but also against temperature changes, droughts or oxidative stress (Singh et al., 2015).

## 5 | CONCLUSIONS

Overall, our MLMs against NB analyses supported that general patterns of bacterial transcriptional regulation can discriminate between generalist and specialist lifestyles. Varying correlation strengths of gene regulation levels against NB and stress in the respective categories implied a close covariation of fitness-related traits, but also of HSP genes with fitness levels on the y-axis of the fitness curves (Figure 1). In contrast, gene regulation levels of osmoprotectant transporters were significantly related to NB, but not to stress. Rather, the regulation of osmoprotectant transporters covaried accordingly with the changing environmental conditions displayed on the x-axis of the fitness curves than with fitness (Figure 1). In all cases, taking into account the physiological functioning of the genes in their respective categories, these observations represent meaningful and interpretable responses.

We further propose a list of candidate stress marker genes whose regulation correlated with the stress exposure levels in our study. We suggest that these genes may be tested in future studies to validate their universal applicability to detect stress levels, either in individual populations or in communities that were exposed to changing conditions. The stress exposure of species in a community



has been considered as one of three main environmental axes defining trait distribution in a community (Grime, 1977; Malik et al., 2020) and is therefore a key parameter for community functioning and assembly processes (Romero et al., 2020). The suggested gene-based approach for stress monitoring in microbial communities may accordingly be incorporated into models to predict carbon fluxes, as suggested elsewhere (Malik et al., 2020), and complement an earlier proposed taxon-based approach to define potential bioindicators for stress (Rocca et al., 2019).

The application of transcriptome analyses is an appropriate tool to characterize the expression of traits in microorganisms, while this method may be less useful to assess the functional properties of larger organisms. Still, general ecological rules, such as species-area relationships (Horner-Devine et al., 2004), have often been shown to be valid across all domains of life. We argue that our findings, concerning the negative or positive correlation of the plasticity of either fitness- or adaption-related traits with an organism's NB should be a more general ecological pattern and its application to macroorganisms that have different response timescales remains to be investigated.

#### ACKNOWLEDGEMENTS

The study was supported by a grant from the German Science Foundation (DFG) awarded to S.B. (BE 5937/2-1). A.R.F. was funded by a scholarship from the Chilean National Agency for Research and Development (ANID)/Scholarship Program/DOCTORADO BECAS Chile/2017 - 72180448. The bioinformatic work was supported by the BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) (031A532B, 031A533A, 031A533B, 031A534A, 031A535A, 031A537A, 031A537B, 031A537C, 031A537D, 031A538A). We thank the Bio2mar platform for access and technical support in their molecular biology facilities. We are further thankful to the MARBEC department at Montpellier University for providing laboratory space and facilities for performing initial work on the strains and for the technical support of Corinne Bouvier and Patrice Got.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### AUTHOR CONTRIBUTIONS

The experimental setup for the isolation experiment was discussed and designed by S.B., T.B. and N.M. N.M. and C.B. collected and isolated the bacterial strains. A.R.F. and S.R. performed laboratory work and analysed the data with support of N.M. A.R.F. and S.B. wrote the manuscript and all the authors commented on the manuscript.

#### OPEN RESEARCH BADGES



This article has earned an Open Data, for making publicly available the digitally-shareable data necessary to reproduce the

reported results. The data is available at <http://doi.io-warnemuende.de/10.12754/data-2021-0007>.

#### DATA AVAILABILITY STATEMENT

The sequence data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession no. PRJEB43309 (<https://www.ebi.ac.uk/ena/browser/view/PRJEB43309>). R-scripts that were used for the data analyses have been published on GitHub (<https://github.com/sarabeier/Strains.NB>).

#### ORCID

Angel Rain-Franco <https://orcid.org/0000-0002-0389-7154>

Nicolas Mouquet <https://orcid.org/0000-0003-1840-6984>

Thierry Bouvier <https://orcid.org/0000-0002-7151-5680>

Sara Beier <https://orcid.org/0000-0003-3707-4487>

#### REFERENCES

- Alekshun, M. N., & Levy, S. B. (1999). The mar regulon: Multiple resistance to antibiotics and other toxic chemicals. *Trends in Microbiology*, 7(10), 410–413. [https://doi.org/10.1016/S0966-842X\(99\)01589-9](https://doi.org/10.1016/S0966-842X(99)01589-9)
- Andrews, E. S. V., & Arcus, V. L. (2020). PhoH2 proteins couple RNA helicase and RNase activities. *Protein Science*, 29(4), 883–892. <https://doi.org/10.1002/pro.3814>
- Aseev, L. V., Koledinskaya, L. S., & Boni, I. V. (2016). Regulation of ribosomal protein operons rplM-rpsI, rpmB-rpmG, and rplU-rpmA at the transcriptional and translational levels. *Journal of Bacteriology*, 198(18), 2494–2502. <https://doi.org/10.1128/JB.00187-16>
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, 19(5), 455–477. <https://doi.org/10.1089/cmb.2012.0021>
- Bartoń, K. (2020). *MuMIn: Multi-Model Inference [R package version 1.43.17]*. Retrieved from <https://CRAN.R-project.org/package=MuMIn>
- Beier, S., Andersson, A. F., Galand, P. E., Hochart, C., Logue, J. B., McMahon, K., & Bertilsson, S. (2020). The environment drives microbial trait variability in aquatic habitats. *Molecular Ecology*, 29(23), 4605–4617. <https://doi.org/10.1111/mec.15656>
- Beier, S., Gálvez, M. J., Molina, V., Sarthou, G., Queroue, F., Blain, S., & Obernosterer, I. (2015). The transcriptional regulation of the glyoxylate cycle in SAR11 in response to iron fertilization in the Southern Ocean. *Environmental Microbiology Reports*, 7(3), 427–434. <https://doi.org/10.1111/1758-2229.12267>
- Beier, S., Holtermann, P. L., Numberger, D., Schott, T., Umlauf, L., & Jürgens, K. (2019). A metatranscriptomics-based assessment of small-scale mixing of sulfidic and oxic waters on redoxcline prokaryotic communities. *Environmental Microbiology*, 21(2), 584–602. <https://doi.org/10.1111/1462-2920.14499>
- Beier, S., Rivers, A. R., Moran, M. A., & Obernosterer, I. (2015). Phenotypic plasticity in heterotrophic marine microbial communities in continuous cultures. *ISME Journal*, 9(5), 1141–1151. <https://doi.org/10.1038/ismej.2014.206>
- Bell, T., Gessner, M. O., Griffiths, R. I., McLaren, J., Morin, P. J., van der Heijden, M., & van der Putten, W. (2009). Microbial biodiversity and ecosystem functioning under controlled conditions and in the wild. In S. Naeem, D. E. Bunker, A. Hector, M. Loreau, & C. Perrings

- (Eds.), *Biodiversity, Ecosystem Functioning, and Human Wellbeing: An Ecological and Economic Perspective*. Oxford University Press.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1), 289–300. JSTOR. Retrieved from JSTOR.
- Bougouffa, S., Radovanovic, A., Essack, M., & Bajic, V. B. (2014). DEOP: A database on osmoprotectants and associated pathways. *Database*, 2014, bau100. doi: <https://doi.org/10.1093/database/bau100>
- Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2020). GTDB-Tk: A toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics*, 36(6), 1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>
- Colles, A., Liow, L. H., & Prinzing, A. (2009). Are specialists at risk under environmental change? Neoecological, paleoecological and phylogenetic approaches. *Ecology Letters*, 12(8), 849–863. <https://doi.org/10.1111/j.1461-0248.2009.01336.x>
- Cottrell, M. T., & Kirchman, D. L. (2016). Transcriptional control in marine copiotrophic and oligotrophic bacteria with streamlined genomes. *Applied and Environmental Microbiology*, 82(19), 6010–6018. <https://doi.org/10.1128/AEM.01299-16>
- Deole, R., & Hoff, W. D. (2020). A potassium chloride to glycine betaine osmoprotectant switch in the extreme halophile *Halorhodospira halophila*. *Scientific Reports*, 10(1), 3383. <https://doi.org/10.1038/s41598-020-59231-9>
- Devictor, V., Clavel, J., Julliard, R., Lavergne, S., Mouillot, D., Thuiller, W., Venail, P., Villéger, S., & Mouquet, N. (2010). Defining and measuring ecological specialization. *Journal of Applied Ecology*, 47(1), 15–25. <https://doi.org/10.1111/j.1365-2664.2009.01744.x>
- Empadinhas, N., & Viète-Vallejo, O. (2008). Osmoadaptation mechanisms in prokaryotes: Distribution of compatible solutes. *International Microbiology*, 11, 151–161. <https://doi.org/10.2436/20.1501.01.55>
- Evans, T. G., & Hofmann, G. E. (2012). Defining the limits of physiological plasticity: How gene expression can assess and predict the consequences of ocean change. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 367(1596), 1733–1745. <https://doi.org/10.1098/rstb.2012.0019>
- Feike, J., Juergens, K., Hollibaugh, J. T., Krueger, S., Jost, G., & Labrenz, M. (2012). Measuring unbiased metatranscriptomics in suboxic waters of the central Baltic Sea using a new in situ fixation system. *ISME Journal*, 6(2), 461–470. <https://doi.org/10.1038/ismej.2011.94>
- García, F. C., Bestion, E., Warfield, R., & Yvon-Durocher, G. (2018). Changes in temperature alter the relationship between biodiversity and ecosystem functioning. *Proceedings of the National Academy of Sciences of the United States of America*, 115(43), 10989. <https://doi.org/10.1073/pnas.1805518115>
- Gifford, S. M., Becker, J. W., Sosa, O. A., Repeta, D. J., & DeLong, E. F. (2016). Quantitative transcriptomics reveals the growth- and nutrient-dependent response of a streamlined marine methylotroph to methanol and naturally occurring dissolved organic matter. *MBio*, 7(6), e01279-16. <https://doi.org/10.1128/mBio.01279-16>
- Giovannoni, S. J. (2017). SAR11 bacteria: The most abundant plankton in the oceans. *Annual Review of Marine Science*, 9(1), 231–255. <https://doi.org/10.1146/annurev-marine-010814-015934>
- Gravel, D., Bell, T., Barbera, C., Bouvier, T., Pommier, T., Venail, P., & Mouquet, N. (2011). Experimental niche evolution alters the strength of the diversity-productivity relationship. *Nature*, 469(7328), 89–U1601. <https://doi.org/10.1038/nature09592>
- Grime, J. (1977). Evidence for existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *American Naturalist*, 111(982), 1169–1194. <https://doi.org/10.1086/283244>
- Grimm, N. B., Chapin, F. S., Bierwagen, B., Gonzalez, P., Groffman, P. M., Luo, Y., Melton, F., Nadelhoffer, K., Pairis, A., Raymond, P. A., Schimel, J., & Williamson, C. E. (2013). The impacts of climate change on ecosystem structure and function. *Frontiers in Ecology and the Environment*, 11(9), 474–482. <https://doi.org/10.1890/120282>
- Häuser, R., Pech, M., Kijek, J., Yamamoto, H., Titz, B., Naeve, F., Tovchigrechko, A., Yamamoto, K., Szafarski, W., Takeuchi, N., Stellberger, T., Diefenbacher, M. E., Nierhaus, K. H., & Uetz, P. (2012). RsfA (YbeB) proteins are conserved ribosomal silencing factors. *Plos Genetics*, 8(7), e1002815. <https://doi.org/10.1371/journal.pgen.1002815>
- Hebbali, A. (2020). *olsrr: Tools for Building OLS Regression Models [R package version 0.5.3]*. Retrieved from <https://CRAN.R-project.org/package=olsrr>
- Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D. M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., & Wardle, D. A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75(1), 3–35. <https://doi.org/10.1890/04-0922>
- Horner-Devine, M. C., Lage, M., Hughes, J. B., & Bohannon, B. J. (2004). A taxa-area relationship for bacteria. *Nature*, 432(7018), 750–753.
- Hoshino, T., Doi, H., Uramoto, G.-I., Wörmer, L., Adhikari, R. R., Xiao, N., Morono, Y., D'Hondt, S., Hinrichs, K.-U., & Inagaki, F. (2020). Global diversity of microbial communities in marine sediment. *Proceedings of the National Academy of Sciences of the United States of America*, 117(44), 27587–27597. <https://doi.org/10.1073/pnas.1919139117>
- Huey, R. B., & Slatkin, M. (1976). Cost and benefits of lizard thermoregulation. *The Quarterly Review of Biology*, 51(3), 363–384. <https://doi.org/10.1086/409470>
- Hutchinson, G. E. (1957). Concluding remarks. *Cold Spring Harbor Symposia on Quantitative Biology*, 22, 415–427. doi:<https://doi.org/10.1101/SQB.1957.022.01.039>
- Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11(1), 119. <https://doi.org/10.1186/1471-2105-11-119>
- Jasmin, J.-N., & Kassen, R. (2007). On the experimental evolution of specialization and diversity in heterogeneous environments. *Ecology Letters*, 10(4), 272–281. <https://doi.org/10.1111/j.1461-0248.2007.01021.x>
- Jessup, C. M., Kassen, R., Forde, S. E., Kerr, B., Buckling, A., Rainey, P. B., & Bohannon, B. J. M. (2004). Big questions, small worlds: Microbial model systems in ecology. *Trends in Ecology & Evolution*, 19(4), 189–197. <https://doi.org/10.1016/j.tree.2004.01.008>
- Joshi, N., & Fass, J. (2001). *Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33) [Software]*. Available at <https://github.com/najoshi/sickle>
- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T., Kawashima, S., Okuda, S., Tokimatsu, T., & Yamanishi, Y. (2007). KEGG for linking genomes to life and the environment. *Nucleic Acids Research*, 36(Database), D480–D484. <https://doi.org/10.1093/nar/gkm882>
- Kazanov, M. D., Vitreschak, A. G., & Gelfand, M. S. (2007). Abundance and functional diversity of riboswitches in microbial communities. *BMC Genomics*, 8(1), 347. <https://doi.org/10.1186/1471-2164-8-347>
- Kellermann, V., van Heerwaarden, B., Sgrò, C. M., & Hoffmann, A. A. (2009). Fundamental Evolutionary Limits in Ecological Traits Drive *Drosophila* Species Distributions. *Science*, 325(5945), 1244. <https://doi.org/10.1126/science.1175443>
- Kielbasa, S. M., Wan, R., Sato, K., Horton, P., & Frith, M. C. (2011). Adaptive seeds tame genomic sequence comparison. *Genome Research*, 21(3), 487–493. <https://doi.org/10.1101/gr.113985.110>
- Konopka, A., Lindemann, S., & Fredrickson, J. (2015). Dynamics in microbial communities: Unraveling mechanisms to identify principles. *The ISME Journal*, 9(7), 1488–1495. <https://doi.org/10.1038/ismej.2014.251>

- Kopylova, E., Noe, L., & Touzet, H. (2012). SortMeRNA: Fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics*, 28(24), 3211–3217. <https://doi.org/10.1093/bioinformatics/bts611>
- Krause, S., Le Roux, X., Niklaus, P. A., Van Bodegom, P. M., Lennon, J. T., Bertilsson, S., Grossart, H.-P., Philippot, L., & Bodelier, P. L. E. (2014). Trait-based approaches for understanding microbial biodiversity and ecosystem functioning. *Aquatic Microbiology*, 5, 251. <https://doi.org/10.3389/fmicb.2014.00251>
- Kültz, D. (2003). Evolution of the cellular stress proteome: From monophyletic origin to ubiquitous function. *Journal of Experimental Biology*, 206(18), 3119–3124. <https://doi.org/10.1242/jeb.00549>
- Ladau, J., & Eloe-Fadrosh, E. A. (2019). Spatial, temporal, and phylogenetic scales of microbial ecology. *Trends in Microbiology*, 27(8), 662–669. <https://doi.org/10.1016/j.tim.2019.03.003>
- Lambert, S., Tragin, M., Lozano, J.-C., Ghiglione, J.-F., Vaulot, D., Bouget, F.-Y., & Galand, P. E. (2019). Rhythmicity of coastal marine picoeukaryotes, bacteria and archaea despite irregular environmental perturbations. *The ISME Journal*, 13(2), 388–401. <https://doi.org/10.1038/s41396-018-0281-z>
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4), 357–359. <https://doi.org/10.1038/NMETH.1923>
- Lecompte, O., Ripp, R., Thierry, J. C., Moras, D., & Poch, O. (2002). Comparative analysis of ribosomal proteins in complete genomes: An example of reductive evolution at the domain scale. *Nucleic Acids Research*, 30(24), 5382–5390. <https://doi.org/10.1093/nar/gkf693>
- Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7), 923–930. <https://doi.org/10.1093/bioinformatics/btt656>
- Lin, J., Liang, H., Yan, J., & Luo, L. (2017). The molecular mechanism and post-transcriptional regulation characteristic of *Tetragenococcus halophilus* acclimation to osmotic stress revealed by quantitative proteomics. *Journal of Proteomics*, 168, 1–14. <https://doi.org/10.1016/j.jprot.2017.08.014>
- Lindquist, S., & Craig, E. A. (1988). The heat-shock proteins. *Annual Review of Genetics*, 22(1), 631–677. <https://doi.org/10.1146/annurev.ev.ge.22.120188.003215>
- Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 550. <https://doi.org/10.1186/s13059-014-0550-8>
- Lozupone, C. A., & Knight, R. (2007). Global patterns in bacterial diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 104(27), 11436–11440. <https://doi.org/10.1073/pnas.0611525104>
- Lynch, M., & Gabriel, W. (1987). Environmental tolerance. *American Naturalist*, 129(2), 283–303. <https://doi.org/10.1086/284635>
- MacArthur, R. H. (1972). *Geographical Ecology*. Princeton University Press.
- Malik, A. A., Martiny, J. B. H., Brodie, E. L., Martiny, A. C., Treseder, K. K., & Allison, S. D. (2020). Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. *The ISME Journal*, 14(1), 1–9. <https://doi.org/10.1038/s41396-019-0510-0>
- Marie, D., Simon, N., Guillou, L., Partensky, F., & Vaulot, D. (2000). Flow Cytometry Analysis of Marine Picoplankton. In R. A. Diamond, & S. Demaggio (Eds.), *Living Color: Protocols in Flow Cytometry and Cell Sorting* (pp. 421–454). Springer. [https://doi.org/10.1007/978-3-642-57049-0\\_34](https://doi.org/10.1007/978-3-642-57049-0_34)
- Matias, M. G., Combe, M., Barbera, C., & Mouquet, N. (2013). Ecological strategies shape the insurance potential of biodiversity. *Frontiers in Microbiology*, 3, 432. <https://doi.org/10.3389/fmicb.2012.00432>
- Ohtsu, I., Kawano, Y., Suzuki, M., Morigasaki, S., Saiki, K., Yamazaki, S., Nonaka, G., & Takagi, H. (2015). Uptake of L-cystine via an ABC transporter contributes defense of oxidative stress in the L-cystine export-dependent manner in *Escherichia coli*. *PLoS One*, 10(4), e0120619. <https://doi.org/10.1371/journal.pone.0120619>
- Ottesen, E. A., Young, C. R., Eppley, J. M., Ryan, J. P., Chavez, F. P., Scholin, C. A., & DeLong, E. F. (2013). Pattern and synchrony of gene expression among sympatric marine microbial populations. *Proceedings of the National Academy of Sciences of the United States of America*, 110(6), E488–E497. <https://doi.org/10.1073/pnas.1222099110>
- P. Grime, J., & Pierce, S. (2012). *The Evolutionary Strategies that Shape Ecosystems*. Wiley-Blackwell.
- Park, C. B., & Lee, S. B. (2000). Effects of exogenous compatible solutes on growth of the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Journal of Bioscience and Bioengineering*, 89(4), 318–322. [https://doi.org/10.1016/S1389-1723\(00\)88952-5](https://doi.org/10.1016/S1389-1723(00)88952-5)
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team (2020). *nlme: Linear and Nonlinear Mixed Effects Models [R package version 3.1-148]*. Retrieved from <https://CRAN.R-project.org/package=nlme>
- Planton, S., Déqué, M., Chauvin, F., & Terray, L. (2008). Expected impacts of climate change on extreme climate events. *Comptes Rendus Geoscience*, 340(9), 564–574. <https://doi.org/10.1016/j.crte.2008.07.009>
- Prosser, J. I. (2015). Dispersing misconceptions and identifying opportunities for the use of “omics” in soil microbial ecology. *Nature Reviews Microbiology*, 13(7), 439–446. <https://doi.org/10.1038/nrmicro3468>
- Prosser, J. I., & Martiny, J. B. H. (2020). Conceptual challenges in microbial community ecology. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 375(1798), 20190241. <https://doi.org/10.1098/rstb.2019.0241>
- Rao, M. R. R., & Stokes, J. L. (1953). Utilization of ethanol by acetic acid bacteria. *Journal of Bacteriology*, 66(6), 634–638. <https://doi.org/10.1128/jb.66.6.634-638.1953>
- Rocca, J. D., Simonin, M., Blaszczyk, J. R., Ernakovich, J. G., Gibbons, S. M., Midani, F. S., & Washburne, A. D. (2019). The microbiome stress project: Toward a global meta-analysis of environmental stressors and their effects on microbial communities. *Frontiers in Microbiology*, 9, 3272. <https://doi.org/10.3389/fmicb.2018.03272>
- Rojas-Jimenez, K., Araya-Lobo, A., Quesada-Perez, F., Akerman-Sanchez, J., Delgado-Duran, B., Ganzert, L., & Grossart, H.-P. (2021). Variation of bacterial communities along the vertical gradient in Lake Issyk Kul, Kyrgyzstan. *Environmental Microbiology Reports*, 13(3), 337–347. doi: <https://doi.org/10.1111/1758-2229.12935>
- Romero, F., Acuña, V., & Sabater, S. (2020). Multiple stressors determine community structure and estimated function of river biofilm bacteria. *Applied and Environmental Microbiology*, 86(12), e00291-20. <https://doi.org/10.1128/AEM.00291-20>
- Roncarati, D., & Scarlato, V. (2017). Regulation of heat-shock genes in bacteria: From signal sensing to gene expression output. *FEMS Microbiology Reviews*, 41(4), 549–574. <https://doi.org/10.1093/femsre/fux015>
- Satinsky, B. M., Gifford, S. M., Crump, B. C., & Moran, M. A. (2013). Use of internal standards for quantitative metatranscriptome and metagenome analysis. *Methods in Enzymology*, 531, 237–250. <https://doi.org/10.1016/B978-0-12-407863-5.00012-5>
- Seneviratne, S., Nicholls, N., Easterling, D., Goodess, C., Kanae, S., Kossin, J., Zwiers, F. W. (2012). Changes in climate extremes and their impacts on the natural physical environment. In C. B. Field, V. Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G. - K. Plattner, S. K. Allen, M. Tignor & P. M. Midgley (Eds.), *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC) (pp. 109–230). Cambridge University Press. <https://doi.org/10.7916/d8-6nbt-s431>

- Sévin, D. C., Stählin, J. N., Pollak, G. R., Kuehne, A., & Sauer, U. (2016). Global metabolic responses to salt stress in fifteen species. *PLoS One*, 11(2), e0148888. <https://doi.org/10.1371/journal.pone.0148888>
- Sexton, J. P., Montiel, J., Shay, J. E., Stephens, M. R., & Slatyer, R. A. (2017). Evolution of ecological niche breadth. *Annual Review of Ecology, Evolution, and Systematics*, 48(1), 183–206. <https://doi.org/10.1146/annurev-ecolsys-110316-023003>
- Shen, D., Juergens, K., & Beier, S. (2018). Experimental insights into the importance of ecologically dissimilar bacteria to community assembly along a salinity gradient. *Environmental Microbiology*, 20(3), 1170–1184. <https://doi.org/10.1111/1462-2920.14059>
- Singh, M., Kumar, J., Singh, S., Singh, V. P., & Prasad, S. M. (2015). Roles of osmoprotectants in improving salinity and drought tolerance in plants: A review. *Reviews in Environmental Science and Bio/Technology*, 14(3), 407–426. <https://doi.org/10.1007/s11157-015-9372-8>
- Slatyer, R. A., Hirst, M., & Sexton, J. P. (2013). Niche breadth predicts geographical range size: A general ecological pattern. *Ecology Letters*, 16(8), 1104–1114. <https://doi.org/10.1111/ele.12140>
- Sørensen, J. G., Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and ecological role of heat shock proteins. *Ecology Letters*, 6(11), 1025–1037. <https://doi.org/10.1046/j.1461-0248.2003.00528.x>
- Stead, M. B., Agrawal, A., Bowden, K. E., Nasir, R., Mohanty, B. K., Meagher, R. B., & Kushner, S. R. (2012). RNAsnap (TM): A rapid, quantitative and inexpensive, method for isolating total RNA from bacteria. *Nucleic Acids Research*, 40(20), e156. <https://doi.org/10.1093/nar/gks680>
- Sultan, S. E. (2001). Phenotypic plasticity for fitness components in polygonum species of contrasting ecological breadth. *Ecology*, 82(2), 328–343.
- Szekely, A. J., Berga, M., & Langenheder, S. (2013). Mechanisms determining the fate of dispersed bacterial communities in new environments. *ISME Journal*, 7(1), 61–71. <https://doi.org/10.1038/ismej.2012.80>
- Thuiller, W., Lavorel, S., & Araújo, M. B. (2005). Niche properties and geographical extent as predictors of species sensitivity to climate change. *Global Ecology and Biogeography*, 14(4), 347–357. <https://doi.org/10.1111/j.1466-822X.2005.00162.x>
- Tischer, K., Zeder, M., Klug, R., Pernthaler, J., Schattenhofer, M., Harms, H., & Wendeberg, A. (2012). Fluorescence in situ hybridization (CARD-FISH) of microorganisms in hydrocarbon contaminated aquifer sediment samples. *Systematic and Applied Microbiology*, 35(8), 526–532. <https://doi.org/10.1016/j.syapm.2012.01.004>
- Van Buskirk, J. (2002). A comparative test of the adaptive plasticity hypothesis: Relationships between habitat and phenotype in anuran larvae. *The American Naturalist*, 160(1), 87–102. <https://doi.org/10.1086/340599>
- Weidenhaupt, M., Rossi, P., Beck, C., Fischer, H.-M., & Hennecke, H. (1996). Bradyrhizobium japonicum possesses two discrete sets of electron transfer flavoprotein genes: FixA, fixB and etfS, etfL. *Archives of Microbiology*, 165(3), 169–178. <https://doi.org/10.1007/BF01692858>
- Welsh, D. T. (2000). Ecological significance of compatible solute accumulation by micro-organisms: From single cells to global climate. *FEMS Microbiology Reviews*, 24(3), 263–290. <https://doi.org/10.1111/j.1574-6976.2000.tb00542.x>
- Yachi, S., & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis. *Proceedings of the National Academy of Sciences of the United States of America*, 96(4), 1463–1468. <https://doi.org/10.1073/pnas.96.4.1463>

#### SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Rain-Franco, A., Mouquet, N., Gougat-Barbera, C., Bouvier, T., & Beier, S. (2022). Niche breadth affects bacterial transcription patterns along a salinity gradient. *Molecular Ecology*, 31, 1216–1233. <https://doi.org/10.1111/mec.16316>