



HAL
open science

Medicinal Macrofungi as Cosmeceuticals: a Review

Susanna Badalyan, Anush Barkhudaryan, Sylvie Rapior

► To cite this version:

Susanna Badalyan, Anush Barkhudaryan, Sylvie Rapior. Medicinal Macrofungi as Cosmeceuticals: a Review. *International Journal of Medicinal Mushrooms*, 2022, 24 (4), pp.1-13. <10.1615/IntJMedMushrooms.2022043124>. <hal-03633501>

HAL Id: hal-03633501

<https://hal.umontpellier.fr/hal-03633501v1>

Submitted on 7 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Short title: Mushrooms as Cosmeceuticals

Medicinal Macrofungi as Cosmeceuticals: A Review

Susanna M. Badalyan,^{a,*} Anush Barkhudaryan,^b & Sylvie Rapior^c

^aLaboratory of Fungal Biology and Biotechnology, Institute of Pharmacy, Department of Biomedicine, Yerevan State University, Yerevan, Armenia; ^bDepartment of Cardiology, Clinic of General and Invasive Cardiology, University Hospital № 1, Yerevan State Medical University, Yerevan, Armenia; ^cCEFE, CNRS, Univ Montpellier, EPHE, IRD, Laboratory of Botany, Phytochemistry and Mycology, Faculty of Pharmacy, 15 Avenue Charles Flahault, 34093 Montpellier Cedex 5, France

*Address all correspondence to: Susanna M. Badalyan, Laboratory of Fungal Biology and Biotechnology, Institute of Pharmacy, Department of Biomedicine, Yerevan State University, Yerevan, Armenia, E-mail: s.badalyan@ysu.am

ABSTRACT: Macrofungi (mushrooms) have been previously studied for their nutritional value and medicinal properties. However, there is still slow progress in the biotechnological application of macrofungi in cosmetic industry as cosmeceuticals and nutricosmetics. Currently, the cosmetic industry is in a constant search for valuable natural ingredients or extracts with relevant bioactive properties (anti-aging, anti-collagenase, anti-elastase, anti-hyaluronidase, anti-pigmentation, anti-inflammatory, antioxidant and anti-tyrosinase) to design formulations. Edible medicinal mushrooms are unlimited source of nutraceuticals and pharmaceuticals. They can be used as a source of ingredients to develop organic cosmeceuticals, nutraceuticals and nutricosmetics for topical and oral administration.

KEY WORDS: anti-aging, bioactive compounds, cosmeceuticals, enzyme inhibitors, medicinal mushrooms and macrofungi, nutricosmetics, nutricosmetics

ABBREVIATIONS: **ABA**, antibacterial activity; **ABTS**, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid); **AD**, atopic dermatitis ; **AIA**, anti-inflammatory activity ; **AMA**, antimicrobial activity; **AOA**, antioxidant activity; **CWE**, cold-water extract; **EPS**, exopolysaccharide; **FIPs**, fungal immunomodulatory proteins; **FMG**, fucomannogalactan; **HaCaT**, human keratinocytes; **HWE**, hot-water extracts; **MRSA**, methicillin-resistant *Staphylococcus aureus*; **NF-κB**, nuclear factor-κB; **NO**, nitric oxid; **PUFA**, polyunsaturated fatty acids; **ROS**, reactive oxygen species; **SIRT1**, Sirtuin 1; **SPF**, sun protection factor; **TCM**, traditional Chinese medicine; **UV**, ultraviolet

I. INTRODUCTION

The natural process of aging, ultraviolet (UV) radiation and hormonal alterations affect the skin and other body parts, which result in the generation of reactive oxygen species (ROS) causing oxidative stress.¹ Cosmetic skin and hair care formulations based on naturally derived active ingredients, including plants and fungi, are currently preferred by consumers for their antioxidant effects.

Currently, the cosmetic industry is in a constant search for bioactive ingredients to design novel cosmeceuticals possessing lower toxicity.² Cosmeceuticals are products between cosmetics and pharmaceuticals containing bioingredients with anti-aging, anti-inflammatory, antioxidant, and photoprotective effects. They can be extracted from different organisms or obtained biotechnologically (cultivation, fermentation, enzymatic synthesis, etc.).

Recent progress in the biology and biotechnology of macrofungi (commonly referred to as mushrooms), genomics, proteomics and myco-pharmacology has contributed to their usage in medicine and food industries.³ Macrofungi are also considered a valuable source of multi-functional cosmetic ingredients to be exploited as cosmeceuticals.^{4,5,6} They are mainly used in

skin and hair care products, and the trend for their industrial usage is expected to increase over the years.⁷

Wild or cultivable, edible or inedible, medicinal or poisonous macrofungi are incorporated in many cosmetic products as producers of bioactive compounds (phenolics, glucans and other polysaccharides and terpenoids) with anti-aging, antibacterial, antifungal, anti-inflammatory, antioxidant, hypocholesterolemic, hypoglycemic, immune-modulating as well as anti-collagenase, anti-elastase, anti-hyaluronidase, anti-tyrosinase, wound-healing and other activities.^{5,8-11} However, there are numerous undescribed or not yet cultivated mushroom species with a potential cosmetic application.^{6,12,13}

Edible mushrooms are considered well-known untapped source of valuable food and nutraceuticals which can improve human health and quality of life. However, inedible or poisonous species (i.e., *Amanita muscaria*, *Ganoderma lucidum*, *Inonotus obliquus*, *Phellinus igniarius*, *Ramaria formosa*, etc.) represent a source of bioactive secondary metabolites to develop myco-pharmaceuticals (dietary supplements or nutraceuticals) with health-enhancing and therapeutic properties.^{5,9} Macrofungi-derived nutraceuticals and nutraceuticals are obtained from mycelia and/or fruiting bodies and administered *per os* in the form of capsules, tablets, or extracts.^{14,15}

Currently, macrofungi-derived numerous commercial products, such as cosmeceuticals (applied topically, i.e., creams, lotions, and ointments) and nutracosmetics (administered *per os*) are available in the market. Their usage is significantly high due to minimal regulation and safety compared to traditional drugs.^{16,17}

The current review reports the main genera and species of macrofungi presently used, or patented to be used, in formulations of cosmeceuticals and nutracosmetics, such as *Agaricus*

subrufescens, *A. bisporus*, *Choiromyces maeandriiformis*, *Ophiocordyceps sinensis*, *Ganoderma lingzhi*, *G. lucidum*, *Grifola frondosa*, *Hypsizygus ulmarium*, *Inonotus obliquus*, *Lentinula edodes*, *Polyporus* spp., *Phellinus* spp., *Schizophyllum commune*, *Trametes versicolor*, *Tremella fuciformis*, and *Tuber* spp. (Table 1).

We also describe the main ranges of human cosmetic and cosmeceutical products incorporating macrofungi or their extracts as anti-aging, antioxidant, hair and skin care, anti-acne or perfumers. Cosmetic brands include Bliss (Hut.com Ltd., Cheshire, UK), La Roche (F. Hoffmann-La Roche Ltd., Basel, Switzerland), Nu-Derm (Obagi Medical Products Inc., Irvine, CA, USA), SensiClear (Mission Scientific Skincare Inc., Gold River, CA, USA) and others (Table 2).

II. MACROFUNGI-DERIVED BIOACTIVE COMPOUNDS AS POTENTIAL COSMECEUTICALS

Over the past years, growing consumers' demand for natural low toxicity organic products with effective skin and hair repairing effects have caused cosmetic industry to search for alternative ingredients. Macrofungi are considered sources of bioactive compounds increasingly used as cosmeceutical ingredients. Scientific data showed that medicinal, edible and inedible mushrooms are resources of raw extracts, fractions of extracts, bioactive molecules, i.e. fatty acids, phenolics, peptides, polysaccharides, terpenoids, vitamins and volatile organic compounds, some of them being also enzyme-inhibitory compounds; that large spectrum of bioactive natural sources possess significant anti-aging, anti-inflammatory, antioxidant, anti-pigmentation, antimicrobial, anti-wrinkle, moisturizing, and skin-whitening effects, may act as inhibitors of collagenase, elastase, hyaluronidase and be used to regulate tyrosinase activity; consequently, they may be potentially used as natural cosmeceuticals or nutracosmetics.

However, the mechanisms behind the mentioned bioactivities are still unknown.

Antioxidants against harmful free radicals and cellular damage are the most demanded cosmeceuticals since their topical administration is essential for protection of aging skin. Skin hyperpigmentation is caused by several factors that upregulate melanogenesis and controlling this unwanted process is a major challenge in dermatology and cosmetology to develop safe and effective anti-melanogenic products. Macrofungi-derived compounds, such as *p*-coumaric acid, can be used to reduce oxidative stress and inflammation, as anti-aging, and skin whitening agents.¹⁸ Previous studies have shown that *p*-coumaric acid inhibited melanin synthesis in murine melanoma cells, human epidermal melanocytes, showed efficient transdermal delivery and functional efficacy in reducing erythema development and skin UV-induced pigmentation, therefore has a potential to be used as a skin-lightening ingredient.¹⁹

The bioactive peptides with plant and animal origins can have a beneficial effect on multiple physiological pathways in the skin which makes them ideal candidates for development of cosmeceuticals. Significant antioxidant and dermatoprotective effects of these peptides after topical application or *per os* administration were revealed which suggests that they can be used in skin care formulations as functional ingredients. However, the mechanisms underlying the skin-protective effect have not been completely elucidated.²⁰

Phenolic compounds and tocopherols derived from edible mushrooms *Boletus edulis*, *Lentinula edodes* and *Xerocomus badius* are the most responsible for their AOA.²¹ The exopolysaccharide (EPS-1) isolated from fermentation broths of rare, valuable edible medicinal polypore fungus *Abortiporus biennis* also possesses prominent radical-scavenging and macrophage-stimulatory activities *in vitro* and could be explored as a functional ingredient with potential applications in medical and cosmetic industries.²² Sclerotia of medicinal polyporoid

species *Polyporus umbellatus* have been used in traditional Chinese medicine (TCM) to treat oedema due to its diuretic effect. It contained bioactive compounds responsible for AOA and AIA and was used as an ingredient in several biotech products.²³

Polysaccharides, triterpenes, and fungal immunomodulatory proteins (FIPs) found in *Ganoderma* species have been exploited for potential ingredients in the cosmetic industry to develop new cosmeceuticals and nutracosmetics with antioxidant, antibacterial, anti-inflammatory, and melanin-inhibitory effects. However, numerous questions need to be addressed before FIPs can be accepted and used in cosmetic products.²⁴

Tyrosinase is an enzyme which has a key role in the formation of melanin. *Ganoderma lucidum* extract was found to have potent anti-tyrosinase activity and is part of several cosmetic, skin-lightening beauty products in China.^{25,26} *G. lucidum* also contains α/β -D-glucans, alkaloids, triterpenoids (ganoderic acids, ganoderenic acids, ganoderol, ganoderiol, lucidenic acids), sterols (ergosterol), proteins (LZ-8, LZ-9), nucleosides (adenosine, inosine, uridine), and nucleotides (adenine, guanine) with different pharmacological potential.⁹ Various marketed products of *G. lucidum* with nutritional and cosmeceutical value are already available in the market.²⁷

AOA and ABA of secondary metabolites of *Pleurotus* species have also been reported.²⁸ *In vitro* safety of ethanolic extracts derived from *G. lucidum* and *P. ostreatus* incorporated into base cosmetic creams was evaluated. The absence of toxicity for keratinocytes and fibroblasts in a concentration-dependent manner was revealed. Protocatechuic and syringic acids were the only compounds permeating from *G. lucidum* extract. These results have shown the safety of studied extracts as skin care ingredients.²⁹

Extracellular polysaccharides derived from medicinal mushroom *Grifola frondosa* (GF-EPS) were applicable in the development of functional cosmeceuticals. The GF-EPS, together

with mycelial extract of *G. frondosa* (GF-MPS), showed AOA, stimulatory effect on collagen biosynthesis and anti-melanogenesis activities, without significant cytotoxicity. GF-EPS and GF-MPS were shown to be promising cosmetic ingredients.³⁰ Chemical constituents isolated from medicinal bracket fungus *Phellinus igniarius* could also be used as ingredients to develop anti-inflammatory cosmetic products.³¹

Edible medicinal cultivated mushroom *Schizophyllum commune* contains phenolics and β -glucans beneficial to skin health. However, data on the development of *Sch. commune* as a cosmeceutical ingredient is lacking. The optimal water extraction conditions at 30°C for 1 h were suggested for effective antioxidant and anti-tyrosinase properties of β -glucans in *Sch. commune*.³² The β -glucans isolated from ethanol extracts of *Sch. commune* also showed potent AOA and may be used as cosmeceuticals.³³

Edible mushroom *Tremella fuciformis* has attracted increasing attention due to its various bioactivities. The polysaccharides extracted from *T. fuciformis*, mainly consisting of mannose and uronic acid, could efficiently reduce water and collagen losses of the skin, and inhibit the increase of glycosaminoglycans. Moreover, after oral treatment with polysaccharides a histopathological study showed that, UV-induced structural alterations of the skin were alleviated, as well as repairing endogenous collagen breakdown and increasing AOA compared to the irradiated control group (without treatment). Thus, polysaccharides from *T. fuciformis* can be used as a skin-protective cosmetic ingredient.³⁴

Atopic dermatitis (AD) is an inflammatory skin disease usually accompanied by *Staphylococcus aureus* (MRSA) infection due to cutaneous barrier-function damage. Benzenoid aromatic compound, coenzyme Q₀ isolated from *Antrodia cinnamomea*, possesses ABA and AIA. The strongest anti-MRSA activity was achieved by inhibiting DNA polymerase,

topoisomerases, reducing the ribosomal proteins, as well as downregulating the expression of interleukin (IL)-6, chemokine (C-C motif) ligand (CCL)5, and CCL17 in HaCaT cells, respectively. CoQ₀ at 0.5 µg/mL could recover the filaggrin decreased by HaCaT activation to the normal control and is effective in mitigating AD symptoms associated with bacterial infection.³⁵

Yellow strain *Flammulina velutipes* (FVY) has become popular among customers due to its distinct texture different from *F. velutipes* white strain (FVW). The physicochemical properties and AOA of FVY and FVW strains were evaluated. The results showed that FVY-80 possesses the highest AOA based on 2,2-diphenyl-1-picrylhydrazyl (DPPH) (IC₅₀ = 2.22 mg/mL) and 2,2'-azino-bis-3-ethylbenzthiazoline-6-sulphonic acid (ABTS) radical assay (IC₅₀ = 2.04 mg/mL). FVY-80 significantly reduced the ROS content in L929 cell by 55.96% and was suggested as the best source for cosmeceutical ingredient.³⁶

The study of ABA, AOA and sun protection factor (SPF) of ethyl acetate extracts of mycelia of selected edible mushrooms revealed ABA against *Staphylococcus epidermidis*, *Escherichia coli*, *Micrococcus luteus* and *Bacillus megaterium*, high SPF activity (34.02 at 200 µg/mL) and AOA, particularly of cultural broth extracted from *F. velutipes*, *P. florida*, and *P. ostreatus*. The results showed the presence of bioactive ingredients in these fungi with promising application in cosmeceuticals.³⁷

Anti-melanogenetic activity and AIA of hot water extracts (HWE) of culinary-medicinal mushrooms *Agaricus bisporus* (white and brown), *G. lucidum*, *Hypsizygus marmoreus*, *Pleurotus floridanus*, and *P. pulmonarius* were evaluated. Brown *A. bisporus* reduced intracellular melanin content to the largest extent (up to 57.05 ± 3.90 %) without a cytotoxic effect on B16F10 melanoma cells, as well as showed anti-tyrosinase activity to 17.93 ± 2.65 %.

The extract obtained from brown *A. bisporus* has AIA decreasing the levels of NO and TNF- α and can be used to develop skin-whitening and anti-inflammatory products. Other mushroom extracts showed moderate activity.³⁸

The ethanolic extract from *Agaricus blazei* (= *A. brasiliensis*) (ABM) was incorporated into a semi-solid base cosmetic cream (safety tests for nutraceutical applications) and cell viability effects of the extract, and of the final cream formulation, on a keratinocyte cell line (HaCaT) (safety tests for cosmeceutical applications) were studied. The study detected proteins, carbohydrates, and a low-fat content. Overall, 22 fatty acids including 53% of polyunsaturated fatty acids (PUFA) were estimated the most abundant fraction. The cell viabilities of Caco-2 and HT29 cell lines were maintained up to 100 μ g/mL. After incorporation into the base cream, a formulation with a pale-yellow colour and favourable pH was obtained. The viability of HaCaT cells in the presence of the extract and the final cream formulation was maintained in a concentration-dependent manner, which indicates the safety of this extract for use in cosmetic industry. The results suggest that ABM residues can be used as an inexpensive and sustainable source of nutraceutical and cosmeceutical ingredients.²⁵

Bioactive compounds extracted from macrofungi showed that the extract of *Trametes versicolor* suppressed the UV-induced cellular senescence in HaCaT cells possibly through augmenting Sirtuin 1 (SIRT1) protein expression.³⁹ Phenolic compounds (cinnamic acid, *p*-coumaric acid, *p*-hydroxybenzoic acid, pyrogallol) with strong AOA and ABA against *Bacillus subtilis*, *E. coli*, *Listeria monocytogenes*, *Salmonella typhimurium*, *S. aureus* were detected in *Macrocybe lobayensis*. As a natural antioxidant and antibacterial agent, this fungus was suggested to be used in pharmaceutical, cosmetic or food industries.⁴⁰

Macrofungi are also producers of bioactive compounds with anti-aging tyrosinase,

hyaluronidase, collagenase and elastase enzymes inhibitory effects which are incorporated in many cosmetic products (Tables 1 and 2).^{5,8,9,11}

Ascomycetous *Cordyceps* (= *Ophiocordyceps*) species is a unique source of bioactive cordycepin with anti-aging property and represents a potential ingredient for cosmeceutical applications.⁴¹

Evaluation of skin care benefits of ethyl acetate, ethanol and hot water mycelial extracts from valuable medicinal fungus *O. sinensis* exhibited *in vitro* moderate or significant anti-collagenase and photoprotective activity with a SPF up to 25. The ethyl acetate extracts showed the highest anti-tyrosinase and anti-elastase activity at 0.14 – 0.47 mg/mL, comparable to arbutin and epigallocatechin gallate. These results revealed the potential of *O. sinensis* as a source of cosmetic bioingredients for skin-care products.⁴²

The tyrosinase-inhibitory effect, AIA, AOA and ABA of ethanolic extracts obtained from *A. bisporus*, *P. ostreatus*, and *L. edodes* were reported and their cosmeceutical potential has been evaluated. The phenolic acids and ergosterol were suggested as basic ingredients for cosmetic creams. The formulations presented 85 - 100% of these compounds can further be exploited as cosmeceuticals.⁸

AOA of mushroom-derived biomolecules could provide a skin-lightening effect by inhibition of tyrosinase, while anti-collagenase and anti-elastase activities can help to restore skin elasticity. Three different polysaccharide extracts obtained from *G. lucidum* were screened for their antioxidant, anti-collagenase and anti-elastase activities. In this study, smaller quantities of proteins, phenols and flavonoids were detected, as well. None of the analysed extracts showed overt toxicity to HaCaT cells at concentrations up to 2 mg/mL and displayed superior scavenging ability on ABTS and DPPH radicals. The hot water crude polysaccharide (HWCP) extract and

partially purified (HWPP) fraction were found to be effective inhibitors of lipid peroxidation (LPx) with a two-fold higher inhibition of LPx compared with ascorbic acid ($EC_{50} = 1.65 \pm 0.08$ mg/mL), a common supplement in cosmeceutical formulations. HWCP showed the strongest anti-tyrosinase activity and inhibition of skin extracellular matrix degradation enzymes. Thus, *G. lucidum* may be considered as a promising source of natural cosmeceutical ingredients. The ethanolic extracts obtained from *A. bisporus* and *P. ostreatus* can be used as a source of cosmeceutical ingredients since they possess antioxidant, anti-tyrosinase and antimicrobial effects. All extracts may be used in free or encapsulated forms in cosmeceutical formulations. The encapsulated forms will provide a controlled release of the cosmeceutical ingredients resulting bioactivity control.^{43,44}

The chemical content and cosmetic properties, such as inhibition of tyrosinase and hyaluronidase activity, SPF values of mycelial extracts derived from *Ganoderma applanatum*, *Laetiporus sulphureus*, and *T. versicolor* have been reported.⁴⁵ The total amount of phenolic acids in the extracts ranged from 2.69 mg/100 g (*G. applanatum*) to 10.30 mg/100 g (*T. versicolor*) dry weight, the total amount of sterols at 48.40 mg/100 g (*T. versicolor*) to 201.04 mg/100 g (*L. sulphureus*) dry weight, and indoles at 2.90 mg/100 g (*G. applanatum*) to 16.74 mg/100 g (*L. sulphureus*) dry weight. Kojic acid was revealed in the extracts of *L. sulphureus* and *G. applanatum*. *L. sulphureus* extract caused a dose-dependent anti-hyaluronidase activity, while all other extracts inhibited tyrosinase activity. The extract of *G. applanatum* exhibited an SPF value of ~ 9. The screened species may be used as bioingredients in the formulation of cosmeceuticals.⁴⁵

The assessment of HWE and cold-water extract (CWE) from fruiting bodies of cultivated macrofungi *Auricularia polytricha*, *P. ostreatus*, *G. lucidum*, and *Sch. commune* revealed their

huge cosmeceutical potential. Total phenolic, polysaccharide and glucan contents were determined, the AOA was estimated. Anti-hyaluronidase activity was used as an indicator of anti-aging and anti-inflammatory properties, while anti-tyrosinase activity was evaluated to assess the whitening property of these extracts. The total polysaccharide content of *P. ostreatus* extracts was the highest, while extracts from *G. lucidum* contained the lowest level of glucan (10.12 – 10.67%). CWE from *Sch. commune* exhibited 98.15% anti-tyrosinase activity and 94.82% AOA. *G. lucidum* HWE revealed the strongest anti-hyaluronidase activity (72.78%). The cosmeceutical properties of these mushrooms can be mainly attributed to the combination of polysaccharides and phenolics. Overall, CWE of *Sch. commune* and HWE of *G. lucidum* revealed the best outcome and suggested them for further study as potential cosmeceuticals.⁴⁶

An culinary-medicinal mushroom *A. bisporus* is rich in nutrients and minerals, and possesses potential anticancer, antioxidant, anti-obesity, and anti-inflammatory properties. The revealed anti-tyrosinase activity and ergothioneine with skin-whitening properties have shown that this fungus is not only a source of nutraceuticals and pharmaceuticals, but also cosmeceutical products.¹

Many commercially available macrofungi-derived cosmetic products have replaced synthetic compounds that have long-term adverse effects.⁴⁷ However, the use of macrofungi and their extracts obtained from fruiting bodies or mycelia as cosmeceuticals need to undergo several steps of biotechnological cultivation and production (extraction optimization, assessment of efficacy and safety) to be produced as a cosmetic product. Previous reports regarding stability, compatibility, and toxicological studies of macrofungi also need to be considered.⁴⁸

III. CONCLUSIONS

Macrofungi (mushrooms) have been regarded as a traditional source of natural low- and

high-molecular-weight secondary metabolites and used for centuries to prevent and treat many diseases; however, over the past decade, they are being exploited for potential ingredients in the cosmetic industry. Several species and their extracts derived from fruiting bodies and/or mycelia are either used or patented to be used as cosmeceuticals and nutracosmetics for their anti-aging, anti-wrinkle, skin-whitening and moisturizing, wound-healing, anti-collagenase, anti-elastase, anti-hyaluronidase, anti-tyrosinase, anti-hyperpigmentation, anti-inflammatory, and antioxidant effects. Among known biodiversity, only a few taxonomically identified and cultivated species are utilized as cosmeceuticals. Further evaluation of natural resources of macrofungi will promote their exploitation in cosmetic industry. Advances in fungal biology and biotechnology, genomics, proteomics, metabolomics, as well as systems pharmacology and molecular cosmetology, study of the molecular mechanisms of medicinal and cosmetic effects of macrofungi will support the discovery of new species to be used in the formulation of organic cosmetic products and cosmeceuticals.

Currently, different mushroom-derived cosmeceuticals and nutracosmetics with various properties are available in the multi-billion-dollar cosmetic market. Despite the availability of a number of products, the industry is largely unregulated, and consumers should consult a doctor prior to the use of these products. Recent developments in cosmeceutical biotechnology, including nano-systems and nano-delivery technologies, have been introduced to improve the efficacy and safety of usage of organic products in commercial production of anti-aging cosmeceuticals. Further realization of clinical studies, manufacturing of high-quality standardized products, and sustainable production of cosmeceuticals under controlled conditions are warranted.

ACKNOWLEDGMENT

The work was supported by the Science Committee of Republic of Armenia, in the frames of the thematic research project № 21T-1F228.

The authors declare to have no conflict of interest

REFERENCES

1. Usman M, Murtaza G, Ditta A. Nutritional, medicinal, and cosmetic value of bioactive compounds in Button Mushroom (*Agaricus bisporus*): A review. *Appl Sci*. 2021; 11:5943.
2. Kaur IP, Agrawal R. Nanotechnology: A new paradigm in cosmeceuticals. *Rec Pat Drug Deliv Formul*. 2007;1(2):171–82.
3. Niego AG, Rapior S, Thongklang N, Raspé O, Jaidee W, Lumyong S, Hyde KD. Macrofungi as a nutraceutical source: Promising bioactive compounds and market value. *J Fungi*. 2021; 7:397.
4. Wu Y, Choi MH, Li J, Yang H, Shin HJ. Mushroom cosmetics: the present and future. *Cosmetics*. 2016;3(3):22.
5. Badalyan SM, Zambonelli A. Biotechnological exploitation of macrofungi for the production of food, pharmaceuticals and cosmeceuticals. In: Sridhar KR and Deshmukh SK, editors. *Advances in Macrofungi: Diversity, Ecology and Biotechnology*. Boca Raton: CRC Press, 2019, p 199–230.
6. Angelini P, Girometta CE, Venanzoni R, Bertuzzi G. Micoterapici per la medicina estetica sistemica: i funghi contro l'invecchiamento cutaneo. *Natural*. 2021;1:44–53.
7. Pareek A, Jain V, Ratan Y, Sharma S, Jain PK, Dave V. Mushrooming of herbals in new emerging markets of cosmeceuticals. *Int J Adv Res Pharm Boil Sci*. 2012;2(4):473–80.
8. Taofiq O, González-Paramás AM, Martins A, Barreiro MF, Ferreira ICFR. A mushroom extracts and compounds in cosmetics, cosmeceuticals and nutricosmetics. A review. *Ind*

Crops Prod. 2016;15;90:38–48.

9. Badalyan SM, Gharibyan NG, Kocharyan AE. Perspectives in the usage of bioactive substances of medicinal mushrooms in pharmaceutical and cosmetic industries. *Int J Med Mushrooms*. 2007;9(3-4):275.
10. Badalyan SM, Barkhudaryan A, Rapior S. Recent progress in research on the pharmacological potential of mushrooms and prospects for their clinical application. In: Agrawal DC, Dhanasekaran M, editors. *Medicinal mushrooms: Recent Progress in Research and Development*. Singapore: Springer Nature, 2019, p 1–70.
11. Sharifi-Rad J, Butnariu M, Ezzat SM, Adetunji CO, Imran M, Sobhani SR, Tufail T, Hosseinabadi T, Ramírez-Alarcón K, Martorell M, Maroyi A, Martins N. Mushrooms-rich preparations on wound healing: from nutritional to medicinal attributes. *Front Pharmacol*. 2020; 11:567518.
12. Hyde KD, Bahkali AH, Moslem MA. Fungi - an unusual source for cosmetics. *Fungal Divers*. 2010;43(1):1–9.
13. Lindequist U. Macrofungi in pharmacy, medicine, cosmetics and nutrition - An appraisal. In: Sridhar KR, Deshmoukh SK, editors. *Advances in Macrofungi: Pharmaceuticals and Cosmeceuticals*. Boca Raton: CRC Press, 2021, p 1-6.
14. Chang ST, Wasser SP. Current and future research trends in agricultural and biomedical applications of medicinal mushrooms and mushroom products (Review). *Int J Med Mushrooms*. 2018;20(12):1121–33.
15. Wong KH, Cheng PG. Mushroom nutraceuticals in Malaysia. *Int J Complem Alt Med*. 2018;11(2):75–6.
16. Brandt FS, Cazzaniga A, Hann M. Cosmeceuticals: current trends and market analysis.

- Semin Cutan Med Surg. 2011;30(3):141–3.
17. Elkhateeb WA. What medicinal mushroom can do? Chem Res J. 2020;5(1):106–118.
 18. Alvarez E, Villa R, Nieto S, Donaire A, García-Verdugo E, Luis SV, Lozano P. The suitability of lipases for the synthesis of bioactive compounds with cosmeceutical applications. Mini-Reviews Org Chem. 2021;18(4):515–28.
 19. Boo YC. p-Coumaric acid as an active ingredient in cosmetics: A review focusing on its anti-melanogenic effects. Antioxidants. 2019;8(8):275.
 20. Aguilar-Toalá JE, Hernández-Mendoza A, González-Córdova AF, Vallejo-Cordoba B, Liceaga AM. Potential role of natural bioactive peptides for development of cosmeceutical skin products. Peptides. 2019;122:170170.
 21. Heleno SA, Ferreira RC, Antonio AL, Queiroz MJRP, Barros L, Ferreira ICFR. Nutritional value, bioactive compounds and antioxidant properties of three edible mushrooms from Poland. Food Biosci. 2015;11:48–55.
 22. Long Z, Liu H, Li J, Sun J, Xue Y, Hu Z, Su Z, Xu C, Yan JK. Preliminary characterization of exopolysaccharides produced by *Abortiporus biennis* in submerged fermentation. Sains Malays. 2019;48(12):2633–40.
 23. Bandara AR, Rapior S, Bhat DJ, Kakumyan P, Chamyuang S, Xu J, Hyde KD. *Polyporus umbellatus*, an edible-medicinal cultivated mushroom with multiple developed health-care products as food, medicine and cosmetics: a review. Cryptogam Mycol. 2015;36(1):3–42.
 24. Li LD, Mao PW, Shao KD, Bai XH, Zhou XW. Ganoderma proteins and their potential applications in cosmetics. Appl Microbiol Biotechnol. 2019;103(23-24):9239–50.
 25. Jiang L. *Ganoderma lucidum* (Reishi Mushroom): potential application as health supplement and cosmeceutical ingredient. Global J Res Analysis (GJRA). 2015;4:124–5.

26. Chien CC, Tsai ML, Chen CC, Chang SJ, Tseng CH. Effects of tyrosinase activity by the extracts of *Ganoderma lucidum* and related mushrooms. *Mycopathologia*. 2008;166:117–20.
27. Ahmad R, Riaz M, Khan A, Aljamea A, Algheryafi M, Sewaket D, Alqathama A. *Ganoderma lucidum* (reishi) an edible mushroom; a comprehensive and critical review of its nutritional, cosmeceutical, mycochemical, pharmacological, clinical, and toxicological properties. *Phytother Res*. 2021;35(11):6030–6062.
28. Cardoso RVC, Carocho M, Fernandes Â, Pinela J, Stojković D, Soković M, Zied DC, Cobos JDV, González-Paramás AM, Ferreira ICFR, Barros L. Antioxidant and antimicrobial influence on oyster mushrooms (*Pleurotus ostreatus*) from substrate supplementation of calcium silicate. *Sustainability*. 2021;13:5019.
29. Taofiq O, Rodrigues F, Barros L, Barreiro MF, Ferreira ICFR, Oliveira MB. Mushroom ethanolic extracts as cosmeceuticals ingredients: Safety and ex vivo skin permeation studies. *Food Chem Toxicol*. 2019;127: 228–36.
30. Kim SW, Hwang HJ, Lee BC, Yun JW. *Grifola frondosa* polysaccharides as cosmeceuticals, *Food Technol Biotechnol*. 2007;45(3):295–305.
31. Jiang Z, Jin M, Zhou W, Li R, Zhao Y, Jin X, Li G. Anti-inflammatory activity of chemical constituents isolated from the willow bracket medicinal mushroom *Phellinus igniarius* (Agaricomycetes). *Int J Med Mushrooms*. 2018;20(2):119–28.
32. Abd Razak DL, Fadzil NH, Jamaluddin A, Abd Rashid NY, Sani NA, Manan MA. Effects of different extracting conditions on anti-tyrosinase and antioxidant activities of *Schizophyllum commune* fruit bodies. *Biocatal Agric Biotechnol*. 2019; 19:101116.
33. Phuket SR, Sangkaew T, Chanapan P, Techaoei S. Biological activity of β -glucans from edible mushroom, *Schizophyllum commune* in Thailand. *Int J Appl Pharm*. 2019;11(5):110–

2.

34. Wen L, Gao Q, Ma CW, Ge Y, You L, Liu RH, Fu X, Liu D. Effect of polysaccharides from *Tremella fuciformis* on UV-induced photoaging. *J Funct Foods*. 2016;20:400–10.
35. Chou WL, Lee TH, Huang TH, Wang PW, Chen YP, Chen CC, Chang ZY, Fang JY, Yang SC. Coenzyme Q0 from *Antrodia cinnamomea* exhibits drug-resistant bacteria eradication and keratinocyte inflammation mitigation to ameliorate infected atopic dermatitis in mouse. *Front Pharmacol*. 2019;10:ID1445.
36. Hu YN, Sung TJ, Chou CH, Liu KL, Hsieh LP, Hsieh CW. Characterization and antioxidant activities of yellow strain *Flammulina velutipes* (Jinhua Mushroom) polysaccharides and their effects on ROS content in L929 cell. *Antioxidants*. 2019;8(8):298.
37. Saraswat A, Mathur P, Sanyal D. Assessing cosmeceuticals properties of some macrofungi for improved skin care. *Int J Pharm Pharm Sci*. 2020;12(2):15–19.
38. Saad HM, Sim KS, Tan YS. Antimelanogenesis and anti-inflammatory activity of selected culinary-medicinal mushrooms. *Int J Med Mushrooms*. 2018;20(2):141–53.
39. Chong Z, Matsuo H, Kuroda M, Yamashita S, Parajuli GP, Manandhar HK, Shimizu K, Katakura Y. Mushroom extract inhibits ultraviolet B-induced cellular senescence in human keratinocytes. *Cytotechnology*. 2018;70(3):1001–8.
40. Khatua S, Acharya K. Functional ingredients and medicinal prospects of ethanol extract from *Macrocybe lobayensis*. *Pharmacogn J*. 2018;10(6):1154–1158.
41. Kunhorm P, Chaicharoenaudomrung N, Noisa P. Enrichment of cordycepin for cosmeceutical applications: culture systems and strategies. *Appl Microbiol Biotechnol*. 2019;103(4):1681–91.
42. Cheng WY, Wei XQ, Siu KC, Song AX, Wu JY. Cosmetic and skincare benefits of

- cultivated mycelia from the Chinese caterpillar mushroom, *Ophiocordyceps sinensis* (Ascomycetes). *Int J Med Mushrooms*. 2018;20(7):623–36.
43. Taofiq O, Heleno SA, Calhelha RC, Fernandes IP, Alves MJ, Barros L, González-Paramás AM, Ferreira ICFR, Barreiro MF. Mushroom-based cosmeceutical ingredients: Microencapsulation and in vitro release profile. *Ind Crops Prod*. 2018;124:44–52.
44. Kozarski M, Klaus A, Jakovljević D, Todorović N, Wan-Mohtar WAAQI, Nikšić M. *Ganoderma lucidum* as a cosmeceutical: antiradical potential and inhibitory effect on hyperpigmentation and skin extracellular matrix degradation enzymes. *Arch Biol Sci*. 2019;71(2):253–64.
45. Sułkowska-Ziaja K, Grabowska K, Apola A, Kryczyk-Poprawa A, Muszyńska B. Mycelial culture extracts of selected wood-decay mushrooms as a source of skin-protecting factors *Biotechnol Lett*. 2021; 43:1051–61.
46. Abd Razak DL, Jamaluddin A, Abd Rashid NY, Sani NA, Manan MA. Assessment of cosmeceutical potentials of selected mushroom fruit body extracts through evaluation of antioxidant, anti-hyaluronidase and anti-tyrosinase activity. *J*. 2020; 3:329–42.
47. Sujarit K, Suwannarach N, Kumla J, Lomthong T. Mushrooms: Splendid gifts for the cosmetic industry. *Chiang Mai J Sci*. 2021;48(3):699–725.
48. Taofiq O, Barreiro MF, Ferreira ICFR. The role of bioactive compounds and other metabolites from mushrooms against skin disorders - A systematic review assessing their cosmeceutical and nutricosmetic outcomes. *Curr Med Chem*. 2020;27(41):6926–65.
49. Ruthes AC, Carbonero ER, Machado Córdova M, Baggio CH, Lanzi Sasaki G, Gorin PAJ, Soares Santos AR, Iacomini M. Fucomannogalactan and glucan from mushroom *Amanita muscaria*: structure and inflammatory pain inhibition. *Carbohydr Polym*. 2013;98(1):761–9.

50. Kalaras MD, Richie JP, Calcagnotto A, Beelman RB. Mushrooms: A rich source of the antioxidants ergothioneine and glutathione. *Food Chem.* 2017;233:429-33.
51. Vaz JA, Barros L, Martins A, Santos-Buelga C, Vasconcelos MH, Ferreira ICFR. Chemical composition of wild edible mushrooms and antioxidant properties of their water soluble polysaccharidic and ethanolic fractions. *Food Chem.* 2011;126(2):610–16.
52. Kim YH, Ryoo IJ, Choo SJ, Xu GH, Lee S, Seok SJ, Bae K, Yoo ID. Clitocybin D, a novel human neutrophil elastase inhibitor from the culture broth of *Clitocybe aurantiaca*. *J Microbiol Biotechnol.* 2009;19(10):1139–41.
53. Lee JE, Lee IS, Kim KC, Yoo ID, Yang HM. ROS scavenging and anti-wrinkle effects of clitocybin A isolated from the mycelium of the mushroom *Clitocybe aurantiaca*. *J Microbiol Biotechnol.* 2017; 27:933–8.
54. Wu JY, Siu KC, Geng P. Bioactive ingredients and medicinal values of *Grifola frondosa* (Maitake). *Foods.* 2021; 10:95.
55. Ito T, Kato M, Tsuchida H, Harada E, Niwa T, Osawa T. Ergothioneine as an anti-oxidative/anti-inflammatory component in several edible mushrooms. *Food Sci Technol Res.* 2011;17(2):103–10.
56. Liu K, Xiao X, Wang J, Chen CYO, Hu H. Polyphenolic composition and antioxidant, antiproliferative, and antimicrobial activities of mushroom *Inonotus sanghuang*. *LWT - Food Sci Technol.* 2017; 82:154–61.
57. Berg A., Reiber K., Dörfelt H., Walther G., Schlegel B., Gräfe U. Laccaridiones A and B, new protease inhibitors from *Laccaria amethystea*. *J Antibiot Res.* 2000;53(11):1313–6.
58. Chen HP, Zhao ZZ, Li ZH, Huang Y, Zhang SB, Tang Y, Yao JN, Chen L, Isaka M, Feng T, Liu JK. Anti-proliferative and anti-inflammatory lanostane triterpenoids from the Polish

- edible mushroom *Macrolepiota procera*. *J Agric Food Chem*. 2018;66:12, 3146–54.
59. Ishihara A, Ide Y, Bito T, Ube N, Endo N, Sotome K, Maekawa N, Ueno K, Nakagiri A. Novel tyrosinase inhibitors from liquid culture of *Neolentinus lepideus*. *Biosci Biotechnol Biochem*. 2018;82(1):22–30.
60. Souilem F, Fernandes Â, Calhella RC, Barreira JCM, Barros L, Skhiri F, Martins A, Ferreira ICRF. Wild mushrooms and their mycelia as sources of bioactive compounds: Antioxidant, anti-inflammatory and cytotoxic properties. *Food Chem*. 2017;230:40–8.
61. Lee IS, Ryoo IJ, Kwon KY, Ahn JS, Yoo ID. Pleurone, a novel human neutrophil elastase inhibitor from the fruiting bodies of the mushroom *Pleurotus eryngii* var. *ferulae*. *J Antibiot*. 2011;64(8):587–9.
62. Abreu H, Fogagnoli Simas F, Ribeiro Smiderle F, Sovrani V, Dallazen JL, Maria-Ferreira D, Werner MF, Cordeiro LMC, Iacomini M. Gelling functional property, anti-inflammatory and antinociceptive bioactivities of β -D-glucan from the edible mushroom *Pholiota nameko*. *Int J Biol Macromol*. 2019; 122:1128–35.
63. Kim KC, Lee IS, Yoo ID, Ha BJ, Sesquiterpenes from the fruiting bodies of *Ramaria formosa* and their human neutrophil elastase inhibitory activity. *Chem Pharm Bull*. 2015;63(7):554–7.
64. Ma K, Han J, Bao L, Wei T, Liu H. Two sarcoviolins with antioxidative and α -glucosidase inhibitory activity from the edible mushroom *Sarcodon leucopus* collected in Tibet. *J Nat Prod*. 2014;77(4):942–7.
65. Thomas P, Elkhateeb W, Daba G. Industrial applications of truffles and truffle-like fungi. In: Sridhar KR, Deshmukh SK, editors. *Advances in Macrofungi: Industrial Avenues and Prospects*. Boca Raton: CRC Press, 2021, p 82-8.

66. Lima ATM, Santos MN, de Souza LAR, Pinheiro TS, Paiva AAO, Dore CMPG, Costa MSSP, Santos ND, Baseia YG, Araújo RM, Leite EL. Chemical characteristics of a heteropolysaccharide from *Tylophilus ballouii* mushroom and its antioxidant and anti-inflammatory activities. *Carbohydr Polym.* 2016;144:400–9.

TABLE 1: Medicinal macrofungi-derived bioactive compounds and their cosmetic effects

Mushroom species	Bioactive compounds	Cosmetic effects	Refs.
<i>Abortiporus biennis</i> (Bull.) Singer	Exopolysaccharides	AOA	22
<i>Agaricus bisporus</i> (J.E. Lange) Imbach	Caffeic, chlorogenic, cinnamic, ferulic, gallic, <i>p</i> -hydroxybenzoic, <i>p</i> -coumaric and protocatechuic acids, alkaloids, polyphenols	AOA, AIA, ABA, anti-tyrosinase	1,8,3 8,43, 54
<i>Agaricus subrufescens</i> Peck	Polysaccharides, benzenoid aromatic compound	AOA, anti-pigmentation	29
<i>Amanita muscaria</i> (L.) Lam.	β -D-Glucan, FMG	AIA	49
<i>Antrodia cinnamomea</i> T.T. Chang & W.N. Chou.	Benzenoid aromatic compound coenzyme Q ₀	ABA, AIA	35
<i>Auricularia polytricha</i> (Mont.)Sacc.	Phenolics, glucans	Anti-hyaluronidase, anti-tyrosinase, AIA	46
<i>Boletus edulis</i> Bull.	Ergothioneine, glutathione	AOA	50
<i>Calocybe gambosa</i> (Fr.) Donk	Ascorbic acid (vitamin C)	Anti-radiation, stimulating collagen synthesis	51
<i>Clitocybe aurantiaca</i> (Wulfen) Studer-Steinhäuslin [= <i>Hygrophoropsis aurantiaca</i> (Wulfen)	Clitocybin A, D	Anti-elastase, stimulation procollagen synthesis, AOA	52,53

Maire]			
<i>Coprinus comatus</i> (O.F. Muller)	Tocopherols (vitamin E)	Reducing UV damage to skin	51
<i>Ophiocordyceps sinensis</i> (Berkeley) G.H. Sung et al.	Cordycepin	Anti-pigmentation, anti-collagenase, anti-tyrosinase, anti-elastase, photoprotective	41,42
<i>Cyclocybe aegerita</i> (V. Brig.) Vizzini	Ergothioneine, glutathione	AOA	50
<i>Flammulina velutipes</i> (Curtis) Singer	Glycosides	AIA, AOA	4,36, 37
<i>Ganoderma lucidum</i> (Curtis) P. Karst.	Polysaccharides, triterpenes, FIPs	Anti-tyrosinase, melanin inhibitory, AIA	9,25, 26,27 ,29,3 8,44
<i>Grifola frondosa</i> (Dicks.) Gray	Extracellular polysaccharides, β -glucan, gallic and tannic acids	AOA, stimulation of collagen biosynthetic, anti-melanogenesis	30,54
<i>Grifola gargal</i> Singer	Ergothioneine	Suppressed TNF- α -mediated activation of NF- κ B	55
<i>Inonotus sanghuang</i> Sheng H. Wu, T. Hatt. & Y. C. Dai	Chlorogenic acid, icarisid II, isorhamnetin, quercetin, quercitrin, rutin	AOA	56
<i>Laccaria amethystina</i> Cooke	Laccaridiones A, B	Anti-collagenase	57
<i>Laetiporus sulphurous</i> (Bull.) Murrill	Phenolic acids	Anti-hyaluronidase, anti-tyrosinase	45

<i>Lentinula edodes</i> (Berk.) Pegler	Polysaccharides, β -glucan lentinan, eritadenine, phenolics, tocopherols	Anti-tyrosinase, AIA, AOA, ABA	8, 21
<i>Macrocybe lobayensis</i> (R. Heim) Pegler & Lodge	Phenolic compounds (cinnamic acid, p-coumaric acid, p-hydroxybenzoic acid, pyrogallol)	AOA, ABA	40
<i>Macrolepiota procera</i> (Scop.) Singer	Lanostane triterpenoids	Inhibition of NO production	58
<i>Neolentinus lepideus</i> (Fr.) Redhead & Ginns	1,3- Dihydroisobenzofuran- 4,5,7-triol,5-methoxy-1,3- dihydroisobenzofuran-4,7-diol, lepidepyrone	Anti-tyrosinase, anti- hyaluronidase	59
<i>Pleurotus eryngii</i> (DC.) Quél.	Cinnamic, p-coumaric and p- hydroxy-phenylacetic acids, ergosterol	AOA	60
<i>Pleurotus eryngii</i> var. <i>ferulae</i> (Lanzi) Saccardo	Ergosterol, nicotinic acid, pleurone, (24E)-3 β - hydroxycucurbita-5,24-diene- 26-oic acid	Anti-elastase	37,61
<i>Pleurotus ostreatus</i> var. <i>florida</i>	Phenolics, gallic acid	ABA, AOA, anti- pigmentation	37
<i>Pleurotus ostreatus</i> (Jacq.) P. Kumm.	Alkaloids, phenolics, gallic acid	AMA, AOA, AIA, anti- pigmentation, anti- tyrosinase	4,8,2 8,29, 37
<i>Phellinus igniarius</i> (L.) Quél.	Nepetidin, betulic acid	NF- κ B inhibitory	31

<i>Pholiota nameko</i> (T. Itô) S. Ito & S. Imai	β -D-Glucan	AIA	62
<i>Polyporus umbellatus</i> (Pers.) Fr.	Polysaccharides, steroids, ergosteol, terpenoids	AIA, ABA, AOA	23
<i>Ramaria formosa</i> (Pers.) Quél.	Axinysonone A, <i>ent</i> -aristolone, nambinone A, ramarin A, B	Anti-elastase	63
<i>Sarcodon leucopus</i> (Pers.) Maas Geest. & Nannf.	Sarcoviolin β , episarcoviolin β	AOA	64
<i>Schizophyllum commune</i> Fr.	Phenolics, α - and β -glucans	Anti-pigmentation, anti-tyrosinase, AOA	32,33
<i>Suillus bellinii</i> (Inzenga) Kuntze	Cinnamic, <i>p</i> -hydroxyphenylacetic and protocatechuic acids, ergosterol	AOA	60
<i>Trametes versicolor</i> (L.) Lloyd	Protein SIRT1, phenolic acids	Anti-tyrosinase, anti-hyaluronidase, AOA	39,45
<i>Tremella fuciformis</i> Berk.	Mannose, polysaccharides, uronic acid	Reduce the water and collagen losses, inhibit the increase of glycosaminoglycans, skin protection, AOA	34
<i>Tuber</i> spp.	Polysaccharides, volatile compounds	AMA, AIA, wound-healing	65
<i>Tylopilus ballouii</i> (Peck) Singer	FMG	Inhibition of superoxide and hydroxyl radicals	66
<i>Volvariella bombycina</i> (Schaeff.)	Isodeoxyhelicobasidin,	Anti-elastase, AMA	4

Singer	alkaloids	AIA, AOA	
--------	-----------	----------	--

ABA, antibacterial activity; **AIA**: anti-inflammatory activity; **AMA**, antimicrobial activity; **AOA**, antioxidant activity; **FIPs**, fungal immunomodulatory proteins; **FMG**, fucomannogalactan; **NF-κB**, nuclear factor-κB; **SIRT1**, Sirtuin 1; **SPF**, sun protection factor; **UV**, ultraviolet.

TABLE 2: Medicinal macrofungi incorporated in different cosmetic products

Mushroom species	Cosmetic Product	Cosmetic Effect
<i>Agaricus subrufescens</i>	Vitamega Facial Moisturizing Mask	Renews and revitalizes skin
<i>Albatrellus confluens</i> (Alb. & Schw. ex Fr.) Kotl.	REN Clean Skincare Evercalm Ultra Comforting Rescue Mask	Skin feels smooth with even tone
<i>Ophiocordyceps sinensis</i>	Kosé Sekkisei Cream	Moisturizer, suppress melanin production
<i>Ganoderma lucidum</i>	Eminence Organic Skin Care Birch Water Purifying Essence	Anti-aging, provides hydration
	Yves Saint Laurent Temps Majeur Elixir De Nuit	Anti-aging
	Dr. Andrew Weil for Origins™ Mega-Mushroom Skin-Relief Micellar Cleanser	Anti-aging
	Shiseido: Ultimune Power Infusing Concentrate	Anti-aging, anti-inflammatory
	Tela Beauty Organics Encore Dri Styling Cream	Hair protection

<i>Inonotus obliquus</i> (Fr.) Pilát	Supermood - The Perfect Day Cream	Antioxidant, skin moisturizer
	Root Science, Reborn Organic Face Mask	Anti-inflammatory, soothes irritated skin
<i>Lentinula edodes</i>	Griffin+Row Naturally Effective Skincare	Anti-inflammatory, skin brightening
	Aveeno Positively Ageless, Daily Exfoliating Cleanser	Anti-aging, skin lifting
	One Love Organics Vitamin D Moisture Mist	Skin lightener, moisturizer and toner
<i>Pleurotus ostreatus</i>	Hankook Sansim Hyoum Jin Firming Cream	Enhances elasticity, skin tight and vitalize
<i>Polyporus umbellatus</i>	iS Clinical Active Serum	Brightens and revitalizes skin
<i>Schizophyllum commune</i>	Alqvimia Eternal Youth Maximum Recovery Face Cream	Brightens, unifies skin tone
	Alqvimia Eternal Youth Cream Facial Máxima Regeneración	Anti-aging, skin lifting
	Sulwhasoo Hydro-Aid, cream	Hydrates, revitalizes skin and promotes clear
<i>Tremella fuciformis</i>	Eminence Organic Skin Care Birch Water Purifying Essence	Antioxidant, enhances skin elasticity

	Dr. Dennis Gross Skincare Alpha Beta® Pore Perfecting Cleansing Gel	Enhances skin elasticity, skin firming
	La Prairie Advanced Marine Biology Night Solution	Moisturizes, nourishes, revitalizes
<i>Tuber spp.</i>	Black Orchid Tom Ford, Jo Malone Red Truffle 21, Estée Lauder Re-Nutriv skin cream, Perigold truffle magnetic face mask	Perfumes with truffles' aroma, moisturizes, nourishes