

Occurrence of Cyclusphaera scabrata in Achiri (late middle-early late Miocene?, Bolivian Altiplano): Paleogeographical implication

Mercedes B Prámparo, Pierre-Olivier Antoine, Laurent Marivaux, Rubén Andrade Flores, Marcos Fernández-Monescillo, Alberto Boscaini, Bernardino Mamani Quispe, Séverine Fauquette, Colline Bonnet, Philippe Münch, et al.

▶ To cite this version:

Mercedes B Prámparo, Pierre-Olivier Antoine, Laurent Marivaux, Rubén Andrade Flores, Marcos Fernández-Monescillo, et al.. Occurrence of Cyclusphaera scabrata in Achiri (late middle-early late Miocene?, Bolivian Altiplano): Paleogeographical implication. Journal of South American Earth Sciences, 2022, 119, pp.103990. 10.1016/j.jsames.2022.103990. hal-03789913

HAL Id: hal-03789913 https://hal.umontpellier.fr/hal-03789913

Submitted on 28 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Occurrence of *Cyclusphaera scabrata* in Achiri (late middleearly late Miocene?, Bolivian Altiplano): paleogeographical implication

Mercedes B. Prámparo^{1, 2}, Pierre-Olivier Antoine³, Laurent Marivaux³, Rubén Andrade Flores⁴, Marcos Fernández-Monescillo^{2, 5} Alberto Boscaini^{2, 6}, Bernardino Mamani Quispe⁴, Séverine Fauquette³, Colline Bonnet³, Philippe Münch⁷, François Pujos^{1, 2}

- ¹ Instituto Argentino de Nivología Glaciología y Ciencias Ambientales (IANIGLA), CONICET-UNCUYO-Mendoza, Avda.Ruiz Leal s/n, Parque Gral. San Martín, 5500 Mendoza, Argentina. <u>mprampar@mendoza-conicet.gob.ar</u>
- ² Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Godoy Cruz 2290, C1425FQB, Ciudad Autónoma de Buenos Aires, Argentina
- ³ Institut des Sciences de l'Évolution de Montpellier, cc 64, Univ Montpellier, CNRS, 18 IRD, place Eugène Bataillon, F-34095 Montpellier Cedex 05, France
- ⁴ Unidad de Paleontología, Museo Nacional de Historia Natural, Calle 26 s/n, Cota Cota, La Paz, Plurinational State of Bolivia
- ⁵ Cátedra y Museo de Paleontología, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba. Vélez Sarsfield 1611, (X5016GCA) Córdoba, Argentina
- ⁶ Instituto de Ecología, Genética y Evolución de Buenos Aires (IEGEBA CONICET), Departamento de Ecología, Genética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Int. Guiraldes 2160, C1428EGA, Ciudad Autónoma de Buenos Aires, Argentina
- ⁷ Géosciences Montpellier, Université de Montpellier, CNRS, F-34095 Montpellier, France

Abstract

A palynological assemblage including the extinct araucarian dispersed pollen, Cyclusphaera scabrata, was analysed from the Achiri locality, Bolivian Altiplano. The fossil-yielding locality of Achiri is located in the Pacajes Province, La Paz Department, in the northwestern Altiplano, Bolivia. The Achiri locality is very well known by the fossil vertebrate faunas recovered from the Mauri Formation (M6). The palynological samples come from fine-grained and plant-rich carbonated lacustrine levels lower in the same unit. An age between late middle to early late Miocene? was inferred for the palynological association studied based on local stratigraphy and radioisotopic ages obtained higher in the series. The retrieved association is mainly represented by angiosperms; although gymnosperms, bryophytes, ferns, and fresh-water algae are also documented. Compositae are the most abundant group within angiosperms, and Polylepis also occurred but in low percentages. The presence of C. scabrata in the Bolivian Altiplano expands the taxon distribution outside northern South America and provides additional evidence to infer timing and distribution routes of conifers, restricted toward the tropics during the latest Neogene. Its occurrence supports a paleoelevation of no more than 2000 m for the northern area of the Altiplano in agreement with previous results. Compositae of Fenestrites type and Polylepis occur earlier in Bolivia and C. scabrata later, comparing their occurrence in miocene microfloras from low to mid- latitudes of South America.

Keywords: Neogene palynomorphs; araucarian dispersed pollen; northwestern Altiplano; Mauri Formation.

1. Introduction

Nowadays, the Bolivian Altiplano as a whole is characterized by Puna vegetation, but the vegetation likely has considerably shifted in the Miocene and onwards. The precise timing and pattern of this shift is not well-constrained. In addition, Neogene palynological records from the Plurinacional State of Bolivia are pretty scarce. Graham et al. (2001) described a Miocene-Pliocene palynoflora originally presented by Berry (1922a) as the Pislepampa flora, Eastern Cordillera, Bolivia, with the following recognized microfossils: Isoetes, Lycopodium, Cnemidaria, Cyathea, Grammitis, Hymenophyllum, Pteris, trilete fern spores, Danaea, monolete fern spores, Podocarpus, Gramineae, Arecaceae, Ilex, cf. Oreopanax, Cavanillesia, cf. Pereskia, Compositae (three types), Ericaceae, and Tetrorchidium. Based on sedimentological, plant microfossil and macrofossil evidence, Graham et al. (2001) recovered evidence of a lacustrine environment bordered by elements of the Amazonian rainforest, with a cloud forest growing on the surrounding slopes. Gregory-Wodzicki (2002) analysed a new collection of the late Miocene Jakokkota flora (the upper Jakokkota flora) from a lacustrine unit, located above the lower Jakokkota flora, from a fluvial unit, first documented by Berry (1922b) and further described by Gregory-Wodzicki et al. (1998). These fossil leaf impressions, located in the northern Altiplano of Bolivia, in the Member 6 of the Mauri Formation, correspond to 24 fossil-species, mainly represented by Anacardiaceae, Berberis, Myrtaceae, Zizyphus, Polylepis, and several leguminosae. Based on leaf morphology for the combined flora (both lower and upper Jakokkota floras), the mean annual temperature was inferred to be 21.5±2.0°C and the mean annual precipitation of 550±180 mm. The most accurate age based into radioisotopic method (Sanidine from ash falls) of these floras is around 10.66 ± 0.06 Ma (Gregory-Wodzicki et al.2002). The inferred elevation at that time (Late Miocene) was no more than 1600-1200 m, whereas today is almost 4000 m above sea level (ASL; Gregory-Wodzicki, 2002). Thus, based on the study of these floras, Gregory-Wodzicki (2002) suggested that a large portion of the Central Andes was covered by subtropical-dry forest with nearly half of the modern Altiplano elevation reached since the early late Miocene interval.

In this contribution, we document for the first time a palynological assemblage including the extinct araucarian dispersed pollen, *Cyclusphaera scabrata*, recovered from the northern Altiplano (Achiri locality: ACH-26 locus Cerro Pisakeri) and we discuss its correlation with palynological zonations from different sections of northern South America. The data presented here partially bridges the gap for Neogene floras from Bolivia, providing useful information to reveal the plant distribution during this period in the Central Andes region. All this, especially taking into account the difficulty of finding favorable lithologies for palynological studies in the Neogene of the Altiplano region.

1.1. Geological and stratigraphical contexts

The fossil-yielding locality of Achiri is located in the Pacajes Province, La Paz Department, in the northwestern Altiplano, Bolivia (Fig. 1). The Altiplano is an extensive intramontane basin formed during the Cenozoic era, starting with the Eastern Cordillera uplifting (Suárez-Soruco, 2000). The Altiplano plateau of western South America lies within the Central Andes above the subducting Nazca plate. The extensive sedimentary archive accumulated in this area is of special interest as it reflects the Altiplano Neogene paleoclimate and surface uplift history (Garzione et al., 2006; 2014). In the western Altiplano, the late Oligocene-Miocene Mauri Fm. unconformable overlies the Berenguela Fm. (Evernden et al., 1977). The Mauri Fm. comprises six members (M1 to M6) which include volcanoclastic rocks interbedded with volcanic flows (Sirvas and Torres, 1966). This unit was deposited in a wide fluvial and lacustrine plain in a backarc basin. The Achiri locality is well known by the fossil vertebrate faunas recovered from the Mauri Fm, M6 (e.g., Villarroel 1974; Villarroel and Marshall, 1983; Marshall et al., 1983; Hoffstetter, 1986; Saint-André, 1993, 1996; Pujos et al., 2012; Fernández-Monescillo et al., 2019; Gaudin et al., 2022). The fossiliferous levels of the Achiri locality were dated between 10.5 Ma (K-Ar, at Hako-kota; Evernden et al., 1977) and 8 Ma (K-Ar, west of San Andrés, Evernden et al., 1977). In the same area, Fernández-Monescillo et al. (2019) described a partial pathological skeleton of the notoungulate mesotheriid *Plesiotypotherium achirense* from the Cerro Pisakeri (Fig. 1) in the locus ACH-27 (Fig. 2), immediately bracketed by two 15mdistant ash layers for which they performed step-heating 40Ar/39Ar experiments on feldspar micro-populations and obtained plateau ages of 10.42 ± 0.09 Ma (ACH-TUF3, below) and 9.42 ± 0.1 Ma (ACH-TUF4, above), respectively. Accordingly, they confirmed the late Miocene age (Tortonian or late Mayoan-early Chasicoan from South American Land Mammal Ages [SALMAs] for the corresponding locus (ACH-27) and surrounding assemblages, and provided a sedimentation rate approximating 15 m per million years in the concerned time interval. The presence of several successive volcanoclastic beds, occurring throughout the sequence cropping out on the Cerro Pisakeri, suggests an almost continuous volcanic activity during the late Miocene in this area.

Extensive field explorations conducted by our team during the last decade also resulted in the discovery of older fossil-bearing levels, at the bottom of the available stratigraphic section (Fig. 2). These fine-grained and plant-rich carbonated levels (ACH-25 and ACH-26) testify to the presence of lacustrine settings in the area, in which the palynological samples were collected (ACH-25, ACH- 26; Fig. 2). Interestingly, ACH-26 has yielded amphibian, bird, rodent and plant remains (Douillez et al., 2018). A late middle to early late Miocene? age can be inferred for ACH-26 and thus for ACH-25 when considering a roughly constant and low sedimentation rate throughout the whole Cerro Pisakeri section (Fig. 2).

1.2. Achiri vertebrate faunas

The late Miocene mammalian assemblage of Achiri is, together with that of Quebrada Honda (Croft, 2007), one of the most abundant and diversified faunal association during the Neogene in the Plurinacional State of Bolivia. This fauna includes several notoungulates such as the mesotheriid Plesiotypotherium achirense, the small hegetotheriid cf. Hemihegetotherium, and the toxodontid Hoffstetterius imperator, a macraucheniid litoptern, the sparassodontan metatherian Borhyaenidium altiplanicum, and a wide array of xenarthrans with the glyptodont Trachycalyptoides achirense, several Dasypodidae, a pampatheriid, and ground sloths. The latter group is documented by a megatheriine, an unidentified mylodontid, the scelidotheriine cf. Proscelidodon sp. and the small megatherioid "Xyophorus" villarroeli (e.g., Marshall et al., 1983; Saint-André, 1996; Pujos et al., 2012). Recently, new discoveries have permitted to: i) extend our knowledge of Plesiotypotherium achirense (Fernández-Monescillo et al., 2018, 2019), ii) suggest that the extinct sloth "Xyophorus" spp. (including X. villarroeli) could correspond to an early-diverging nothrothere for which the new genus Mcdonaldocnus is proposed (Pujos et al., 2021; Gaudin et al., 2022), and iii) identify several rodents including the cavioids Prodolichotis prisca and Cardiomys sp., the chinchilloids Tetrastylus sp. and Lagostomus sp., and an octodontoid of uncertain affinities (Pérez et al., 2021).

2. Material and methods

Five samples were collected and processed for palynology:ACH-25, ACH-26 (PN1,2,3), ACH 14. Only two yielded palynomorphs. The two productive samples come from ACH-25 (S17°11.261', W68°00.153') and ACH-26 (S17°14.459', W69°00.255') in a dry canyon at the NW base of the Cerro Pisakeri. ACH-26 corresponds to lacustrine grey siltstone with abundant plant detritus and leaf fragments. The physical and chemical extractions of palynomorphs were performed in the Paleopalynology Laboratory at the Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA) located in Mendoza, Argentina. The samples were treated with hydrochloric and hydrofluoric acids following the palynological processing techniques according to Volkheimer and Melendi (1976). No oxidation was needed. The residues were sieved with a 10 µm filter and the final slides were mounted in glycerin jelly. The specimens were examined using an Olympus BX 50 microscope. Location of the specimens on the slides, and in figure references is given by ESF (England slide finder) coordinates. The slides are housed at the Laboratory of Palynology of the IANIGLA under the catalogue numbers: 10566 (ACH-25) and 10567 (ACH-26) MPLP (Mendoza-Paleopalinoteca- Laboratorio de Paleopalinología). The palynostratigraphic analysis was based mainly on the zonation presented by Hoorn (1993) and Jaramillo et al. (2011) for the Neogene of Amazonia and Northern South America palynology.

3. Results

3.1. Palynology

The palynological association from both levels ACH-25 and ACH-26 comprises pollen grains, spores, and fresh-water algae. However, the palynomorphs recovered from the lower level (ACH 25) were very scarce, thus preventing statistical counts. This sample yielded only some psilate and verrucate trilete spores, a few angiosperms of compositae type (type 1 and 2, see below to which correspond each) and the tricolporate *Psilatricolporites crassoexinatus*.

The material recovered at the upper level (ACH-26) was quite abundant and well preserved instead, allowing to make a detailed taxonomic and relative abundance analysis based on 202 specimens counted (Fig. 3). Angiosperms are the most abundant group (90%) in the association corresponding to ACH-26, but gymnosperms, bryophytes and ferns are also documented. The fern spores retrieved included mainly laevigate monolete and trilete spores (*Laevigatosporites* sp. and *Deltoidospora* sp.; Figs. 4 A, B) and aquatic ferns (*Azolla* and *Salvinia*). Liverwort spores (Marchantiales) of *Riccia* type, similar to *Reboulisporites fuegiensis* Zamaloa and Romero (Fig. 4 C), are recorded.

Gymnosperms are represented by pollen grains with coniferous affinities: *Cyclusphaera scabrata* (1%) and *Podocarpidites* sp. (0,5%). *Cyclusphaera* is a taxon with a typical Gondwanan distribution during the Mesozoic era. It corresponds to a biaperturate pollen grain with affinity to the Araucariaceae, based on early Cretaceous cones found with pollen *in situ* in Patagonia, Argentina (Del Fueyo and Archangelsky, 2005). The general distribution of the genus was reduced during the Cenozoic, and its geographical range shifted mostly toward the tropics. *Cyclusphaera* was a common element of tropical floras in South America and Africa during this time, and it became restricted to western South America by mid-Neogene times (Jaramillo et al., 2013).

Cyclusphaera scabrata was erected by Jaramillo and Dilcher (2001) from the Paleocene–early Eocene Interval of Colombia. Subsequently, Jaramillo et al. (2013) expanded the original description of the species adding new data of the exine structure obtained by transmission electron and scanning electron microscope. The Bolivian specimens (eight measured from ACH-26) have circular to elliptical amb (55 to 72 μ m) with two symmetrical circular apertures with regular edges (always regular) (20 to 37 μ m) occupying nearly one-third of the total grain diameter (Figs. 4 E-H). One of these specimens (Fig. 4 F) is corroded showing the granulae at the innermost portion of the sexine. The area between the aperture and equatorial margin of the Bolivian specimens is 17 to 20 μ m. The aperture membrane is not present in the studied specimens. The equatorial exine thickening, of 2 μ m, of *C. scabrata* given by Jaramillo and Dilcher (2001) is observed in none of the Bolivian specimens. The thickness of the exine (1 μ m) is equal between the equatorial zone and the aperture. The specimens from Achiri locality (ACH-26) are slightly larger than those from the Paleogene of Colombia, in agreement with the size increase-upward trend, proposed by Jaramillo et al. (2013) in

their stratigraphic section. These authors gave a complete discussion of the worldwide spatial and temporal distributions of this taxon. The distribution of *Cyclusphaera scabrata* was even more restricted during the late Miocene–early Pliocene, recording both a drastic increase in aridity in the tropics and the expansion of C4 grasses and savannas (Zachos et al., 2001; Edwards et al., 2010).

In the Neotropics, Parra et al. (2020) described the palynology from a Neogene sequence in the Marañon Basin, north of Peru and mentioned the presence of *C. scabrata* in Zone Mar-B, corresponding to the Burdigalian stage (17.71–16.1 Ma). Sá and Carvalho (2017) and D'Apolito et al. (2021) recorded the same fossil-taxon in Miocene deposits of the Solimões Fm. in the Solimões Basin (western Brazilian Amazonia), as well as Hoorn et al. (2017) in the middle–late Miocene interval of Well 2 of Foz do Amazonas Basin (offshore Pará, Brazil). The latter authors recognize *C. scabrata* as a component of the tropical lower montane forest. The youngest record of *Cyclusphaera* (only one specimen) was reported from the Choco region in westernmost Colombia, at *ca*. 1.5°N, dated at 5.5 Ma (Jaramillo et al., 2013).

Podocarpaceae (represented in the association by *Podocarpidites* sp. Fig. 4D) is a southern hemisphere conifer family, which comprises evergreen, coniferous trees, and shrubs. Today, it is still a conspicuous component in angiosperm-dominated tropical forests. Tropical podocarps are most abundant in mid- to high elevation forests but also occasionally at low elevation (Cernuzak et al., 2010). The pattern of dispersal or extinction of this family during the Cenozoic was related to the availability of areas of wet climate and equable mesothermal temperatures (Morley, 2011). Thus, its presence in the palynoflora is considered as indicative of upland vegetation (mountain slopes, Andean Forest). Jaramillo et al. (2013) consider its distribution a better analogue to the pattern seen in *Cyclusphaera*.

Compositae are the most abundant group within angiosperms in the Achiri association (77.2% in ACH-26). They are represented by four morphological types similar to Asteraceae: i) prolate tricolporoidate grains (19 to 26 μ m, polar axe), echinae short (2μ m) (Figs. 4, I-J) Compositae type 1 similar to *Senecio*); ii) roughly similar to C. type 1 but differs in the size of the echinae which are more than 5 μ m long (Fig. 4, L Compositae type 2 similar to *Baccharis*); iii) tricolporate, echinate, fenestrate larger than 30 μ m (Fig. 4, K Compositae type 3 similar to *Taraxacum*) and iv) small fenestrate pollen grains (20–24 μ m) *Gomphrena* type and *Vernonia* type (similar to the morphotype *Fenestrites*; Figs. 4, M, N). Compositae first appear in South America in the late Oligocene–early Miocene interval and do not become frequent or diverse until the middle to late Miocene (Graham et al., 2001). Barreda et al. (2010) based on the broad distribution patterns of basal Asteraceae and allied families during the Oligocene in the Southern Hemisphere, suggested an Eocene origin of the clade. A significant diversification of Asteraceae occurred during the Early Miocene (Barreda et al., 2010).

Tricolporate pollen grains corresponding to *Polylepis/Acaena* group occurred (Fig. 4 O, P). They have an endoaperture with a zone of $35-42 \mu m$ (diameter); the sexine (ectexine) covers the endopore, forming a chamber with prominent flaps (pouchlike

extensions); and the exine is 1.5 µm thick microverrucate. The genus Polylepis consists of shrubs or trees native to the mid- and high-elevation tropical Andes (Simpson, 1979). Polylepis appears to be the highest natural occurring arborescent angiosperm genus in the world, since some members of the genus are components of the upper montane forest, but others occur in woodlands at elevations as high as 5200 m (Simpson, 1979). High pollen values of Polylepis/Acaena and Compositae, are considered to be diagnostic of a sub-puna shrubland (Hansen et al. 1984). Today Polylepis is commonly found in Bolivia in the Yungas (Paramo), southern Puna and northern Puna (Ortuño et al. 2011). In the late Miocene of the Bolivian Altiplano, Gregory-Wodzicki (1998, 2001) recorded *Polylepis* as one of the most abundant forms among fossil dicot leaves from Jakokkota flora. Martínez et al. (2020) recorded the presence of Polylepis from El Descanso Fm (Central Andean Plateau, Peru) in both studied Members (B and C). Going to east, Hoorn et al. (2017) recorded the first occurrence of *Polylepis-Acaena* in much younger strata from a well in the Foz do Amazonas Basin (5.8 Ma; Brazil). The presence of this pollen type, despite its low abundance (ca. 2%), points to the onset of subpuna vegetation (following Hansen et al. 1984) since the late Middle to early late Miocene in Bolivia.

Psilatricolporites crassoexinatus Hoorn (1993), an angiosperm of unknown affinity, is also present (Fig. 4 S). It was previously recorded in the Miocene–Pliocene Solimões Fm. (Hoorn, 1993; Nogueira et al., 2013) and the Miocene–Pliocene Novo Remanso Fm. from central Amazonia (Guimaraes et al., 2015). In addition, Polycolporate endocingulate pollen grains occur (Fig. 4 Q), corresponding to Polygalaceae (similar to *Xanthophyllum*). Polygalaceae have a cosmopolitan distribution and include trees, shrubs, lianas, herbs, and parasites (Banks et al., 2008). The pollen assemblage also includes tricolpate reticulate pollen grains (probably of Brassicaceae affinity, Fig. 4R) and a reticulate tricolporate of unknown affinity (Fig. 4T).

Freshwater algae referable to *Botryococcus* are retrieved (6%), further testifying to a lacustrine environment.

Finally, abundant organic matter mainly conformed by biostructured phytoclasts, cuticles, and abundant woody remains (bigger than 100 μ m in length), complete the palynological association of both levels (ACH-25 and ACH-26), giving account of local vegetation which was much more diversified at that time in terms of types.

3.2. Palynostratigraphy

We correlate our data from Achiri to the palynological zonation from the Llanos Basin in Colombia (Jaramillo et al., 2011) and coeval floras from low to mid- latitudes of South America. The palynostratigraphic scheme from Colombia has been applied also to correlate outcrops and cores in the Miocene of Peru, southern Colombia, and Brazil (Boonstra et al., 2015; Hoorn et al., 2017; Jaramillo et al., 2017, Parra et al., 2020). The Achiri association can be correlated with a part of the *Fenestrites spinosus* Zone T16 (12.7 to 7.1 Ma), from the Cenozoic of Los Llanos, Colombia (Jaramillo et al., 2011)

due to the occurrence of this taxon in both associations. Nevertheless, *Cyclusphaera* scabrata had thus far its last appearance datum (LAD) in this scheme in Zone T-13 of Los Llanos, Colombia, i.e. in the late early Miocene (17.7 to 16.1 Ma). However, Jaramillo et al. (2013), based on unpublished data from western Colombia, mentioned a youngest record of this taxon as young as 5 Ma-old. In Achiri, *C. scabrata* occurs 58 meters below the tuff ACH-TUF-3 (Fig. 2) dated as 10.42 ± 0.09 Ma (Fernández-Monescillo et al., 2019). The sedimentary dynamics seems to be fairly steady and constant over the complete sequence measured at Cerro Pisakeri (Fig. 2), with fluviolacustrine and floodplain deposits, accumulated with a mean sedimentation rate of 15 m per million years at the top (between ACH-TUF-3 and ACH-TUF4). This in turn allows for hypothesizing a probable age between late middle to early late Miocene? for the palynological association studied.

The presence of *C. scabrata* in the Neogene of Bolivia expands the known distribution of the taxon outside northern South America and corresponds to the southernmost record of the specie during the Miocene (a complete distribution of the taxon is available in Jaramillo et al., 2013).

Martínez et al. (2020) studied the plant fossil record from the Descanso-Yauri Basin, Southern Peru including palynological (pollen and spores) and macrofossil samples from two lithostratigraphic units (Members B and C) of more than 88 localities. The members were previously dated using radiometric methods: Member B, dated to ca. 18.7 to 9.1 Ma (early to late Miocene), and Member C, dated to ca. 4.8 to 3.9 Ma (early Pliocene). The palynoflora of Member B is composed of common montane forest indicators such as Podocarpus, Hedyosmum, Bocconia, Mutisia, and Cyatheaceae, as well as higher elevation taxa such as Calamagrostis, Polylepis, and Valeriana. Part of Member B correlates to the Achiri palynoflora, having in common Podocarpaceae and Polvlepis. Parra et al. (2020) carried out a detailed palynostratigraphic research of the Marañon Basin (western Amazonia, Peru) based on the identification of qualitative changes in palynomorphs through the intervals sampled in six exploration wells. They recognized six palynozones spanning the early Miocene- early Pliocene time interval (Mar- A-F). Accordingly, the Achiri palynoflora may be coeval to Mar-C (latest Burdigalian to late Langhian) and part of Mar-D (late Serravallian) from Marañon Basin. However, there are no common species in both associations. Tcheumeleu et al. (2020), recorded palynomorphs, fossil plants, microfossils (charophyte oogonia and ostracod fragments), mollusks, and vertebrates from the Middle Ipururo Fm, at the the Shumanza Section, Subandean Zone of Northern Peru. The Peruvian association palynostratigraphic markers Grimsdalea magnaclavata, includes as Palaeosantalaceaepites cingulatus, Echitricolporites spinosus, and *Fenestrites* longispinosus, pointing to a late Miocene-early Pliocene age for the TAR-27 locality (10.06–3.72 Ma). They concluded the persistence of a steady lowland rainforest (with absence of characteristic high-altitude taxa such as Podocarpus, Hedyosmum, Jamesonia, or Polylepis) close to the Andes between 10 and 8 Ma. The marker taxa recovered in Peru were not recognized in the Achiri palynoflora with the exception of Compositae of *Fenestrites* type, present in both palynofloras but with different species.

The Asteraceae zone which corresponds to the middle Tortonian–late Messinian period is very uncommon in the Brazilian Amazon palynofloras and there are only three records of the Asteraceae zone coming from three boreholes of the southwestern part of the Amazonas state (Leandro et al., 2019). In summary, we can conclude that an association of *Compositae-Polylepis-Cyclusphaera* characterizes the Achiri palynoflora which can be dated from the late middle to early late Miocene?. In this sense, Compositae of *Fenestrites* type and *Polylepis* occur earlier in Bolivia and *Cyclusphaera scabrata* later, comparing their occurrence in other miocene microfloras of the region.

4. Conclusions

The presence of Cyclusphaera scabrata at the bottom of the Achiri section (late middle to early late Miocene?) greatly increases the southernmost distribution of this taxon during the Neogene, farther to the South. This data also provides a useful additional contribution to infer the timing and distribution routes of conifers towards a more restricted location in the tropics during the latest Neogene. The occurrence of Cyclusphaera, of Araucariaceae affinities, together with Podocarpaceae pollen confirms the effective presence of conifers as components of the upland vegetation (Andean Forest) in the northwestern Bolivian Altiplano, during the late middle to early late Miocene. We assume for the extinct Cyclusphaera taxon similar environmental conditions of closely-related extant species of Araucaria. In this sense, Araucaria angustifolia grows today at a similar latitude (20°S) at high elevations between 1500 and 1800 m near Rio de Janeiro, Brazil (Ledru and Stevenson, 2012) and in adjacent areas of eastern Paraguay and northeastern Argentina. This taxon generally occurs at 800 m to 1800 m in the north and at 500 m to 1200 m near its southern limits (Veblen et al. 1995). Thus, considering that conifers need high levels of rainfall in temperate regions, and that they barely tolerate to drought, we can suggest an elevation of no more than 2000 m for the northern area of the Altiplano in agreement with the results of Gregory-Wodzicki (2002) and Boschman (2021). Besides, the occurrence of abundant Compositae pollen associated with incipient presence of Polylepis in the Achiri microflora are indicative of the sub-puna shrubland vegetation. Botryococcus and aquatic fern spores (Azolla and Salvinia) embody the freshwater communities related to the lacustrine environment. We hope that these first results will trigger new field efforts. in order to better bridge the existing gaps in the Neogene palynological record of the Bolivian Altiplano.

Acknowledgements

Thanks to the editor, to Carlos Jaramillo and two anonymous reviewers for providing constructive comments that improved the manuscript. Fieldwork in the Bolivian Altiplano has been carried out in the framework of an ongoing cooperation agreement (N°864/2014) between the MNHN-Bol (La Paz, Bolivia), the *Institut des Sciences de l'Evolution de Montpellier* (France) and the CCT-CONICET (Mendoza, Argentina). M. Prámparo thanks CONICET (PIP 11220200101738CO) for supporting her research.

This research was also supported by an "*Investissements d'Avenir*" grant managed by the French *Agence Nationale de la Recherche* (ANR) in the framework of the LabEx CEBA (ANR-10-LABX-25-01). This is a contribution of the Paltiplano project that has received financial support from the CNRS through the MITI interdisciplinary programs.

References

- Banks, H., Klitgaard, B.B., Claxton, F., Forest, F., Crane, P., 2008. Pollen morphology of the family Polygalaceae (Fabales). Bot. J. Linnean Soc., 156, 253–289.
- Barreda, V.d., Palazzesi, L., Telleria, M.C., Katinas, L., Crisci, J.V., 2010. Fossil pollen indicates an explosive radiation of basal Asteracean lineages and allied families during Oligocene and Miocene times in the Southern Hemisphere. Rev.Palaeobot.Palynol., 160, 102-110. DOI:10.1016/j.revpalbo.2010.02.004
- Berry, E.W., 1922a. Pliocene fossil plants from eastern Bolivia. John Hopkins Univ. Stud. Geol. 4, 145–202.
- Berry, E.W., 1922b. Late Tertiary plants from Jancocota, Bolivia. Johns Hopkins Univ. Stud. Geol. 4, 205–221.
- Boonstra, M., Ramos, M.I.F., Lammertsma, E.I., Antoine, P.-O., Hoorn, C., 2015. Marine connections of Amazonia: evidence from foraminifera and dinoflagellate cysts (early to middle Miocene, Colombia/Peru). Palaeogeogr. Palaeoclimatol. Palaeoecol, 417, 176–194.
- Boschman, L.M., 2021. Andean mountain building since the Late Cretaceous: a paleoelevation reconstruction. Earth-Sci. Reviews, 220, 103640.
- Cernusak L, H., Adie, P., Bellingham, E., Biffin, T., Brodribb, P., Coomes, J., Dalling, et al., 2011. Podocarpaceae in tropical forests: a synthesis. In: L Cernusak, B Turner, (Eds), Ecology of the Podocarpaceae in tropical forests. Smithsonian Institution, Washington, DC, 189–196.
- Croft, D.A., 2007. The middle Miocene (Laventan) Quebrada Honda Fauna, Southern Bolivia and a description of its Notoungulates. Palaeontology, 50, 277–303.
- D'Apolito, C., Jaramillo, C., Harrington, G., 2021. Miocene palynology of the Solimões Formation (Well 1-AS-105-AM), Western Brazilian Amazonia. Smithsonian Contributions to Paleobiology, 105, 1–134.
- Del Fueyo, G., Archangelsky, S., 2005. A new araucarian pollen cone with in situ *Cyclusphaera* Elsik from the Aptian of Patagonia, Argentina. Cretac. Res., 26, 757–768.
- Douillez, S., Münch, P., Merzeraud, G., Fernández-Monescillo, M., Andrade Flores, R., Antoine, P.-O., Boscaini, A., Mamani Quispe, B., Marivaux, L., Prámparo, M.B., Pujos, F., 2018. A multidisciplinary study of late Miocene terrestrial deposits from the Corque basin, central Altiplano, Bolivia. International Palaeontological Congress 5, 8-13 juillet 2018, Paris.
- Edwards, E, C., Osborne, C., Stromberg, S., Smith, W., Bond, P., Christin, A., Cousins, et al., 2010. The origins of C4 grasslands: integrating evolutionary and ecosystem science. Science, 328, 587–591.

- Evernden, J.F., Stanislav, J.K., Cherroni, M., 1977. Potassium-Argon ages of some Bolivian rocks. Economic Geology, 72, 1042–1061.
- Fernández-Monescillo, M. Bernardino Mamani Quispe, François Pujos, Pierre-Olivier Antoine, 2018. Functional anatomy of the Forelimb of *Plesiotypotherium* achirense (Mammalia, Notoungulata, Mesotheriidae) and Evolutionary Insights at the Family Level. J. Mamal. Evol., 24, 197-211
- Fernández-Monescillo, M., Antoine, P.-O., Mamani Quispe, B., Münch, P., Andrade Flores, R., Marivaux, L., Pujos, F., 2019. Multiple skeletal and dental pathologies in a late Miocene mesotheriid (Mammalia, Notoungulata) from the Altiplano of Bolivia: palaeoecological inferences. Palaeogeogr, Palaeoclimatol, Palaeoecol, 534, 109297.
- Garzione, C.N., Molnar, P., Libarkin, J.C., MacFadden, B.J., 2006. Rapid late Miocene rise of the Bolivian Altiplano, Evidence for removal of mantle lithosphere. Earth Planet. Sci. Lett. 241, 543–556. http://dx.doi.org/10.1126/science.1119365
- Garzione, C.N., Auerbach, D.J., Jin-Sook Smith, J., Rosario, J.J., Passey, B.H., Jordan, T., Eiler, J.M., 2014. Clumped isotope evidence for diachronous surface cooling of the Altiplano and pulsed surface uplift of the Central Andes. Earth Planet. Sci. Lett., 393, 173–181. http:// dx.doi.org/10.1016/j.epsl. 2014.02.029
- Gaudin,T.J., Boscaini, A., Mamani Quispe, B., Andrade Flores, R., Fernández-Monescillo, M., Marivaux, L., Antoine, P.O., Münch, P., Pujos, F., 2022.
 Recognition of a new nothrotheriid genus (Mammalia, Folivora) from the early late Miocene of Achiri (Bolivia) and the taxonomic status of the genus Xyophorus, Hist. Biol, DOI: 10.1080/08912963.2022.2075744
- Graham, A., Gregory-Wodzicki, K.M., Wright, K.L., 2001. Mio-Pliocene palynoflora from the Eastern Cordillera, Bolivia: implications for the uplift history of the Central Andean Plateau. Amer. J. Bot., 88, 1545–1557.
- Gregory-Wodzicki, K. M., 2002. A late Miocene subtropical-dry flora from the northern Altiplano, Bolivia. Palaeogeogr. Palaeoclimatol. Palaeoecol., 180, 331–348.
- Gregory-Wodzicki, K.M., McIntosh, W.C., Velasquez, K., 1998. Climatic and tectonic implications of the late Miocene Jakokkota flora, Bolivian Altiplano. J. S. Am. Earth Sci., 11, 533–560.
- Guimaraes J. T., Rodrigues Nogueira, A.C., Bandeira Cavalcante Da Silva Junior, J., Lima Soares, J., Alves, R., Kern, A.K., 2015. Palynology of the Middle Miocene-Pliocene Novo Remanso Formation, Central Amazonia, Brazil. Ameghiniana, 52, 107–134.
- Hansen, B.C.S., Wright, H.E., Bradbury, J.P., 1984. Pollen studies in the Junín area, central Peruvian Andes. Geol. Soc. Am. Bull., 95, 1454–1465.
- Hoffstetter, R., 1986. High andean mammalian faunas during the Plio-Pleistocene. In: High altitude tropical Biogeography. Vuilleumier, F., Monasterio, M., (Eds) Oxford University Press, Oxford, 219–245.
- Hoorn, C., 1993. Marine incursions and the influence of Andean tectonics on the Miocene depositional history of northwestern Amazonia: results of a palynostratigraphic study. Palaeogeogr. Palaeoclimatol. Palaeoecol., 105, 267– 309.

- Hoorn, C., Bogotá, G.R., Romero-Baez, M., Lammertsma, E.I. Flantua, S.G.A., Dantas, E.L., Dino, R., do Carmo, D.A., Chemale Jr., F., 2017. The Amazon at sea: onset and stages of the Amazon River from a marine record, with special reference to Neogene plant turnover in the drainage basin. Global and Planetary Change, 153, 51–65.
- Jaramillo, C., Dilcher, D., 2001. Middle Paleogene palynology of central Colombia, South America: a study of pollen and spores from tropical latitudes. Palaeontographica Abt B 258, 87–213.
- Jaramillo, C., Rueda, M., Torres, V., 2011. A palynological zonation for the Cenozoic of the Llanos Foothills of Colombia. Palynology, 35, 46–84.
- Jaramillo, C., Zavada, M., Ortiz, J., Pardo, A., Ochoa, D., 2013. The Biogeography of the Araucarian Dispersed Pollen *Cyclusphaera*. Int. J. Plant Sci., 174, 489–498
- Jaramillo C, Romero I, D'Apolito C, Bayona G, Duarte E, Louwye S, Escobar J, Luque J, Carrillo-Briceño JD, Zapata V, Mora A. 2017. Miocene flooding events of western Amazonia. Sci. Advances 3, e1601693.
- Leandro, L.M., Vieira, C.E.L., Santos, A., Fauth, G., 2019. Palynostratigraphy of two Neogene boreholes from the Northwestern portion of the Solimões Basin, Brazil. J. S. Am. Earth Sci., 89, 211–218.
- Ledru, M.-P., Stevenson, J., 2012. The rise and fall of the genus *Araucaria*: a Southern Hemisphere climatic connection. In: Haberle, S., David, B. (Eds), Peopled Landscapes: Archaeological and Biogeographic Approaches to Landscapes. Terra Australis, 34, 241–253.
- Marshall, L.G., Hoffstetter, R., Pascual, R., 1983. Mammals and stratigraphy: geochronology of the continental mammal-bearing Tertiary of South America. Palaeovertebrata, 1–93.
- Martínez, C., Jaramillo, C., Correa-Metrío, A., Crepet, W., Moreno, J.E., Aliaga, A. Moreno, F., Ibañez-Mejia, M., Bush, M.B., 2020. Neogene precipitation, vegetation, and elevation history of the Central Andean Plateau. Sci. Advances, 6, 1–9.
- Morley, J., 2011. Dispersal and paleoecology of tropical podocarps. In: Cernusak, L.A., Turner, B.L. (eds.), Ecology of the Podocarpaceae in Tropical Forests. Smithsonian Institution, Washington, DC., 21–42.
- Nogueira, A.C.R., Silveira, R., Guimarães, J.T.F., 2013. Neogene-Quaternary sedimentary and paleovegetation history of the eastern Solimões Basin, Central Amazon region. J. S. Am. Earth Sci., 46, 89–99.
- Ortuño, T., Ledru, M.-P., Cheddadi, R., Kuentz, A., Favier, C., Beck, S., 2011. Modern pollen-rain, vegetation and climate in Bolivian ecoregions. Rev.Palaeobot.Palynol., 165, 61–74. doi:10.1016/j.revpalbo.2011.02.004
- Parra, F.J., Navarrete, R.E., di Pasquo, M.M., Roddaz, M., Calderon, Y., Baby, P., 2020. Neogene palynostratigraphic zonation of the Maranon Basin, Western Amazonia, Perú. Palynology, 44, 675–695.
- Pérez, M.E., Candela, A.M., Mamani Quispe, B., Abello, A., Adnet, S., Andrade Flores,
 R., Billet, G., Boscaini, A., Fernández-Monescillo, M., Marivaux, L., Münch, P.,
 Prámparo, M.B., Antoine, P.-O., Pujos, F., 2021. Los roedores caviomorfos

(Rodentia, Hystricognathi) del Mioceno tardío de Achiri (Bolivia). 34° JAPV, p.75–76.

- Pujos, F., Antoine, P.-O., Mamani Quispe, B., Abello, A., Andrade Flores, R., 2012. The Miocene vertebrate faunas of Achiri, Bolivia. J. Vertebr. Paleontol., 32 (Supp. to 3), p. 159.
- Pujos, F., Gaudin, T.J., Boscaini, A., Abello, M.A., Andrade Flores, R., Fernández-Monescillo, M., Mamani Quispe, B., Marivaux, L., Prámparo, M.B., Antoine, P.-O., Münch, P., 2021. The megatherioid sloth "*Xyophorus*" from the Late Miocene of Achiri (Bolivia). *Libro de Resúmenes* – 14° CAPA, p.197–198.
- Sá, N.P., Carvalho, M.A., 2017. Miocene fern spores and pollen grains from the Solimões Basin, Amazon region, Brazil. Acta Bot. Bras., 31, 720–735. http://10.1590/010 2-33062017abb0160.
- Saint-André, P.-A., 1993. *Hoffstetterius imperator* n. g., n. sp. du Miocène supérieur de l'Altiplano bolivien et le statut des Dinotoxodontinés (Mammalia, Notoungulata).
 C. R. Acad. Sci. Paris, 316 (Série 2), 539–545.
- P-A., 1996. Deux nouveaux Édentés (Mammalia, Saint-André. Xenarthra) Trachycalyptoides achirense nov. gen. (Glyptodontidae, et nov. sp. Sclerocalyptinae) et *Xvophorus* villarroeli nov. (Megatheriidae, sp. Nothrotheriinae) du Huayquérien (Miocène supérieur) de l'Altiplano bolivien. Bulletin du Muséum National d'Histoire Naturelle, 4, 79-106.
- Simpson, B.B., 1979. A revision of the genus *Polylepis* (Rosaceae: Sanguisorbeae). Smithsonian Contributions to Botany, 43, 1–62
- Sirvas-Carranza, F., Torres, E., 1966. Consideraciones geológicas de la zona noroeste de la Provincia Pacajes del depto. La Paz. Boletín del Instituto Boliviano del Petróleo, 66, 54–64.
- Suárez-Soruco, R., 2000. Subdivisión tectono-estratigráfica de las rocas del Ciclo Cordillerano de Bolivia. Memorias del 14° Congreso Geológico de Bolivia, La Paz, Bolivia, 142–145.
- Tcheumeleu, A. F., Fauquette, S., Aliaga Castillo, A., Martínez, C., Moreno, F., Navarrete, R., Parra, F., Wesselingh, F., Salas-Gismondi, R., Varas-Malca, R., Roddaz, M., Antoine, P-O., 2020. Depositional environments and landscapes of the upper Miocene Ipururo Formation at Shumanza, Subandean Zone, northern Perú. Palaeobiodiversity and Palaeoenvironments, 10.1007/s12549-019-00400-8. hal-02355089v2
- Veblen, T.T., Burns, B. R., Kitzberger, T., Lara, A., Villalba, R., 1995. The ecology of the conifers of southern South America. In: Ecology of Southern Conifers. Enright, N.J., Hill, R.S. (Eds). Melbourne University Press, Melbourne, pp. 120 –155.
- Villarroel, C., 1974. Un mésothériiné nouveau (Notongulata, Mammalia) dans le Miocène supérieur de Bolivie. C.R. Acad. Sci. Paris, 279(D), 551–554.
- Villarroel, C., Marshall, L.G., 1983. Two New Late Tertiary Marsupials (Hathlyacyninae and Sparassocyninae) from the Bolivian Altiplano. J. Paleontol., 57, 1061–1066.

- Volkheimer, W., Melendi, D., 1976. Palinomorfos como fósiles guía. Tercera parte: Técnicas de Laboratorio palinológico. Rev. Min. Geol. Mineral, 34, 19–30.
- Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. Science, 292, 686–693
- Zamaloa, M.del C., Romero, E.J., 1990. Some spores and pollen from the Cullen Formation (upper Eocene to middle Oligocene), Tierra del Fuego, Argentina, Palynology, 14:1, 123–133. DOI: 10.1080/01916122.1990.9989376

Figure captions

- Figure 1. Location map of the studied area around Achiri, in the La Paz Department, Pacajes Province, Bolivia. General map of the Plurinacional State of Bolivia showing Achiri locality (present contribution); Jakokkota flora (Berry 1922b; Gregory-Wodzicki et al., 1998; 2002) and Pislepampa flora (Berry 1922a; Graham et al., 2001) (right side). Detailed map showing location of the sampled points for palynology and ash levels (left side), in the north of Cerro Pisakeri.
- Figure 2. (A) Lithologic column of Member 6 of Mauri Fm. at Cerro Pisakeri, Achiri locality, north-western Altiplano, Bolivia. Location of the fossil and volcaniclastic levels. (B)(C) photographs showing levels sampled for palynology; ACH-25 and ACH-26 locations respectively (and details of each of them). Abbreviation: ACH-TUF, volcanic ash (tuff) samples; PN, palynological samples. ACH-14 unproductive palynological sample; ACH-11 vertebrate remains, ACH-27 locus of the partial pathological skeleton of *Plesiotypotherium achirense* described by Fernández-Monescillo et al. (2019).
- Figure 3. (A) Bar diagram showing relative percentages of different groups of palynomorphs in the ACH-26; (B) relative percentage of the different groups among angiosperms from ACH-26.
- **Figure 4.** Palynomorphs from Achiri locality. (A) *Deltoidospora* sp. 10567B: J 29/3 (25 μm); (B) *Laevigatosporites* sp. 10567C: E32/0 (36 μm); (C) *Reboulisporites fuegiensis* Zamaloa and Romero 1990, 10567A: T30/2 (45 μm); (D) *Podocarpidites* sp. 10567B: D42/2 (55 μm); (E-H) *Cyclusphaera scabrata* Jaramillo and Dilcher 2001, (E) 10567B: V24/2 (72 μm), (F) 10567A: W29/2 (55 μm), (G) 10567B: T26/0 (70 μm each specimen); (H) 10567B: V27/4 (60 μm); (I) Compositae type 1,10567C: J42/1 (22 μm); (J) Compositae type 1, 10567A: W44/0 (22 μm); (K) Compositae type 3, 10567C: O22/0 (37 μm); (L) Compositae type 2,10567C: F20/1 (23 μm); (M-N) *Fenestrites* sp. 10567B: M32/3 (20μm); (O-P) *Polylepis* sp. (O) 10567B: U38/1 (39 μm), (P) 10567B: U43/0 (42 μm); (Q) Polygalaceae 10567C: T20/3 (37 μm); (R) Tricolpate reticulate indet. 10567C: S37/4 (42 μm) (S) *Psilatricolporites crassoexinatus* Hoorn 1993, 10567 A*: E32/3 (35 μm) (T) reticulate tricolporate 10567A: M40/1 (22 μm). Scale bars= 10μm.









PALYNOMORPHS%	ACH-26
Bryophytes/Pteridophytes	2.5
Gymnosperms	1.5
Angiosperms	90
Fresh-water algae	6

100%

ANGIOSPERMS %	ACH-26
Compositae type 1-2	61.5
Compositae type 3	22.6
Fenestrites type	1.6
Psilatricolporites crassoexinatus	9.9
Polylepis/Acaena	2.2
Schinus type	2.2
	100%