



HAL
open science

Mesophotic reefs are not refugia for neither taxonomic nor functional diversity of reef fishes

Nicolas Loiseau, Sébastien Villéger, Charles Le Bozec, Marie Gimenez, Seiji
Léo Kawahara, Thomas Claverie

► **To cite this version:**

Nicolas Loiseau, Sébastien Villéger, Charles Le Bozec, Marie Gimenez, Seiji Léo Kawahara, et al..
Mesophotic reefs are not refugia for neither taxonomic nor functional diversity of reef fishes. Coral
Reefs, 2022, 10.1007/s00338-022-02311-1 . hal-03860627

HAL Id: hal-03860627

<https://hal.umontpellier.fr/hal-03860627v1>

Submitted on 31 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Coral Reefs**

2

3 **Mesophotic reefs are not refugia for neither taxonomic nor functional**
4 **diversity of reef fishes**

5

6 *Authors* : Nicolas Loiseau¹, Sébastien Villéger¹, Charles Le Bozec², Marie Gimenez², Seiji Léo
7 Kawahara², Thomas Claverie^{1,2}

8 *Institutions*:

9 1 MARBEC, University of Montpellier, CNRS, IFREMER, IRD, Montpellier, France

10 2 CUFR of Mayotte, Dembeni, France

11

12 *Mailing addresses*

13 Nicolas Loiseau: *nicolas.loiseau@cnrs.fr*

14 UMR MARBEC MONTPELLIER

15 UNIVERSITE DE MONTPELLIER - FACULTE DES SCIENCES

16 Place Eugène Bataillon - bât 24 - CC093

17 34 095 MONTPELLIER Cedex 5

18

19 Sébastien Villéger: *sebastier.villeger@cnrs.fr*

20 Charles Le Bozec: *charles.lebozec1@gmail.com*

21 Marie Gimenez: *mariegimenez34@yahoo.fr*

22 Seiji Léo Kawahara: *leo.kawahara@gmail.com*

23 Thomas Claverie: *tclaverie@gmail.com*

24

25 *Keywords*: Coral reef, functional traits, beta diversity, video surveys, Western Indian Ocean

26

27 *Data availability*: All code and data will be provided on <https://github.com/LoiseauN/Bubot>

28

29 On behalf of all authors, the corresponding author states that there is no conflict of interest.

30 **ABSTRACT**

31 Mesophotic reefs, defined as reefs between 30 m - 150m, have been less investigated than
32 their shallower counterparts. Yet, more protected from human activities and bleaching events, they
33 could be refuge zones allowing the resilience of shallow reefs. However, it is currently unknown
34 if their role as refugia for species diversity also extends to functional diversity and thus potentially
35 ecosystem functioning. Using an extensive video field surveys from Mayotte Island in the Western
36 Indian Ocean, we analyzed how taxonomic diversity and functional diversity of coral reef fishes
37 changed along a depth gradient from 10 to 100 m. We found that shallow and mesophotic reefs
38 host different species, but that these dissimilar assemblages share similar richness combinations
39 of traits values. However, when accounting for biomass of species, it appears that functional
40 structure of assemblages differs along the depth gradient, with, for instance, herbivorous fishes
41 being very scarce in deep reefs while large-bodied carnivorous are more abundant. Therefore,
42 surveys of mesophotic fishes should assess not only species, but also trait composition, abundance
43 and biomass. Overall, our results support the claim that extending marine protected areas to
44 mesophotic reefs is required to maximize the protection of both species and functional diversities
45 to eventually maintain coral reef ecosystem functioning and associated services across the broader
46 seascape.

47
48 *Keywords: Coral reef, functional traits, beta diversity, video surveys, Western Indian Ocean*

49
50 **INTRODUCTION**

51 Coral reefs cover close to 300,000 km², which is just under 0.1% of the oceans' surface area
52 (Spalding et al. 2001). Despite this restricted cover, coral reefs are one of the world's most diverse
53 and productive ecosystems, supporting prodigious biodiversity and providing important ecosystem
54 services to many millions of people. In fact, over 4,000 species of teleost fishes are associated with
55 coral reefs (Sale 2013). Fishes are central to reef ecosystems, providing essential ecological
56 functions such as biomass production, regulation of foodwebs, nutrient cycling and sand formation
57 among others (Villéger et al. 2017; Brandl et al. 2019). Fishes also provide food security to
58 hundreds of millions of people (Sala et al. 2021). However, coral reef fishes are severely affected
59 by overexploitation, habitat loss and climate change, especially through marine heatwaves (Eddy
60 et al. 2021). In response to this “coral reef crisis,” studies around the world have examined the
61 social, institutional, and environmental conditions that influence fish biodiversity and ecosystem
62 functions (Cinner et al. 2016, 2020). However, most of these studies focused on shallow (< 30m)
63 reef communities (Rocha et al. 2018). As a result, the distribution, abundance and function of fish
64 assemblages that occupy the deeper range of reefs are less known compared to shallow reefs. In
65 other words, variation of the large diversity of biological traits (e.g. size, diet, mobility, behavior;
66 (Villéger et al. 2017), and their participation in core processes (Brandl et al. 2019) along depth
67 gradients remains largely unknown (see however Cure et al. 2021 and Carrington et al. 2021). Yet,
68 unraveling the distribution of fishes and of their traits across depth is critical for effective
69 ecosystem-based fisheries management (Lindfield et al. 2016).


70
71 Mesophotic coral ecosystems occur in the deeper parts of the ocean’s photic zone, with
72 depth ranging from 30 to 150 m (Eyal et al. 2021), experience reduced temperatures and light, and
73 may be protected from localized anthropogenic stressors such as pollution and fishing. This
74 suggests that cooler conditions in deeper coral reefs could buffer the impacts of the strongest

75 stressors affecting fishes in shallow water. The deep reef refugia hypothesis emerged from genetic
76 connectivity research (Lesser et al. 2009; Bongaerts et al. 2010; Assis et al. 2016) and was
77 reinforced by the fact that many coral reef species occur across a wide depth range (Rocha et al.
78 2018; Medeiros et al. 2021). However, the increasing frequency and severity of marine heatwaves
79 (Oliver et al. 2018; Venegas et al. 2019), and accumulating reports of impacts reaching mesophotic
80 depths (Jackson et al. 2001), cast doubt on the ability of depth refuges to provide protection to fish
81 communities over the long-term (Rocha et al. 2018).

82
83 A refugium is defined as an “habitat where some components of biodiversity retreat to, persist in
84 and can potentially expand from under changing environmental conditions” (sensu, Keppel et al.
85 2012). The role of refugia to protect species from extinction has been assessed worldwide for
86 different taxa (Pellissier et al. 2014; Brown et al. 2020). However, biodiversity is multifaceted and
87 includes other components in addition to the number of species. A largely unexplored dimension
88 of the role of refugia is their effects on functional diversity, despite the importance of functional
89 diversity for community and ecosystem functioning. In other words, the ability of refugia to fully
90 encapsulate diversity has been poorly investigated, especially as it relates to their capacity to
91 reduce the loss of ecological function (Ordóñez and Svenning 2015). This is particularly true for
92 mesophotic reefs. Indeed, despite the hypothesis suggesting that mesophotic reefs could be refugia
93 for reef biodiversity (Jankowski et al. 2015; Lindfield et al. 2016; Coleman et al. 2018; Rocha et
94 al. 2018; Stefanoudis et al. 2019), there is still a critical lack of evaluation of their role as refugia
95 for species and trait diversity of fishes (Carrington et al. 2021; Cure et al. 2021).

96
97 Thus, considering the multifaceted definition of biodiversity, the mesophotic reef refugia
98 hypothesis could follow several expectations, depending on similarity in species and trait diversity
99 between shallow and mesophotic reefs (Figure 1): (1) Enriched full refugia: mesophotic reefs host
100 the same species and trait composition as shallow reefs, plus host unique species and traits not
101 observed in shallow reefs; (2) Full refugia: mesophotic reefs host the same species and traits
102 composition than shallow reef while maintaining the same biomass distribution among species;
103 (3) Functional refugia: mesophotic reefs host different species than shallow reefs but these distinct
104 species have the same trait values; (4) Not refugia: mesophotic and shallow reefs host distinct
105 species and traits composition.

106
107 Here, we tested the potential refugia role of mesophotic ecosystems for multiple facets of
108 fish diversity using a video-based survey in coral reefs from Mayotte (an island in the Western
109 Indian Ocean), along a depth gradient from 7.8 to 99.7 m. More specifically, we analyzed how
110 taxonomic and functional diversity and composition of coral reef fishes changed along the depth
111 gradient.



	Taxonomic		Functional	
	Δ Richness	Dissimilarity	Δ Richness	Dissimilarity
A	0	1	0	1
B	< 0	≈ 0.5	< 0	≈ 0.5
C	0	≈ 0.5	0	≈ 0.5
D	0	1	0	0
E	0	0	0	0
F	> 0	≈ 0	> 0	≈ 0

112
 113 *Figure 1: A multifaceted approach to test whether deep reefs are refugia for taxonomic (circle)*
 114 *and functional (square) facets of biodiversity. The size of a circle or square represents the*
 115 *taxonomic or functional richness of an assemblage while overlapping represents similarity in*
 116 *species or functional composition between assemblages from mesophotic and shallow reefs. The*
 117 *associated table shows the expected value of four diversity metrics for each scenario. Δ Richness*
 118 *is the difference in richness between mesophotic and shallow and dissimilarity is the proportion*
 119 *of richness not shared by shallow and mesophotic reefs. A: No refugia, mesophotic and shallow*
 120 *reefs host highly distinct species and trait combinations. B & C: Partial refugia, shallow and*
 121 *mesophotic reefs share a subset of species and traits but also a proportion of unique species and*
 122 *traits, in B, shallow reefs host more species and trait combinations than mesophotic reef. D:*
 123 *Functional refugia, mesophotic reef host dissimilar species communities from shallow reefs, but*
 124 *similar traits composition. E: Full refugia, mesophotic reefs host the same composition of species*
 125 *and hence of traits as shallow reefs. F: Enriched full refugia, mesophotic reefs host all the species*
 126 *and trait combinations present in shallow reefs, plus host unique species which have unique traits*
 127 *(absent in shallow reefs).*
 128

129 **MATERIALS AND METHODS**


130 **Underwater video surveys of coral reef fishes**

131 Mayotte is a volcanic island (360 km²) in the northern Mozambique Channel and is
132 surrounded by one of the world's largest lagoons (area of ~1500 km² surface). The coral reefs of
133 Mayotte hosts more than 700 fish species (Wickel et al. 2014), but recent and rapid human
134 development (annual growth rate higher than 4%, Marie et al. 2018) is driving fast-paced change
135 across the coral reef communities.

136 The choice of study sites was made using two complementary tools: observation of the
137 habitat using rosette cameras and a 3D reconstruction of the site using data collected by the boat's
138 sounder. The rosette camera system consisted of four cameras oriented in four different directions.
139 This system was manually lowered to the bottom over the point of interest using a rope and the
140 videos were viewed afterwards. For the 3D reconstruction, latitude, longitude and depth
141 coordinates were recorded every second using a GPS plotter sonar (Lowrance, HDS10). These
142 traces were then used to reconstruct the 3D profile of the substrate using the *raster* (Hijmans et al.
143 2015), *gstat* (Pebesma and Heuvelink 2016) and *sp* (Bivand et al. 2008) R packages (Figure 2).
144 Using to this information, we selected six sites along the west coral reef barrier of the island with
145 as much similarity as possible in coral cover and complexity within each four different depths,
146 ranging from 7.8 to 99.7 m deep (Figure 2).

147 Fish communities were surveyed using a non-baited remote underwater stereovideo
148 system, which is a cost-effective technique for surveying reef fish across a range of depths and
149 habitats and beyond the limits of regular scientific diving (Mallet and Pelletier 2014). We chose
150 non-baited video systems to avoid over-representation of piscivore species. The stereovideo
151 system consisted of two digital cameras (GoPro Hero 4+ and Hero 3+) inside 150m-rated housings
152 mounted on an aluminum bar, 80 cm apart and each oriented 8° inward using 3D printed mounts
153 to maximize pictures overlap 5 m away (Letessier et al. 2015). Four PVC legs (length 60 cm), each
154 weighted with 1 kg, were attached under the bar to make a quadripod system. The recording was
155 made using the cameras' medium view setting (84° of horizontal field of view in water) and at full
156 HD definition (1080p) with a frame rate of 30 fps. Calibration of the system was performed using
157 a 2x2x1 m black cubic frame with white dots of known coordinates situated 4 m from the system
158 underwater. Distortion of the cameras was calibrated with an 80x50 cm chessboard situated 1 m
159 from the cameras underwater (Neuswanger et al. 2016). Measurement accuracy was verified by
160 filming a graduated ruler underwater with each system.

161 Each of the 6 sites was surveyed several times (between 9 and 64 times, 20.2 ± 21.6 , mean
162 \pm sd) from 26-Feb to 23-Mar 2019. All surveys were performed between 9:00 and 17:00. Weather
163 ranged from warm and calm sunny days to rainy and windy days with as much as 1.5 m waves in
164 that period. Current did not exceed 0.8 knots. To minimize fishes moving between drops, and
165 potential spatial pseudo-replication, the four depths within a site were simultaneously surveyed,
166 and video systems within a depth level were dropped at least 100m apart from each other (Langlois
167 et al. 2020). Videos recording lasted from 55 minutes to 171 minutes (98.9 ± 36.3).



168
 169 *Figure 2: Location of surveyed sites around Mayotte Island in the SW Indian Ocean. On the left,*
 170 *the 3D reconstruction of each site and on the right each red dot represents a drop of a remote*
 171 *underwater camera.*
 172

173 **Video analyses**

174 We analyzed data from 121 drops (i.e., assemblages with fewer than five species were not
 175 included). For each video, the first two minutes were not analyzed to limit disturbance due to the
 176 video system deployment. As the total duration of video recorded varied among sites and depth
 177 level, and as analyzing stereovideos is a demanding task we analyzed five, one-minute non-
 178 overlapping sequences randomly selected within each recording. Hence, we keep a constant
 179 sampling effort across all replicates for estimating abundance of fish species, even if this sampling
 180 effort did not allow to reach the asymptote of the species accumulation curve (Supplementary
 181 materials, Figure S1). For each of these sequences, all fishes up to 5 m from the camera were
 182 counted and identified to the lowest possible taxonomic level. Abundance of each species was
 183 estimated as the maximum number of individuals of each species observed in a single frame during
 184 each one-minute sequence (MaxN; Ellis 1995). A total of 12,891 individuals (50%) belonging to
 185 182 species were identified. 2206 individuals were identified at the genus level (44 genera) and
 186 10,382 (40%) at the family level (26 families). We also did not consider cryptobenthic species
 187 (*Apogonidae* [7 individuals], *Gobiidae* [33 individuals] and *Chondrichthyans* [30 individuals]).
 188 Because a high proportion of individuals were not identified at the species levels, we also tested
 189 the sensitivity of our analyses to the taxonomic resolution by rerunning our analyses at the genus
 190 level. Size calibration calculation and 3D fish measurements were all performed using the open-
 191 source software VidSync (www.vidsync.org).
 192

193 **Biomass estimation**

194 Measuring manually each individual for the 121 videos was impractical given their number. Thus,
 195 we measured all individuals for 45 videos (for a total of 7h18min) at one of the six sites. Then, for
 196 each individual in the present study, we assigned the measured average size at the lowest
 197 taxonomic level (species, genus, family) possible according to available data. This process may
 198 reduce the dissimilarity in species size between shallow and mesophotic reefs. However,
 199 comparison of the distribution in size between shallow reef (from 0 to 30m) and mesophotic reef
 200 (from 30 to 99m) for the most measured common species (37) show that body size did not

201 significantly vary across the depth gradient (Supplementary materials, Figure S2). Some species
202 or genera were not observed at the selected site for fish measurement (59%). Thus, we associated
203 the average size for the given taxonomic level from the Reef Life Survey database (highly
204 correlated with the average size from our dataset, $\text{cor} = 0.78$, $\text{p-value} < 0.001$,
205 <https://reeflifesurvey.com/>, Edgar et al. 2020).

206

207 **Functional traits**

208 Five traits linked to the key functions performed by fishes were selected: Diet, Position in
209 the water column, Gregariousness, Activity, Mobility, Size. These traits linked to food acquisition,
210 locomotion and behaviour are a proxy of the ability of each species to impact ecosystem processes.
211 Values were taken from the dataset used by (Mouillot et al. 2014). Diet characterized each species'
212 main food source and was coded using seven categories: herbivore and detritivore, macro-algae
213 herbivore, sessile invertebrate feeder, mobile invertebrate feeder, planktivore, piscivore and
214 omnivore. Position in the water column was coded using four ordered categories: benthic
215 (sedentary on the bottom), demersal (swimming near the bottom), pelagic site-attached (swimming
216 off the bottom within a reef), and pelagic mobile (swimming off the bottom between reefs).
217 Mobility was coded using three ordered categories: sedentary, mobile within a reef, and mobile
218 between reefs. Active period was coded as either diurnal (active during the day), nocturnal (active
219 during the night) or both. Gregariousness was coded using five ordered categories from solitary to
220 large schooling. Body size was coded using six ordered categories: 0–7 cm, 7.1–15 cm, 15.1–30
221 cm, 30.1–50 cm, 50.1–80 cm, and >80 cm. For individuals identified at the genus or family level,
222 we used the average modality of each trait for the given genus or family. Functional distance
223 between all pairs of species was computed using Gower's metric (Gower and Legendre 1986).

224

225 **Taxonomic and functional diversity**

226 To evaluate the influence of depth on reef fish diversity of each assemblage we applied a
227 framework based on a generalization of Hill numbers to measure taxonomic and functional
228 diversity in all assemblages (Chao et al. 2019). Hill numbers provide a parametric family of
229 diversity indices, differing by a parameter “ q ” that determines their sensitivity to species relative
230 abundances, all expressed in the same unit as “equivalent number of species” (*sensu* species with
231 even dominance and even functional distance).

232

233 Taxonomic diversity was described using two indices: species richness (i.e., $q = 0$ indicates that
234 all species have equal weight) and taxonomic entropy (i.e., $q=1$, as exponential of the Shannon
235 index):

$$236 \quad \text{TD} = e^{-\sum_{i=1}^S p_i \log \log p_i} \quad \text{Eq (1)}$$

237

238 where i is the number of species in an assemblage and p_i is the relative dominance (abundance or
239 biomass) of the i th species.

240

241 To estimate functional diversity, we also used the framework from Chao et al. (2019)
242 setting the parameter τ to the mean functional distance over all pairs of species:

243

$$244 \quad \text{FD} = (\sum_{i=1}^S p_i (\sum_{j=1}^S [1 - f(d_{ij}(\tau))] p_j)^{q-1})^{1/(1-q)} \quad \text{Eq (2)}$$

245

246 Functional richness ($q = 0$) was computed with species presence/absence (i.e., all p_i equals
247 to $1/S$):

$$248 \quad FD = \sum_{i=1}^S (\sum_{j=1}^S [1 - f(d_{ij}(\tau))])^{-1} \quad \text{Eq (3)}$$

249
250 Functional entropy ($q = 1$) was computed on species biomass:

$$251 \quad FD = e^{-\sum_{i=1}^S p_i \log(\sum_{j=1}^S [1 - f(d_{ij}(\tau))] p_j)} \quad \text{Eq (4)}$$

252
253
254 This metric is a generalization of the Shannon entropy with functional distance having equal
255 importance as species dominance.

256 257 **Taxonomic and functional dissimilarities**

258 The dissimilarity in species and functional entities among assemblages was also computed
259 using the framework of Chao et al. (2019) with, as for alpha-diversity, $q=0$ (i.e., dissimilarity in
260 composition of species or functional entities) and $q=1$ (i.e., dissimilarity in dominance of species
261 or functional entities). The dissimilarity indices are bounded between 0 and 1, with 0 indicating
262 identical assemblages (in terms of species or trait values) and values close to 1 indicating a strong
263 dissimilarity in species or functional entities. We applied a depth decay analysis (similar to the
264 distance decay in biogeography, Nekola and White 1999) by comparing the dissimilarity between
265 0-20m assemblages and each of all other deeper assemblages.

266
267 Changes in functional richness and structure through depth were illustrated in a functional
268 space built using Principal Coordinates Analysis (PCoA) computed on functional distances
269 between species. Indices and functional space were computed and drawn using the *mFD* package
270 (Magneville et al. 2022).

271 272 **Habitat predictors**

273 To include the environment of each assemblage as covariate, we measured variables related to
274 abiotic and biotic habitat recorded by the video systems. It is however important to note that hard
275 substrate was mainly targeted and any results on variation among habitat should be interpreted
276 with caution. For the abiotic variables, we recorded the substrate type (soft, mix, hard),
277 topography, habitat complexity (from 0 to 3), temperature and depth. For temperature and depth
278 measurements, we used a custom-made data logger built by the company Syera
279 (<https://syera.fr/wp/glog/>) and using the Bar30 temperature-depth sensor from the company blue
280 robotics (<https://bluerobotics.com/>). Moreover, categorical abundance of dominant coral biota,
281 sponge, non-scleractinian coral and scleractinian coral were recorded. We performed a principal
282 component analysis of a mixture of numerical and categorical variables (Factor analysis of mixed
283 data, FAMD, using *FactoMineR*). Depth was not included in this analysis. The first four
284 orthogonal PCA axes explained 62% of the total variance in habitat predictors among assemblages
285 and were retained as explanatory variables in the analyses to test environmental influence on fish
286 assemblages (Supplementary materials, Figure S3).

287 288 **Determinants of fish diversity**

289 We used generalized linear mixed models (GLMMs) to investigate the effect of depth while
290 controlling for environmental differences in the four diversity indices and biomass hypothesized

291 to influence composition, entropy and traits of reef fish communities. Variables included the first
292 four axes of the environmental FAMD and depth while accounting for the site and date of the
293 sampling as random effects. For each model, we computed marginal R^2 providing the variance
294 explained only by fixed effects and conditional R^2 providing the variance explained by the entire
295 model, i.e., both fixed effects and random effects. Finally, we performed visreg analyses (Breheny
296 and Burchett 2017) to isolate the exclusive effect of depth on diversity after accounting for
297 environmental confounding factors.

298

299 **Determinants of fish dissimilarity**

300 To test if depth was the best predictor of taxonomic and functional dissimilarities, we applied
301 generalized dissimilarity modelling (*GDM*, Ferrier et al. 2007). *GDM* examines the dissimilarities
302 among pairs of assemblages as a non-linear multivariate function of the depth and environment of
303 those assemblages. *GDM* thus overcomes two potential problems: non-linearity in dissimilarity
304 between assemblages and environmental dissimilarities, and uneven rates of dissimilarities along
305 environmental gradients (Ferrier et al. 2007; Manion et al. 2016). We selected the three I-spline
306 basis function options and accounted for spatial autocorrelation by including the geographic
307 distance between pairs of assemblages as a predictor variable. To assess the impact of each
308 environmental variable, multiple *GDMs* were run by removing one variable at a time. Percentage
309 contribution of each variable was determined by comparing deviance of the full model to deviance
310 of models without the variable of interest. Variables included the first four axes of the
311 environmental PCA and depth.

312


313 **RESULTS**

314


315 **Diversity of fishes along depth gradient**

316 Total fish biomass and taxonomic richness ($q = 0$) significantly decreased with depth
317 (Figure 3 & 4, p -value < 0.001 , Table S1). On average, $29.3 (\pm 10.7 \text{ sd})$ species were observed
318 between 0-20 m, and $8.4 (\pm 3.6 \text{ sd})$ beyond 80 m. Taxonomic entropy ($q = 0$) also decreased with
319 depth, but less steeply than taxonomic richness, indicating that decreasing species richness was
320 associated with an increase in species dominance (p -value = 0.002).

321 Functional richness values varied from 1.7 to 6.2 among assemblages (4.3 ± 1.0 , Figure 4).
322 Functional entropy varied from 1.1 to 4.8 among assemblages (2.8 ± 0.7 , Figure 4). Unlike
323 taxonomic diversity, functional richness and functional entropy were not significantly influenced
324 by depth (p -value = 0.134, 0.498, respectively, Table S1).



325
 326 *Figure 3 Partial regression plot showing changes in biomass of fish along the depth gradient*
 327 *while accounting for environmental variables. Statistical analyses were carried out using*
 328 *generalized linear mixed models and related statistics are reported in the Results.*
 329



330
 331
 332 *Figure 4: Partial regression plot showing changes in taxonomic richness ($q=0$, A) and entropy*
 333 *($q=1$, B) and functional richness ($q=0$, τ = mean functional distance, C) and entropy ($q=1$, τ =*
 334 *mean functional distance, D) of fish communities along the depth gradient, while accounting for*


335 *environmental variables. Statistical analyses were carried out using generalized linear mixed*
 336 *models and related statistics are reported in the Results.*

337

338 **Dissimilarity of fish assemblages along depth gradient**

339 The dissimilarity in taxonomic composition and structure between assemblages was high overall,
 340 ranging from 0.4 to 1 (0.86 ± 0.09) and 0.1 to 1 (0.77 ± 0.2), respectively. The highest dissimilarity
 341 values were observed between shallow (0-20 m) and deep assemblages (>60 m) (i.e., high value
 342 in the difference in depth axis, Figure 5). Limited dissimilarity in functional composition was
 343 observed between assemblages (0.13 ± 0.13). The dissimilarity in functional structure was higher
 344 than functional composition (0.46 ± 0.25), highlighting an important change in the distribution of
 345 biomass between functional entities. Both taxonomic and functional dissimilarities increased with
 346 an increase in depth difference (all p-values < 0.001 and R^2 ranged between 0.30 and 0.47; Figure
 347 5).

348 The diversity patterns described above were very similar to those computed only with the
 349 genus presence and dominance (Supplementary materials, Figure S4 and S5).




350

351 *Figure 5: Depth decay of fish diversity, mean dissimilarity (whiskers representing sd) in taxonomic*
 352 *and functional composition and structure between 0-20m assemblages (coded as being at 0m) and*
 353 *each all other deeper assemblages. The first row shows dissimilarity in species composition ($q =$*
 354 *0) and structure ($q = 1$). The second row dissimilarity in functional composition ($q = 0, \tau =$ mean*
 355 *functional distance) and structure ($q = 0, \tau =$ mean functional distance). Yellow lines are linear*

356 regressions (with associated statistics) between the difference in depth and beta diversity indices.
357 Shaded areas represent the 95% confidence interval of the regression.

358
359 Finally, the projection of all species along the first two axes of the PCoA confirmed that
360 mesophotic and shallow reefs mostly overlap in functional space but distribution of biomass within
361 this shared functional volume markedly changed (Figure 6).
362



363
364 *Figure 6: Changes in functional composition and structure along the depth gradient. The traits*
365 *space represented is made of the first two axes (Principal Components of a PCoA, relative*
366 *importance: 32 and 30%, respectively). Species are represented by circles whose size is*
367 *proportional to the biomass of the species. The gray area represents the portion of the trait space*
368 *filled by all species present across all depths and colored areas represented the functional*
369 *richness, i.e., trait space filled by species present in each depth range. Vertical and horizontal*
370 *lines illustrate the biomass-weighted average position of species from each depth range.*

371
372 *Environmental drivers of taxonomic and functional diversity*

373 Despite differences in habitat among sites, depth was the most important variable in
374 explaining biomass, number of species and taxonomic entropy, followed by PC1 of the habitat
375 data (Table S1). This axis is driven by the shift from scleractinian corals to non-scleractinian corals
376 with depth (Supplementary materials, Figure S3). For taxonomic richness and taxonomic entropy,
377 the models explain 66% (conditional $R^2 = 0.66$ and marginal $R^2 = 0.45$) and 22% (conditional R^2
378 $= 0.22$ and marginal $R^2 = 0.19$) of the variance, respectively. Visreg confirmed that depth, after
379 controlling for environmental conditions, significantly explained the biomass, species richness and
380 entropy pattern between assemblages (Figure 3 and 4). Conversely, environmental conditions and
381 depth poorly explain variation of functional richness and entropy (25 and 12%, respectively;
382 conditional $R^2 = 0.25$, 0.12 and marginal $R^2 = 0.21$ and 0.09, respectively).
383 The GDM explained 41 and 44% of the variation in species composition and structure,
384 respectively. Depth was the most important explanatory variable for changes in species
385 composition (Supplementary materials, Figure S6), contributing to 16% and 21% of the explained
386 deviance, respectively, with a lower effect of environmental PC1 (11% of the deviance explained
387 for both models). Variation in functional composition was poorly explained by differences in
388 environmental variables and depth (9% of deviance explained). The GDM explained 41% of the
389 variation in functional structure, and depth was the greatest driver, contributing to 23% of the
390 deviance explained.

391 392 **DISCUSSION**

393 Here, thanks to a video survey, we documented how taxonomic and trait diversity of reef
394 fishes changed along a depth gradient from 7.8 to 99.7m around Mayotte Island. Deeper
395 ecosystems exhibit reduced species richness as already reported (Piacenza et al. 2015). However,
396 our results demonstrate that shallow reefs and mesophotic reefs contain similar levels of functional
397 richness with a partial overlap in trait space despite the difference in species composition. While
398 such convergence has been observed at biogeographic scale (Loiseau et al. 2017; Olivier et al.
399 2018; McLean et al. 2021), ours is a novel example showing trait convergence in marine fish
400 communities driven by taxonomically different species.

401 Thus, while the taxonomic composition of the fish fauna varied over depth, species were
402 often replaced by another with similar functional roles such that only minor differences in
403 functional entities occurred across depth ranges. For instance, small planktivorous *Chromis* spp.
404 were progressively replaced by *Pseudanthias* spp. another small planktivorous species, with a ratio
405 of two *Pseudanthias* spp. per 100 *Chromis* spp. between 0 and 20 m versus a ratio of 49 per 100
406 beyond 60 m. This trait similarity across the depth gradient could limit the competition between
407 species that have different distributions along the depth gradient and thus promote coexistence of
408 fishes on the seascape (Goetze et al. 2021; Bosch et al. 2021).

409 Thus, functional space remains relatively constant over the entire water column. This
410 finding is similar to Carrington et al.'s (2021) findings that temperate shallow-water and deep-sea
411 ecosystems have similar functional composition. The limited shift in trait composition could be
412 explained by the relative homogeneity in the temperature of the water column from our sampling
413 area, ranging only from 24.6 to 31°C. Even if biologically significant, this 6°C degree difference
414 may not be strong enough to limit the establishment of adapted shallow reef functional entities at
415 greater depths. For instance, in the Bay of La Paz, across the same range of depth temperature
416 decreased by 11°C (Hollarsmith et al. 2020). However, conversely to our study Medeiros et al.
417 (2021) found a divergence in both species and functional composition with depth along the
418 Northeastern Brazilian subprovince characterized by a temperature range similar to ours (7°C).

419 However, despite the functional richness of shallow and mesophotic reefs remaining
420 relatively similar, traits dependent on light energy, like herbivory, do not occur beyond 60m depth
421 (Supplementary materials, Figure S7). Such a decrease is expected to be due to harsher
422 environmental conditions in deep habitat that filter out some species present on shallow reefs
423 (Price et al. 1999; Costello and Chaudhary 2017). Moreover, the overlap in trait space (i.e.,
424 functional similarity) is accompanied by changes in the structure of the traits with a significant
425 shift in biomass between functional entities. Indeed, the increase in the dissimilarity of functional
426 structure with depth highlights the fact that dominant and rare functional entities are different
427 between shallow reefs and deep reefs (Figure 5). For instance, 75% of *Chaetodontidae*,
428 corallivores, were observed between 0 and 34 m and only 5% were observed beyond 48 m.
429 Conversely, 50% of *Carangidae*, which are highly mobile and piscivorous, were recorded beyond
430 48 m. Overall, variation in trait structure was characterized by a greater relative biomass and
431 abundance of piscivores on deep reefs, while herbivorous fishes were relatively more abundant on
432 shallow reefs (Supplementary materials, Figure S7). In fact, piscivorous species represent 8% of
433 the total biomass in shallow reefs (0-20m), but represent 48% of the biomass beyond 60 m. The
434 biomass of piscivorous and mobile fishes within reefs, such as grouper, is even higher beyond
435 60 m ($538g \pm 859$, mean \pm sd) than between 0 and 20 m ($191g \pm 222$).

436 Coral reef ecosystems have been heavily impacted by temperature rise over the last few
437 decades, with documented increases in intense heatwaves leading to massive bleaching events
438 (Genevier et al. 2019). Synergistically, fisheries strongly impact coral reef fish communities
439 (MacNeil et al. 2015). Deep-sea ecosystems have been described as stable, with the incidence and
440 intensity of heatwaves, occurrence of storms and fishing activities lower in mesophotic depths than
441 in shallow water (Kahng et al. 2017). This “stability” suggests that mesophotic reefs could provide
442 protection to fish communities (Bongaerts et al. 2010). In the same vein, we found that the upper
443 mesophotic (30-60 m) zone holds the greatest potential to aid in reef recovery following
444 disturbance due to the functional overlap with the shallow reef, suggesting that the upper
445 mesophotic could be a “functional refugia”. However, the limited similarity in species composition
446 between mesophotic assemblages and shallow reef assemblages suggests that only a relatively
447 small number of deeper species may be able to live on shallow reefs. Moreover, the strong decrease
448 in biomass with depth supports the hypothesis that mesophotic reefs are a limited functional
449 refugia in the face of climate change (i.e. they host dissimilar species, but similar traits while
450 exhibiting a decrease and a shift in biomass distribution across traits). Lastly, despite the minimal
451 shift in functional trait composition, there is a marked shift from dominance of herbivorous fishes
452 to piscivorous fishes with increasing depth. Herbivorous species can have positive effects on coral
453 cover by controlling macroalgae populations, which can prevent coral mortality and promoting
454 resilience of coral reefs in the face of anthropogenic disturbance and heatwaves (Burkepile and
455 Hay 2008). Maintaining high herbivore biomass and diversity is an important goal for sustaining
456 coral reef functioning (Lefcheck et al. 2019) that cannot be supported by mesophotic reefs deeper
457 than 60m. However, our results demonstrate the importance of mesophotic reefs to fishery-targeted
458 species, since the biomass of some piscivorous fishes that are targets of fisheries increase with
459 depth (Lindfield et al. 2016; Weijerman et al. 2019; Bosch et al. 2021). Even low fishing pressure
460 on shallow reefs of Mayotte could also create a decreasing gradient of biomass of targeted
461 piscivorous (Doherty et al. 2015).

462 While we documented the convergence of trait diversity over depth, we are unable to
463 explain the potential consequences for the ecosystem functioning, since studies linking fish traits
464 to ecosystem functioning are still lacking (Villéger et al. 2017; Bellwood et al. 2018; Bosch et al.

465 2021b). However, since biomass distribution across traits determines the influence of fish on
466 ecosystems functioning, we can easily assume that shallow and mesophotic reef fish communities
467 do not have the same influence of reef ecosystems. As with all traits-based studies, trait choices
468 ultimately influence research conclusions. Our results are limited to the set of traits used and to
469 the depth range scales examined. Future studies should therefore attempt to identify the influence
470 of depth on physiologic traits such as reproduction and direct functions such as nutrient cycling
471 (Allgeier et al. 2016) and productivity (Morais and Bellwood 2020) to better understand the
472 functional complementarity of mesophotic and shallow reefs. Understanding how these functions
473 vary along depth gradients is critically important to predict how mesophotic reefs may respond to
474 future environmental change and for their management and conservation. Finally, fishes associated
475 with mesophotic reefs may perform vertical movements over seasonal and diel time frames
476 (Bacheler et al. 2021; Brazo et al. 2021). This behavior could increase the similarity between
477 shallow and mesophotic reefs.

478
479 Preserving marine ecosystems is a crucial priority, as they support the livelihood and well-being
480 of millions of citizens (human food provisioning, water purification, disease regulation, cultural
481 values, etc.) (Cabral et al. 2020). Despite their importance, and although the percentage of the coral
482 reef surface devoted to protected areas has markedly increased globally (Maxwell et al. 2020),
483 deep reefs are rarely the focus of conservation strategies. Fishing has historically impacted shallow
484 coral reefs more than the mesophotic part, but as technology continues to improve, the exploitation
485 of the deeper areas has been increasing (Audzijonyte and Pecl 2018). Thus, in the context of the
486 ongoing rise in the severity of threats, mesophotic reefs are an important habitat for conserving
487 coral reef fish diversity. The unique species composition hosted by mesophotic reefs supports the
488 recommendation that protected areas should be representative of a broad range of habitats across
489 depths. Expanding marine protected areas to adjacent mesophotic reefs is required to protect a
490 mosaic of habitat, maximising the number of species and functional entities protected, critical to
491 prevent future local population extirpations and maintain coral reef ecosystem functioning and
492 associated services at all depths. Sustainable management of deep reef ecosystems will also have
493 strong impacts on the management of fishery resources by providing shelters for targeted species.

494 495 **Acknowledgement**

496 This research was funded by the CUFR of Mayotte, by the second Research Consortium Iles eparse
497 2017-2020 (project MAPOR), by the French Foundation for Research on Biodiversity and its
498 partners (FRB - www.fondationbiodiversite.fr) and by ANR (The French National Research
499 Agency) under the “investissement d’avenir” program with the reference ANR-16-IDEX-0006.
500 The authors thank Priscilla Dupont and Julien Wikel for helping with fish ID on videos and Yann
501 Mercky and Elliott Sucre for their help on field operation.

502 **REFERENCES**

- 503
- 504 Allgeier JE, Valdivia A, Cox C, Layman CA (2016) Fishing down nutrients on coral reefs. *Nat*
505 *Commun* 7:12461
- 506 Assis J, Coelho NC, Lamy T, Valero M, Alberto F, Serrão EÁ (2016) Deep reefs are climatic
507 refugia for genetic diversity of marine forests. *Journal of biogeography* 43:833-844
- 508 Audzijonyte A, Pecl GT (2018) Deep impact of fisheries. *Nature ecology & evolution* 2:1348-
509 1349
- 510 Bachelier NM, Shertzer KW, Runde BJ, Rudershausen PJ, Buckel JA (2021) Environmental
511 conditions, diel period, and fish size influence the horizontal and vertical movements of
512 red snapper. *Scientific Reports* 11:1-16
- 513 Bellwood DR, Streit RP, Brandl SJ, Tebbett SB (2019) The meaning of the term ‘function’ in
514 ecology: A coral reef perspective. *Functional Ecology*, 33:948-961.
- 515 Bivand RS, Pebesma EJ, Gómez-Rubio V, Pebesma EJ (2008) *Applied spatial data analysis with*
516 *R*. Springer
- 517 Bongaerts P, Ridgway T, Sampayo E, Hoegh-Guldberg O (2010) Assessing the ‘deep reef
518 refugia’ hypothesis: focus on Caribbean reefs. *Coral reefs* 29:309-327
- 519 Bosch NE, Monk J, Goetze J, Wilson S, Babcock RC, Barrett N, ... & Langlois, TJ (2022) Effects
520 of human footprint and biophysical factors on the body-size structure of fished marine
521 species. *Conservation Biology*, 36(2), e13807.
- 522 Bosch NE, McLean M, Zarco-Perello S, Bennett S, Stuart-Smith RD, Vergés A, ... & Wernberg,
523 T (2022) Persistent thermally driven shift in the functional trait structure of herbivorous
524 fishes: Evidence of top-down control on the rebound potential of temperate seaweed
525 forests?. *Global Change Biology*, 28:2296-2311.
- 526 Brandl SJ, Rasher DB, Côté IM, Casey JM, Darling ES, Lefcheck JS, Duffy JE (2019) Coral reef
527 ecosystem functioning: eight core processes and the role of biodiversity. *Frontiers in*
528 *Ecology and the Environment* 17:445-454
- 529 Brazo A, Marques R, Zimmermann M, Aspillaga E, Hereu B, Saragoni G, Mercière A, Crec’Hriou
530 R, Mercader M, Verdoit-Jarraya M (2021) Seasonal influence on the bathymetric
531 distribution of an endangered fish within a marine protected area. *Scientific Reports* 11:1-
532 16
- 533 Breheny P, Burchett W (2017) Visualization of regression models using visreg. *R J* 9:56
- 534 Brown SC, Wigley TM, Otto-Bliesner BL, Rahbek C, Fordham DA (2020) Persistent Quaternary
535 climate refugia are hospices for biodiversity in the Anthropocene. *Nature Climate Change*
536 10:244-248
- 537 Burkepile DE, Hay ME (2008) Herbivore species richness and feeding complementarity affect
538 community structure and function on a coral reef. *Proceedings of the National Academy of*
539 *Sciences* 105:16201-16206
- 540 Cabral RB, Bradley D, Mayorga J, Goodell W, Friedlander AM, Sala E, Costello C, Gaines SD
541 (2020) A global network of marine protected areas for food. *Proceedings of the National*
542 *Academy of Sciences* 117:28134-28139
- 543 Carrington VG, Papa Y, Beese CM, Hall J, Covain R, Horn P, Ladds MA, Rogers A, Yeo D (2021)
544 How functionally diverse are fish in the deep? A comparison of fish communities in deep
545 and shallow-water systems. *Diversity and Distributions*

546 Chao A, Chiu CH, Villéger S, Sun IF, Thorn S, Lin YC, Chiang JM, Sherwin WB (2019) An
547 attribute-diversity approach to functional diversity, functional beta diversity, and related
548 (dis) similarity measures. *Ecological Monographs* 89:e01343

549 Cinner JE, Huchery C, MacNeil MA, Graham NA, McClanahan TR, Maina J, Maire E, Kittinger
550 JN, Hicks CC, Mora C (2016) Bright spots among the world's coral reefs. *Nature* 535:416-
551 419

552 Cinner JE, Zamborain-Mason J, Gurney GG, Graham NA, MacNeil MA, Hoey AS, ... & Mouillot
553 D. (2020). Meeting fisheries, ecosystem function, and biodiversity goals in a human-
554 dominated world. *Science*, 368(6488), 307-311.

555 Coleman RR, Copus JM, Coffey DM, Whitton RK, Bowen BW (2018) Shifting reef fish
556 assemblages along a depth gradient in Pohnpei, Micronesia. *PeerJ* 6:e4650

557 Costello MJ, Chaudhary C (2017) Marine biodiversity, biogeography, deep-sea gradients, and
558 conservation. *Current Biology* 27:R511-R527

559 Cure K, Currey-Randall L, Galaiduk R, Radford B, Wakeford M, Heyward A (2021) Depth
560 gradients in abundance and functional roles suggest limited depth refuges for herbivorous
561 fishes. *Coral Reefs* 40:365-379

562 Doherty B, Herfaut J, Le Manach F, Harper S, Zeller D (2015) Reconstructing domestic marine
563 fisheries in Mayotte from 1950–2010. *Fisheries Catch Reconstructions in the Western*
564 *Indian Ocean*, eds F Le Manach and D Pauly (Vancouver: University of British
565 Columbia):53-65

566 Eddy TD, Lam VW, Reygondeau G, Cisneros-Montemayor AM, Greer K, Palomares MLD, Bruno
567 JF, Ota Y, Cheung WW (2021) Global decline in capacity of coral reefs to provide
568 ecosystem services. *One Earth* 4:1278-1285

569 Edgar GJ, Cooper A, Baker SC, Barker W, Barrett NS, Becerro MA, Bates AE, Brock D,
570 Ceccarelli DM, Clausius E (2020) Reef Life Survey: Establishing the ecological basis for
571 conservation of shallow marine life. *Biological Conservation* 252:1-14

572 Ellis D (1995) Evaluation of video camera technique for indexing abundances of juvenile pink
573 snapper *Pristipomoides filamentosus*, and other Hawaiian insular shelf fishes. *Fishery*
574 *Bulletin* 93:67-77

575 Eyal G, Laverick JH, Bongaerts P, Levy O, Pandolfi JM (2021) Mesophotic coral ecosystems of
576 the Great Barrier Reef are understudied and underexplored. *Frontiers in Marine Science*
577 8:83

578 Ferrier S, Manion G, Elith J, Richardson K (2007) Using generalized dissimilarity modelling to
579 analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity*
580 *and distributions* 13:252-264

581 Geneviev LG, Jamil T, Raitos DE, Krokos G, Hoteit I (2019) Marine heatwaves reveal coral reef
582 zones susceptible to bleaching in the Red Sea. *Global change biology* 25:2338-2351

583 Goetze JS, Wilson S, Radford B, Fisher R, Langlois TJ, Monk J, Knott NA, Malcolm H, Currey-
584 Randall LM, Ierodiaconou D (2021) Increased connectivity and depth improve the
585 effectiveness of marine reserves. *Global Change Biology*

586 Gower JC, Legendre P (1986) Metric and Euclidean properties of dissimilarity coefficients.
587 *Journal of classification* 3:5-48

588 Hijmans RJ, Van Etten J, Cheng J, Mattiuzzi M, Sumner M, Greenberg JA, Lamigueiro OP, Bevan
589 A, Racine EB, Shortridge A (2015) Package 'raster'. R package 734

590 Jackson JB, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH,
591 Cooke R, Erlandson J, Estes JA (2001) Historical overfishing and the recent collapse of
592 coastal ecosystems. *science* 293:629-637

593 Hollarsmith JA, Ramirez-Ortiz G, Winqvist T, Velasco-Lozano M, DuBois K, Reyes-Bonilla H,
594 Neumann KC, Grosholz ED (2020) Habitats and fish communities at mesophotic depths in
595 the Mexican Pacific. *Journal of Biogeography*, 47:1552-1563.

596 Jankowski M, Graham N, Jones G (2015) Depth gradients in diversity, distribution and habitat
597 specialisation in coral reef fishes: implications for the depth-refuge hypothesis. *Marine*
598 *Ecology Progress Series* 540:203-215

599 Kahng S, Copus JM, Wagner D (2017) Mesophotic coral ecosystems. *Marine animal forests: the*
600 *ecology of benthic biodiversity hotspots*:185-206

601 Keppel G, Van Niel KP, Wardell-Johnson GW, Yates CJ, Byrne M, Mucina L, Schut AG, Hopper
602 SD, Franklin SE (2012) Refugia: identifying and understanding safe havens for
603 biodiversity under climate change. *Global Ecology and Biogeography* 21:393-404

604 Langlois T, Goetze J, Bond T, Monk J, Abesamis RA, Asher J, ... & Harvey ES (2020) A field and
605 video annotation guide for baited remote underwater stereo-video surveys of demersal fish
606 assemblages. *Methods in Ecology and Evolution* 11: 1401-1409.

607 Lefcheck JS, Innes-Gold AA, Brandl SJ, Steneck RS, Torres RE, Rasher DB (2019) Tropical fish
608 diversity enhances coral reef functioning across multiple scales. *Science advances*
609 5:eaav6420

610 Lesser MP, Slattery M, Leichter JJ (2009) Ecology of mesophotic coral reefs. *Journal of*
611 *Experimental Marine Biology and Ecology* 375:1-8

612 Letessier TB, Juhel J-B, Vigliola L, Meeuwig JJ (2015) Low-cost small action cameras in stereo
613 generates accurate underwater measurements of fish. *Journal of Experimental Marine*
614 *Biology and Ecology* 466:120-126

615 Lindfield SJ, Harvey ES, Halford AR, McIlwain JL (2016) Mesophotic depths as refuge areas for
616 fishery-targeted species on coral reefs. *Coral Reefs* 35:125-137

617 Loiseau N, Legras G, Kulbicki M, Mérigot B, Harmelin-Vivien M, Mazouni N, Galzin R, Gaertner
618 J-C (2017) Multi-component β -diversity approach reveals conservation dilemma between
619 species and functions of coral reef fishes. *Journal of Biogeography* 44:537-547

620 MacNeil MA, Graham NA, Cinner JE, Wilson SK, Williams ID, Maina J, Newman S, Friedlander
621 AM, Jupiter S, Polunin NV (2015) Recovery potential of the world's coral reef fishes.
622 *Nature* 520:341-344

623 Magneville C, Loiseau N, Albouy C, Casajus N, Claverie T, Escalas A, Leprieur F, Maire E,
624 Mouillot D, Villéger S (2022) mFD: an R package to compute and illustrate the multiple
625 facets of functional diversity. *Ecography* 2022

626 Mallet D, Pelletier D (2014) Underwater video techniques for observing coastal marine
627 biodiversity: a review of sixty years of publications (1952–2012). *Fisheries Research*
628 154:44-62

629 Manion G, Lisk M, Ferrier S, Nieto-Lugilde D, Fitzpatrick M (2016) gdm: Functions for
630 generalized dissimilarity modeling. Rpackage version 1

631 Marie C-V, Breton D, Crouzet M (2018) More than half of all adults living in Mayotte were born
632 elsewhere. *Population Societies*:1-4

633 Maxwell SL, Cazalis V, Dudley N, Hoffmann M, Rodrigues AS, Stolton S, Visconti P, Woodley
634 S, Kingston N, Lewis E (2020) Area-based conservation in the twenty-first century. *Nature*
635 586:217-227

636 Medeiros APM, Ferreira BP, Alvarado F, Betancur RR, Soares MO, Santos BA (2021) Deep reefs
637 are not refugium for shallow-water fish communities in the southwestern Atlantic. *Ecol*
638 *Evol* 11:4413-4427

639 Morais RA, Bellwood DR (2020) Principles for estimating fish productivity on coral reefs. *Coral*
640 *Reefs* 39:1221-1231

641 Mouillot D, Villéger S, Parravicini V, Kulbicki M, Arias-González JE, Bender M, Chabanet P,
642 Floeter SR, Friedlander A, Vigliola L (2014) Functional over-redundancy and high
643 functional vulnerability in global fish faunas on tropical reefs. *Proceedings of the National*
644 *Academy of Sciences* 111:13757-13762

645 Nekola JC, White PS (1999) The distance decay of similarity in biogeography and ecology. *Journal*
646 *of biogeography* 26:867-878

647 Neuswanger JR, Wipfli MS, Rosenberger AE, Hughes NF (2016) Measuring fish and their
648 physical habitats: versatile 2D and 3D video techniques with user-friendly software.
649 *Canadian Journal of Fisheries and Aquatic Sciences* 73:1861-1873

650 Olivier D, Loiseau N, Petatán-Ramírez D, Millán OT, Suárez-Castillo AN, Torre J, Munguia-Vega
651 A, Reyes-Bonilla H (2018) Functional-biogeography of the reef fishes of the islands of the
652 Gulf of California: integrating functional divergence into marine conservation. *Global*
653 *Ecology and Conservation* 16:e00506

654 Ordonez A, Svenning JC (2015) Geographic patterns in functional diversity deficits are linked to
655 glacial-interglacial climate stability and accessibility. *Global Ecology and Biogeography*
656 24:826-837

657 Oliver EC, Donat MG, Burrows MT, Moore PJ, Smale DA, Alexander LV, Benthuisen JA, Feng
658 M, Gupta AS, Hobday AJ, Holbrook NJ, Perkins-Kirkpatrick SE, Scannell HA, Straub SC,
659 Wernberg, T. (2018). Longer and more frequent marine heatwaves over the past century.
660 *Nature communications*, 9(1), 1-12.

661 Pebesma E, Heuvelink G (2016) Spatio-temporal interpolation using gstat. *RFID Journal* 8:204-
662 218

663 Pellissier L, Leprieur F, Parravicini V, Cowman PF, Kulbicki M, Litsios G, Olsen SM, Wisz MS,
664 Bellwood DR, Mouillot D (2014) Quaternary coral reef refugia preserved fish diversity.
665 *Science* 344:1016-1019

666 Piacenza SE, Barner AK, Benkwitt CE, Boersma KS, Cerny-Chipman EB, Ingeman KE,
667 Kindinger TL, Lee JD, Lindsley AJ, Reimer JN (2015) Patterns and variation in benthic
668 biodiversity in a large marine ecosystem. *PLoS One* 10:e0135135

669 Price A, Keeling M, O'callaghan C (1999) Ocean-scale patterns of 'biodiversity' of Atlantic
670 asteroids determined from taxonomic distinctness and other measures. *Biological Journal*
671 *of the Linnean Society* 66:187-203

672 Roberts TE, Bridge TC, Caley MJ, Madin JS, Baird AH (2019) Resolving the depth zonation
673 paradox in reef-building corals. *Ecology* 100:e02761

674 Rocha LA, Pinheiro HT, Shepherd B, Papastamatiou YP, Luiz OJ, Pyle RL, Bongaerts P (2018)
675 Mesophotic coral ecosystems are threatened and ecologically distinct from shallow water
676 reefs. *Science* 361:281-284

677 Sala E, Mayorga J, Bradley D, Cabral RB, Atwood TB, Auber A, Cheung W, Costello C, Ferretti
678 F, Friedlander AM (2021) Protecting the global ocean for biodiversity, food and climate.
679 *Nature* 592:397-402

680 Sale PF (2013) *The ecology of fishes on coral reefs*. Elsevier

- 681 Spalding M, Spalding MD, Ravilious C, Green EP (2001) World atlas of coral reefs. Univ of
682 California Press
- 683 Stefanoudis PV, Gress E, Pitt JM, Smith SR, Kincaid T, Rivers M, Andradi-Brown DA, Rowlands
684 G, Woodall LC, Rogers AD (2019) Depth-dependent structuring of reef fish assemblages
685 from the shallows to the rariphotic zone. *Frontiers in Marine Science* 6:307
- 686 Venegas RM, Oliver T, Liu G, Heron SF, Clark SJ, Pomeroy N, Young C, Eakin CM, Brainard
687 RE (2019) The rarity of depth refugia from coral bleaching heat stress in the western and
688 central Pacific Islands. *Scientific reports* 9:1-12
- 689 Villéger S, Brosse S, Mouchet M, Mouillot D, Vanni MJ (2017) Functional ecology of fish: current
690 approaches and future challenges. *Aquatic Sciences* 79:783-801
- 691 Weijerman M, Grüss A, Dove D, Asher J, Williams ID, Kelley C, Drazen JC (2019) Shining a
692 light on the composition and distribution patterns of mesophotic and subphotic fish
693 communities in Hawai 'i. *Marine Ecology Progress Series* 630:161-182
- 694 Wickel J, Jamon A, Pinault M, Durville P, Chabanet P (2014) Composition et structure des
695 peuplements ichthyologiques marins de l'île de Mayotte (sud-ouest de l'océan Indien).
696 *Cybium: Revue Internationale d'Ichtyologie* 38:179-203