



HAL
open science

Trajectory of a coastal social-ecological system: analyzing co-evolution and regime shifts in the Thau lagoon (Mediterranean Sea, France), 1970–2018

Valérie Derolez, Rémi Mongruel, Helene Rey-Valette, Nicole Lautrédou-Audouy

► **To cite this version:**

Valérie Derolez, Rémi Mongruel, Helene Rey-Valette, Nicole Lautrédou-Audouy. Trajectory of a coastal social-ecological system: analyzing co-evolution and regime shifts in the Thau lagoon (Mediterranean Sea, France), 1970–2018. *Regional Environmental Change*, 2023, 23 (2), pp.68. <10.1007/s10113-023-02061-y>. <hal-04092807>

HAL Id: hal-04092807

<https://hal.umontpellier.fr/hal-04092807v1>

Submitted on 12 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Trajectory of a coastal social-ecological system: analyzing co-evolution and regime shifts in the Thau lagoon (Mediterranean Sea, France), 1970-2018

Valérie Derolez^{1,*}, Rémi Mongruel², Hélène Rey-Valette³, Nicole Lautrédou-Audouy³

¹MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France

²AMURE, Ifremer, Univ Bretagne Occidentale, CNRS, Brest, France

³CEE-M, Université de Montpellier, CNRS, INRAE, InstitutAgro, Montpellier, France

* Corresponding author: valerie.derolez@ifremer.fr; (+33)4.99.57.32.79

Abstract

As coastal social-ecological systems (SES) are used for many different purposes and subject to different stressors, they are also among the most vulnerable and change the most rapidly. To achieve resilience, coastal areas require adaptive management based on integrated interdisciplinary approaches, requiring a deep understanding of SES dynamics, i.e., the ecological and social changes that comprise synchronicities and regime shifts.

For the purpose of analyzing co-evolution and regime shifts, we propose an original and operational methodology combining: (i) a chronosystemic timeline approach and (ii) time series analyses based on indicators from the “Drivers-Pressures-State-Impacts-Responses” framework (14 indicators including population growth, eutrophication status, ecosystem services, management actions and climate change indicators). The methodology was tested on a SES located on the French Mediterranean coast, the Thau lagoon, whose trajectory we studied over a 5-decade period (1970-2018).

Our results show that the dynamics of co-evolution and regime shifts of the SES were driven by a succession of interactions between sanitary or ecological crises and management adaptations. During the study period, this typical Mediterranean coastal SES appears to have evolved toward the multiplication of uses, improvement of water quality and more integrative management. Our methodology also highlighted three contrasting periods characterized by stable emerging properties and identified several cause-and-effect relationships, which is rare in historical analyses of SES. These cause-and-effect relationships concern major sanitary and

29 ecological crises that have led to a reorganization of governance mechanisms and to technical
30 remediation actions.

31 Our approach based on long-term trajectories will help identify the conditions required for
32 increasing the resilience of coastal SES and the best management objectives to face new
33 challenges related to global climate change.

34 **Keywords:** social-ecological system (SES); resilience; regime shift; timelines; trajectories;
35 coastal lagoon.

36

37 1. INTRODUCTION

38

39 Coastal ecosystems are known to provide many ecosystem services that contribute to human
40 wellbeing (Newton et al. 2014). As they are used for multiple purposes and subject to many
41 different stresses, they are also among the most vulnerable (Halpern et al. 2008). It is
42 therefore increasingly acknowledged that to achieve resilience, coastal zones require adaptive
43 management and development policies based on interdisciplinary and integrated approaches
44 (Lloyd et al. 2013). Stemming from ecology, resilience theory has become a powerful
45 transdisciplinary tool to understand the dynamics and trajectories of social-ecological systems
46 (Gunderson and Holling 2002). The idea of explicitly coupling the dynamics of ecological
47 and social systems as a whole was extensively developed for the first time by Berkes and
48 Folke (1998). These authors used the term social-ecological system (SES) to stress that the
49 delineation between social and ecological systems is artificial and arbitrary: basically, the SES
50 perspective analyzes how ecological and social processes interact at different spatial and time
51 scales, and to consider this complex dynamic system as capable of self-organization and
52 perpetual adaptations (Folke 2006). In practice, dealing with the diversity of processes
53 involved in SES dynamics has given rise to numerous conceptual and methodological
54 frameworks requiring interdisciplinary approaches.

55 Many frameworks have been proposed to analyze a whole complex SES, which often overlap
56 or refer to each other (Binder et al. 2013). Some are purely conceptual, such as the theory of
57 adaptive cycles and its extension to panarchy proposed by Gunderson and Holling (2002).
58 Panarchy analyzes how SES evolve and re-organize at different spatial and temporal scales: a
59 panarchy is defined as a set of nested adaptive cycles organized into a hierarchy, which
60 connects adaptive cycles at small scales to adaptive cycles at large scales (Gunderson and
61 Holling 2002). Other approaches appear to be more appropriate to understand the dynamics of
62 a local SES while encompassing the wide range of the issues and processes at stake in coastal
63 zones, such as the approaches developed in the three frameworks below.

64 First, the framework developed by Ostrom (2009), based on an institutional analysis, is the
65 one that presents the most thorough analysis of the social components of an SES. Secondly,
66 more and more frameworks include ecosystem services (ES) as a boundary object between
67 disciplines (Steger et al. 2018). Considering both the potential of ecosystems to maintain
68 useful ecological functions and the benefits people derive from it, the ES framework is able to
69 describe the complexity of ecosystem-society interactions in an understandable and

70 comprehensive way (Pendleton et al. 2015). Finally, many scientists involved in the
71 management of coastal zones use the Drivers-Pressures-State-Impacts-Responses (DPSIR)
72 analytical framework because it can be an efficient tool to simultaneously organize
73 sophisticated empirical scientific research and transdisciplinary knowledge at an appropriate
74 level to build knowledge on coastal SES (Lewison et al. 2016; Newton et al. 2014). The
75 European Environment Agency (EEA 2003) defines each category of the DPSIR framework,
76 allowing to analyze the causality of interactions. The causal links start with Driving forces,
77 continue through Pressures to the State of the environment and Impacts on ecosystem
78 functions and Human welfare, while Responses may apply to all the other four categories.
79 DPSIR focuses on the factors of change (drivers and pressures) that people are capable of
80 managing.

81 However, these frameworks also have certain shortcomings that need to be overcome. The
82 DPSIR framework is structured around an adaptive management feedback loop designed to
83 address a particular issue, and fails to account for certain dimensions of the SES that are not
84 considered *per se*, such as ecological functions, ecosystem services or institutions. For these
85 reasons, the DPSIR approach should be combined with the ecosystem services and the
86 institutional analysis approaches to be sure to take all the ecological, social and governance
87 issues raised by the SES dynamics into consideration. Ecological and social dynamics are at
88 the core of SES research (Holland 1992; Costanza et al. 1993; Levin 1998; Gunderson &
89 Holling 2002). However, few SES conceptual frameworks have produced an empirical
90 methodology capable of systematically analyzing changes over time and of providing a
91 comprehensive view of the SES trajectory (Binder et al. 2013). Most applications of the SES
92 approach to case studies end with narratives that identify the conditions required to achieve
93 resilience (see for instance Lloyd et al. 2013), but fail to provide clear evidence of changes
94 through quantitative assessments, especially as regards institutions and governance. Several
95 authors recently highlighted the need to use historical or dynamic approaches to study
96 changes in ecosystem services, but few applications are available to date (Lavorel et al. 2015;
97 Renard et al. 2015; Tomscha et al. 2016).

98 In this article, we hypothesize that long-term trajectories of coastal SES, which are exposed to
99 high anthropogenic pressure and environmental changes and are therefore marked by
100 synchronicities between ecological and social evolution, can be split into contrasted periods
101 separated by regime shifts involving both ecological and social dynamics. Indeed, most


102 current SES approaches consider that ecological and social changes are made of
103 synchronicities and regime shifts (Norgaard 1995). However, although regime shifts in
104 ecological sciences are based on well-documented theories (Scheffer and Carpenter 2003) and
105 have been widely studied (Möllemann et al. 2011; Tomczak et al. 2022), their operational
106 definition at the scale of SES is still the subject of debate (Biggs et al., 2018). We use the
107 definition of SES regime shift proposed by Biggs et al. (2018): “large systemic changes that
108 are policy relevant, in that they affect ecosystem services, and where there are established or
109 at least proposed changes in feedbacks that make the systemic changes difficult to reverse”.
110 To test our hypothesis, we developed and applied an operational methodology to understand
111 the trajectory of a local SES and to highlight synchronicities and regime shifts. We argue that
112 an efficient methodology should combine qualitatively and quantitatively oriented approaches
113 to account for the key processes involved in SES dynamics.

114 Several recent operational methods proposed for resilience assessment recommend building
115 the first stages of the study of an SES based on timelines and historical profiling and on
116 interviews with key stakeholders (Resilience Alliance 2020; Enfors-Kautsky et al. 2021;
117 Biggs et al. 2021). The qualitative part of the methodology proposed in the present paper is
118 adapted from the “chronosystemic timeline” approach (Bergeret et al. 2015; Spiegelberger et
119 al. 2018), which consists in an interdisciplinary and holistic analysis of the trajectory of local
120 territories based on literature searches and interviews with stakeholders. The quantitative part
121 of the assessment used time-series analyses based on DPSIR indicators, including ES
122 assessments, as proposed by Santos-Martín et al. (2019) to scrutinize long-term social-
123 ecological dynamics at national scale. We applied our methodology to the Thau lagoon, an
124 SES located on the French Mediterranean coast, and focused our analysis on the adaptation of
125 local societies to changes in the state of the Thau lagoon (South of France) over a 5-decade
126 period (1970-2018).

127 **2. METHODS**

128

129 The original approach we used to design our analytical framework and for the organization
130 and classification of the information available on the SES led us to distinguish several steps
131 corresponding to the qualitative (steps 2a-3a) and quantitative (steps 2b-3b) approaches (Fig.
132 1). The two approaches were combined by matching variables (steps 2a and 2b) and looking
133 for regime shifts (steps 3a and 3b), to provide a synthesis of contrasting periods (step 4).



134

135 **Fig. 1** Methodological process developed to analyze the long-term trajectory of a coastal social-
136 ecological system (SES), combining a chronosystemic timeline approach (steps 2a-3a) and a Drivers-
137 Pressures-State-Impacts-Responses (DPSIR) approach (steps 2b-3b). The two approaches were
138 combined to analyze synchronicities and regime shifts (step 4)

139

140 2.1. Step 1 - Study site: Thau lagoon

141 Thau lagoon is a euhaline coastal lagoon, the largest and deepest in the Gulf of Lions
142 (appendix, Fig. 4). Our study focused on the water body of the Thau lagoon, which covers
143 about 6,790 ha (Fiandrino et al. 2017).

144 The population of the watershed is 103,500 inhabitants, half of whom live in the municipality
145 of Sète (INSEE, 2016). Thau lagoon is the scene of traditional fishing and shellfish farming.
146 The shellfish species reared today are *Mytilus galloprovincialis* (*M. galloprovincialis*)
147 mussels but mainly the Pacific oyster *Crassostrea gigas* (*C. gigas*), whose production is
148 estimated at about 10% of the total oyster production in France (Gangnery et al. 2003).

149 2.2. Qualitative approach: chronosystemic timelines

150 The full implementation of the “chronosystemic timeline” approach presupposes the
151 formation of a multidisciplinary team of researchers (Bergeret et al. 2015; Spiegelberger et al.
152 2018). This work was conducted by a multidisciplinary team of researchers (one ecologist,
153 three social scientists) as recommended for the methodology, but due to the limited number of
154 researchers, we applied a simplified version (Fig. 1, steps 2a-3a).

155 2.2.2. Step 2a - Identification of processes and components

156 Because we focused on the adaptation of local societies to changes in the state of the Thau
157 lagoon, we depicted the trajectory of our coastal SES through four main processes: the state of
158 the ecosystem, human activities, management of the SES and sanitation. Each of these

159 processes is depicted in a thematic narrative timeline (i.e. descriptive chronology) including
160 components of an economic, political, environmental or social nature, with a particular
161 attention paid to institutions and their changes. The following components were considered:

- 162 • **state of the ecosystem:** anoxic crises, appearance of invasive macroalgae, sanitary
163 crises or ban on the sale of shellfish due to microbiological contamination or toxic
164 phytoplankton, chemical or ecological problems that impact shellfish;
- 165 • **human activities:** fishing, shellfish farming, recreational and spa activities, tourism;
- 166 • **management of the SES:** European and national regulations and funds, regional
167 water management schemes, creation of local structures, a local charter and decree,
168 management tools and contracts for the use of the SES;
- 169 • **sanitation:** chemical wastewater treatment plant or lagooning treatment, changes in
170 the treatment system: transformation into lagooning/chemical process, connection to
171 the sanitation network, hydraulic, rehabilitation or resizing works.

172 **2.2.3. Step 3a - Analysis of documents and identification of milestones**

173 Several documents and data sources were used to inform the components and to identify
174 significant events to be reported in the first version of the timelines:

- 175 • gray literature on ecology, shellfish farming and management of coastal SES over the
176 period 1970-2018, representing 418 digitized files;
- 177 • search for scientific gray literature and publications in Web of Science databases;
- 178 • documents in the departmental archives and on regional and local websites;
- 179 • phone and email surveys of associations and administrative services in 2019 and 2020
180 to assess the importance of some tourist and recreational activities.

181 Data were mainly qualitative but quantitative data were also used to inform some components
182 (see appendix, Table 2 for the list of indicators and the data sources).

183 To validate and possibly clarify the interpretations resulting from our document analysis, we
184 interviewed nine organizations representative of the range of local institutions, involved in
185 water management and use and identified as key-actors by previous studies on the Thau
186 lagoon (Bouleau et al. 2009; La Jeunesse et al. 2015) (appendix, Table 3). One person per
187 organization was selected for the interviews, according to its seniority in the territory. Only
188 for the Syndicate of the Thau Lagoon (SMBT), two people were interviewed to represent the

189 broad spectrum of activities carried out by this management structure. In that case,
190 quantitative data were averaged and qualitative information were merged.

191 In a semi-structured interview lasting 90 minutes and transcribed, the stakeholders were asked
192 about their perception of changes in the SES between 1970 and 2018 concerning
193 environmental and socioeconomic problems, human activities and SES management
194 (appendix, Table 4). They were given two blank chronological timelines and asked to position
195 the 'major events' of the SES trajectory in relation to (1) the ecological and sanitary state of
196 the ecosystem, (2) uses and management of the SES. The 'major events' were defined as
197 events that marked the evolution of the characteristics and management policies of the lagoon
198 (environmental and/or socio-economic issues). We listed the major events mentioned by
199 stakeholders in the timelines and also during interviews (questions 2, 7 and 8 in appendix,
200 Table 4). A second version of the four narrative timelines was then produced, highlighting the
201 'milestones' corresponding to the major events cited by more than five stakeholders among
202 the nine interviewed, representing 55.5% of the stakeholders interviewed.

203 **2.3. Quantitative approach: DPSIR framework**

204 We adapted the methodology proposed by Santos-Martín et al. (2019) to analyze time series
205 of the coastal SES based on DPSIR indicators. From a SES perspective, we expanded the
206 scope of DPSIR indicators to include environmental drivers of change (Fig. 1, steps 2b-3b).

207 **2.3.1. Step 2b - Synthetic indicators**

208 To summarize the SES dynamics according to the DPSIR approach, we selected a set of
209 synthetic indicators using the standard evaluation criteria recommended by Tam et al. (2017).
210 Availability and reliability of data was evaluated according to the following criteria: 1) data
211 available for a large part of the study period (1970-2018); 2) data of medium reliability
212 (qualitative data supported by several sources or quantitative data with relatively high
213 uncertainty) or high reliability (quantitative data delimited in space and time). This resulted in
214 the selection of 14 indicators (appendix, Table 2). We selected population growth as the
215 driving force (D) linked to human activities and as for a coastal SES. As climate change
216 cannot be considered as a result of local pressures, rainfall and temperature were included as
217 environmental drivers of change. As waste waters are known to be the most impactful
218 anthropogenic pressure for marine waters (Pedreschi et al. 2019) and especially
219 Mediterranean lagoons (Souchu et al. 2010; Derolez et al. 2019), the proportion of households
220 not connected to the sanitation system was the variable chosen to illustrate the anthropic

221 pressure (P) caused by urban nutrient inputs. Then, two environmental variables illustrating
222 the eutrophication status of the ecosystem and used for Water Framework Directive diagnoses
223 (MTES 2018) were selected as state indicators (S) and corresponding to regulation ecosystem
224 services. Five variables illustrating ecosystem services, corresponding to provisioning and
225 cultural services, were selected as impact indicators (I). The selection of the ES was drawn up
226 in a multidisciplinary manner by scientists specialized in the study area, based on the
227 Common International Classification of Ecosystem Services (Haines-Young and Potschin
228 2018). Finally, three variables illustrating management actions were selected as response
229 indicators (R). The eight variables including missing values (NA) (appendix, Table 2) were
230 interpolated using the algorithm of Stineman (*imputeTS* package in R).

231 **2.3.2. Step 3b - PCA analysis and clustering**

232 Data from 1970 to 2018 on the 12 environmental and social variables selected using the
233 DPSIR approach were analyzed through an explanatory Principal Component Analysis (PCA)
234 (*ade4* package in R). The two environmental drivers of change were considered as illustrative
235 variables and not included in the construction of the PCA space. K-means cluster analysis was
236 then applied to identify years with similar patterns of environmental and social variables
237 (*vegan* package in R). The optimal number of clusters was estimated based on the simple
238 structure index (SSI) (Dolnicar et al. 1999).

239 **2.4. Step 4 - Periodization: synchronicities and regime shifts**

240 Synchronicities were characterized with the help of PCA analyses showing correlations
241 between DPSIR indicators. We considered that the periods were separated by a regime shift,
242 using the definition given by Biggs et al. (2018).

243 To highlight contrasting periods and regime shifts, we compared the results of the two
244 approaches: (i) milestones identified using the chronosystemic timelines approach and (ii)
245 groups of years with similar patterns identified by the DPSIR approach. A regime shift
246 separating contrasting periods corresponds to phases of rupture: (i) highlighted by clustering
247 and identified as a milestone by the stakeholders; and (ii) that led to the reorganization of
248 actor strategies in a context of regulatory renewal and governance mechanisms. Emergent
249 properties of the SES at each period were also identified by both PCA and the narrative
250 timelines. These results were used to build a synthetic timeline to describe the contrasted
251 periods identified along the SES trajectory.

252 **3. RESULTS**

253 The analysis of the synchronicity of factors makes it possible to identify trajectories and to
254 distinguish different contrasting periods in terms of modes of organization on a local scale,
255 and on a wider scale in terms of societal and institutional developments. Several timelines
256 specific to each field were therefore produced (Section 3.1) and then combined to characterize
257 the global trends and the factors of change characteristic of the different periods (Section 3.2).

258 **3.1. CHRONOSYSTEMIC TIMELINE APPROACH USED TO ANALYZE THE** 259 **TRAJECTORY OF COASTAL SES**

260 The trends and milestones found with document analysis and mentioned by stakeholders on
261 the first two processes - (i) the sanitary status of the waters, the shellfish, and the ecological
262 and chemical status of the lagoon; (ii) and human activities - evidenced a succession of
263 environmental crises and a strong vocation for fishing and shellfish farming for the Thau
264 lagoon during the period 1970-2018 (see the timelines in appendix, Fig. 5 and 6).

265 The milestones identified by stakeholders concerning the management measures and the
266 wastewater treatment system (timelines in appendix, Fig. 7 and 8) show a change from: i)
267 sectoral management, hierarchical organization and sanitation works (1970s-1980s), to ii) a
268 structuring of watershed management (1990s) and then to iii) integrated management (from
269 the 2000s).

270


271 **3.2. SYNCHRONICITIES AND REGIME SHIFTS IN THE COASTAL SES** 272 **TRAJECTORY**

273

274 **3.2.2. DPSIR approach**

275 PCA was performed on 12 active environmental and social variables selected for the DPSIR
276 approach, with two illustrative variables corresponding to environmental drivers of change
277 (changes observed between 1970 and 2018 in each variable are available in appendix, Fig. 9).

278 The first two axes of the PCA explained 87.8% of total variability (axis 1 explaining 75.0%
279 and axis 2 explaining 12.8%; the correlation circle of the PCA is available in appendix, Fig.
280 10). The first axis (horizontal) revealed a clear temporal structure, contrasting the earliest and
281 the most recent years (respectively on the left and right in Fig. 2).



282

283 **Fig. 2** Results of PCA of environmental and social variables from 1970 to 2018. Years on the plane
 284 defined by the first and second axes and ellipses representing the 3 groups of years identified by
 285 cluster analysis. The variables are positioned on the plane according to their correlations with axes 1
 286 and 2 (correlation circle is available in appendix, Fig. 10; meaning of the acronyms available in
 287 appendix, Table 2), and colored according to their DPSIR category. The first and second axes account
 288 for 87.8% of total variability

289 This axis, driven by many variables, illustrates the synchronicity of several processes in the
 290 SES trajectory. This axis is negatively correlated mainly with the number of fishers, and to a
 291 lesser extent with the population not connected to the wastewater treatment plant and the
 292 Dissolved Inorganic Phosphorus (DIP) levels in water (on the left in Fig. 2). The first axis is
 293 positively correlated mainly with the number of tourist beds, water recreational and cultural
 294 associations, of management structures and tools and with population (on the right in Fig. 2).
 295 The first axis is correlated to a lesser extent with the number of scientific publications, the
 296 surface area of the lagoon protected by Natura 2000 and with temperature. The second axis
 297 (vertical) is mainly defined by shellfish production (at the top in Fig. 2). Chlorophyll *a*
 298 concentration is correlated with both axis 1 and 2, but to a lesser extent.

299 Among the three clusters identified by the simple structure index index in the PCA, the first
 300 cluster, located on the left in Fig. 2, groups the first 22 years of the study period (1970 to
 301 1991). This first cluster is associated with the variables negatively correlated with the first
 302 axis. On the opposite side, the third cluster groups the last 14 years of the study period (2005

303 to 2018) and is associated with the variables positively correlated with the first axis (on the
 304 right in Fig. 2). In the middle of Fig. 2, the second cluster isolates the period 1992 to 2004,
 305 among which the years 2001 to 2003 were characterized by high levels of shellfish
 306 production.

307 3.2.3. Periodization: synchronicities and regime shifts

308 Among the events listed in the four narrative timelines, nine were identified as milestones by
 309 the majority of stakeholders (more than 5 of 9, Table 1). Five milestones concerned the
 310 ecosystem status of the lagoon, two milestones referred to shellfish farming and two
 311 milestones concerned lagoon management. The events related to sanitation works were not
 312 identified as milestones by the stakeholders even though they contributed to significant
 313 changes in water quality.


314 **Table 1** Summary of the approaches used to identify the regime shifts: i) milestones mentioned by the majority
 315 of stakeholders through the chronosystemic timeline approach (left column), ii) years identified by cluster
 316 analysis in the DPSIR approach (right column). Rows shaded in grey represent the regime shifts considered to
 317 separate the 3 contrasting periods. *The colors represent different components: blue: ecological and sanitary
 318 status, orange: shellfish farming, green: management tools

Chronosystemic timeline approach Milestones mentioned by the majority of stakeholders*	DPSIR approach Years identified with the cluster analysis
1970-1972: reorganization of shellfish farming area	
1989: « <i>Salmonella</i> crisis »	1991-1992
1995: Sea Development Plan (French acronym 'SMVM')	
2003: anoxic crisis	
2004: sanitary downgrading	2004-2005
2005: creation of the Syndicate of the Thau Lagoon (SMBT)	
2006: anoxic crisis	
2008: Oyster spat mortalities (OsHV-1)	
2018: "Green water crisis"	

319
 320 Among these milestones, two periods of environmental crises can be considered as regime
 321 shifts because (i) they led to reorganization of actor strategies in a context of regulatory
 322 renewal and governance mechanisms; (ii) they were identified by the clustering analysis
 323 applied in the DPSIR approach (Table 1). The first regime shift occurred in 1989 with the
 324 "Salmonella crisis", which caused an economic crisis and led to the involvement of all the

325 local actors in the problem of the quality of the water in the lagoon, and to the first
 326 Management Contract in 1990. The sequence of these events constitutes a first cause-and-
 327 effect relationship identified with our analyses. The DPSIR approach also identified a shift
 328 between clusters of years in this period, with clusters 1 and 2 separating in 1991-1992. The
 329 second regime shift occurred in 2004 with sanitary downgrading of the lagoon that took place
 330 between two major anoxic crises, one in summer 2003 and the other in summer 2006. These
 331 environmental crises occurred in a context of changes in regulations related to water
 332 management (Water Framework Directive in 2000) and transformation of the local
 333 institutional organization that favored the creation of the Syndicate of the Thau Lagoon in
 334 2005, a major actor in the management of Thau SES. The creation of this syndicate, dedicated
 335 mainly to the protection of water quality, following the ecological crises of 2003 and 2006
 336 constitutes the second cause-and-effect relationship highlighted by our analyses. The year
 337 2008, which the stakeholders identified as a milestone because of the oyster spat mortalities,
 338 was not identified by PCA clustering. However, as this event in 2008 did not lead to
 339 reorganization of actor strategies or governance, we did not consider it as a regime shift.

340 The two regime shifts identified with our combined approach distinguished three contrasting
 341 periods in the trajectory of Thau SES: period 1 (1970-1989), period 2 (1990-2004) and period
 342 3 (2005-2018). The emerging properties of these periods are highlighted by the two axes of
 343 the PCA and can be interpreted as follows (i) axis 1 discriminates periods 1 and 3 (from left to
 344 right), as a gradient of oligotrophication, change of uses (from the domination of traditional
 345 uses to recreational and diversified uses), development of management tools, demographic
 346 growth and climate warming; ii) axis 2 discriminates period 2 as a gradient of shellfish
 347 production. Fig. 3 is a synthetic timeline of Thau SES that summarizes the regime shifts, the
 348 periods and their emergent properties.



350 **Fig. 3** Synthetic timeline of trajectory of Thau SES from 1970 to 2018, showing the regime shifts that
351 separate 3 contrasting periods with their emergent properties concerning the state of the ecosystem,
352 human activities, management of the SES, sanitation and environmental drivers of change (climate).
353 Milestones identified by the stakeholders are shown in red

354 The main characteristics of the three periods are listed below:

355 • Period 1 (1970-1989)

356 The beginning of this period was marked by the reorganization of shellfish farming that
357 allocated more space to shellfish farming encroaching on areas traditionally dedicated to
358 fishing. Both this event and the parallel disappearance of the Portuguese oyster can be
359 considered as a regime shift in the trajectory of the SES. During this period, the lagoon was
360 the scene of few recreational activities apart from recreational fishing, hydrotherapy,
361 swimming and boating. Management of these activities was essentially sectoral, with
362 considerable intervention by the State as regulator. This period was marked by the installation
363 of the first wastewater treatment plants in the lagoon watershed. The status of the lagoon was
364 seriously damaged by excessive inputs of nutrients - causing a eutrophic state and recurring
365 anoxic crises - and by regular microbiological contamination leading to measures banning the
366 sale of shellfish.

367 • Period 2 (1990-2004)

368 The "*Salmonella* crisis" in 1989 triggered the establishment of partnership management by the
369 stakeholders in the territory and corresponded to the start of a series of Management Contracts
370 for the Thau lagoon. The work to improve the sanitation system resulting from these contracts
371 improved the status of the lagoon in terms of eutrophication and reduced the intensity of
372 anoxic episodes. However, anoxic crises continued and microbiological contamination of
373 shellfish still frequently occurred. The Sea Development Plan (1995) reaffirmed the priority
374 given to traditional activities such as shellfish farming and fishing in the lagoon. This period
375 also saw the development of tourism centered on the lagoon and the emergence of new
376 recreational and leisure uses.

377 • Period 3 (2005-2018)

378 The beginning of the third period was marked by the sanitary downgrading of the lagoon to
379 category B, meaning the shellfish had to be purified before being sold. The creation of
380 Syndicate of the Thau Lagoon in 2005 marked the establishment of an integrated management

381 policy, implemented in collaboration with professional actors, municipalities and State
382 services. The Syndicate of the Thau Lagoon pursued a dual objective of reconciling uses and
383 preserving the good status of the lagoon. The period that then began was characterized by the
384 diversification of uses of the lagoon and significant development of tourism and new
385 recreational activities. This period was also marked by weakening of shellfish farming activity
386 from 2008 (mortality of OsHV spat) and a decline in fishing. Despite the ongoing
387 oligotrophication of the lagoon, anoxic crises occurred during summer heatwaves (2003,
388 2006, 2018), and the “Green water crisis” in 2018, a sign of the growing pressure of global
389 warming on the Thau lagoon. This unprecedented dystrophic crisis can be seen as a new
390 regime shift in the trajectory of Thau SES.

391

392 **4. DISCUSSION**

393 **4.1. Summary of the coastal SES trajectory**

394 The methodology we used to analyze the long-term trajectory of a coastal SES enabled us to
395 show a succession of sanitary or ecological crises and management adaptations. Similar
396 successions of crises leading to adaptations of local management methods have been
397 highlighted in other estuarine systems in Europe and in Pacific regions (Daniell et al. 2020).
398 Examples of coastal SES regime shifts in the literature are generally associated with
399 degradations (de Jonge et al. 2002; Biggs et al. 2021), but few examples are available to
400 illustrate the recovery or coastal lagoon ecosystems (Derolez et al. 2019), and even less at the
401 SES scale (De Wit et al. 2020; Erostate et al. 2022). Our dynamic study over a 5-decade
402 period showed that the SES evolved towards the improvement of water quality, multiplication
403 of uses and more integrative management. Our results revealed a shift in a system originally
404 dedicated to food production towards a system with diversified uses, with a particular focus
405 on the development of tourism and recreational activities. This trend reflects the growing
406 attraction of the coast for leisure activities (Corlay 1995). The three contrasting periods we
407 identified in this study of the trajectory of the coastal SES are almost identical (within one or
408 two years) to those identified in a study based only on the ecological changes (Derolez et al.
409 2020). This study showed successive periods in ecosystem functioning during the
410 oligotrophication process: a eutrophic period (1970-1992), a transition phase (1993-2003) and
411 a regime shift (2003-2006) followed by a period towards oligotrophy (2006-2018). The
412 similarity of the periods could be explained by the low inertia of estuaries and coastal lagoons
413 in terms of ecological functioning in response to reduced nutrient loads following remediation

414 measures (Boynnton et al. 2013; Lie et al. 2021). In parallel, the social system is very
415 responsive to water and shellfish quality issues, which play a major role in mobilizing local
416 actors when they are threatened (Barone 2016). However, the management measures we
417 identified in our study were driven by more general issues, given that the Thau basin was
418 selected as a national pilot site for the implementation of the Integrated Coastal Zone
419 Management (GESAMP 1996). Likewise, the changes in management methods should be
420 considered in the broader context of decentralization in France in the 1980s, and their
421 promotion by European Union recommendations and French laws in the 1990s designed to
422 promote sustainable development (French Law 1999; Parliamentary Assembly 1999). The
423 principles of « New public management » were reinforced in France in the 2000s, and
424 decentralization was reinforced to the benefit of local authorities.

425 Despite improvements in water quality and the development of an integrated management
426 policy in the most recent period, several problems still threaten the stability of the SES. The
427 recent ecological crises associated with the increasing pressure of global warming (Lagarde et
428 al. 2021) greatly weakened shellfish farmers and point to a new regime shift in the trajectory
429 of the SES. Shellfish diseases continue to affect production (Pernet et al. 2018) and fishing is
430 also declining. Finally, the average age of today's fishers and shellfish farmers is going up, so
431 these traditional uses do not suggest stability or future growth. However, for those involved in
432 water policy, the strict water quality standards required by the traditional fishing and shellfish
433 farming activities, who readily present themselves as "sentinels of the coastal environment",
434 can be considered as a driving force for the territory towards exemplarity and innovation in
435 scientific activities and environmental management (Barone 2016; Chesbrough 2003; Daudin
436 et al. 2021).

437 **4.2. Perspectives for the dynamic analysis of the SES**

438 The original and operational methodology used to depict the trajectory of a coastal SES
439 proposed in this paper, which combines qualitative and quantitative information, proved to be
440 particularly appropriate to reveal the dynamics of coevolution of the social and ecological
441 components of a complex coastal system along with regime shifts. As recommended by
442 Bergeret et al. (2015) and Spiegelberger et al. (2018), the trajectory of a SES must be studied
443 using a holistic approach, calling on interdisciplinary expertise and using timelines as
444 boundary objects between the natural sciences and social sciences. This holistic approach is in
445 accordance with the theories developed by researchers studying complex systems (Holland

446 1992; Levin 1998). Moreover, our study enabled the identification of regime shifts, in
447 accordance with the concepts developed by several authors in the system analysis perspective
448 (Scheffer and Carpenter 2003; Biggs et al. 2018) and illustrated in several terrestrial or
449 aquatic SES (Kermagoret et al. 2019; Lavorel et al. 2015). Other recent studies analyzed the
450 long-term dynamics of SES from the same theoretical perspective, to name a few examples: a
451 study of the management of the Everglades wetlands (Florida, USA) (Gunderson et al. 2002),
452 the trajectory of Lake Taihu (China) (Lin et al. 2019) or of a Corsican lagoon (Erostate et al.
453 2022).

454 The holistic and combined approach we developed allowed us to identify several cause-effect
455 relationships and stable emerging properties characterizing contrasted periods, but needs to be
456 completed by more detailed analyses of causal links and to better account for the complexity
457 of systems (Costanza et al. 1993). For example, more precise characterization of the
458 ecological and social determinants of the stability or instability of the SES is required to fully
459 understand its dynamics. To this end, our study could be supplemented by a study of historical
460 changes in ecosystem services (ES) as recommended by several authors (Renard et al. 2015;
461 Tomscha et al. 2016). Other authors have analyzed the past trajectory of a mountain SES
462 through changes in nature's contributions to people and resulting adaptations (Bruley et al.
463 2021). A dynamic approach is crucial to explore the threshold effects and possible
464 irreversibilities and to identify the synergies and antagonisms between ES that require trade-
465 offs or compromises (Renard et al. 2015; Tomscha et al. 2016). In addition, the distinction
466 between demands, flows and potentials of ES is particularly useful to assess the sustainability
467 of uses and thus to support environmental management (Burkhard et al. 2014; Villamagna et
468 al. 2013).

469
470 Our study of historical trajectories will also be useful to identify the conditions that trigger
471 crises or favor the ability of SES to adapt in the long term. In the context of global change and
472 faced with the multiplicity of uses and the complexity of the issues, stakeholders need more
473 knowledge and management scenarios to increase resilience and guarantee the sustainability
474 of SES activities (Yorque et al. 2002; van Staveren and van Tatenhove 2016). Such models
475 have already been applied to Thau lagoon to test the influence of different nutrient inputs and
476 shellfish stock scenarios on oyster production, carrying capacity, and Water Framework
477 Directive ecological status (Pete et al. 2020). However, the lack of reliable data on fish and

478 shellfish production is an obstacle to the calibration of such models and the necessary
479 management tools need to be co-developed by scientists and stakeholders (Byron et al. 2011).
480 More knowledge on how local users, inhabitants, managers perceive ecosystem and how it is
481 changing is needed to better define restoration goals and to account for the notion of the
482 “desired state” of the environment, whose meaning may differ among local actors and is
483 sometimes far removed from ecological standards (de Wit et al. 2020; Baker and Eckerberg
484 2016). Some authors designed a participatory diagnosis approach for fisheries and
485 demonstrated its potential to guide contextually informed improvements to management in
486 practice, including transitions to contemporary forms of governance (Eriksson et al. 2016).
487 Finally, the results we obtained using the empirical analysis of the long-term trajectory of a
488 coastal SES highlighted the key role of episodic disturbances in triggering reorganization of
489 governance. These results should now be reanalyzed using the concept of ecological resilience
490 to provide the critical insights needed to identify the role of social dynamics during
491 alternating periods of stability and instability in SES. Such an approach could help identify
492 the structures and processes best suited to shift governance towards sufficiently flexible forms
493 to meet the challenge of mitigating the impacts of global climate change (Chaffin and
494 Gunderson 2016).

495 **5. CONCLUSION**

496

497 We propose an innovative analytical framework to study the long-term trajectory of coastal
498 social-ecological systems. Our methodology, which combines qualitative and quantitative
499 information, was applied to the Thau lagoon case study and revealed co-evolution and regime
500 shifts caused by a succession of sanitary or ecological crises and highlighted the key role of
501 episodic disturbances in triggering reorganization of governance. The contrasting periods
502 shown in the 5-decade trajectory of the SES were almost identical to those based on
503 ecological changes alone. This confirmed the importance of water quality and of the
504 traditional uses of shellfish farming as well as their role in mobilizing local actors when
505 threatened.

506 Our method provides the first clues to emergent properties that may favor resilience at the
507 coastal SES scale and guide future management options to protect ecosystem functions and
508 services in times of global changes. The analysis of past changes makes it possible to
509 characterize the processes at the origin of the observed adaptations. These result from
510 interactions between spontaneous responses, both ecological and socio-economic, and the

511 responses of regulatory systems, which are the responsibility of local governance, but also of
512 multi-level decisions due to the interweaving of governance systems. The synchronicities
513 observed lead us to advocate the reactivity of regulatory systems, but also the evolution of
514 paradigms according to the periods. Moreover, this type of analytical framework, conducted
515 in collaboration with the stakeholders, illustrates the interest of close collaboration between
516 scientists and managers to better understand the complexity of interactions. The aim is to
517 better understand the interactions between the various components, but also between cyclical
518 and structural determinants. This understanding of past trajectories facilitates the analysis of
519 the conditions of adaptation of the SES and thus its long-term resilience, which must be
520 dynamic and adaptive (Sgrò et al. 2010; Davoudi et al. 2012).

521 **Acknowledgments:** This study was carried out in the framework of the project
522 RETROSCOPE funded by Ifremer. The authors would like to thank all RETROSCOPE
523 project members; Syla Benidiri for her contribution to the survey of stakeholders; Jean-Louis
524 Guillou (Ifremer); Ludovic Cesmat, Béatrice Pary, Romain Pete and Stéphane Roumeau
525 (SMBT); Anaïs Giraud and Anahi Barrera (AERMC); Matthew Hebert and Cécile Mablouké
526 (Cépralmar); Denis Régler and Adeline Perrignon (CRCM); Fabrice Jean (Prud'homie de
527 Mèze); Emmanuel Liberti (CRPMEM); Emilie Varraud and Florian Martel (CPIE); Frédéric
528 Le Gal (DML34).

529 LITERATURE CITED

- 530
- 531 Baker S, Eckerberg K (2016) Ecological restoration success: a policy analysis understanding. *Restor Ecol*
532 24:284-290. [doi:10.1111/rec.12339](https://doi.org/10.1111/rec.12339)
- 533 Barone S (2016) Building a narrative on environmental policy success. Reflections from a watershed
534 management experience. *Crit Policy Stud* 12:135-148. [doi:10.1080/19460171.2016.1194765](https://doi.org/10.1080/19460171.2016.1194765)
- 535 Bergeret A, George-Marcelpoil E, Delannoy J, Piazza-Morel D (2015) L'outil-frise : une expérimentation
536 interdisciplinaire. Comment représenter des processus de changements en territoires de montagne ?
- 537 Berkes F, Folke C (Eds.) (1998) Linking social and ecological systems: management practices and social
538 mechanisms for building resilience. Cambridge University Press.
- 539 Biggs R, Peterson GD, Rocha JC (2018) The regime shifts database: A framework for analyzing regime shifts in
540 social-ecological systems. *Ecology and Society* 23. [10.5751/ES-10264-230309](https://doi.org/10.5751/ES-10264-230309)
- 541 Biggs R, de Vos A, Preiser R, Clements H, Maciejewski K et al (2021) The routledge handbook of research
542 methods for social-ecological systems. [10.4324/9781003021339](https://doi.org/10.4324/9781003021339)

- 543 Binder CR, Hinkel J, Bots PW, Pahl-Wostl C (2013) Comparison of frameworks for analyzing social-ecological
544 systems. *Ecology and Society* 18(4). doi:10.5751/ES-05551-180426
- 545 Boynton WR, Hodgkins CLS, O'Leary CA, Bailey EM, Bayard AR, Wainger LA (2013) Multi-decade
546 Responses of a Tidal Creek System to Nutrient Load Reductions: Mattawoman Creek, Maryland USA.
547 *Estuaries and Coasts* 37, 111-127. doi:10.1007/s12237-013-9690-4
- 548 Bruley E, Locatelli B, Vendel F, Bergeret A, Elleaume N et al (2021) Historical reconfigurations of a social-
549 ecological system adapting to economic, policy and climate changes in the French Alps. *Reg Environ*
550 *Change* 21, 34. <https://doi.org/10.1007/s10113-021-01760-8>
- 551 Burkhard B, Kandziora M, Hou Y, Müller F (2014) Ecosystem service potentials, flows and demands-concepts
552 for spatial localisation, indication and quantification. *Landsc Online* 34:1-32. doi:10.3097/LO.201434
- 553 Byron C, Bengtson D, Costa-Pierce B, Calanni J (2011) Integrating science into management: Ecological
554 carrying capacity of bivalve shellfish aquaculture. *Mar Policy* 35:63-370. doi:10.1016/j.marpol.2010.10.016
- 555 Chaffin BC, Gunderson LH (2016) Emergence, institutionalization and renewal: Rhythms of adaptive
556 governance in complex social-ecological systems. *Journal of Environmental Management* 165:81-87.
557 doi:10.1016/j.jenvman.2015.09.003
- 558 Chesbrough H (2003) *Open innovation: the new imperative for creating and profiting from technology*. Boston,
559 MA: Harvard Business.
- 560 Corlay JP (1995) Géographie sociale, géographie du littoral. *Norois* 165:247-265.
561 doi.org/10.3406/noroi.1995.6623
- 562 Costanza R, Wainger L, Folke C, Mäler K (1993) Modeling Complex Ecological Economic Systems. Towards
563 an evolutionary, dynamic understanding of people and nature. *Bioscience* 43:545-555
- 564 Daniell KA, Plant R, Pilbeam V, Sabinot C, Paget N et al (2020) Evolutions in estuary governance? Reflections
565 and lessons from Australia, France and New Caledonia. *Marine Policy* 112.
566 doi.org/10.1016/j.marpol.2019.103704
- 567 Daudin K, Weber C, Colin F, Cernesson F, Maurel P et al (2021) The Collaborative Process in Environmental
568 Projects, a Place-Based Coevolution Perspective. *Sustainability* 13(15), 8526. doi:10.3390/su13158526
- 569 Davoudi S, Shaw K, Haider LJ, Quinlan AE, Peterson GD, Wilkinson C, Fünfgeld H, McEvoy D, Porter L
570 (2012) Resilience: a bridging concept or a dead end? Reframing resilience: challenges for planning theory
571 and practice. *Planning theory and practice* 13, 299-333. doi:10.1080/14649357.2012.677124
- 572 De Wit R, Leruste A, Le Fur I, Sy MM, Bec B et al. (2020). A multidisciplinary approach for restoration
573 ecology of shallow coastal lagoons, a case study in South France. *Front Ecol Evol - Conserv*
574 doi:10.3389/fevo.2020.00108

- 575 Derolez V, Bec B, Munaron D, Fiandrino A, Pete R et al (2019). Recovery trajectories following the reduction
576 of urban nutrient inputs along the eutrophication gradient in French Mediterranean lagoons. *Ocean Coast*
577 *Manag* 171:1-10. doi:10.1016/j.ocecoaman.2019.01.012
- 578 Derolez V, Malet N, Fiandrino A, Lagarde F, Richard M et al (2020). Fifty years of ecological changes: regime
579 shifts and drivers in a coastal Mediterranean lagoon during oligotrophication. *Science of the Total*
580 *Environment* 732. doi.org/10.1016/j.scitotenv.2020.139292
- 581 Dolnicar S, Grabler K, Mazanec JA (1999) A tale of three cities: perceptual charting for analyzing destination
582 images. In: Woodside A. et al (ed) *Consumer psychology of tourism, hospitality and leisure*, CAB
583 International, New York, pp 39-62
- 584 EEA (2003) *Environmental Indicators: Typology and Use in Reporting*. European Environment Agency
- 585 Enfors-Kautsky E, Järnberg L, Quinlan AL, Ryan P (2021) Wayfinder: a new generation of resilience practice.
586 *Ecology and Society* 26(2):39. <https://doi.org/10.5751/ES-12176-260239>
- 587 Eriksson H, Adhuri DS, Adrianto L, Andrew NL, Apriliani T et al (2016) An ecosystem approach to small-scale
588 fisheries through participatory diagnosis in four tropical countries. *Glob Environ Chang* 36:56-66.
589 [doi:10.1016/j.gloenvcha.2015.11.005](https://doi.org/10.1016/j.gloenvcha.2015.11.005)
- 590 Erostate M, Ghiotti S, Huneau F, Jouffroy D, Garel E et al. (2022) The challenge of assessing the proper
591 functioning conditions of coastal lagoons to improve their future management. *Science of the Total*
592 *Environment* 803:150052. [doi:10.1016/j.scitotenv.2021.150052](https://doi.org/10.1016/j.scitotenv.2021.150052)
- 593 Fiandrino A, Ouisse V, Dumas F, Lagarde F, Pete R et al (2017). Spatial patterns in coastal lagoons related to
594 the hydrodynamics of seawater intrusion. *Mar Pollut Bull* 119:132-144.
595 doi:10.1016/j.marpolbul.2017.03.006
- 596 Folke C (2016). Resilience (republished). *Ecology and Society*, 21(4). [doi:10.5751/ES-09088-210444](https://doi.org/10.5751/ES-09088-210444)
- 597 French Law (1999) Loi n° 99-533 du 25 juin 1999 d'orientation pour l'aménagement et le développement
598 durable du territoire et portant modification de la loi n° 95-115 du 4 février 1995 d'orientation pour
599 l'aménagement et le développement du territoire.
600 <https://www.legifrance.gouv.fr/loda/id/LEGITEXT000005628113/>. Accessed 24 August 2022
- 601 Gangnery A, Chabirand JM, Lagarde F, Le Gall P, Oheix J et al (2003) Growth model of the Pacific oyster,
602 *Crassostrea gigas*, cultured in Thau Lagoon (Méditerranée, France). *Aquaculture* 215:267-290.
603 doi:10.1016/S0044-8486(02)00351-4
- 604 Garmestani A, Craig RK, Gilissen HK, McDonald J, Soininen N et al (2019) The role of social-ecological
605 resilience in coastal zone management: A comparative law approach to three coastal nations. *Frontiers in*
606 *ecology and evolution*, 7(410). [doi:10.3389/fevo.2019.00410](https://doi.org/10.3389/fevo.2019.00410)

- 607 GESAMP (1996) The contributions of science to coastal zone management. Reports and studies GESAMP, 61.
608 Food and Agricultural Organization of the United Nations
- 609 Gunderson LH, Holling CS (2002) Panarchy. Understanding transformations in human and natural systems.
610 Island Press
- 611 Halpern BS, Walbridge SKA, Selkoe CV, Kappel F, Micheli C et al (2008) A global map of human impacts on
612 marine ecosystems. *Science* 319:948-952
- 613 Haines-Young R, Potschin M (2018) Common International Classification of Ecosystem Services (CICES)
614 V5.1. Guidance on the Application of the Revised Structure.
615 <https://cices.eu/content/uploads/sites/8/2018/01/Guidance-V51-01012018.pdf>
- 616 Holland JH (1992) Complex adaptative systems. *Daedalus* 121:17-30
- 617 INSEE (2016) National institute of statistics and economic studies. Population in year.
618 <https://www.insee.fr/en/accueil>. Accessed 24 August 2022
- 619 Kermagoret C, Claudet J, Derolez V, Nugues MM, Ouisse V et al (2019) How does eutrophication impact
620 bundles of ecosystem services in multiple coastal habitats using state-and-transition models. *Ocean Coast*
621 *Manag* 174:144-153. doi:10.1016/j.ocecoaman.2019.03.028
- 622 La Jeunesse I, Cirelli C, Sellami H, Aubin D, Deidda R et al (2015). Is the governance of the Thau coastal
623 lagoon ready to face climate change impacts? *Ocean Coast Manag* 1-13.
624 doi:10.1016/j.ocecoaman.2015.05.014
- 625 Lagarde F, Atteia A, Gobet A, Richard M, Mostajir B et al (2021). Phénomène d'Eaux Vertes à *Picochlorum* en
626 lagune de Thau pendant les années 2018 et 2019, Observations environnementales.
627 <https://archimer.ifremer.fr/doc/00689/80087/>
- 628 Lavorel S, Colloff MJ, McIntyre S, Doherty MD, Murphy HT et al (2015) Ecological mechanisms underpinning
629 climate adaptation services. *Glob Chang Biol* 21-12-31. doi:10.1111/gcb.12689
- 630 Levin SA (1998) Ecosystems and the biosphere as complex adaptative systems. *Ecosystems* 1:431-436
- 631 Lewison RL, Rudd MA, Al-Hayek W, Baldwin C, Beger M et al (2016) How the DPSIR framework can be
632 used for structuring problems and facilitating empirical research in coastal systems. *Environmental Science*
633 *& Policy* 56-110-119. doi:10.1016/j.envsci.2015.11.0011462-9011/
- 634 Lie AAY, Wong CK, Lam JYC, Liu JH, Yung YK (2011) Changes in the nutrient ratios and phytoplankton
635 community after declines in nutrient concentrations in a semi-enclosed bay in Hong Kong. *Mar. Environ.*
636 *Res.* 71, 178-188. doi:10.1016/j.marenvres.2011.01.001
- 637 Lin Q, Zhang K, Shen J, Liu E (201) Integrating long-term dynamics of ecosystem services into restoration and
638 management of large shallow lakes. *Sci Total Environ* 671:66-75. doi:10.1016/j.scitotenv.2019.03.307

- 639 Lloyd MG, Peel D, Duck RW (2013) Towards a social–ecological resilience framework for coastal planning.
640 Land Use Policy 30(1):925-933. [doi:10.1016/j.landusepol.2012.06.012](https://doi.org/10.1016/j.landusepol.2012.06.012)
- 641 MTES (Ministère de la Transition Ecologique et Solidaire) (2018) Arrêté du 27 juillet 2018 modifiant l'arrêté
642 du 25 janvier 2010 relatif aux méthodes et critères d'évaluation de l'état écologique, de l'état chimique et du
643 potentiel écologique des eaux de surface pris en application des articles R. 212-10, R. 212-11 et R. 212-18 du
644 code de l'environnement.
- 645 Möllmann C, Conversi A, Edwards M (2011) Comparative analysis of European wide marine ecosystem shifts:
646 A large-scale approach for developing the basis for ecosystem-based management. *Biology Letters* 7:484-
647 486. [10.1098/rsbl.2010.1213](https://doi.org/10.1098/rsbl.2010.1213)
- 648 Newton A, Icely J, Cristina S, Brito A, Cardoso AC et al (2014) An overview of ecological status, vulnerability
649 and future perspectives of European large shallow, semi-enclosed coastal systems, lagoons and transitional
650 waters. *Estuar Coast Shelf Sci* 140:95-122. [doi:10.1016/j.ecss.2013.05.023](https://doi.org/10.1016/j.ecss.2013.05.023)
- 651 Norgaard RB (1995) Beyond Materialism: A Coevolutionary Reinterpretation of the Environmental Crisis. *Rev*
652 *Soc Econ* LIII:475-492
- 653 Ostrom E (2009) A general framework for analyzing sustainability of social-ecological systems. *Science*
654 325(5939):419-422. [doi:10.1126/science.1172133](https://doi.org/10.1126/science.1172133)
- 655 Parliamentary Assembly (1999) Recommendation 1431. Future action to be taken by the Council of Europe in
656 the field of environment protection. Report of the Committee on the Environment, Regional Planning and
657 Local Authorities. <https://assembly.coe.int/nw/xml/XRef/Xref-XML2HTML-en.asp?fileid=16731&lang=en>.
658 Accessed 24 August 2022
- 659 Pedreschi D, Bouch P, Moriarty M, Nixon E, Knights AM et al (2019) Integrated ecosystem analysis in Irish
660 waters; Providing the context for ecosystem-based fisheries management. *Fisheries Research* 209:218-229.
661 <https://doi.org/10.1016/j.fishres.2018.09.023>
- 662 Pendleton L, Mongrue R, Beaumont N, Hooper T, Charles M (2015) A triage approach to improve the
663 relevance of marine ecosystem services assessments. *Mar Ecol Prog Ser* 530:183-193.
664 [doi:10.3354/meps11111](https://doi.org/10.3354/meps11111)
- 665 Pernet F, Fuhrmann M, Petton B, Mazurié J, Bouget JF et al (2018) Determination of risk factors for herpesvirus
666 outbreak in oysters using a broad-scale spatial epidemiology framework. *Sci Rep* 8:1-11.
667 [doi:10.1038/s41598-018-29238-4](https://doi.org/10.1038/s41598-018-29238-4)
- 668 Pete R, Guyondet T, Bec B, Derolez V, Cesmat L et al (2020) A box-model of carrying capacity of the Thau
669 lagoon in the context of ecological status regulations and sustainable shellfish cultures. *Ecol Modell*
670 [doi:10.1016/j.ecolmodel.2020.109049](https://doi.org/10.1016/j.ecolmodel.2020.109049)
- 671 Renard D, Rhemtulla JR, Bennett EM (2015) Historical dynamics in ecosystem service bundles. *PNAS*
672 112:13411-13416. [doi:10.1073/pnas.1502565112](https://doi.org/10.1073/pnas.1502565112)

- 673 Resilience Alliance (2020) Assessing Resilience in Social-Ecological Systems: Workbook for Practitioners.
674 Version 2.0. <http://www.resalliance.org/3871.php>
- 675 Santos-Martín F, García-Mon BG, González JA, Iniesta-Arandia I, García-Llorente M et al (2019) Identifying
676 past social-ecological thresholds to understand long-term temporal dynamics in Spain. *Ecol Soc* 24.
677 doi:10.5751/ES-10734-240210
- 678 Scheffer M, Carpenter SR (2003) Catastrophic regime shifts in ecosystems: linking theory to observation.
679 *Trends Ecol Evol* 18:648-656. doi:10.1016/j.tree.2003.09.002
- 680 Sgrò CM, Lowe AJ, Hoffmann AA (2010) Building evolutionary resilience for conserving biodiversity under
681 climate change. *Evolutionary Applications* 4, 326-337. doi:10.1111/j.1752-4571.2010.00157.x
- 682 Souchu P, Bec B, Smith VH, Laugier T, Fiandrino A et al (2010) Patterns in nutrient limitation and chlorophyll
683 *a* along an anthropogenic eutrophication gradient in French Mediterranean coastal lagoons. *Canadian Journal*
684 *of Fisheries and Aquatic Sciences* 67(4):743-753. doi:10.1139/F10-018
- 685 Spiegelberger T, Bergeret A, Crouzat E, Tschanz L, Piazza-Morel D et al (2018) Interdisciplinary construction
686 of a socio-ecological vulnerability trajectory based on the Quatre Montagnes (Isère, France) area from 1950
687 to 2016. doi:10.4000/rga.5046
- 688 Steger C, Hirsch S, Evers C, Branoff B, Petrova M et al (2018) Ecosystem services as boundary objects for
689 transdisciplinary collaboration. *Ecological economics* 143:153-160
- 690 Tam JC, Link JS, Rossberg AG, Rogers SI, Levin PS et al (2017) Towards ecosystem-based management:
691 identifying operational food-web indicators for marine ecosystems, *ICES Journal of Marine Science*
692 74(7):2040-2052. <https://doi.org/10.1093/icesjms/fsw230>
- 693 Tomczak MT, Müller-Karulis B, Blenckner T, Ehrnsten E, Eero M et al (2022) Reference state, structure,
694 regime shifts, and regulatory drivers in a coastal sea over the last century: The Central Baltic Sea case.
695 *Limnology and Oceanography* 67:S266-S284. 10.1002/lno.11975
- 696 Tomscha ST, Sutherland IJ, Renard D, Gergel SE, Rhemtulla JM et al (2016) Guide to Historical Data Sets for
697 Reconstructing Ecosystem Service Change over Time. *Bioscience* 66:747-762
- 698 Tscherning K, Helming K, Krippner B, Sieber S, y Paloma SG (2012) Does research applying the DPSIR
699 framework support decision making? *Land use policy* 29(1):102-110
- 700 van Staveren MF, van Tatenhove JPM (2016) Hydraulic engineering in the social-ecological delta:
701 understanding the interplay between social, ecological, and technological systems in the Dutch delta by
702 means of “delta trajectories”. *Ecology and Society* 21(1):8. doi:10.5751/ES-08168-210108
- 703 Villamagna AM, Angermeier PL, Bennett EM (2013) Capacity, pressure, demand, and flow: A conceptual
704 framework for analyzing ecosystem service provision and delivery. *Ecol Complex* 15:114-121.
705 doi:10.1016/j.ecocom.2013.07.004

This version of the article has been accepted for publication, after peer review but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s10113-023-02061-y>. Use of this Accepted Version is subject to the publisher's Accepted Manuscript terms of use <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

706 Yorque R, Walker B, Holling CS, Gunderson LH, Folke C et al (2002) Toward an Integrative Synthesis. In:
707 Gunderson LH, Holling CS (ed) Panarchy. Understanding transformations in human and natural systems,
708 Island Press, pp 419-438

709

This version of the article has been accepted for publication, after peer review but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s10113-023-02061-y>. Use of this Accepted Version is subject to the publisher's Accepted Manuscript terms of use <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

711 Supplementary materials

712 **Table 2** List of the 12 variables (and 2 environmental drivers of change) selected to analyze the SES trajectory according to the DPSIR approach. The process addressed, using the chronosystemic
 713 timeline approach, is listed in the second column: state of the ecosystem, human activities, management of the SES, sanitation. The category and number of ecosystem services according to CICES
 714 typology are given for impact indicators. The period covered by the data is given in column 5. Column 6 lists the data sources and the variables for which interpolation was applied to replace the
 715 missing data (*). The last column indicates the level of data reliability

Category in the DPSIR approach	Process addressed using the chronosystemic timeline approach	Service category (CICES Class)	Indicator (unit) "Name on the PCA"	Data availability period	Data sources ⁽¹⁾ (details on the data) The symbol (*) indicates that interpolation was used to replace NA	Data reliability: (1) low, (2) moderate, (3) high
Driver	Environmental driver of change	/	Annual cumulative rainfall (mm) "Rainfall"	1970-2018	Sète Météo-France weather station (n°34301002)	3
	Environmental driver of change	/	Annual mean air temperature (°C) "Temperature"	1970-2018	Sète Météo-France weather station (n°34301002)	3
	Human activities	/	Watershed population (inhabitants) "Population"	1975-2018	(*) La Jeunesse 2001; INSEE 2006-2015, 2018 (14 cities in the watershed: Balaruc-les-Bains, Balaruc-le-Vieux, Bouzigues, Gigean, Loupian, Marseillan, Mèze, Montagnac, Montbazin, Pinet, Pomerols, Poussan, Sète, Villeveyrac)	3
Pressure	Sanitation	/	Watershed population not connected to the sanitation system and discharging into the lagoon "Population_no_WWTP"	1975-2015	(*) La Jeunesse (2001); INSEE (2006-2015, 2018); Couton et al (2007) (14 cities in the watershed)	3
State	State of the ecosystem	Regulation (2.2.5.2)	DIP concentration in water (summer median, µM) "DIP_summer"	1972-2018	(*) Souchu et al (1998); Derolez et al (2020a)	3
	State of the ecosystem	Regulation (2.2.4.2)	Chlorophyll <i>a</i> concentration in water (annual median, µg L ⁻¹) "CHLA_annual"	1976-2017	(*) Vaumat and Frisoni (1986); Péna (1989); Collos et al (2009); Derolez et al (2020a)	2
Impact	Human activities	Provisioning (1.1.4.1)	Total shellfish production (t.y-1) "Shellfish_prod"	1970-2018	(*) Affaires Maritimes de Sète (1970-1989); Comps (2000); DML (2011-2019)	3
	Human activities	Provisioning (1.1.6.1)	Number of active fishers "N° Fishers"	1973-2018	(*) Mazouni et al (1999); Cejpa (1985); Les Ecologistes de l'Euzière (1988); Affaires Maritimes de Sète (2004, 2005); Cépralmar (2006); SMBT (2010, 2014); Chaboud et al (2015); SIH (2018, 2019).	2
	Human activities	Cultural (3.1.1.1)	Number of associations linked to water recreational activities (sailing, diving, kitesurf, water skiing, rowing) "Water_recreat_asso"	1970-2018	Survey on the web (French Yellow Pages) and phone or e-mail contacts by N. Lautrédou-Audouy (associations with headquarters located in the Thau watershed)	2
	Human activities	Cultural (3.1.1.2)	Number of tourist beds "Tourist_beds"	1975-2016	(*) La Jeunesse (2001); SMBT (2010); Pagès (2003); Rey-Valette (2006); data SMBT 2016 (comm. pers. L.Cesmat) (14 cities of the watershed)	2
	Human activities	Cultural (3.1.2.3)	Number of associations for the protection and enhancement of cultural and linguistic heritage "Cultural_asso"	1970-2018	(*) Répertoire National des Associations de l'Hérault (1970-2006, 2018): Search of city websites (data available only for Sète and Balaruc-les-Bains)	2
Response	Management of the SES	/	Number of publications citing the Thau lagoon "Scientific_pub"	1975-2018	Search of Web of Science (search terms = "Thau lagoon" or "étang de Thau")	3
	Management of the SES	/	Part of the lagoon protected as a Natural Area of Ecological Floristic and Fauna Interest under Natura 2000 (ha) "N2000_surf"	1970-2018	SMBT 2010	3
	Management of the SES	/	Number of management tools & structures "Management"	1970-2018	Gray literature ⁽¹⁾	3

716 **(1) Data sources for step 2b (gray and local literature):**

- 717 Affaires Maritimes de Sète (1970 to 2005) Statistiques des Affaires Maritimes. Monographies conchylicoles
- 718 Andral B, Sargian P (2010) Directive Cadre Eau. District "Rhône et Côtiers Méditerranéens". Contrôles de surveillance/opérationnel
- 719 Bec B, Derolez V, Soudant D, Cesmat L, Pete R et al (2018) Projet CAPATHAU : Capacité Trophique de la lagune de Thau. Evolution temporelle
720 de l'état écologique de la lagune de Thau et des performances des coquillages en élevage au regard de la réduction des apports issus du bassin
721 versant et des changements météo
- 722 Bouchoucha M, Derolez V, Munaron D, Gonzalez JL, Cimiterra N et al (2019) Directive Cadre sur l'Eau Bassin Rhône Méditerranée Corse -
723 Année 2018
- 724 Cejpa AM (1985) Les métiers de la pêche pratiqués sur l'étang de Thau - Premiers éléments d'informations. Rapport Ifremer-Cépralmar
- 725 Cépralmar (2006) Prud'homies du Languedoc-Roussillon. Données de suivi de la pêche aux petits métiers
- 726 Cesmat L (2018) Note d'identification des diverses crises et évaluation de la vulnérabilité de l'activité conchylicole sur la lagune de Thau
- 727 Chaboud C, Cazalet B, Reyes N, Rubin A (2015) Enquête sur les petits métiers de la pêche en Languedoc-Roussillon. Programme POLYPECHE
- 728 Collos, Y., Bec, B., Jauzein, C., Abadie, E., Laugier, T., Lautier, J., Pastoureaud, A., Souchu, P., Vaquer, A., 2009. Oligotrophication and
729 emergence of picocyanobacteria and a toxic dinoflagellate in Thau lagoon, southern France Regional index term: Mediterranean. J. Sea Res. 61,
730 68-75.
- 731 Comps MA (2000) La qualité bactériologique des coquillages de l'étang de Thau. Analyse des données de colimétrie recueillies entre 1971 et
732 1998. Relations avec l'état de l'assainissement des communes du bassin versant
- 733 Couton P, Marty D, Lequette C, Serais O, Derolez V et al (2007) Inventaire et hiérarchisation des sources de pollution microbiologique du bassin
734 de Thau - OMEGA Thau étape 1
- 735 DML (Délégation à la Mer et au Littoral, Direction Départementale des Territoires et de la Mer 34) (2011 to 2019) Bilan de la production
736 conchylicole dans le département de l'Hérault
- 737 DREAL Languedoc-Roussillon, SMBT (2011) Document d'Objectifs des sites Natura 2000 de Thau. Tome 1. Sète
- 738 Farrugio H, Le Corre G (1985) Les pêcheries de lagune en Méditerranée. Définition d'une stratégie d'évaluation. Conv. CEE XIV-B-1-84/2/M03 P1
- 739 Fauvel Y (1986) L'étang de Thau : compétition dans l'exploitation, une redite
- 740 Gervasoni E, Giffon C (2016) La conchyliculture en Occitanie. Résultats d'une enquête menée auprès de 135 entreprises conchylicoles
- 741 Gallois D (1973) Etude de deux Veneridae de l'Etang de Thau : *Venerupis decussata* et *Venerupis aurea*. PhD Thesis, Univ. Orléans
- 742 Ifremer (2008) Réseau de Suivi Lagunaire du Languedoc-Roussillon. Bilan des résultats 2007. <https://archimer.ifremer.fr/doc/00118/22923/>
- 743 INSEE (2006 to 2015, 2018) National institute of statistics and economic studies. Population in year. <https://www.insee.fr/en/accueil>
- 744 La Jeunesse I. (2001) Etude intégrée dynamique du phosphore dans le système bassin-versant - lagune de Thau (Mer Méditerranée, Hérault).
745 Université d'Orléans. PhD Thesis
- 746 Les Écologistes de l'Euzière (1988) Etang de Thau
- 747 Maurel P (2012) Signes, Données et Représentations Spatiales : Des éléments de sens dans l'élaboration d'un projet de territoire intercommunal.
748 Application au territoire de Thau. Université du Sud - Toulon, Var
- 749 Mazouni N, Rey-Valette H, Valarié P, Sauvagnargues JC, Berthou P et al (1999) Gestion d'une ressource naturelle exploitée. Cas de la Palourde
750 (*Ruditapes decussatus*) dans la lagune de Thau. Diagnostic pluridisciplinaire
- 751 Mazouni N, Rey-Valette H, Valarié P, Despres E, Dedieu O et al (2000) Complexité des espaces mobilisés par un système halieutique. Les Espac
752 l'Halieutique 497-515
- 753 Munaron D, Derolez V, Foucault E, Cimiterra N, Tapie et al (2020) OBSLAG - Volet Pesticides. Bilan 2017-2019 du suivi des lagunes
754 méditerranéennes. Rapport final. <https://archimer.ifremer.fr/doc/00656/76769/>
- 755 Pagès S (2003) Etude sectorielle du tourisme et des activités récréatives du Bassin de Thau. Mémoire de DESS, Univ. Montpellier
- 756 Péna G. (1989) Sels nutritifs et micropolluants métalliques dans un écosystème lagunaire : l'étang de Thau. PhD Thesis
- 757 Rey-Valette H (2001) Les lagunes méditerranéennes : revue des questions et des travaux de sciences sociales. In: Sourisseau E, Rey-Valette H.
758 (ed) Women in Fisheries Sector, France
- 759 Rey-Valette H. (2006) Fiches synthétiques des principales filières du territoire de Thau
- 760 Richard M (2019) Conséquences des mortalités massives de juvéniles d'huitres dans la lagune de Thau. In: Les dossiers D'Agropolis International
761 (ed) Compétences de la communauté Scientifique en Région Occitanie 24
- 762 Serais O, Derolez V, Fiandrino A (2007) Synthèse des résultats REMI - Etang de Thau - Période 1997-2006 - OMEGA Thau étape 1.
763 <https://archimer.ifremer.fr/doc/00042/15338/>
- 764 SIH (Système d'Informations Halieutiques de l'Ifremer) (2018) Activité des navires de pêche 2017 : quartier maritime Sète.
765 <https://archimer.ifremer.fr/doc/00490/60144/>
- 766 SIH (Système d'Informations Halieutiques) (2019) Quartier maritime Sète. 2018. Activité des navires de pêche.
767 <https://archimer.ifremer.fr/doc/00608/72051/>
- 768 SMBT (2010) Natura 2000 Thau. DOCUMENT d'Objectifs des sites FR 9112018 «étang de Thau et Lido de Sète à Agde» FR 9101411 «herbiers
769 de l'étang de Thau». PHASE 1. Etat des lieux
- 770 SMBT (2014) DLAL FEAMP 2014-2020. Thau et sa bande côtière de Frontignan à Agde. Annexes au dossier de candidature. Volume 1

771 Souchu P, Gasc A, Collos Y, Vaquer A, Tournier H et al (1998) Biogeochemical aspects of bottom anoxia in a Mediterranean lagoon (Thau,
 772 France). Mar Ecol Prog Ser 164:135-146. doi:10.3354/meps164135

773 Vault D, Frisoni GF (1986) Phytoplankton productivity and nutrients in five Mediterranean lagoons. Oceanol Acta 9(5)

774

775 **Table 3** Type of stakeholders interviewed during the study and date of the interviews

776

Stakeholder category	Name of the organization	Date of the interviews
Local management organization	Syndicate of the Thau Lagoon (French acronym 'SMBT')	June 21 2019
State representatives	Delegation for the Sea and Coast (French acronym 'DML')	June 19 2019
	Rhône-Mediterranean-Corsica Water Agency (French acronym 'AERMC')	June 27 2019
Representative of the Region for shellfish and fishing	Center for the Promotion of Lagoon and Maritime Activities (French acronym 'Cépralmar')	August 2 2019 and March 24 2020
Research institute	French Research Institute for the Exploitation of the Sea (French acronym 'Ifremer')	February 7 2019
Environmental association	Center for the Promotion of Environmental Initiatives (French acronym 'CPIE')	June 28 2019 and April 2020
Representative of shellfish and fisheries	Regional Mediterranean Shellfish Committee (French acronym 'CRCM')	April 4 2019
	Mèze 'Prud'homie'	October 7 2019
	Regional Committee for Maritime Fisheries in the Mediterranean (French acronym 'CRPMEM')	July 22 2019

777

778

779

780


781

782

I. Environmental and socio-economic problems related to the Thau lagoon	II. Uses and benefits of the Thau lagoon	III. Management of the lagoon
1. What does the Thau lagoon represent for your structure? Or how would you define the Thau lagoon?	4. Can you tell us how the weight of the different activities and uses linked to the lagoon has evolved from 1970 to today? (diversity of uses: shellfish farming, fishing, tourism, thermalism, recreational activities, etc.)	8. How has the management of the lagoon evolved in terms of objectives and actors involved? What have been the main milestones in this area? Who are these new actors? When did they appear? And why?
2. What are the major events that have marked the evolution of the lagoon's characteristics and management policies (environmental and/or socio-economic problems)? <i>A timeline is presented to be completed with these events. Additional questions will be asked when the actor establishes the timeline in order to know the causes and consequences of these events.</i>	5. How do you rate the level of ecosystem services provided by the lagoon from 1970 to today? <i>To do this, we present a list of services in a table and propose to give a score to each service on a scale of 1 to 5 over 3 different time periods from 1970 to today.</i>	9. What are your organization's commitments to lagoon management? Are these commitments the same since the beginning or have they evolved or changed? What are the strategic choices in managing the lagoon to meet user needs? From 1970 to the present?
3. In your opinion, what are the threats to the lagoon today?	6. Has the lagoon always had the same expectations of the services it provides? If so, what are the new forms of demand from 1970 to the present?	10. Are there conflicts between the user community and/or between the lagoon management community? If so, over what periods from 1970 to the present? What do they consist of? And how are they managed?
	7. In your opinion, what are the contributions of the lagoon to the territory? Are they the same (from 1970 to today)? Or has the relationship between these contributions changed over this period?	11. How do you see the future of the lagoon? Will the lagoon be able to ensure the sustainability of its uses?


787

788 **Fig. 4** Geographical location of the Thau lagoon. Shellfish aquaculture concessions are located in three rearing areas: Bouzigues, Mèze and Marseillan. The main exchanges of sea water take place via the Sète Canals
789



790

791 **Fig. 5** Timeline of the main events concerning the ecological, chemical and sanitary status of the Thau lagoon (1970-2018)




792

793


794

795 **Fig. 6** Timeline of the main events concerning human activities centered on the Thau lagoon (1970-2018)



796

797 **Fig. 7** Timeline of the main events concerning the management of the Thau lagoon (1970-2018)




798

799

800

801 **Fig. 8** Timeline of the main events concerning the installation of wastewater treatment plants and the main works on the
802 wastewater treatment system of the Thau lagoon from 1970 to 2018 (La Jeunesse 2001; pers. comm. L. Cesmat, R.
803 Pete/SMBT, A. Barrera/AERMC)




804


805

806
807
808
809
810
811
812
813
814
815
816
817
818

819
820
821
822
823
824
825


Fig. 9 Changes observed between 1970 and 2018 in the 12 variables selected by the PCA in the DPISR approach and environmental drivers of change (in parentheses, code of the variables according to Table A1). Drivers: 1) Population of the watershed ("Population"); Pressures: 2) Population not connected to WWTP ("Population no WWTP"); State: 3) DIP concentration in water ("DIP summer") and 4) Chlorophyll a concentration in water ("CHLA annual"); Impacts: 5) Shellfish production ("Shellfish prod"), 6) Number of fishers ("N° fishers"), 7) Number of tourist beds ("Tourist beds"), 8) Number of cultural associations ("Cultural assoc"), 9) Number of water recreational associations ("Water recreat asso"); Responses: 10) Surface area protected by Natura 2000 ("N2000 surf"), 11) Cumulative number of scientific publications ("Scientific pub"), 12) Number of management tools and structures ("Management"); Environmental drivers of change: 13) Annual mean temperature ("Temperature") and 14) Annual cumulative rainfall ("Rainfall"). Interpolated data are in grey





829

830 **Fig. 10** Results of PCA of environmental and social variables from 1970 to 2018. First and second axes showing the 12 active
831 variables colored according to their DPSIR category (+2 illustrative variables in grey). The first and second axes account for
832 87.8% of total variability



833

834

835

836