



HAL
open science

Multilocus Sequence-Based Analysis Delineates a Clonal Population of *Agrobacterium (Rhizobium) radiobacter (Agrobacterium tumefaciens)* of Human Origin

Fabien Aujoulat, Estelle Jumas-Bilak, Agnès Masnou, Fanny Sallé, Denis Faure, Christine Segonds, Hélène Marchandin, Corinne Teyssier

► **To cite this version:**

Fabien Aujoulat, Estelle Jumas-Bilak, Agnès Masnou, Fanny Sallé, Denis Faure, et al.. Multilocus Sequence-Based Analysis Delineates a Clonal Population of *Agrobacterium (Rhizobium) radiobacter (Agrobacterium tumefaciens)* of Human Origin. *Journal of Bacteriology*, 2011, 193 (10), pp.2608-2618. <10.1128/jb.00107-11>. <hal-04315786>

HAL Id: hal-04315786

<https://hal.umontpellier.fr/hal-04315786v1>

Submitted on 30 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Multilocus Sequence-Based Analysis Delineates a Clonal Population of *Agrobacterium (Rhizobium) radiobacter (Agrobacterium tumefaciens)* of Human Origin[∇]

Fabien Aujoulat,¹ Estelle Jumas-Bilak,^{1*} Agnès Masnou,¹ Fanny Sallé,¹ Denis Faure,² Christine Segonds,³ Hélène Marchandin,^{1,4} and Corinne Teyssier¹

Université Montpellier 1, Unité Mixte de Recherche 5119 Ecosym, Equipe Pathogènes et Environnements, Faculté de Pharmacie, 15 Avenue Charles Flahault, BP 14491, 34093 Montpellier Cedex 5, France¹; Centre National de la Recherche Scientifique, Institut des Sciences du Végétal, UPR2355, 1 Avenue de la Terrasse, 91198 Gif-sur-Yvette, France²; Centre Hospitalier Universitaire de Toulouse, Hôpital de Purpan, Laboratoire de Bactériologie-Hygiène, 31059 Toulouse Cedex 9, France³; and Centre Hospitalier Universitaire de Montpellier, Hôpital Arnaud de Villeneuve, Laboratoire de Bactériologie, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5, France⁴

Received 24 January 2011/Accepted 3 March 2011

The genus *Agrobacterium* includes plant-associated bacteria and opportunistic human pathogens. Taxonomy and nomenclature within the genus remain controversial. In particular, isolates of human origin were all affiliated with the species *Agrobacterium (Rhizobium) radiobacter*, while phytopathogenic strains were designated under the synonym denomination *Agrobacterium tumefaciens*. In order to study the relative distribution of *Agrobacterium* strains according to their origins, we performed a multilocus sequence-based analysis (MLSA) on a large collection of 89 clinical and environmental strains from various origins. We proposed an MLSA scheme based on the partial sequence of 7 housekeeping genes (*atpD*, *zwf*, *trpE*, *groEL*, *dnaK*, *glnA*, and *rpoB*) present on the circular chromosome of *A. tumefaciens* C58. Multilocus phylogeny revealed that 88% of the clinical strains belong to genovar A7, which formed a homogeneous population with linkage disequilibrium, suggesting a low rate of recombination. Comparison of genomic fingerprints obtained by pulsed-field gel electrophoresis (PFGE) showed that the strains of genovar A7 were epidemiologically unrelated. We present genetic evidence that genovar A7 may constitute a human-associated population distinct from the environmental population. Also, phenotypic characteristics, such as culture at 42°C, agree with this statement. This human-associated population might represent a potential novel species in the genus *Agrobacterium*.

Members of the genus *Agrobacterium* are well-known plant-associated bacteria. *Agrobacterium tumefaciens*, *Agrobacterium rhizogenes*, *Agrobacterium rubi*, *Agrobacterium vitis*, and *Agrobacterium larrymoorei* may be phytopathogens, causing diverse tumors in plants (4, 45). Since *A. tumefaciens* is able to transform vegetal cells, it has been used to obtain genetically engineered plants (32, 40). In the past 2 decades, *Agrobacterium (Rhizobium) radiobacter* has been recognized as an opportunistic human pathogen responsible for nosocomial infections, mainly bacteremia, peritonitis, and urinary tract infections (1, 3, 10, 28). Bacteremias are usually secondary to the use of intravenous devices (3, 11, 17). *A. radiobacter* strains have also been found in the respiratory tracts of cystic fibrosis patients (5).

The genus *Agrobacterium* belongs to the family *Rhizobiaceae* in subphylum alpha of the proteobacteria (16). 16S rRNA gene-based phylogeny grouped the genus *Agrobacterium* and some fast-growing nitrogen-fixing bacteria of the genus *Rhizobium* in the same clade (12). The paraphyly of the two genera has led to diverse revisions of the nomenclature (12, 42, 43,

44), but expert controversies have received little attention in practice. Currently, *A. tumefaciens*, *A. radiobacter*, and *R. radiobacter* have authority to denominate the same bacterium, even if two type strains named *A. tumefaciens* CFBP 2413^T and *A. radiobacter* CFBP 2414^T, are deposited and validated. As a rule, but without clear argumentation, clinical strains of the genus *Agrobacterium* are usually affiliated with the species *A. radiobacter*, whereas *A. tumefaciens* includes environmental phytopathogenic or nonphytopathogenic strains. In this study, *Agrobacterium (Rhizobium)* strains will be named only *Agrobacterium*.

The strains of the genus *Agrobacterium* can be grouped into three biovars on the basis of biochemical and physiological tests and into 11 genomospecies defined by DNA-DNA hybridization (30). Finally, relationships could not be established among biovars, genomospecies, and species. In particular, biovar 1 corresponds to 9 genomospecies and contains the type strains of *A. tumefaciens* and *A. radiobacter*.

Previously reported clinical strains were assigned to the genus *Agrobacterium* and to the species *A. radiobacter* (*R. radiobacter*) on the basis of conventional methods used for the routine identification of oxidase-positive, nonfastidious, nonfermenting Gram-negative bacilli or, though less frequently, on the 16S rRNA gene sequence. However, these methods are not reliable for identification to the species level due to the absence of species other than *A. radiobacter* in the commercial systems' phenotypic databases and to low interspecific poly-

* Corresponding author. Mailing address: UMR 5119 Ecosym, Equipe Pathogènes et Environnements, Laboratoire de Bactériologie, Faculté de Pharmacie, 15 Avenue Charles Flahault, BP 14491, 34093 Montpellier Cedex 5, France. Phone: 33 4 67 63 54 26. Fax: 33 4 67 63 45 11. E-mail: ebilak@univ-montp1.fr.

[∇] Published ahead of print on 11 March 2011.

TABLE 1. Characteristics of *Agrobacterium* species strains of clinical origin^{a,g}

Strain ^b	Genovar within clade A	ST	Allelic profile ⁱ							Growth (42°C) ^f	3-Ketolactose production ^f	PDA CaCO ₃ ^{f,h}	Origin of strain	Place, yr of isolation
			<i>atpD</i>	<i>dnaK</i>	<i>glnA</i>	<i>groEL</i>	<i>rpoB</i>	<i>trpE</i>	<i>zwf</i>					
AGR01	A7	3	3	5	7	3	31	2	5	+	-	-	Scalp wound	Montpellier, Fr, 2003
AGR35	A7	3	3	5	7	3	31	2	5	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR38	A7	3	3	5	7	3	31	2	5	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR40	A7	3	3	5	7	3	31	2	5	-	-	-	Blood	Montpellier, Fr, 2009
AGR02	A7	4	3	6	10	4	1	3	6	+	-	-	Urine	Montpellier, Fr, 2004
AGR03	A7	5	18	7	10	5	2	5	7	+	+	-	Urine	Montpellier, Fr, 2004
AGR04	A7	5	18	7	10	5	2	5	7	+	+	-	Undocumented ^e	Nimes, Fr, 2004
AGR17	A7	5	18	7	10	5	2	5	7	+	+	-	Sputum ^d	Montpellier, Fr, 2006
AGR24	A7	5	18	7	10	5	2	5	7	+	+	-	Sputum ^d	Montpellier, Fr, 2008
AGR33	A7	5	18	7	10	5	2	5	7	-	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR41	A7	5	18	7	10	5	2	5	7	+	+	-	Glans penis	Montpellier, Fr, 2009
AGR05	A7	6	3	6	12	4	1	3	6	+	-	-	Fiberscope rinse fluid ^e	Toulouse, Fr, 1997
AGR07	A7	6	3	6	12	4	1	3	6	+	+	-	Undocumented ^e	Toulouse, Fr, 1998
AGR09	A7	6	3	6	12	4	1	3	6	+	-	-	Collyre ^e	Montpellier, Fr, 1995
AGR15	A7	6	3	6	12	4	1	3	6	+	-	-	Sputum ^d	Montpellier, Fr, 2005
AGR20	A7	6	3	6	12	4	1	3	6	+	-	-	Sputum ^d	Montpellier, Fr, 2007
AGR21 A	A7	6	3	6	12	4	1	3	6	+	-	-	Sputum ^d	Montpellier, Fr, 2007
AGR21 B	A7	6	3	6	12	4	1	3	6	+	-	-	Sputum ^d	Montpellier, Fr, 2007
CCUG 48648	A7	6	3	6	12	4	1	3	6	+	+	-	Blood	Stockholm, Sweden, 2004
AGR06	A7	7	4	8	5	6	3	22	8	+	+	-	Culture medium ^e	Toulouse, Fr, 1997
AGR10	A7	8	5	9	9	7	4	9	9	+	+	-	Sputum ^d	Montpellier, Fr, 2005
AGR18	A7	8	5	9	9	7	4	9	9	+	-	-	Sputum ^d	Montpellier, Fr, 2006
AGR11	A7	9	10	10	9	8	5	6	6	+	-	-	Sputum	Nimes, Fr, 2005
AGR16	A7	9	10	10	9	8	5	6	6	+	-	-	Feces	Montpellier, Fr, 2006
AGR26	A7	9	10	10	9	8	5	6	6	+	-	-	Sputum	Montpellier, Fr, 2008
AGR27	A7	9	10	10	9	8	5	6	6	+	ND	ND	Sputum	Montpellier, Fr, 2008
AGR37	A7	9	10	10	9	8	5	6	6	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR39	A7	9	10	10	9	8	5	6	6	+	-	-	Sputum	Montpellier, Fr, 2009
CCUG 49619	A7	9	10	10	9	8	5	6	6	+	-	-	Bronchus brush rinse	Göteborg, Sweden, 2004
AGR12	A7	10	3	5	7	3	32	2	5	+	-	-	Toe wound	Montpellier, Fr, 2005
AGR14	A7	11	6	11	6	8	7	8	11	+	-	-	Sputum ^d	Montpellier, Fr, 2006
AGR23	A7	11	6	11	6	8	7	8	11	+	-	-	Blood	Toulouse, Fr, 2006
AGR43	A7	11	6	11	6	8	7	8	11	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR19	A7	12	7	5	11	16	5	10	12	+	+	-	Blood	Montpellier, Fr, 2007
CCUG 12509	A7	13	4	22	12	16	14	5	21	+	+	-	Lesion	Texas, 1974
AGR25	A7	13	4	22	12	16	14	5	21	+	-	-	Feces	Montpellier, Fr, 2008
AGR32	A7	15	8	25	8	19	3	9	23	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR34	A7	16	9	8	5	8	28	8	5	+	-	-	Sputum ^d	Montpellier, Fr, 2009
AGR08	A7	18	3	3	7	3	33	1	5	+	-	-	Sputum ^d	Montpellier, Fr, 2005
LMG 361.1	A7	37	30	32	24	25	27	29	11	+	+	-	Sputum	Missouri, NA
LMG 378	A7	38	7	5	9	16	5	10	12	+	+	-	Blood	Georgia, NA
LMG 399	A7	39	9	8	5	26	28	8	7	+	+	-	Infected eye	Louisiana, NA
LMG 355	A7	48	3	5	7	3	36	2	5	-	+	-	Bronchial washings	Hawaii, NA
AGR28	A2	14	11	23	13	17	15	11	22	-	+	-	Blood	Montpellier, Fr, 2008
AGR29	A2	14	11	23	13	17	15	11	22	-	+	-	Blood	Montpellier, Fr, 2008
AGR30	A2	14	11	23	13	17	15	11	22	-	+	-	Blood	Montpellier, Fr, 2008
AGR13	A4	2	17	4	3	9	6	7	10	-	+	-	Sputum ^d	Montpellier, Fr, 2005
AGR36	A4	17	17	4	3	9	23	7	10	-	+	-	Sputum ^d	Montpellier, Fr, 2009
LMG 227 ^c	A3	28	16	16	18	10	12	16	4	-	+	-	Blood	Denmark, before 1972

^a Results are presented by genovar within clade A and then by ST.
^b All reference strains were purchased as *Agrobacterium (Rhizobium) radiobacter (A. tumefaciens)* strains from the corresponding collection.
^c Biovar 1.
^d From cystic fibrosis patients.
^e Probable human origin.
^f +, positive for the test; -, negative for the test; ND, not determined.
^g CCUG, Culture Collection University of Göteborg; LMG, Laboratorium voor Microbiologie Universiteit Gent; Fr, France; NA, not available.
^h PDA CaCO₃, differential production of acid from a glucose-containing medium associated with differential dissolution of CaCO₃.
ⁱ For each locus, each different allele was assigned an arbitrary number.

morphism within the 16S rRNA gene. Considering the taxonomic problems associated with the absence of suitable tools for species identification, the distribution of the clinical strains among the general population of *Agrobacterium* has never been described.

The availability of the complete genomic sequences of *A. tumefaciens* strain C58 (18), *Rhizobium rhizogenes* strain K84 (35) (formerly *A. radiobacter*, formerly *A. rhizogenes* [41]), and *A. vitis* strain S4 (35) allowed the use of multilocus based-methods to study the genetic diversity in *Agrobacterium* species populations. The aim of the present study was to analyze a

large collection of clinical and environmental *Agrobacterium* strains using multilocus genetics and phenotypic traits in order to determine whether clinical isolates displayed specific characteristics.

MATERIALS AND METHODS

Bacterial strains, culture, and phenotype. A total of 89 strains of *Agrobacterium* spp., including 49 clinical and 40 environmental isolates, were analyzed (Tables 1 and 2). Clinical strains were sampled over a 37-year period in four countries in Europe, North America, and Oceania. However, most of them were recovered over a 14-year period from patients admitted to three French univer-

TABLE 2. Characteristics of *Agrobacterium* species strains of environmental origin^{a,f}

Strain/biovar ^b	Genovar within clade A	ST	Allelic profiles ^c										Growth (42°C) ^d	3-Ketolactose production ^e	PDA CaCO ₃ ^e	Phytopathogenicity ^{c,d}	Origin of strain	Place, yr of isolation ^f
			<i>atpD</i>	<i>dnaK</i>	<i>glnA</i>	<i>groEL</i>	<i>rhoB</i>	<i>tpoE</i>	<i>zvf</i>									
181	A3	26	21	14	23	12	10	24	15	15	16	-	-	-	+	Populus, tumor	New York, UD	
RTC2	A3	27	15	15	17	10	11	15	15	16	16	-	+	-	-	<i>Lycopersicon esculentum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTe4	A3	27	15	15	17	10	11	15	15	16	16	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTB12	A3	27	15	15	17	10	11	15	15	16	16	-	+	-	-	<i>Lycopersicon esculentum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
LMG 232/bv1	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Beta vulgaris</i> , rhizosphere	United Kingdom, 1963	
RTB1	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTB8	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Lycopersicon esculentum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTP7	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTP1	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTP2	A3	28	16	16	18	10	12	16	16	4	4	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTB7	A3	30	24	18	16	10	12	18	18	18	18	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
RTA10	A3	31	22	19	17	10	12	19	19	19	19	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
LMG 142/bv1	A3	34	22	19	17	10	12	19	29	29	29	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere soil	Netherlands or Germany, 1904	
LMG 197/bv1	A3	35	28	12	19	10	8	13	13	13	13	-	+	-	-	<i>Malus</i> sp., tumor	Wisconsin, 1935	
LMG 175/bv1	A3	35	28	12	19	10	8	13	13	13	13	-	+	-	-	<i>Prunus cerasifera</i> , tumor	North Bulgaria, 1959	
LMG 90/bv1	A3	40	31	15	25	23	29	27	30	30	30	-	+	-	-	<i>Rosa</i> sp. cv. Golden Rapture, tumor	New York, UD	
A6	A3	41	23	12	19	10	8	13	13	13	13	-	+	-	-	apple tree, tumor	United States, UD	
R10	A3	41	23	12	19	10	8	13	13	13	13	-	+	-	-	NA, tumor	United States, UD	
LMG 167/bv1	A3	45	35	16	18	10	12	16	4	4	4	-	+	-	-	<i>Prunus</i> sp., tumor	Gembloux, Belgium, UD	
LMG 303/bv1	A3	45	35	16	18	10	12	16	4	4	4	-	+	-	-	<i>Chrysanthemum frutescens</i> , tumor	Germany, 1927	
C58	A5	20	1	1	20	1	20	1	1	1	1	-	+	-	-	<i>Prunus</i> sp. cv. Montmorency tumor	New York, 1958	
CCUG 50385B	A5	22	27	30	20	22	25	26	28	28	28	-	+	-	-	Soil	Chile, 2004	
T37	A5	25	12	13	20	11	9	14	14	14	14	-	+	-	-	<i>Juglans regia</i> , tumor	California, 1926	
H100	A5	25	12	13	20	11	9	14	14	14	14	-	+	-	-	<i>Humulus lupulus</i> , tumor	United States, UD	
AB35-9	A4	2	17	4	3	9	6	7	10	10	10	-	+	-	-	Nematode	Avignon, Fr, 2007	
B10-2	A4	2	17	4	3	9	6	7	10	10	10	-	+	-	-	Nematode	Avignon, Fr, 2007	
RTP8	A4	32	13	20	4	14	6	20	20	20	20	-	+	-	-	<i>Nicotiana tabacum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
CFBP 2414T	A1	23	20	21	15	15	13	21	32	32	32	-	+	-	-	Ditch water	Netherlands, before 1927	
LMG 306/bv1	A1	36	29	31	22	24	26	28	31	31	31	-	+	-	-	<i>Prunus</i> sp., tumor	Saitama, Japan, 1956	
RTH4	A6	29	14	17	14	13	18	17	17	17	17	-	+	-	-	<i>Lycopersicon esculentum</i> , rhizosphere	Gif-sur-Yvette, Fr, 2006	
<i>A. vitis</i> S4		1	25	26	2	20	22	25	24	24	24	-	+	-	-	<i>Vitis vinifera</i> , tumor	Hungary, UD	
LMG 109/bv2		19	34	35	28	30	21	32	2	2	2	-	+	-	-	<i>Rosa</i> sp., tumor	Texas, UD	
LMG 29/bv2		21	33	38	27	28	35	31	33	33	33	-	+	-	-	<i>Rosa xanthina</i> , tumor	United Kingdom, UD	
LMG 253/bv2		24	34	37	28	30	21	4	2	2	2	-	+	-	-	<i>Prunus persica</i> , tumor	Greece, UD	
K84/bv2		33	2	2	1	2	21	4	2	2	2	-	+	-	-	<i>Prunus persica</i> , soil around tumor	Australia, 1972	
LMG 341/bv2		42	34	37	28	29	21	4	2	2	2	-	+	-	-	<i>Prunus dulcis</i> , tumor	Israel, UD	
LMG 63/bv2		43	34	34	28	29	21	4	2	2	2	-	+	-	-	<i>Malus sylvestris</i> , hairy root	United States, UD	
LMG 99/bv2		44	34	35	29	30	21	4	2	2	2	-	+	-	-	<i>Rosa</i> sp., tumor	Pennsylvania, UD	
LMG 229/bv2		46	36	36	28	30	21	32	2	2	2	-	+	-	-	<i>Rosa</i> sp., tumor	South Africa, UD	
<i>A. vitis</i> LMG 8750T		47	37	27	30	31	34	33	34	33	34	-	+	-	-	<i>Vitis vinifera</i> , tumor	Australia, 1977	

^a Results are presented by genovar within clade A and/or by ST.

^b All type and reference strains were purchased as *Agrobacterium (Rhizobium) radiobacter (A. tumefaciens)* strains from the corresponding collection, except for *A. vitis* S4, *A. vitis* LMG 8750T, and *A. rhizogenes* LMG 63.

^c Phytopathogenicity data for collection strains were obtained from the corresponding collection databases and from references 6, 22, 24, and 30.

^d +, positive for the test; -, negative for the test; ND, not determined.

^e PDA, CaCO₃, differential production of acid from a glucose-containing medium associated with differential dissolution of CaCO₃.

^f LMG, Laboratorium voor Microbiologie Universiteit Gent; CCUG, Culture Collection University of Göteborg; CFBP, Collection Française des Bactéries Phytopathogènes; bv, biovar when available for *Agrobacterium (Rhizobium)* strains from LMG collection; Fr, France; UD, undocumented.

^g For each locus, each different allele was assigned an arbitrary number.

TABLE 3. Primers used for gene amplification and sequencing

Locus	Function	Putative gene product	Locus position ^a	Gene size (bp)	Primers ^b	Primer sequence 5'-3'	Sequence length (bp)
<i>atpD</i>	Energy metabolism	F ₀ -F ₁ ATP synthase subunit beta	2604717	1,454	800F 1350R	GGCCAGGACGTTCTGTTCTT CTTGAAGCCCTTGATCGTGT	465
<i>dnaK</i>	Stress response	Heat shock protein, 70 kDa	126205	1,901	720F 1400R	GAAGACTTCGACATGCGTCT GCCGAGCAGCTTGTGTGTC	480
<i>glnA</i>	Amino acid biosynthesis	Glutamine synthetase	196923	1,358	144F 1340R 900R	GTCATGTTTCGACGGCTCCT CGCATGACTTCCTGCATCT CCTTGGCATGCTTGATGAT	474
<i>groEL</i>	Stress response	Heat shock protein, 60 kDa	676328	1,634	100F 1240R 760R	GTGGTGATCAGCAGCGAAG AGGCCAAGGCCAAGAAGAT CTGGAAGACATCGCCATCT	504
<i>rpoB</i>	Transcription	Beta subunit RNA polymerase	1927198	4,136	2040F 3150R 2718R	GAAAACGACGACGCCAAC TGGACCTTTTCGACCTTGTC GCGCAGAAGCTTTCTTCC	534
<i>trpE</i>	Amino acid biosynthesis	Anthranilate synthase	2262145	2,189	890F 2090R 1630R	CGCCCTATTCTTCTTCATC ATCGATTCCGGGTGGAAGT GAAATAATTCGCCAGCGTGT	510
<i>zwf</i>	Pentose phosphate pathway	Glucose-6-phosphate 1-dehydrogenase	585849	1,475	530F 950R	AGATCTTCCGCATCGACCA CTTGATGGCGACGAAGTT	384 ^c

^a Gene start codon position on the circular chromosome sequence of *A. tumefaciens* C58 (accession number AE007869).

^b F, forward primer; R, reverse primer. Primers in boldface were used for gene sequencing in both directions. The primer denominations correspond to their hybridization regions in the gene according to the complete genome sequence of *A. tumefaciens* C58.

^c The size of the *zwf* sequences of the two strains of *A. vitis* was 381 bp.

sity hospitals. Nonclinical or nonhuman strains were collected in 14 countries and four continents over a 103-year period. They were representative of the 3 biovars of *Agrobacterium* and originated from soil, plants, or nematodes. *A. tumefaciens* C58, *A. rhizogenes* K84 (formerly *A. radiobacter* K84), *A. radiobacter* CFBP 2414^T, and *A. vitis* LMG 8750^T were included as the reference and type strains. All isolates were cultivated on nutrient glucose agar (NGA) at 28°C for 24 h. They were identified to the genus level by routine phenotypic tests, including Gram stain, oxidase production, and a API 20NE system (bioMérieux, Marcy l'Etoile, France). Genus identification was confirmed by 5'-end sequencing of the 16S rRNA gene as previously described (37). The ability to grow at 35°C and 42°C was tested on NGA incubated for 24 h. Two of the assays usually used for biovar identification (3-ketolactose production from lactose and differential acid production) were performed as described previously (2). The strains from the LMG collection identified at the biovar level in the LMG database ($n = 16$) were used for the validation of the assays based on the following criteria: 3-ketolactose production positive in biovar 1 strains and negative in biovar 2 strains and acid production from glucose positive only in biovar 2 strains (2).

PFGE-restriction fragment length polymorphism (RFLP). Genomic DNA was prepared in agarose plugs as previously described (39) and digested at 25°C with 40 U of SmaI (New England BioLabs, Hertfordshire, United Kingdom). SmaI fragments were separated by pulsed-field gel electrophoresis (PFGE) using a CHEF-DRII apparatus (Bio-Rad Laboratories, Hercules, CA) in a 1% agarose gel in 0.5× Tris-borate-EDTA (TBE) buffer at 150 V and at 10°C. Pulse ramps were 100 to 160 s for 35 h, followed by 40 to 80 s for 15 h. The gel was stained with ethidium bromide and photographed under UV light. PFGE profiles were interpreted visually.

Gene amplification and sequencing. The complete genomic sequences of *A. tumefaciens* C58 (AE007869), *R. rhizogenes* K84 (formerly *A. radiobacter*, formerly *A. rhizogenes*) (CP000628), and *A. vitis* S4 (CP000633) were used as references for gene selection and primer design. The primers are shown in Table 3. Genomic DNA was extracted using the MasterPure DNA Purification kit (Epicentre, Madison, WI). PCR was carried out in 50 µl of reaction mixture containing 200 nM (each) primer (Sigma Genosys), 200 µM (each) desoxynucleoside triphosphate (dNTP) (Euromedex, Mundolsheim, France), 2.5 U of *Taq* DNA polymerase (Promega, Madison, WI) in the appropriate reaction buffer, and 50 ng of genomic DNA as the template. The amplification conditions were as follows: initial denaturation for 3 min at 95°C, followed by 35 cycles with 1 min

at 94°C; 1 min at 60°C (*trpE* and *atpD*), 65°C (*dnaK*, *glnA*, *groEL*, and *zwf*), or 68°C (*rpoB*); and 2 min 30 s (*dnaK*, *glnA*, *groEL*, *trpE*, and *rpoB*) or 1 min 30 s (*zwf* and *atpD*) at 72°C. The final extension was carried out at 72°C for 10 min. PCR products and a molecular weight marker (phage phiX DNA digested with HaeIII; New England BioLabs) were separated in a 1.5% (wt/vol) agarose gel in 0.5× TBE buffer. The amplification products were sequenced in both directions using forward and reverse amplification primers (Table 3) on an Applied Biosystems automatic sequencer (Beckman Coulter Genomics).

Multilocus sequence-based phylogeny. Gene sequences were codon aligned using CLUSTALW after translation with TRANSLATE (<http://www.expasy.org>). The sizes of the codon-aligned sequences used for further analyses are indicated in Table 3. Phylogenetic analyses were performed for each of the seven gene sequences and for the manually concatenated sequence. Evolutionary distance was analyzed using the Phylip package v3.66 (13) by neighbor joining (NJ) after distance matrix construction using DNADIST (with F84 as a substitution model). Bootstrap values were calculated using SEQBOOT and CONSENSE after 1,000 reiterations. Maximum-likelihood (ML) analysis was performed using phylogenetic analysis at <http://www.phylogeny.fr> (9). The general time-reversible (GTR) model plus gamma distribution and invariant sites was used as a substitution model. ML bootstrap support was computed after 100 reiterations. The sequences of *Ochrobactrum anthropi* ATCC 49188^T (CP000758) were used as outgroup sequences in order to place an artificial tree root. The seven gene sequences from complete genomic sequences of *A. vitis* S4 (CP000633), *Sinorhizobium fredii* NGR234 (CP001389), *Rhizobium etli* CFN 42^T (CP000133), and *Rhizobium leguminosarum* bv. *trifolii* WSM2304 (CP001191) were also used for phylogenetic analysis.

Multilocus sequence analysis (MLSA). For each locus, each different allele was assigned to a different arbitrary number using a nonredundant database program available at <http://linux.mlst.net/nrdb/nrdb.htm>. The combination of allele numbers for each isolate defined the sequence type (ST). A distance matrix in nexus format was generated from the set of allelic profiles using a Web version of the program SplitsTree (<http://www.pubmlst.org>) and then used for decomposition analyses with SplitsTree 4.0 software (21). The program LIAN 3.1 (19) was used to calculate the standardized index of association (sIA) and to test the null hypothesis of linkage disequilibrium, as well as to determine the mean genetic diversity (H) and genetic diversity at each locus (h_i). The numbers of synonymous (dS) and nonsynonymous (dN) substitutions per site were deter-




FIG. 1. Maximum-likelihood tree based on concatenated sequences of the seven housekeeping gene fragments indicating the relative placement of clinical (boldface) and environmental (lightface) strains in the genus *Agrobacterium*. The horizontal lines show genetic distance. The numbers at the nodes are support values estimated with 100 bootstrap replicates. Only bootstrap values of >0.5 are indicated. The scale bar indicates the number of substitutions per nucleotide position. *O. anthropi* ATCC 49188^T was used as the outgroup organism. The sequences of

mined on codon-aligned sequences using SNAP online software (<http://www.hiv.lanl.gov>).

Nucleotide sequence accession numbers. The nucleotide sequences determined in this study have been deposited in the GenBank data library under accession numbers HM238283 to HM238884.

RESULTS

MLSA scheme design. Seven genes encoding housekeeping proteins involved in transcription (*rpoB*), stress response (*dnaK* and *groEL*), amino acid biosynthesis (*glnA* and *trpE*), and energy metabolism (*atpD* and *zwf*) were selected from the complete genome sequence of *A. tumefaciens* C58 (Table 3). They were dispersed on the large circular chromosome of the bipartite genome of *A. tumefaciens* strain C58 (minimal distance between loci, 90 kb) in order to minimize the influence of localized linkage disequilibrium. Partial sequences of the seven loci were obtained for all 89 strains studied, showing that the MLSA scheme was suitable for analysis of strains belonging to the 3 biovars in the genus *Agrobacterium*, biovar 1, biovar 2, and *A. vitis* (biovar 3). The sequence size at each locus is given in Table 3.

Multilocus sequence-based phylogeny. We applied distance and ML phylogenetic approaches to the concatenated sequences (3,351 nucleotides) of the seven loci. The two methods gave congruent trees, and the ML tree is presented in Fig. 1. Phylogenetic analysis clustered *Agrobacterium* strains in a major clade, clade A. Clade A included 79 of the 89 strains distributed into seven subclades, named genovars A1 to A7, and all the isolates of human origin (Fig. 1). The main genovars, A7 and A3, grouped 54% ($n = 43$) and 27% ($n = 21$) of the isolates of clade A, respectively. The 43 strains found in genovar A7 were clinical isolates, representing 88% (43 out of 49 strains) of the strains of human origin in this study. *Agrobacterium* sp. strains AGR28, AGR29, and AGR30, isolated from the same patient, presented the same sequence for the seven loci and formed, by themselves, genovar A2. Strains AGR13 and AGR36 belonged to genovar A4, together with three environmental strains, and *A. radiobacter* strain LMG 227 was the only clinical isolate among the 21 strains of genovar A3. Other minor genovars, A1, A5, and A6, did not include clinical strains. Outside of clade A, we observed *Agrobacterium* strains grouping together with *Rhizobium* sp. strains. This group, including only environmental *Agrobacterium* strains, was not well enough represented ($n = 10$), and therefore, we focused on clade A for further analyses.

For each locus, phylogenetic trees reconstructed using NJ or ML methods were compared to the NJ or ML trees reconstructed from the seven concatenated sequences (data not shown). The genovars determined in the concatenated tree were also found in individual trees. Despite some conflicting topologies in terms of branching order observed among the trees, the distribution of strains in each genovar was not mod-

ified according to the gene used or the phylogenetic method, except for strain AGR08, which belonged to genovar A5 after *trpE* locus analysis and was placed in an intermediate position between genovars A5 and A7 by *rpoB* locus analysis.

Genomic species have been previously described in the genus *Agrobacterium* by determination of DNA-DNA hybridization (7, 8, 30). These data are available for 13 strains included in this study and are shown in Fig. 1. The population studied here contained strains representative of the genomic species G1, G2, G4, and G8 according to the nomenclature of Mougel et al. (26). We observed concordance between MLSA clusters and genomic groups for members of clade A: the eight strains of group G1 belonged to genovar A3, while strains of groups G4 and G8 were found in genovars A1 and A5, respectively. The only strain representative of group G2, *A. radiobacter* CCUG 12509, was located in the “human” genovar, A7. It was noteworthy that this clinical strain, isolated in the United States in 1974, had the same sequence for the seven loci as the clinical strain AGR25, isolated in France in 2008. Genovars A2, A4, and A6 in clade A did not contain strains previously affiliated with a genomic group, and therefore, the relationship between these genovars and genomic species could not be established.

Multilocus and genomic typing. Forty-eight unique sequence types (STs) were observed; 33 of them (69%) were identified only once, suggesting an overall high level of genetic diversity among the studied population (Tables 1 and 2). In clade A, the largest STs were ST6 (8 isolates); ST9 and ST28, each comprising 7 isolates; and ST5 (6 isolates). These major STs included 35% of the clade A strains. Within the “human” genovar A7, 58% of the strains were grouped in 4 STs (ST3, ST5, ST6, and ST9). In each of these STs, strains appeared epidemiologically unrelated, i.e., they were isolated over a large period of time (from 5 to 10 years) and in some cases from geographically distant places. No obvious relationships between the STs and the types of clinical samples were observed, except for ST9, which comprised 6 of the 7 isolates from the human respiratory tract. Outside of genovar A7, only ST28 in genovar A3 comprised several strains ($n = 7$) with different lifestyles; these isolates were sampled over a 43-year period in three different countries (Tables 1 and 2).

Fifty-six strains sharing 15 identical STs, i.e., 37 clinical and 19 environmental isolates of clade A, were further analyzed by PFGE in order to detect the occurrence of genomically identical strains in the corresponding population. Genomic macrorestriction with the endonuclease *SwaI* produced PFGE patterns suitable for strain comparison and comprising an average of 10 bands (data not shown). A high level of genomic diversity was observed: the 56 strains studied showed 51 pulsotypes, and 74 distinguishable isolates were demonstrated among the 79 strains of clade A based on both multilocus and PFGE analysis results. However, the levels of genomic diversity in each ST

S. fredii NGR234 (CP001389), *R. etli* CFN 42^T (CP000133), and *R. leguminosarum* bv. *trifolii* WSM2304 (CP001191) were also included. *Agrobacterium* strains with identical concatenated sequences are presented on collapsed branches and are listed on the right of the triangular representations. The names of the clades and genovars discussed in the text are shown at the corresponding roots, indicated by bold lines. The node with an asterisk varied depending on the ML trees reconstructed with each of the seven loci. Genomic species G1 to G9 are indicated in gray boxes after the names of the corresponding strains. The data are from references 7, 8, and 30.

TABLE 4. PFGE analysis of *Agrobacterium* strains belonging to the same ST

Parameter	Value														
	ST2 (n = 3) ^c	ST3 (n = 4)	ST5 (n = 6)	ST6 (n = 8)	ST8 (n = 2)	ST9 (n = 7)	ST11 (n = 3)	ST13 (n = 2)	ST14 (n = 3)	ST25 (n = 2)	ST27 (n = 3)	ST28 (n = 7)	ST35 (n = 2)	ST41 (n = 2)	ST45 (n = 2)
Clinical strains (%) ^a	33	100	100	100	100	100	100	100	100	0	0	17	0	0	0
Phytopathogen strains (%) ^a	0	0	0	0	0	0	0	0	0	100	0	0	100	100	100
Other strains (%) ^a	66	0	0	0	0	0	0	0	0	0	100	83	0	0	0
Shared bands (%) ^b	42	22	40	10	75	31	21	9	93	95	35	29	100	86	100

^a Percentage among all strains of an ST.

^b Percentage of shared bands obtained from the PFGE patterns of strains belonging to the same ST.

^c n, number of strains.

varied greatly, with from 9% to 100% of shared bands obtained from the patterns of the same ST (Table 4). A relationship between the diversity of PFGE patterns in an ST and the lifestyle of the strains could be observed (Table 4). Identical or very similar patterns with more than 95% shared bands were observed for pairs of phytopathogenic strains that in other respects were unrelated on the basis of date and/or site of isolation, such as T37 and H100 (ST25), LMG 175 and LMG 197 (ST35), and LMG 167 and LMG 303 (ST45) (Table 4). Identical or closely related patterns were also observed for three clinical strains isolated from blood cultures from the same patient within a 5-day period (strains AGR28, AGR29, and AGR30 in ST14), between strains from two patients attending the same clinical ward 1 year apart (AGR10 and AGR18 in ST8), and between strains from two patients hospitalized in two different hospitals in Montpellier on the same date (AGR26 and AGR27 in ST9). The other clinical strains studied show no more than 40% shared bands (Table 4). An intermediate genomic-diversity level was observed for nonphytopathogenic environmental strains, except for strains RTP1 and RTP2 (ST28), originating from the same experimental field, which appeared identical after PFGE analysis.

Phenotypic traits in genovars and sequence types. Among the 88 strains tested, including 49 clinical and 39 environmental isolates, all grew at 35°C but only 41 strains grew at 42°C (Tables 1 and 2). Among the latter, 40 isolates were of clinical origin and belonged to genovar A7; the remaining strain was *A. radiobacter* LMG 90 (genovar A3) from *Rosa* sp. galls. In other words, only 3 out of the 43 clinical strains of genovar A7 (AGR33, AGR40, and LMG 355) did not grow at 42°C. Clinical strains belonging to other genovars than A7 did not grow at 42°C, and only one strain of environmental origin showed significant growth at this temperature.

The production of 3-ketolactose from lactose and differential acid production, which are major characteristics commonly used for determining biovar affiliation, were studied for the 74 genomically distinguishable isolates belonging to clade A. The assays gave results congruent with biovar identifications for the LMG control strains tested. Forty-four strains out of 74 (59%) were positive for production of 3-ketolactose (Tables 1 and 2). Genovars A1, A2, A5, and A6 contained only 3-ketolactose-producing strains, while in other genovars, the ability to produce 3-ketolactose varied among strains. For example, genovar A7 contained 36% (15 out of the 42 strains tested) 3-ketolactose-producing strains, and production was also found to be variable among strains sharing the same ST (ST5, ST6, ST8,

and ST13). In clade A, all the 3-ketolactose nonproducers were isolated from animal hosts (human or nematode), except for strain 181 from *Populus* sp.

Whatever their clinical or environmental origins, all clade A isolates are unable to produce large amounts of acid from glucose and, consequently, to dissolve CaCO₃ from a glucose-containing medium: they were negative in the differential acid production assay.

Genetic statistics and recombination. Genetic analysis of the population was performed on the 74 genomically distinguishable isolates of clade A. Genetic statistics were also calculated for the two main genovars, A3 and A7 (Table 5). Within clade A, a total of 544 single-nucleotide polymorphisms (SNPs) in the 7 loci were observed. This corresponded to a range of 13.1% to 23.4% polymorphic sites, depending on the gene (Table 5). The number of different alleles for the seven loci ranged from 22 (*glnA* and *groEL*) to 28 (*atpD*) (Table 5) and did not depend on the size of the sequence studied. All loci had equivalent mol% G+C contents, from 59.6% to 61.1%, with a mean value of 60.5%, which was similar to the mean mol% G+C contents of the *A. tumefaciens* C58 chromosomes (59%) (18). The mean genetic diversity (*H*) and the genetic diversity at each locus (*h*) for clade A and genovars A3 and A7 indicated a high level of genetic diversity both inside clade A and inside the main genovars, A3 and A7 (Table 5). The locus *tpE* displayed the highest percentage of polymorphic sites, while *groEL* appeared to be the most conserved in both genovars A3 and A7. For instance, *groEL* displayed only 6 polymorphic sites among the 18 strains of genovar A3 tested. In contrast, the *groEL* gene was the locus for which a higher rate of nonsynonymous SNPs versus synonymous SNPs (dN/dS ratio) was observed at the clade level. Nevertheless, inside each of the genovars A3 and A7, no nonsynonymous mutations were observed at this locus. The dN/dS ratio for the other six loci was found to be weak (Table 5), indicating that these loci were not subjected to strong positive selective pressure. The nonsynonymous mutations did not correspond to any premature stop codon.

Evidence in favor of clonal or recombining population structure can be obtained by assessing the levels of linkage between alleles at different loci around the chromosome. We assessed the linkage between alleles from the 7 loci by determination of the sIA value. The sIA value is expected to be zero when a population is at linkage equilibrium, i.e., when free recombination occurs. Analyses were carried out using one isolate from each ST in order to minimize any bias due to a possible epi-

TABLE 5. Sequence analysis of the seven loci

Locus	Clade or genovar	No. of alleles	No. (%) of polymorphic sites	Genetic diversity (<i>h</i>)	No. of nonsynonymous codons	dN/dS ^a ratio
<i>atpD</i>	Clade A	28	68 (14.6)	0.9716	3	0.02
	Genovar A7	10	13 (2.8)	0.8824	0	
	Genovar A3	9	17 (3.7)	0.9778	1	0.08
<i>dnaK</i>	Clade A	26	67 (14)	0.9730	2	0.02
	Genovar A7	11	15 (3.1)	0.9085	0	
	Genovar A3	6	14 (2.9)	0.9111	0	
<i>glnA</i>	Clade A	22	81 (17.1)	0.9673	6	0.03
	Genovar A7	9	13 (2.7)	0.9085	1	0.06
	Genovar A3	6	16 (3.4)	0.8889	0	
<i>groEL</i>	Clade A	22	67 (13.3)	0.9403	10	0.107
	Genovar A7	9	9 (1.8)	0.9150	0	
	Genovar A3	3	6 (1.2)	0.3778	0	
<i>rpoB</i>	Clade A	27	70 (13.1)	0.9744	5	0.03
	Genovar A7	13	33 (6.2)	0.9608	1	0.015
	Genovar A3	5	8 (1.5)	0.7556	0	
<i>trpE</i>	Clade A	25	101 (19.8)	0.9787	1	0.004
	Genovar A7	10	52 (10.2)	0.9346	0	
	Genovar A3	7	20 (3.9)	0.9333	0	
<i>zwf</i>	Clade A	26	90 (23.4)	0.9730	6	0.02
	Genovar A7	9	15 (3.9)	0.8954	0	
	Genovar A3	8	13 (3.4)	0.9556	1	0.08

^a dN, number of nonsynonymous substitutions per nonsynonymous site; dS, number of synonymous substitutions per synonymous site.

demic population structure. The sIA values were calculated for clade A and for genovars A3 and A7. The sIA values ranged from 0.3425 to 0.3610 and were significantly different from 0 ($P \leq 1.00 \times 10^{-3}$), suggesting that the recombination rates were low.

Linkage disequilibrium in clade A could be present in long-term recombining populations where adaptive clones have emerged over the short term. To explore this hypothesis, we performed decomposition analysis, which depicts all of the shortest pathways linking sequences, including those that produce an interconnected network. The split graph (obtained using the Neighbor-Net method) of all seven loci displayed parallel paths (Fig. 2) corresponding to recombination events. However, the number of events was low and confirmed the low rate of recombination deduced from the determination of the sIA. Recombination events occurred only inside genovars A3, A4, and A7. Finally, the recombination clusters generated by splitting trees were consistent with phylogenetic lineages. This suggested that genetic exchanges occurred inside each genovar but that each genovar formed a lineage genetically isolated from others.

DISCUSSION

Agrobacterium spp. are described as environmental bacteria, and some strains are phytopathogenic by tumorigenesis. The phytopathogenic behavior is related to a particular genomic structure that includes a conjugative plasmid harboring genes involved in tumor formation in plants. Besides these highly specialized bacteria, nonphytopathogenic strains were found in diverse environments not always in association with plants.

Agrobacterium is also a pathogen of human beings by a mechanism probably unrelated to plant tumorigenesis (29), but the characteristics of the pathogenicity of clinical strains remain mostly unknown. Virulence in humans is considered to be low; however, the frequent isolation of *Agrobacterium* in cases of nosocomial infections and in cystic fibrosis patients (3, 5, 23) suggests adaptation of the bacterium to humans, particularly in the context of hospitalization, immunosuppression, or underlying diseases. Typically, mild nosocomial and/or opportunistic pathogens of environmental origin, like *Stenotrophomonas maltophilia* or *Ochrobactrum* spp., displayed a high level of resistance to antibacterial compounds (27, 38). This resistance by itself could explain bacterial adaptation to nosocomial conditions. In the case of *Agrobacterium*, resistance to antibiotics is considered to be lower than that observed for other environmental opportunistic pathogens (3). On the other hand, genetic data suggested heterogeneity within the genus *Agrobacterium* and in the species *A. tumefaciens* and *A. radiobacter* (30, 31). This genetic diversity has been mainly demonstrated by DNA-DNA hybridization and amplified fragment length polymorphism (AFLP) approaches, because 16S rRNA genes displayed low polymorphism in the genus, as was also observed for the entire family *Rhizobiaceae* (42). Adaptation of some strains to human colonization and/or infection, taken together with the genetic heterogeneity in the genus, raises the question of the existence of a genetic subpopulation of *Agrobacterium* that has developed a particular relationship with humans.

In this study, we investigated genetic diversity and some phenotypic traits in the largest collection of clinical isolates of




FIG. 2. SplitsTree decomposition analyses of MLSA data for *A. radiobacter*, *A. tumefaciens*, and *A. vitis* strains. The distance matrix was obtained from allelic profiles of strains. A network-like graph indicates recombination events. A starlike radiation from the central point indicates absence of recombination.

A. radiobacter and *A. tumefaciens* reported so far. The population of strains represents diverse clinical and geographic sources, as well as diverse dates of isolation. Environmental strains were also included in the population studied here. Finally, the overall collection represented different lifestyles encountered in the genus: free living and association with plants (phytopathogenic or not), with nematodes, and with humans. The collection was studied by multilocus analysis and PFGE, two methods applied for the first time to a large population of genomically and biologically diverse *Agrobacterium* isolates.

The multilocus-based analysis proposed here succeeded in analyzing the four represented genomic species defined by DNA-DNA hybridization in biovar 1 of *A. tumefaciens* and *A. radiobacter*, the K84 group (biovar 2), and *A. vitis* (biovar 3). Moreover, MLSA displayed good discriminatory power, defining seven genovars in clade A that roughly corresponded to biovar 1 of *A. tumefaciens* and *A. radiobacter*. The large majority of the clinical strains and isolates (88%) belonged to genovar A7, the only genovar encompassing only clinical strains. The few remaining clinical strains were scattered in three other genovars, either alone or associated with environmental, nonphytopathogenic strains. At first glance, genovar A7 could be considered a “clinical genovar” or a “human-associated” genovar. However, a sampling bias could be suspected in the population studied, since a number of the strains analyzed came from the same region of France (Teaching

Hospitals of Nîmes and Montpellier, 60 kilometers apart in the south of France). For example, ST3 and ST5 in genovar A7 contained only strains from this geographic origin, although they were isolated 5 or 6 years apart. However, PFGE analysis of ST3 and ST5 strains showed genomic patterns differing by at least 8 or 6 bands out of approximately 10, respectively. This suggested that the grouping of strains in STs and then in genovars did not result from the spread of the same clone in a limited geographic zone. This was confirmed for some STs that included strains of remote origin, such as France and Sweden (ST6 and ST9) or France and the United States (ST13). Moreover, in the same ST, the dates of isolation could differ by up to 31 years. For instance, ST13 included strains isolated in 1974 in Texas and in 2008 in Montpellier (France). For these reasons, we considered that genovar A7 was actually a “human-associated” genovar and not a “geographic” genovar. A major phenotypic trait further supported this conclusion, since 91% of the strains belonging to genovar A7 grew at 42°C, whereas the optimal temperature for the genus *Agrobacterium* was defined as between 25°C and 28°C (34). The heat shock protein HSP60 is a chaperone protein involved in stress adaptation, including thermal stresses. The *groEL* gene was included in our MLSA scheme and displayed a higher frequency of nonsynonymous mutations than other loci. This suggested that *groEL* is subjected to particular selective pressure in the population tested. This selective pressure could be the host temperature,

which clearly differs between the human body and plants or other environmental sources. The ability to grow at high temperature should be considered a major trait for adaptation to the human body or to other thermoregulated animals. *Agrobacterium* strains isolated from other warm-blooded animals have recently been described (36), and they should be analyzed in order to decide whether genovar A7 is a real “human-associated” genovar or a more generally “thermophilic” genovar that consequently succeeded in its relationship with the human body.

No differences were observed in epidemiological data, such as geographic origin, clinical site of isolation, medical unit, and date of isolation, between strains in genovar A7 and the few other human strains scattered in other genovars or between strains from different STs in genovar A7. The genomic structure and its dynamics have been related to the lifestyles of bacteria (20). Pathogenic bacteria, which have close relationships with their hosts, generally inhabit a narrow and stable niche and show a low level of genomic polymorphism. Comparison of macrorestriction profiles could provide a snapshot of the genome dynamics. In the alphaproteobacteria, the genome of *Brucella*, which lives in mammalian macrophages, appears highly conserved by diverse comparative methods (14, 15, 25), whereas the genome of its phylogenetic neighbor *Ochrobactrum*, a free-living bacterium and an opportunistic pathogen, is highly fluid and polymorphic (33, 39). We observed differences in the levels of genomic polymorphism assessed by PFGE between the phytopathogenic strains and the clinical strains. The genome of the phytopathogens appeared globally conserved, suggesting the existence of constraints limiting genomic variation and adaptation to a narrow niche, including a specialized virulence mechanism. In contrast, PFGE profiles were highly polymorphic among *A. tumefaciens* (*A. radiobacter*) clinical strains. Variation generated by genome rearrangements has been reported to provide advantages for invading or for merely inhabiting complex environments, a condition that corresponded to the opportunistic behavior of *A. radiobacter*. For epidemiological purposes, PFGE-RFLP appeared more discriminative than MLST, since it was able to differentiate strains belonging to the same multilocus sequence type and is probably a suitable method for the epidemiological follow-up of human outbreaks due to *Agrobacterium*. PFGE previously permitted detection of the nosocomial transmission of *Agrobacterium* isolates from intravenous catheters of two hospitalized patients (17).

Previous phylogenetic studies based on the 16S rRNA gene (42) suggested a high level of recombination in the genus *Agrobacterium*, making its utilization in taxonomy difficult. Our results showed, by the use of a multilocus approach, that the *Agrobacterium* population was structured in robust subpopulations and had a low rate of recombination. The clade A population displayed a basically clonal structure, with no recombination among clones and a low level of recombination within each clone (genovar). The genovars obtained by the multilocus approach seem to be in accordance with the genomic species determined in previous studies (7, 8, 30). Consequently, our multilocus scheme appeared to be a suitable tool for taxonomic studies in the genus and allowed us to suggest that the “human-associated” genovar A7, which contains a strain previously affiliated with genomospecies G2 by DNA-DNA

hybridization (30) and presents particular phenotypic features, such as growth temperature and inconstant production of 3-ketolactose, may represent a new species in the genus *Agrobacterium* (*Rhizobium*). However, further investigations, such as DNA-DNA hybridization between diverse members of genovar A7 and a more complete phenotypic study, remain to be performed.

Agrobacterium spp. are generally mild pathogens, but the existence of a clone adapted to clinical conditions should stress its detection and epidemiological surveillance, as well as its evolution, in order to detect the emergence of virulent and/or resistant strains and to prevent their spread.

ACKNOWLEDGMENTS

We are particularly indebted to the microbiology laboratory team of the Nîmes, Montpellier, and Toulouse, France, academic hospitals for providing some clinical isolates. We also thank CNRS UPR2355 ISV at Gif-sur-Yvette and UMR1133 EMIP of the University Montpellier 2 for providing isolates from plants and nematodes, respectively.

Parts of this study were supported by grants from ADEREMPHA (Sauzet, France).

REFERENCES

- Amaya, R. A., and M. S. Edwards. 2003. *Agrobacterium radiobacter* bacteremia in pediatric patients: case report and review. *Pediatr. Infect. Dis. J.* **22**:183–186.
- Bouzar, H., J. B. Jones, and A. L. Bishop. 1995. Simple cultural tests for identification of *Agrobacterium* biovars. *Methods Mol. Biol.* **44**:9–13.
- Chen, C.-Y., K. S. Hansen, and L. K. Hansen. 2008. *Rhizobium radiobacter* as an opportunistic pathogen in central venous catheter associated bloodstream infection: case report and review. *J. Hosp. Infect.* **68**:203–207.
- Chilton, M. D., et al. 1982. *Agrobacterium rhizogenes* inserts T-DNA into the genomes of the host plant root cells. *Nature* **295**:432–434.
- Coenye, T., J. Goran, T. Spilker, P. Vandamme, and J. J. LiPuma. 2002. Characterization of unusual bacteria isolated from respiratory secretions of cystic fibrosis patients and description of *Inquilinus limosus* gen. nov. sp. nov. *J. Clin. Microbiol.* **40**:2062–2069.
- DeLey, J., M. Bernaerts, A. Rassel, and J. Guilmot. 1966. Approach to an improved taxonomy of the genus *Agrobacterium*. *J. Gen. Microbiol.* **43**:7–17.
- DeLey, J., R. Tytgat, J. De Smedt, and M. Michels. 1973. Thermal stability of DNA:DNA hybrids within the genus *Agrobacterium*. *J. Gen. Microbiol.* **78**:241–252.
- DeLey, J. 1974. Phylogeny of prokaryotes. *Taxon* **23**:291–300.
- Dereeper, A., et al. 2008. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res.* **36**:W465–W469.
- Edmond, M. B., S. A. Riddler, C. M. Baxter, B. M. Wicklund, and A. W. Pascuile. 1993. *Agrobacterium radiobacter* a recently recognized opportunistic pathogen. *Clin. Infect. Dis.* **16**:388–391.
- Erol Cipe, F., F. Dogu, D. Sucuoglu, D. Aysev, and A. Ikinogullari. 2010. Asymptomatic catheter related *Rhizobium radiobacter* infection in a haploidentical hemopoietic stem cell recipient. *J. Infect. Dev. Ctries.* **4**:530–532.
- Farrand, S. K., P. B. van Berkum, and P. Oger. 2003. *Agrobacterium* is a definable genus of the family *Rhizobiaceae*. *Int. J. Syst. Evol. Microbiol.* **53**:1681–1687.
- Felsenstein, J. 1984. Distance methods for inferring phylogenies: a justification. *Evolution* **38**:16–24.
- Foster, J. T., et al. 2009. Whole-genome-based phylogeny and divergence of the genus *Brucella*. *J. Bacteriol.* **191**:2864–2870.
- Gándara, B., A. L. Merino, M. A. Rogel, and E. Martínez-Romero. 2001. Limited genetic diversity of *Brucella* spp. *J. Clin. Microbiol.* **39**:235–240.
- Garrity, G. M., et al. 2007. Part 3, The Bacteria: phylum “Proteobacteria”, class Alphaproteobacteria, p. 52–111. In J. R. Cole et al. (ed.), *The taxonomic outline of Bacteria and Archaea, release 7.7*. Michigan State University, East Lansing, MI. <http://www.taxonomicoutline.org/index.php/toba/article/view/179/212>.
- Giammanco, G. M., et al. 2004. Molecular typing of *Agrobacterium* species isolates from catheter-related bloodstream infections. *Infect. Control Hosp. Epidemiol.* **25**:885–887.
- Goodner, B., et al. 2001. Genome sequence of the plant pathogen and biotechnology agent *Agrobacterium tumefaciens* C58. *Science* **294**:2323–2328.
- Haubold, B., and R. R. Hudson. 2000. LIAN 3.0: detecting linkage disequilibrium in multilocus data. *Linkage analysis. Bioinformatics* **16**:847–848.
- Hughes, D. 8 December 2000. Evaluating genome dynamics: the constraints on rearrangements within bacterial genomes. *Genome Biol.* **1**:reviews0006–reviews0006.8. doi:10.1186/gb-2000-1-6-reviews0006.

21. **Huson, D. H., and D. Bryant.** 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* **23**:254–267.
22. **Kertters, K., J. DeLey, P. H. A. Sneath, and M. Sackin.** 1973. Numerical taxonomic analysis of *Agrobacterium*. *J. Gen. Microbiol.* **78**:227–239.
23. **Lai, C.-C., et al.** 2004. Clinical and microbiological characteristics of *Rhizobium radiobacter* infections. *Clin. Infect. Dis.* **38**:149–153.
24. **Lippincott, J. A., and B. B. Lippincott.** 1969. Tumour-initiating ability and nutrition in the genus *Agrobacterium*. *J. Gen. Microbiol.* **59**:57–75.
25. **Michaux-Charachon, S., et al.** 1997. Genome structure and phylogeny in the genus *Brucella*. *J. Bacteriol.* **179**:3244–3249.
26. **Mougel, C., J. Thioulouse, G. Perriere, and X. Nesme.** 2002. A mathematical method for determining genome divergence and species delineation using AFLP. *Int. J. Syst. Evol. Microbiol.* **52**:573–586.
27. **Nicodemo, A. C., and J. I. Garcia Paez.** 2007. Antimicrobial therapy for *Stenotrophomonas maltophilia* infections. *Eur. J. Clin. Microbiol. Infect. Dis.* **26**:229–237.
28. **Paphitou, N. I., and K. V. Rolston.** 2003. Catheter related bacteremia caused by *Agrobacterium radiobacter* in a cancer patient: case report and literature review. *Infection* **31**:421–424.
29. **Petrunia, I. V., et al.** 2008. *Agrobacterium tumefaciens*-induced bacteraemia does not lead to reporter gene expression in mouse organs. *PLoS One* **3**:e2352.
30. **Popoff, M. Y., K. Kersters, M. Kiredjian, I. Miras, and C. Coynault.** 1984. Position taxonomique de souches d'*Agrobacterium* d'origine hospitalière. *Ann. Microbiol.* **135**:427–442.
31. **Portier, P., et al.** 2006. Identification of genomic species in *Agrobacterium* biovar 1 by AFLP genomic markers. *Appl. Environ. Microbiol.* **72**:7123–7131.
32. **Ream, L. W., and M. P. Gordon.** 1982. Crown gall disease and prospects for genetic manipulation of plants. *Science* **218**:854–859.
33. **Romano, S., et al.** 2009. Multilocus sequence typing supports the hypothesis that *Ochrobactrum anthropi* displays a human-associated subpopulation. *BMC Microbiol.* **9**:267–285.
34. **Sawada, H., H. Ieki, H. Oyaizu, and S. Matsumoto.** 1993. Proposal for rejection of *Agrobacterium tumefaciens* and revised descriptions for the genus *Agrobacterium* and for *Agrobacterium radiobacter* and *Agrobacterium rhizogenes*. *Int. J. Syst. Bacteriol.* **43**:694–702.
35. **Slater, S. C., et al.** 2009. Genome sequences of three *Agrobacterium* biovars help elucidate the evolution of multichromosome genomes in bacteria. *J. Bacteriol.* **191**:2501–2511.
36. **Tabar, M. D., et al.** 2010. Presence of opportunistic bacteria (*Rhizobium* spp.) with potential for molecular misdiagnosis among canine and feline clinical samples. *Can. Vet. J.* **51**:895–897.
37. **Teyssier, C., et al.** 2003. Species identification and molecular epidemiology of bacteria belonging to *Ochrobactrum* genus. *Pathol. Biol.* **51**:5–12.
38. **Teyssier, C., et al.** 2005. Molecular and phenotypic features for identification of the opportunistic pathogens *Ochrobactrum* spp. *J. Med. Microbiol.* **54**:945–953.
39. **Teyssier, C., et al.** 2005. Pulsed-field gel electrophoresis to study the diversity of whole genome organization in the genus *Ochrobactrum*. *Electrophoresis* **26**:2898–2907.
40. **Tzfira, T., and V. Citovsky.** 2006. *Agrobacterium*-mediated genetic transformation of plants: biology and biotechnology. *Curr. Opin. Biotechnol.* **17**:147–154.
41. **Velázquez, E., et al.** 2010. Analysis of core genes supports the reclassification of strains *Agrobacterium radiobacter* K84 and *Agrobacterium tumefaciens* AKE10 into the species *Rhizobium rhizogenes*. *Syst. Appl. Microbiol.* **33**:247–251.
42. **Young, J. M., L. D. Kuykendall, E. Martinez-Romero, A. Kerr, and H. Sawada.** 2001. A revision of *Rhizobium* Frank 1889 with an emended description of the genus and the inclusion of all species of *Agrobacterium* Conn 1942 and *Allorhizobium undicola* deLajudie *et al.* 1998 as new combinations: *Rhizobium radiobacter*, *R. rhizogenes*, *R. rubi*, *R. undicola* and *R. vitis*. *Int. J. Syst. Evol. Microbiol.* **51**:89–103.
43. **Young, J. M., L. D. Kuykendall, E. Martinez-Romero, A. Kerr, and H. Sawada.** 2003. Classification and nomenclature of *Agrobacterium* and *Rhizobium*—a reply to Farrand *et al.* *Int. J. Syst. Evol. Microbiol.* **53**:1689–1695.
44. **Young, J. M., S. R. Pennycook, and D. R. W. Watson.** 2006. Proposal that *Agrobacterium radiobacter* has priority over *Agrobacterium tumefaciens*. Request for an opinion. *Int. J. Syst. Evol. Microbiol.* **56**:491–493.
45. **Zhu, J., et al.** 2000. The bases of crown gall tumorigenesis. *J. Bacteriol.* **182**:3885–3895.